

# UC Irvine

## UC Irvine Previously Published Works

### Title

A Home-Based Telerehabilitation Program for Patients With Stroke

### Permalink

<https://escholarship.org/uc/item/8hq3r5w5>

### Journal

Neurorehabilitation and Neural Repair, 31(10-11)

### ISSN

0888-4390

### Authors

Dodakian, Lucy

McKenzie, Alison L

Le, Vu

et al.

### Publication Date

2017-10-01

### DOI

10.1177/1545968317733818

Peer reviewed



Published in final edited form as:

*Neurorehabil Neural Repair*. 2017 ; 31(10-11): 923–933. doi:10.1177/1545968317733818.

## A Home-Based Telerehabilitation Program for Patients with Stroke

Lucy Dodakian, MA, OTR/L<sup>1</sup>, Alison L. McKenzie, PT, PhD<sup>1,2</sup>, Vu Le, MS<sup>1</sup>, Jill See, MPT<sup>1</sup>, Kristin Pearson-Fuhrhop, PhD<sup>1,3</sup>, Erin Burke Quinlan, PhD<sup>1,3</sup>, Robert J. Zhou, BA<sup>1</sup>, Renee Augsberger, MHA, OTR/L<sup>1</sup>, Xuan A. Tran, BS<sup>1</sup>, Nizan Friedman, PhD<sup>6</sup>, David J. Reinkensmeyer, PhD<sup>3,4,5,6</sup>, and Steven C. Cramer, MD<sup>1,3,4,5</sup>

<sup>1</sup>Dept. Neurology, UC Irvine

<sup>2</sup>Dept. Physical Therapy, Chapman University

<sup>3</sup>Dept. Anatomy & Neurobiology, UC Irvine

<sup>4</sup>Dept. Physical Medicine & Rehabilitation, UC Irvine

<sup>5</sup>Sue & Bill Gross Stem Cell Research Center, UC Irvine

<sup>6</sup>Dept. of Biomedical Engineering, UC Irvine

### Abstract

**Background**—Though rehabilitation therapy is commonly provided after stroke, many patients do not derive maximal benefit due to access, cost, and compliance. A telerehabilitation-based program may overcome these barriers. We designed then evaluated a home-based telerehabilitation system in patients with chronic hemiparetic stroke.

**Methods**—Patients were 3–24 months post-stroke with stable arm motor deficits. Each received 28 days of telerehabilitation using a system delivered to their home. Each day consisted of one structured hour focused on individualized exercises and games, stroke education, and an hour of free-play.

**Results**—Enrollees (n=12) had baseline Fugl-Meyer (FM) score of 39±12 (mean±SD). Compliance was excellent: participants engaged in therapy on 329/336 (97.9%) assigned days. Arm repetitions across the 28 days averaged 24,607±9,934 per subject. Arm motor status showed significant gains (FM change 4.8±3.8 points, p=0.0015), with half of subjects exceeding the minimal clinically important difference. Although scores on tests of computer literacy declined with age (r=-0.92, p<0.0001), neither the motor gains nor the amount of system use varied with computer literacy. Daily stroke education via the telerehabilitation system was associated with a 39% increase in stroke prevention knowledge (p=0.0007). Depression scores obtained in person correlated with scores obtained via the telerehabilitation system 16 days later (r=0.88, p=0.0001). In-person blood pressure values closely matched those obtained via this system (r=0.99, p<0.0001).

**Conclusions**—This home-based system was effective in providing telerehabilitation, education, and secondary stroke prevention to participants. Use of a computer-based interface offers many opportunities to monitor and improve the health of patients after stroke.

### Keywords

stroke; rehabilitation; telehealth; motor; games

---

## Introduction

Rehabilitation therapy is provided to most patients after discharge following acute stroke admission<sup>1</sup>. However, patients may not derive maximal benefits from rehabilitation therapy due to factors such as limited access to appropriate providers, increasing costs that reduce the amount of care provided, difficulty traveling to appointments, and poor patient compliance. Furthermore, even when a patient engages in rehabilitation therapy, doses provided can be strikingly low and fail to approximate those corresponding to favorable preclinical stroke studies<sup>2-4</sup>.

Evidence supports the use of telehealth in the delivery of acute stroke care, but the utility of telehealth in the post-acute treatment of patients with stroke is less clear<sup>5,6</sup>. Telerehabilitation has been defined as “the delivery of rehabilitation services via information and communication technologies”<sup>7</sup>, and therefore can include a broad range of services. This is consistent with the holistic framework outlined by Demiris et al.<sup>8</sup>, who suggested that home-based post-stroke telerehabilitation should include support that spans an array of medical, mental health, and other services; and also addresses many of the specific priorities outlined by the Canadian Stroke Network Consensus Conference on Stroke Rehabilitation and Research<sup>9</sup>. Our long-term goal is to improve patient outcomes after stroke by incorporating telerehabilitation methods that employ these priorities. The short-term goal, addressed in this report, was to develop a telerehabilitation system that provides home-based rehabilitation therapy, focused here on the upper extremity, which builds on standard of care physical/occupational therapy, embeds principles of learning and plasticity, uses a game-based approach, and includes regular videoconference contact between the treatment team and the patient at home.

The current report describes this system and its performance in a pilot study. The main goals of this study were to assess (1) feasibility, measured as patient compliance with assigned sessions, and (2) arm motor gains, measured as the change in arm motor Fugl-Meyer (FM) score from baseline to one-month post-therapy, with this home-based telerehabilitation system. Secondary goals included determining whether compliance and arm motor gains could be achieved independent of the participant’s computer literacy. In addition, as part of the holistic framework, other issues relevant to the long-term care of patients with stroke were examined using this telerehabilitation system, including assessment of risk factors via blood pressure recordings, secondary stroke complications via depression screening, and stroke knowledge via daily education.

## Methods

### Overview of the study and telerehabilitation system

After providing written, informed consent, subjects received 28 days of home-based telerehabilitation, organized as two 14-day blocks of daily telerehabilitation separated by a 1–3 week break; 14 days was selected based on the success of the EXCITE trial of constraint-induced therapy<sup>10</sup>, which showed benefits when making daily demands of patients for a 14-day period. Entry criteria required age >18 years, stroke (ischemic or intracerebral hemorrhage) with onset 3–24 months prior, arm Fugl-Meyer score 22–55, either grip or pinch strength >1 kg, and visual acuity at least 20/40; motor deficits could not be unstable, defined as FM score change >3 points from Visit 1 to Visit 2, which had to be at least one week apart; and subjects could not have deficits in language (NIHSS Q9 <2) or attention (NIHSS Q11 <2) that interfered with study participation.

Subjects made six in-person Visits to the lab for behavioral assessments, from baseline to 1-month post-therapy. At Visit 2, a behavioral contract<sup>10</sup> was signed that included patient goals, duties, and the dates/times of home-based therapy, after which the telerehabilitation system was introduced then delivered by a team member to the patient's home where it was assembled. Subjects were given an instruction manual that covered all devices and their usage, and all games/exercises and their rules, and were encouraged to phone the study team for technical support if needed. In addition, patients took a pill each day one hour prior to therapy, in a double-blind manner, organized as 14 consecutive days of L-Dopa or placebo; there was no significant effect of drug, and these results will be reported elsewhere. Baseline testing included electroencephalography (EEG), the results of which have been reported previously<sup>11</sup>. This study was approved by the Institutional Review Board of UC Irvine.

Design of the telerehabilitation system emphasized patient convenience, ease of use, simple large-font instructions with clear options and easy navigation, feedback, regular interaction with a clinician, and patient autonomy in choosing some therapy details<sup>12</sup>. The approach was built on established principles of learning and plasticity: training was challenging, repeated many times, motivating, interesting, and intensive<sup>12–18</sup>. Game design started with therapists outlining movements to be performed by patients, then USB-based devices were selected to transduce these movements, and finally the device signals were used to drive game play. All telerehabilitation communications between the study team and the patient's home were HIPAA-compliant and used Polycom Converged Management Application client software on both the therapist (in the lab) and patient (in the home) computers.

### Subject treatment

Each of the 28 days of therapy consisted of one required hour of activities selected by the therapist and one optional hour of games selected by the patient. A licensed occupational or physical therapist in the lab reviewed the patient's baseline exam, and then used a Therapist Control Panel program designed for use with this telerehabilitation system to generate the initial content for the required hour of therapy then upload it to the home-based telerehabilitation system. The patient saw the day's 1-hour itinerary and used large buttons and arrows on the mat (Figure 1A) to move through itinerary items. The telerehabilitation

computer software was locked such that patients could only run the telerehabilitation program, and the system would not operate beyond the permitted number of minutes.

During the required hour, patients proceeded through activities that included:

1. Games focused on arm motor therapy. A total of 18 games were available. Therapists could adjust game duration, difficulty, and input device used to drive game play. Games emphasized control of range, speed, timing, and accuracy of arm movements, and spanned categories that included casino, arcade, music, and memory games.
2. Therapeutic arm exercises. A total of 67 upper extremity exercises were also available.
3. Stroke education, consisting of five minutes/day during which patients were presented five of the practiced questions from the stroke knowledge quiz, in pseudo-random order, asked to select the correct answer from four choices, then provided with the correct answer and a brief explanation.
4. Videoconferences three times/week. There were three videoconferences/week between the patient at home and a study team member in the lab (one/week by a licensed therapist and two/week by a research assistant). During their videoconference with the patient at home, the therapist reviewed the prior week's activities with the subject including scores, preferences, and issues in an open-ended manner. This was followed by a structured interview that included specific inquiries related to pain, adverse events, and telerehabilitation system performance. The therapist documented the interaction using standard clinic notes. Game scores and data on telerehabilitation system usage were recorded and transmitted to the lab in real time. Therapists reviewed those data, along with patient responses from questionnaires and patient input during videoconferences, to modify choice, duration, and/or difficulty level of telerehabilitation games/exercises, and then create and upload (silently, at any time of day) revised itineraries for future days. During videoconferences, therapists worked with patients to adjust the challenge level presented during telerehabilitation sessions as appropriate, providing feedback and encouragement to drive motivation, according to standard of care practice.

### Subject testing

Most testing was done live, in the lab. Baseline assessments (Visits 1 and 2) included vital signs, MRI, FM Scale<sup>19</sup>, Box & Blocks (B&B), NIH Stroke Scale (NIHSS), Barthel Index, Geriatric Depression Scale (GDS)-15 question form, Mini-Mental Status Exam (MMSE), Optimization in Primary and Secondary Control Scale (adapted from<sup>20</sup>), The Medical Outcomes Study Social Support Survey<sup>21</sup>, Mental Adjustment to Stroke Scale (Fighting Spirit subscore)<sup>22</sup>, Stroke-Specific Quality Of Life Scale (SS-QOL)<sup>23</sup>, modified functional reach forward displacement, shoulder pain using a visual analog scale from 1–10, gait velocity from 10 meter walk test, and an adaptation of the subjective and objective Computer–Email–Web Fluency Scale tests of computer literacy<sup>24</sup>. The FM was scored at all

six Visits. At Visit 2, patients also took a stroke knowledge quiz with 40 multiple-choice questions (4 choices/question) focused on knowledge about stroke symptoms and risk factors. Of these 40 questions, 20 were subsequently practiced during the 5 minutes of daily telerehabilitation stroke education over the next 14 days, and 20 were not practiced. At Visit 3, two weeks later, the same 40-question quiz was administered. This entire process was repeated across Visits 4 and 5 using a novel 40-question stroke knowledge quiz. Thus from Visit 2 to Visit 5 patients were serially tested on a total of 40 practiced and 40 unpracticed questions.

Other testing was done remotely, via the telerehabilitation system. One set of telerehabilitation-based assessments was used to assess validity. After one therapist assessed depression in person in the lab during Visit 2 using the GDS, as above, a second therapist assessed depression via the videoconference that occurred on the first day of telerehabilitation using the patient health questionnaire (PHQ)-9 and PHQ-2 scales. Also during Visit 2, a therapist measured the patient's systolic (SBP)/diastolic (DBP) blood pressure and pulse using a USB-based wrist blood pressure cuff (ION Health) in person. The patient then measured the same vital signs at the start of each day of telerehabilitation at home using the same cuff. A second set of telerehabilitation-based assessments was used to characterize subjects at baseline. Two measures were obtained in this group, the Stroke Impact Scale (hand subsection), which measures difficulty of hand use<sup>25</sup>, and the Stroke Self-Efficacy Questionnaire<sup>26</sup>. A third set of remote assessments was used to assess patient status on a daily level. There four such assessments: daily blood pressure, Target Bopping game, Wrist Targeting game, and Help The Postman game.

## Hardware

The core telerehabilitation system consisted of a 24"×48" table, bridge chair, Dell Latitude E5420 laptop (with 14" display, internal webcam, and added fisheye lens), Verizon wireless USB modem, USB-based wrist blood pressure cuff, and custom-made USB-based mat that used contact-sensitive switches (Figure 1A). Other input devices, connected via USB and prescribed per therapist judgment, included a wrist splint with a sensor on its hinge to record the wrist flexion/extension position, an accelerometer with a trigger added that was used to move the screen cursor and trigger events, and a Music Glove<sup>27</sup>. The laptop keyboard was covered and in fact no keyboard was used. Instead, patients pressed buttons and arrows on the mat to move through the steps of telerehabilitation assignment. Standard non-electronic exercise equipment, such as a weighted dowel and digit exerciser, was also provided.

## Data analysis

The number of repetitions performed by subjects was assessed by either exact digital recordings or review of scores; this was done for a convenience sample of 12 different sessions (two separate treatment days, for each of six different patients).

Parametric statistical methods were used for measures for which the normality assumption was valid, using raw or transformed values, otherwise non-parametric methods were used. All analyses were two-tailed with  $\alpha=0.05$  and were performed using JMP-8 software. The primary outcome measure was the change in FM score from baseline (mean of Visits 1

and 2) to 1-month post-therapy (Visit 6). For the secondary analyses, Bonferroni correction for multiple comparisons was performed as follows: there were three secondary motor outcomes (B&B, FM proximal subscore, and FM distal subscore), and so  $p < 0.0167$  was used to define significance. Each test of computer literacy was compared with four measures (age, usage, change in FM score, and change in education scores), and so  $p < 0.0125$  defined significance. There were two tests of validation of telehealth methods for depression screening (PHQ-2 and PHQ-9) and so here  $p < 0.025$  was used. There were three tests of validation of telehealth vital signs measurement (pulse, DBP, and SBP), and so  $p < 0.0167$  defined significance.

## Results

### Subjects

A total of 12 patients were enrolled, with no subject dropout. Enrollees were  $54 \pm 17$  years of age (mean  $\pm$  SD) and had moderate-severe motor deficits at baseline, with FM =  $39 \pm 12$ , range 23–55 (Table 1). All but one lived with a caregiver relative. Verizon wireless reception, used to connect the telerehabilitation system to the internet, was present in all patients' homes. Table S1 (Supplement) considers problems, solutions, and lessons learned.

### Patient experience

Compliance was exceptionally good, with subjects engaging in therapy for at least 30 of the required 60 minutes on 329 of 336 (97.9%) assigned days. One patient, with the lowest motor status (FM score = 23) did not complete at least 30 minutes for four sessions, due to fatigue; another patient did not complete at least 30 minutes due to hardware malfunction; and one patient missed two complete sessions due to conflict with other medical appointments. On 94 (28.6%) of the 336 days, the treatment team was contacted by phone for assistance, a rate that decreased over time for each patient across their 28 days of therapy (reflecting patient familiarity with the system; Supplement Figure S1A) and decreased over the seven months during which this study was conducted (reflecting improved performance by the study team; Supplement Figure S1B). Non-electronic devices were used only during a minority of the exercises, accounting for only a very small portion of treatment content, approximately 2–4 minutes/day on average.

Patients completed a questionnaire (Supplement, Table S2) regarding their telerehabilitation experience at the end of all study procedures, 1 month after end of treatment, rating the experience very favorably: when asked "What did you think of the treatment overall?" using a 5-level Likert scale (5 being most favorable), the mean score was 4.4. When asked "Would you recommend this type of treatment to someone with a similar condition after stroke?", the mean score was 4.8. In terms of system components, the lowest rating pertained to some of the hardware that was difficult to use and the highest rating was for the appointment reminder system. Other patient ratings appear in Supplement Table S4.

In terms of specific patient preferences, memory games (e.g., Simon) were not preferred. Patients did not express concerns with the exercise assignments. The most popular specific free-play choices were casino (slot machine, pachinko) and arcade (carnival shooting, duck

hunt) games (Figure 1B/C), but preferences varied widely. Overall, the percent of free-play time spent on music games was  $7.0\pm 6.7\%$ ; on carnival games,  $15.2\pm 7.6\%$ ; on arcade games,  $24.4\pm 19.8\%$ ; on puzzle games,  $24.6\pm 23.4\%$ ; and on casino games,  $28.8\pm 30.5\%$ . The percent of free-play time spent in each free-play category was not related to subject's self-reported interests in these respective categories according to a survey performed at baseline.

Patients underwent daily motor assessments. Each day at home, patients played the Target Bopping game (Supplement, Table S3.14), whereby patients grasped a cylinder to press on different input mat targets according to on-screen instructions. Average scores increased significantly over time ( $r=0.65$ ,  $p=0.0002$ , Pearson's correlation coefficient, Figure 2), by 122% across the 28 days of play of this 3-minute game, from 46 targets successfully bopped during play on day 1 at home to 102 targets on day 28 at home. Day-to-day variability was common. Scores on day 1 correlated significantly with baseline Fugl-Meyer scores ( $r=0.78$ ,  $p<0.003$ , Pearson's correlation coefficient), however, the slope of game scores over the 28 days of home play was not related to the change in Fugl-Meyer score over the same time interval ( $p>0.5$ ). For two other games played daily at home, Wrist Targeting and Help The Postman, reliable data could not be obtained over time, as these two relied on the wrist splint, which proved difficult to don and calibrate consistently.

Active time (time during which games and exercises were performed) averaged  $60\pm 10$  min/day, including an average of  $22\pm 24$  min/day of free-play. Total time (active time plus education questions, measuring blood pressure, reading game instructions, donning devices, and taking breaks between tasks) averaged  $182\pm 61$  min/day. The average number of arm repetitions per day was  $879\pm 355$ , and thus the average number of repetitions per subject across the entire 28 days was  $24,607\pm 9,934$ .

### Therapist experience

Therapists were able to create an initial treatment plan, send it to the telerehabilitation system in the patient's home, review patient scores and performance remotely, and then upload revised therapy settings to the telerehabilitation system in the patient's home without difficulty, at any time of day. Videoconferences were performed as planned, without incident. Of the 48 planned weekly videoconferences with a therapist across all 12 patients, 46 were attended as planned. During these videoconferences, the therapist helped the patient adjust to assigned games, exercises, and equipment, or modified assignments. Common issues that arose were the need to reduce elbow extension, reduce pronation/supination, increase arm elevation, and refer to the Instruction Manual provided by study personnel. Also, patients were asked if any activities were associated with shoulder pain. Responses varied widely across patients and across weeks with no trend for worsening over time, and included pain from the daily target bopping target game, the Simon game, various exercises, and reaching for the furthest targets on the input mat.

### Arm motor gains

FM scores were stable at baseline, changing an average of 0.1 points ( $p>0.9$ ) from Visit 1 to Visit 2, which were  $15\pm 3$  days apart. Regarding the primary outcome measure, arm FM score increased from baseline to one month post-therapy by  $4.8\pm 3.8$  points ( $p=0.0015$ ,



paired t-test, Figure 3A), with the gain in six subjects exceeding the minimal clinically important difference (MCID) of 4.25 points<sup>28</sup>. Gains in proximal and distal FM score subsections are described in the Supplement (Figure S2). None of the measures in Table 1 were significantly correlated with this FM score increase, whether assessed with or without correction for multiple comparisons. Gains in FM were not uniformly distributed across the 33 components comprising the FM score. For example there were no questions where at least 10 of 12 enrollees showed a change of at least 1 point; there were 6 questions where at least 4 people showed an improvement of at least 1 point (shoulder external rotation, hand to lumbar spine, wrist flex/extend with elbow extended, wrist circumduction, palmar prehension, and movement with normal speed); and there were 4 questions where anyone showed a change of 2 points (triceps reflex, hand to lumbar spine, wrist circumduction, and palmar prehension).

The B&B score, a measure of arm function assessed from baseline to end of therapy, also increased significantly, from 13 to 16 blocks transferred over 60 seconds ( $p=0.01$ , paired t-test). Assuming an MCID of 6 blocks<sup>29</sup>, gains in four subjects achieved the MCID for this functional measure.

### **Stroke education**

Patients showed significant gains in stroke prevention knowledge that were specific to facts reinforced in daily quizzes (Figure 3B). For the 40 quiz questions that were practiced on the telerehabilitation system, the number of correct answers increased by 39%, from 22.8 to 31.7 ( $p=0.0007$ , paired t-test). For the 40 quiz questions that were not practiced, scores changed from 25.3 to 26.7 out of 40, a non-significant 5% change ( $p=0.17$ , paired t-test).

### **Findings not dependent on computer skills**

Computer literacy scores declined with age, as expected ( $\rho = -0.92$  and  $-0.90$ ,  $p<0.0001$ , Spearman's rank correlation coefficient, for the subjective and objective portions of the Computer–Email–Web Fluency Scale respectively, Figure 3C left panel). However, neither computer literacy score was significantly related to amount of system usage (Figure 3C right panel), arm motor gains (change in FM score), or stroke education gains ( $p>0.05$ , Spearman's rank correlation coefficient).

### **Validity of depression screening using telehealth methods after stroke**

The PHQ-2 and PHQ-9 were successfully scored in all subjects during the first telerehabilitation videoconference, as planned. GDS scores obtained in person at Visit 1 correlated significantly with these telerehabilitation-derived PHQ-2 scores ( $\rho=0.88$ ,  $p=0.0001$ , Spearman's rank correlation coefficient, Figure 3D) and showed a trend with respect to PHQ-9 ( $\rho=0.57$ ,  $p=0.05$ , Spearman's rank correlation coefficient) scores. At Visit 1, three patients had GDS score of 5–10 (suggestive of depression), and all three had a PHQ-2 score  $>3$  (also suggestive of depression) during the first telerehabilitation session  $16\pm 3$  days later; each was referred for further evaluation.

### Validity of BP measurement using the telerehabilitation system

The subjects recorded their BP and pulse at home using the telerehabilitation system, and the results were then automatically transmitted to the lab, on 329 of 336 (97.9%) assigned days. SBP ( $130\pm 20$  vs.  $130\pm 20$  mmHg), DBP ( $83\pm 16$  vs.  $84\pm 16$  mmHg), and pulse ( $78\pm 11$  vs.  $77\pm 11$ ) recorded by a therapist in the lab at Visit 2 closely matched values obtained by the patient using the same cuff at home during the first telerehabilitation session, which was 1–2 days later (all  $r=0.99$ ,  $p<0.0001$ , Pearson's correlation coefficient).

### Discussion

We designed a home-based telerehabilitation program that provided arm motor therapy while addressing other issues important to the outcome of patients with stroke, such as secondary prevention and screening for complications. We found that (1) patients were highly compliant (97.9% of assigned days) and rated the system favorably; (2) therapists were readily able to review patient performance and revise therapy; (3) videoconferences supported regular communication between the patient and treatment team; (4) arm motor status improved significantly overall and exceeded the MCID in half of the subjects; (5) daily education increased stroke prevention knowledge by 39%; (6) screening for depression was accurate; (7) BP measurement was very accurate; and (8) all of these findings were unrelated to subjects' computer skills. These findings support the feasibility and potential utility of this home-based program for improving outcomes after stroke.

The best treatment program is of little help to patients if they do not adhere to it, and so the telerehabilitation system was designed to maximize compliance. One key feature in this regard was the use of games to drive treatment, an approach known to increase desirability and accessibility and to promote patient participation<sup>30–33</sup>. In addition, patients were asked to sign a behavioral contract<sup>10</sup>, which increases adherence, engagement, and accountability<sup>34</sup>. Our contract listed daily times of therapy, increasing organization, facilitating videoconference planning, and enabling system alerts to begin each day's session on time. Another feature promoting compliance was that patients could pause the system as needed between assigned tasks. Also, in contrast with some commercial systems aiming to drive rehabilitation therapy in the home, our system emphasized variety, with the different modules (therapy, assessment, education, prevention, videoconferencing), numerous games and exercises, and range of different hardware input devices incorporated into individual treatment plans. In addition, patients could phone the lab if a technical question arose, a feature mainly used during subject's initial days of telerehabilitation (Supplement Figure S1A). Also, system design focused on ease of use, including simple, large-font instructions. A key finding in this regard was that computer literacy, while inversely related to age, was not related to treatment gains (Figure 3C), indicating that effective system use did not require computer skills. Patients could only use the telerehabilitation system for the itinerary assigned by the therapist, and in fact the keyboard was inaccessible, making this system more like an appliance than a computer. In general, patient compliance with home-based physical therapy ranges from 23–64%<sup>35</sup>, and after stroke 65.3% of patients report adhering to at least part of a home exercise program<sup>36</sup>. The above-described features likely underlie the superior compliance (97.9%) observed with the current telerehabilitation program.

The primary outcome measure was arm motor gains, assessed by the change in FM score from baseline to 1-month post-therapy. Findings were significant (Figure 3A), with 6 of 12 subjects achieving the MCID. The FM is a measure of body structure/function limitation, so it is worth noting that results using the B&B, a measure of activities limitation, were also significant. The current findings are based on a small cohort in a non-controlled study and so must be viewed with caution; for example, the limited sample size might have increased the risk for ascertainment bias. However, each of the 12 patients was assessed over 28 separate days, for a total of 336 telerehabilitation study days. In addition, a strength of the system that supports the validity of the current findings is the implementation of principles of neuroplasticity in game design, including the use of stimuli that are challenging, repeated many times, intense, fun, and adjustable<sup>12-18</sup>. As part of the research design, the current study limited therapy to 2 hours of therapy/day for 28 days, but greater gains may be possible with a longer duration of intensive therapy<sup>37,38</sup>, which may be pursued in future studies. An important finding is that subjects performed an average of 24,607 over 28 days, based on 879 arm repetitions each day, a value similar to the 466 movements/hour recorded in a prior study of one of the current input devices<sup>27</sup>.

Stroke is a complex chronic disease associated with many risk factors and complications. Using a computer-based interface in the home offers opportunities to address several aspects of patient health in parallel. This type of holistic approach has been advocated<sup>8,9</sup> and was incorporated into the design of the current program in three major ways. First, major depression is a common complication after stroke<sup>1</sup>, and it remains significantly underdiagnosed and poorly treated<sup>39</sup>. The current study validated a home-based telerehabilitation system for measuring depressive symptoms, as PHQ-2 ( $p=0.0001$ , Figure 3D) scores obtained using the telerehabilitation system in the home were significantly related to GDS scores obtained in person, independently, 16 days prior. Second, many patients with stroke have limited knowledge about the causes and effects of this disease; for example, in studies of patients admitted for stroke, 39% could not name one sign or symptom of stroke<sup>40</sup>, while 40% could not identify a single stroke risk factor<sup>41</sup>. Stroke education delivered by the telerehabilitation system increased subjects' knowledge of stroke symptoms and risk factors, with correct scores increasing from 23 out of 40 questions at baseline (58% correct) to 32 out of 40 questions correct at end of therapy (80% correct), a relative increase of 39% ( $p=0.0007$ ). Third, many stroke survivors do not achieve good risk factor control. For example, among 1,252 U.S. survivors of stroke or myocardial infarction, long-term BP control was present in only 53%<sup>42</sup>. Using the current telerehabilitation system, patients recorded their blood pressure at home on 97.9% of the 336 assigned days, and these values closely matched those obtained independently by a therapist in the lab ( $r=0.99$ ,  $p<0.0001$ ). Such findings support the idea that a home-based telehealth system could be used to improve blood pressure control in patients with chronic stroke. Together these findings suggest that, in addition to driving motor rehabilitation, a home-based telerehabilitation system can assess patients for post-stroke complications, educate patients about stroke, and assess risk factor control. Future studies can build on these results to examine how such measures may be used to improve outcomes.

There are a number of strengths to the current study. A holistic framework was implemented, within the convenience of the home, with several different features, and with

no dependency on computer skills. Effective therapist-facing software is as important as the patient-facing program<sup>43</sup> and performed well in this pilot study. Video-based techniques may be a key component to effective telerehabilitation<sup>44</sup> and were a central feature of the system under study. Although the sample size in this pilot study was modest, good compliance with therapy was found in patients with gait velocity < 0.4 m/sec (i.e., mobility limited to the home) or MMSE score as low as 23 (Table 1). There are also important weaknesses to the current study. The sample size was limited. There was no control group, making it difficult to attribute specific outcomes to particular aspects of treatment content. Patients needed substantial support, at least during the initial days at home (Figure S1A). Hopefully, the lessons learned during study performance (Supplement, Table S1) will inform approaches that reduce the need for assistance in future telerehabilitation studies. At baseline (Table 1), patients had high consequence-related motivation for rehabilitation therapy, very good social support, and overall high optimism. It will be important to evaluate compliance with home-based telerehabilitation therapy among subjects with less motivation, social support, and optimism to better understand the extent to which current results generalize across the entire spectrum of stroke survivors. Much of the telerehabilitation was based on tabletop activities, although there were some 3-dimensional components to the games and exercises, and future telerehabilitation-based therapies may wish to expand the amount of time devoted to 3-dimensional functional tasks. A separate control group was not enrolled, however, patients were enrolled after the time when arm motor recovery is generally considered to be at a plateau<sup>45</sup> and confirmation of stable deficits at baseline increased confidence that observed gains in motor status were linked to the telerehabilitation intervention. Finally, this was a pilot study with a limited sample size, unblinded assessors, and no control group; however, results are promising for extension of the current approach to a larger clinical trial.

## Conclusions

The burden of post-stroke disability remains large. Telehealth methods may be useful to improve outcomes, particularly given decreasing lengths of stay for inpatient stroke care<sup>46,47</sup>, and moderate evidence exists for equivalence to conventional rehabilitation<sup>44</sup>. The current study supports the feasibility and utility of a home-based system to effectively deliver telerehabilitation, improve patient education, screen for complications of stroke, and provide patients at home with a means for interaction with medical personnel.

## Supplementary Material

Refer to Web version on PubMed Central for supplementary material.

## Acknowledgments


Support was provided by NIH (K24-HD074722 and UL1-TR000153) and the UC Irvine Dean's Triumvirate Award. Dr. Cramer has served as a consultant for Dart Neuroscience, RAND Corporation, Toyama, MicroTransponder, and Roche, and is cofounder of personalRN.

## References

1. Winstein CJ, Stein J, Arena R, et al. Guidelines for Adult Stroke Rehabilitation and Recovery: A Guideline for Healthcare Professionals From the American Heart Association/American Stroke Association. *Stroke*. 2016; 47(6):e98–e169. [PubMed: 27145936]
2. Lang CE, Macdonald JR, Reisman DS, et al. Observation of amounts of movement practice provided during stroke rehabilitation. *Arch Phys Med Rehabil*. 2009; 90(10):1692–1698. [PubMed: 19801058]
3. Bernhardt J, Chan J, Nicola I, Collier JM. Little therapy, little physical activity: rehabilitation within the first 14 days of organized stroke unit care. *Journal of rehabilitation medicine*. 2007; 39(1):43–48. [PubMed: 17225037]
4. Kimberley TJ, Samargia S, Moore LG, Shakya JK, Lang CE. Comparison of amounts and types of practice during rehabilitation for traumatic brain injury and stroke. *Journal of rehabilitation research and development*. 2010; 47(9):851–862. [PubMed: 21174250]
5. Laver KE, Schoene D, Crotty M, George S, Lannin NA, Sherrington C. Telerehabilitation services for stroke. *Cochrane Database Syst Rev*. 2013; 12:CD010255.
6. Agostini M, Moja L, Banzi R, et al. Telerehabilitation and recovery of motor function: a systematic review and meta-analysis. *Journal of telemedicine and telecare*. 2015; 21(4):202–213. [PubMed: 25712109]
7. Brennan D, Tindall L, Theodoros D, et al. A blueprint for telerehabilitation guidelines. *International journal of telerehabilitation*. 2010; 2(2):31–34. [PubMed: 25945175]
8. Demiris G, Shigaki CL, Schopp LH. An evaluation framework for a rural home-based telerehabilitation network. *J Med Syst*. 2005; 29(6):595–603. [PubMed: 16235812]
9. Bayley MT, Hurdowar A, Teasell R, et al. Priorities for stroke rehabilitation and research: results of a 2003 Canadian Stroke Network consensus conference. *Arch Phys Med Rehabil*. 2007; 88(4):526–528. [PubMed: 17398256]
10. Wolf SL, Winstein CJ, Miller JP, et al. Effect of constraint-induced movement therapy on upper extremity function 3 to 9 months after stroke: the EXCITE randomized clinical trial. *JAMA*. 2006; 296(17):2095–2104. [PubMed: 17077374]
11. Wu J, Quinlan EB, Dodakian L, et al. Connectivity measures are robust biomarkers of cortical function and plasticity after stroke. *Brain*. 2015; 138(Pt 8):2359–2369. [PubMed: 26070983]
12. Jimison, H., Gorman, P., Woods, S., et al. Barriers and Drivers of Health Information Technology Use for the Elderly, Chronically Ill, and Underserved. Rockville, MD: Agency for Healthcare Research and Quality; 2008. Evidence Report/Technology Assessment No. 175. AHRQ Publication No. 09-E004
13. Woldag H, Hummelsheim H. Evidence-based physiotherapeutic concepts for improving arm and hand function in stroke patients: a review. *J Neurol*. 2002; 249(5):518–528. [PubMed: 12021939]
14. Takahashi CD, Der-Yeghiaian L, Le V, Motiwala RR, Cramer SC. Robot-based hand motor therapy after stroke. *Brain*. 2008; 131(Pt 2):425–437. [PubMed: 18156154]
15. Kleim JA, Jones TA. Principles of experience-dependent neural plasticity: implications for rehabilitation after brain damage. *J Speech Lang Hear Res*. 2008; 51(1):S225–239. [PubMed: 18230848]
16. Cramer SC, Sur M, Dobkin BH, et al. Harnessing neuroplasticity for clinical applications. *Brain*. 2011; 134(Pt 6):1591–1609. [PubMed: 21482550]
17. Cramer SC. Repairing the human brain after stroke: I. Mechanisms of spontaneous recovery. *Ann Neurol*. 2008; 63(3):272–287. [PubMed: 18383072]
18. Dobkin BH, Dorsch A. The promise of mHealth: daily activity monitoring and outcome assessments by wearable sensors. *Neurorehabil Neural Repair*. 2011; 25(9):788–798. [PubMed: 21989632]
19. See J, Dodakian L, Chou C, et al. A standardized approach to the fugl-meyer assessment and its implications for clinical trials. *Neurorehabil Neural Repair*. 2013; 27(8):732–741. [PubMed: 23774125]

20. Mackay J, Charles ST, Kemp B, Heckhausen J. Goal striving and maladaptive coping in adults living with spinal cord injury: associations with affective well-being. *Journal of aging and health*. 2011; 23(1):158–176. [PubMed: 20876363]
21. Sherbourne CD, Stewart AL. The MOS social support survey. *Social science & medicine*. 1991; 32(6):705–714. [PubMed: 2035047]
22. Lewis SC, Dennis MS, O'Rourke SJ, Sharpe M. Negative attitudes among short-term stroke survivors predict worse long-term survival. *Stroke*. 2001; 32(7):1640–1645. [PubMed: 11441213]
23. Williams LS, Weinberger M, Harris LE, Clark DO, Biller J. Development of a stroke-specific quality of life scale. *Stroke*. 1999; 30(7):1362–1369. [PubMed: 10390308]
24. Bunz U. The Computer-Email-Web (CEW) Fluency Scale – Development and Validation. *International Journal of Human-Computer Interaction*. 2004; 17(4):479–506.
25. Duncan P, Wallace D, Lai S, Johnson D, Embretson S, Laster L. The stroke impact scale version 2.0. Evaluation of reliability, validity, and sensitivity to change. *Stroke*. 1999; 30(10):2131–2140. [PubMed: 10512918]
26. Jones F, Partridge C, Reid F. The Stroke Self-Efficacy Questionnaire: measuring individual confidence in functional performance after stroke. *Journal of clinical nursing*. 2008; 17(7B):244–252. [PubMed: 18578800]
27. Zondervan DK, Friedman N, Chang E, et al. Home-based hand rehabilitation after chronic stroke: Randomized, controlled single-blind trial comparing the MusicGlove with a conventional exercise program. *Journal of rehabilitation research and development*. 2016; 53(4):457–472. [PubMed: 27532880]
28. Page SJ, Fulk GD, Boyne P. Clinically important differences for the upper-extremity Fugl-Meyer Scale in people with minimal to moderate impairment due to chronic stroke. *Phys Ther*. 2012; 92(6):791–798. [PubMed: 22282773]
29. van der Lee J, Beckerman H, Lankhorst G, Bouter L. The responsiveness of the Action Research Arm test and the Fugl-Meyer Assessment scale in chronic stroke patients. *J Rehabil Med*. 2001; 33(3):110–113. [PubMed: 11482350]
30. Baranowski T, Buday R, Thompson DI, Baranowski J. Playing for real: video games and stories for health-related behavior change. *American journal of preventive medicine*. 2008; 34(1):74–82. [PubMed: 18083454]
31. Brox E, Fernandez-Luque L, Tøllefsen T. Healthy Gaming – Video Game Design to promote Health. *Applied Clinical Informatics*. 2011; 2:128–142. [PubMed: 23616865]
32. Lieberman, D. Designing serious games for learning and health in informal and formal settings. In: Ritterfeld, M., Vorderer, P., editors. *Serious games: Mechanisms and effects*. New York: Routledge; 2009. p. 117-130.
33. Chou, Y. *Actionable Gamification - Beyond Points, Badges, and Leaderboards*. Fremont, CA: Octalysis Media; 2015.
34. Winstein CJ, Miller JP, Blanton S, et al. Methods for a multisite randomized trial to investigate the effect of constraint-induced movement therapy in improving upper extremity function among adults recovering from a cerebrovascular stroke. *Neurorehabil Neural Repair*. 2003; 17(3):137–152. [PubMed: 14503435]
35. Sluijs EM, Kok GJ, van der Zee J. Correlates of exercise compliance in physical therapy. *Phys Ther*. 1993; 73(11):771–782. discussion 783–776. [PubMed: 8234458]
36. Miller KK, Porter RE, DeBaun-Sprague E, Van Puymbroeck M, Schmid AA. Exercise after Stroke: Patient Adherence and Beliefs after Discharge from Rehabilitation. *Top Stroke Rehabil*. 2016:1–7.
37. McCabe J, Monkiewicz M, Holcomb J, Pundik S, Daly JJ. Comparison of robotics, functional electrical stimulation, and motor learning methods for treatment of persistent upper extremity dysfunction after stroke: a randomized controlled trial. *Arch Phys Med Rehabil*. 2015; 96(6):981–990. [PubMed: 25461822]
38. Griffith, V. *A Stroke in the Family*. New York: Delacorte Press; 1970.
39. Herrmann N, Seitz D, Fischer H, et al. Detection and treatment of post stroke depression: results from the registry of the Canadian stroke network. *International journal of geriatric psychiatry*. 2011; 26(11):1195–1200. [PubMed: 21360753]

40. Kothari R, Sauerbeck L, Jauch E, et al. Patients' awareness of stroke signs, symptoms, and risk factors. *Stroke*. 1997; 28(10):1871–1875. [PubMed: 9341687]
41. Zerwic J, Hwang SY, Tucco L. Interpretation of symptoms and delay in seeking treatment by patients who have had a stroke: exploratory study. *Heart & lung: the journal of critical care*. 2007; 36(1):25–34. [PubMed: 17234474]
42. Qureshi AI, Suri MF, Guterman LR, Hopkins LN. Ineffective secondary prevention in survivors of cardiovascular events in the US population: report from the Third National Health and Nutrition Examination Survey. *Arch Intern Med*. 2001; 161(13):1621–1628. [PubMed: 11434794]
43. Putrino D. Telerehabilitation and emerging virtual reality approaches to stroke rehabilitation. *Curr Opin Neurol*. 2014; 27(6):631–636. [PubMed: 25333603]
44. Chen J, Jin W, Zhang X, Xu W, Liu X-N, Ren C-C. Telerehabilitation approaches for stroke patients: systematic review and meta-analysis of randomized controlled trials. *J Stroke Cerebrovasc Dis*. 2015; 24:2660–2668. [PubMed: 26483155]
45. Nakayama H, Jorgensen H, Raaschou H, Olsen T. Recovery of upper extremity function in stroke patients: the Copenhagen Stroke Study. *Arch Phys Med Rehabil*. 1994; 75(4):394–398. [PubMed: 8172497]
46. Ottenbacher KJ, Smith PM, Illig SB, Linn RT, Ostir GV, Granger CV. Trends in length of stay, living setting, functional outcome, and mortality following medical rehabilitation. *Jama*. 2004; 292(14):1687–1695. [PubMed: 15479933]
47. Tong X, Kuklina EV, Gillespie C, George MG. Medical complications among hospitalizations for ischemic stroke in the United States from 1998 to 2007. *Stroke*. 2010; 41(5):980–986. [PubMed: 20203317]



**Figure 1.**


[A] Hardware for the telerehabilitation system included a standard table, chair, laptop with keyboard covered, Verizon wireless modem, mat used for game play and moving through the day's itinerary (via the buttons and arrows), standard rehabilitation equipment, and multiple USB-based input devices to drive game play.

[B] Carnival shooting, a game in which the patient performed supination/pronation movements to move the red cursor then squeezed a trigger using a lateral pinch movement to shoot at the yellow, but not red, ducks.


[C] Slot machine, a game in which the patient had to perform shoulder extension movements to stop each of the three reels from spinning at the correct time to match all three symbols.

[D] Shoulder abduction/adduction, one of the 67 available exercises.





**Figure 2.** Mean±SEM scores on the bopping game, during which subjects had to use a cylinder held in the stroke-affected hand to bop the tabletop mat target that was indicated on the computer screen. Scores increased significantly across the 28 days of therapy ( $r=0.65$ ,  $p=0.0002$ ); note that this increase in scores over time remained significant if values from the first three days were removed.



**Figure 3.**

[A] Arm motor deficits (mean±SEM) were moderate-severe at baseline and increased 1-month post-therapy by 4.8 points ( $p=0.0015$ ), with gains in 6/12 subjects exceeding the minimal clinically important difference. FM=Fugl-Meyer.

[B] Stroke education using the home-based telerehabilitation system improved scores on tests of stroke prevention knowledge. *White bars*: For the 40 questions that were practiced as part of 28 days of daily stroke education, correct answers increased 39%, from 22.8 to 31.7 ( $p=0.0007$ ). *Gray bars*: As a control, 40 questions were not practiced but were also serially tested, and these scores did not change significantly ( $p=0.17$ ).

[C] Computer literacy was inversely related to age ( $r = -0.92$ ,  $p<0.0001$ , left panel) but not system usage (average amount of time subjects spent each day using the telerehabilitation system,  $p>0.5$ , right panel).

[D] Depression score in the lab at Visit 1 using the Geriatric Depression Scale (GDS) correlated significantly ( $r=0.88$ ,  $p=0.0001$ ) with the depression score obtained an average of 16 days later in the home on the telerehabilitation system using the Patient Health Questionnaire (PHQ)-2.

Author Manuscript

Author Manuscript

Author Manuscript

Author Manuscript

**Table 1**

## Baseline subject characteristics

N	12
Age	54±17 (range 26–75)
Gender	6M/6F
Time post-stroke (weeks)	24 [21–36]
Infarct volume (cc)	16 [2–32]
Affected side	6R/6L
Hypertension	58%
Diabetes mellitus	25%
Hyperlipidemia	67%
Fugl-Meyer Scale	
--Total	39.4±12.3 (range 23–56)
--Proximal	23.2±5.7 (range 13.5–31.5)
--Distal	12.1±7.3 (range 1–20.5)
Box & Blocks	13±12
Stroke Impact Scale (hand motor subsection)	2.0±0.8 <sup>a</sup>
NIH Stroke Scale	4 [2–5]
MMSE	27.5 [25–30] (range 23–30)
Barthel Index	92.5 [86–100]
Geriatric Depression Scale	4±2.5 (range 0–8)
Optimization in Primary and Secondary Control Scale	21±1.9 <sup>b</sup>
Medical Outcomes Study Social Support Survey	82±10 <sup>c</sup>
Mental Adjustment to Stroke Scale (Fighting Spirit subsection)	57±5 <sup>d</sup>
SS-QOL	3.7±0.5 <sup>e</sup>
Modified functional reach forward displacement (cm)	36±11
Shoulder pain	4 [0–5.75] <sup>f</sup>
Gait velocity (m/sec)	0.78±0.47 <sup>g</sup>
Stroke Self-Efficacy Questionnaire	46±1 <sup>h</sup>

Values are mean±SD or median [IQR].

<sup>a</sup>Scores five items from 1–5 to measure difficulty of hand use, with higher scores indicating better function. Results indicate substantial difficulty with hand use.

<sup>b</sup>Measures consequence-related motivation for rehabilitation therapy, reflecting the patient's level of dedication to treatment goals. Subjects rated their level of agreement from 1 to 4 for six statements that reflected motivation; maximum score is 24. Results indicate overall high motivation for rehabilitation therapy.

<sup>c</sup>Scores 19 items from 1–5 to measure social support; maximum score is 95. Results indicate very good social support.

<sup>d</sup>Scores 16 items from 1–4 to measure optimism, with higher scores indicating more positive thinking; maximum score is 64. Results indicate high optimism.

<sup>e</sup>Measures quality of life by averaging scores (from 1–5) across 12 domains, with higher scores indicating higher patient-reported quality of life.

<sup>f</sup> Measured using a visual analog scale from 1–10. Value is provided across all subjects; value was 5 [4–6.75] among the 8 reporting any shoulder pain at baseline.

<sup>g</sup> 50% of subjects had gait velocity < 0.8 m/sec and so did not have full mobility in the community, and 25% of subjects were <0.4 m/sec and so mobility was limited to the home.

<sup>h</sup> Patients rate their level of confidence for 13 tasks and functional activities from 1 to 10, with higher scores indicating higher confidence. Results reflect substantial reductions in self-efficacy and confidence.

Author Manuscript

Author Manuscript

Author Manuscript

Author Manuscript