

UCSF

UC San Francisco Previously Published Works

Title

Development and validation of a person-centered abortion scale: the experiences of care in private facilities in Kenya

Permalink

<https://escholarship.org/uc/item/8jn2b144>

Journal

BMC Women's Health, 20(1)

ISSN

1472-6874

Authors

Sudhinaraset, May
Landrian, Amanda
Afulani, Patience A
[et al.](#)

Publication Date

2020-12-01

DOI

10.1186/s12905-020-01071-w

Copyright Information

This work is made available under the terms of a Creative Commons Attribution License, available at <https://creativecommons.org/licenses/by/4.0/>


Peer reviewed

RESEARCH ARTICLE

Open Access



Development and validation of a person-centered abortion scale: the experiences of care in private facilities in Kenya

May Sudhinaraset¹, Amanda Landrian^{1*} , Patience A. Afulani², Beth Phillips², Nadia Diamond-Smith² and Sun Cotter²

Abstract

Background: There is a need for a standardized way to measure person-centered care for abortion. This study developed and validated a measure of person-centered abortion care.

Methods: Items for person-centered abortion care were developed from literature reviews, expert review, and cognitive interviews, and administered with 371 women who received a safe abortion service from private health clinics in Nairobi, Kenya. Exploratory factor analyses were performed and stratified by surgical abortion procedures and medication abortion. Bivariate linear regressions assessed for criterion validity.

Results: We developed a 24-item unifying scale for person-centered abortion care including two sub-scales. The two sub-scales identified were: 1) Respectful and Supportive Care (14 items for medication abortion, 15 items for surgical abortion); and 2) Communication and Autonomy (9 items for both medication and surgical abortion). The person-centered abortion care scale had high content, construct, criterion validity, and reliability.

Conclusions: This validated scale will facilitate measurement and further research to better understand women's experiences during abortion care and to improve the quality of women's overall reproductive health experiences to improve health outcomes.

Keywords: Abortion, Quality of care, Person-centered care, Kenya, Patient experience, Patient-provider communication

Background

Positive patient experience is a critical component in ensuring quality abortion care for women, linked to health outcomes such as decreases in severe abortion complications and seeking timely care [1]. Globally, there are approximately 56 million abortions yearly [2]. In Kenya, changes to the Constitution were passed in 2010, allowing abortions in instances where the health or the life of

the mother is at risk [3]. However, the Kenya law is still restrictive and requires a qualified health professional to determine if the mother's health or life is jeopardized. These restrictions continue to limit women's access to quality care – resulting in abortion providers working in contexts where women experience high levels of fear and uncertainty, mistreatment and discrimination, and consequently women delaying timely care. Given these circumstances, there is evidence of high maternal mortality as a result of unsafe abortions in Kenya [4]. Safe abortion services can help prevent these avoidable deaths, and such services are available even in places like Kenya where laws are restrictive [5]. Women's

* Correspondence: alandrian2@gmail.com; alandrian@ucla.edu

¹Community Health Sciences, University of California, Los Angeles, Jonathan and Karin Fielding School of Public Health, 650 Charles E Young Dr. S, Los Angeles, CA, USA

Full list of author information is available at the end of the article



© The Author(s). 2020 **Open Access** This article is licensed under a Creative Commons Attribution 4.0 International License, which permits use, sharing, adaptation, distribution and reproduction in any medium or format, as long as you give appropriate credit to the original author(s) and the source, provide a link to the Creative Commons licence, and indicate if changes were made. The images or other third party material in this article are included in the article's Creative Commons licence, unless indicated otherwise in a credit line to the material. If material is not included in the article's Creative Commons licence and your intended use is not permitted by statutory regulation or exceeds the permitted use, you will need to obtain permission directly from the copyright holder. To view a copy of this licence, visit <http://creativecommons.org/licenses/by/4.0/>. The Creative Commons Public Domain Dedication waiver (<http://creativecommons.org/publicdomain/zero/1.0/>) applies to the data made available in this article, unless otherwise stated in a credit line to the data.

experiences during these services could help inform women's decisions to seek safe abortions or delay abortion care, particularly for post-abortion complications [6]. Improving women's experiences within broader efforts to improve quality of abortion services is critical in protecting women's reproductive rights and needs.

One overlooked aspect of quality of abortion care is person-centered care, or care that is respectful of and responsive to women's preferences, needs and values [1]. A recent review of person-centered care for abortion services found many instances where women are treated poorly [7], particularly in contexts with restrictive laws. In Brazil, for example, women report discrimination from providers, threats from health facility staff to involve law enforcement, lack of pain management provision, and long waiting lines at the facility [8]. Treating women with respect and dignity during abortion care is important from a human rights perspective, but also because women's experiences can impact outcomes, adherence to post-abortion guidance, and future health-seeking decisions by the patient and women who hear of her experiences [7, 9].

While there is evidence that women experience poor person-centered abortion care around the globe, a significant limitation in the literature is the lack of a standardized scale to assess person-centered abortion care (PCAC). In Kenya, there are few studies that focus on patient experiences of care, with most studies qualitative in nature [4, 10–12]. Standardized measures are important for quality improvement efforts, designing and evaluating health system and policy improvements, identifying the unmet need of the most vulnerable populations, and for advocacy. A recent systematic review maps out measures of quality of abortion services [13], grouping these under structure, process, output, and outcomes. The broad range of indicators evidenced in this review demonstrates that abortion care quality is multidimensional and emphasizes the need for progress in the field to move beyond a clinical-only perspective [13]. As the authors also highlight, a streamlined list of standard indicators to measure abortion quality must be produced to advance program, advocacy, and policy efforts.

This study aims to develop and validate a PCAC scale to assess women's person-centered experiences during abortion care in Kenya. This is the first study to validate a person-centered abortion care scale. We describe the scale development process and procedures to assess content, construct, and criterion validity. Given the unique dimension of abortion care, we examine person-centered care items separately for medication abortions and surgical abortions, acknowledging the important differences in these methods and women's subsequent abortion experiences.

Methods

Scale development

Study activities included a literature review to identify potential items of the scale, expert reviews and cognitive interviews, survey administration with Kenyan women, and psychometric analyses to assess construct and criterion validity (Fig. 1). This study followed a standard sequential procedure to scale development to explore women's experiences of abortion care [14] including: 1) identifying dimensions of PCAC and item generation; 2) conducting expert reviews and cognitive interviews; 3) survey administration; and 4) psychometric analyses.

This study occurred concurrently with the validation of person-centered maternity care (PCMC) scale [15] and person-centered family planning care (PCFP) scale [16], as part of a study on person-centered reproductive health care. Specifically, steps 1–3 occurred simultaneously, and step one informed a framework for equity in person-centered reproductive health care [9].

Step 1: dimension of abortion person-centered care and item generation

This study first identified dimensions, or domains, of abortion person-centered care focused on quality of care for abortion described in detail in another manuscript [9]. Using this literature review, we identified eight domains discussed in full elsewhere [9] and which are shown in Table 1. Using the domains as a guide and based on existing measures drawn from the literature review, we then created a database of potential person-centered care measures for abortion services. In total, we began with 62 items that spanned all domains identified.

Step 2: expert review and cognitive interviews

We conducted expert reviews, in both the United States and Kenya, with reproductive health experts. The purpose of the expert reviews was to assess content validity – the extent to which measures fully describe and represent a construct (i.e. person-centered care) [14]. Additionally, we sent the surveys to individual experts focused on person-centered care for specific feedback, including differences in experiences of women for medication vs. surgical abortion. Expert reviewers gave feedback on the list of measures, suggested measures to include, and commented on item wording and response options. Expert reviewers also suggested including the same questions for both surgical and medication abortions to be able to compare across the two procedures, and because the preparation for medication abortion clients (i.e. ultrasounds, tests, etc.) would warrant the same type of questions as surgical procedures.

Based on these revisions, we conducted cognitive interviews, using an interview guide provided in Additional file 1, with 15 participants in Nairobi, Kenya




Fig. 1 Flowchart of research methodology for the development and validation of a person-centered abortion care scale

in 2017. We recruited women aged 15–24 in the clinic after they had received any type of safe abortion service from one of four clinics. The purpose of cognitive interviews was to assess how important questions are to women, potential problems with survey questions, whether questions are interpreted as intended, whether the questions are contextually appropriate or needed editing, and whether the survey length was appropriate [14]. Interviews were individually conducted, coded, and analyzed by a team of three researchers (specifically, authors MS, NDS, and PA).

Step 3: survey administration

We piloted the full survey with 30 items among a convenience sample of clients ($n = 31$) in three private clinics in Nairobi (Table 1). Based on pilot results, several survey questions were removed to accommodate time constraints and avoid interviewee fatigue. Four items were removed from the initial set of items (facility/bathrooms are clean, the facility is not crowded, electricity, and water are available in facility). The final set of 26 items was administered in the full survey as provided in Additional file 2.

Inclusion criteria for the survey were women who: 1) received a safe abortion-related service at the clinic the day of recruitment (surgical abortion or medication abortion), 2) were at least 18 years old, 3) spoke English or Swahili, and 4) owned a mobile phone where she felt comfortable being contacted by the study team. The study enumerator obtained written consent prior to any study procedures from interested and eligible women. Consented women were then administered a 40-min baseline questionnaire at the clinic in a private space. In total, 371 women were interviewed at six clinics in Nairobi. Participants received airtime equivalent to

approximately \$1.50 as a token of appreciation for their participation.

Step 4: psychometric analyses

Psychometric analyses were conducted to assess construct and criterion validity and reliability. While expert reviewers believed all items would pertain to both medication abortion and surgical abortion procedures, we hypothesized that the extent to which certain behaviors would pertain more for medication abortion vs. surgical abortion procedures might differ. Therefore, all analyses were stratified by abortion type (medication vs. surgical). The following psychometric analyses were performed for both groups.

First, we examined the distribution of each item for heterogeneity. Negative items were reverse coded so that all items could be scored from worst (score of 0 on the item) to best (score of 3 on the item). Responses of “not applicable” were recoded to the highest response category (corresponding to highest PCAC) to obtain a uniform scale for conducting psychometric analyses. After recoding items, a correlation matrix was constructed and the correlations among the items were examined.

Next, we conducted exploratory factor analysis (EFA) using principal factoring. The degree of common variance among items was assessed with the Kaiser-Meyer-Olkin (KMO) test using a standard cutoff of 0.50 indicating that items are sufficiently correlated to warrant principal components analysis [17]. We then began the initial EFA by examining a scree plot of eigenvalues using all 26 scale items to determine the number of factors to retain. We used both Kaiser’s rule of retaining only factors with eigenvalues greater than one and the shape of the plot [18], along with theoretical considerations, to decide on a final number of factors. Multiple rounds of factor analyses were subsequently performed,

Table 1 Items for person-centered abortion care scale in Kenya

| Original Domain | Question | Referred to in text as: | Comment |
|-----------------------------|--|---------------------------|---|
| Dignity/Respect | How did you feel about the amount of time you waited? | Time to care | Retained for both |
| Dignity/Respect | During your time in this clinic did the doctors, nurses, or other health care providers introduce themselves to you when they first came to see you? | Introduce self | Retained for both |
| Dignity/Respect | Did the doctors, nurses, or other health care providers call you by your name? | Called by name | Retained for both |
| Dignity/Respect | Did the doctors, nurses, or other staff at the facility treat you with respect? | Treated with respect | Retained for both |
| Dignity/Respect | Did the doctors, nurses, and other staff at the facility treat you in a friendly manner? | Friendly | Retained for both |
| Dignity/Respect | During your time in the health facility, would you say you were treated differently because of any personal attribute ... like your age, marital status, number of children, your education, wealth, your connections with the facility, or something like that? | Treated differently | Retained for both |
| Dignity/Respect | Did the doctors, nurses, and other staff at the facility show that they cared about you? | Cared | Retained for both |
| Dignity/Respect | Did you feel the doctors, nurses, or other health providers shouted at you, scolded, insulted, threatened, or talked to you rudely? | Verbal abuse | Retained for both |
| Dignity/Respect | Did you feel like you were treated roughly like pushed, beaten, slapped, pinched, physically restrained, or gagged? | Physical abuse | Retained for both |
| Privacy/ Confidentiality | When you were speaking to the doctors, nurses or other staff at the facility, did you feel other people not involved in your care could hear what you were discussing? | Privacy | Retained for both |
| Privacy/ Confidentiality | Do you feel like your health information was or will be kept confidential at this facility? | Record confidentiality | Retained for both |
| Autonomy | Did you feel like the doctors, nurses or other staff at the facility involved you in decisions about your abortion care? | Involvement in care | Retained for both |
| Autonomy | Did the doctors, nurses or other staff at the facility ask your permission/consent before doing procedures on you? | Consent before procedures | Retained for both |
| Communication | Did the doctors, nurses or other staff at the facility speak to you in a language you could understand? | Language understand | Deleted for both |
| Communication | Did the doctors and nurses explain to you why they were doing examinations or procedures on you? | Explain exams | Retained for both |
| Communication | Did the doctors and nurses explain to you why they were giving you any medicine, including pain medicine or medicine to start an abortion? | Explain medicines | Retained for both |
| Communication | Did you feel you could ask the doctors, nurses or other staff at the facility any questions you had? | Ask questions | Retained for both |
| Supportive Care | Did the doctors and nurses at the facility talk to you about how you were feeling? | Ask about feeling | Retained for both |
| Supportive Care | Did the doctors and nurses ask how much pain you were in? | Ask about pain | Retained for both |
| Supportive Care | Do you feel the doctors or nurses did everything they could to help control your pain? | Pain medication given | Retained among SA sub-group; Deleted among MA sub-group |
| Supportive Care | Did you feel the doctors and nurses paid attention to you during your stay in the facility? | Paid attention | Retained for both |
| Supportive Care | During your time at the facility, did any staff at the facility ask you or your family for <i>kitu kidogo</i> [unofficial cost]? | Bribe | Deleted for both |
| Supportive Care | Do you think there was enough health staff in the facility to care for you? | Enough staff | Retained for both |
| Supportive Care | Did you feel the doctors, nurses or other staff at the facility took the best care of you? | Took best care | Retained for both |
| Supportive Care | Did you feel you could completely trust the doctors, nurses or other staff at the facility with regards to your care? | Trust | Retained for both |
| Supportive Care | In general, did you feel safe in the health facility? | Safe | Retained for both |

Adapted from Afulani et al.,[15]

whereby a factor loading cutoff of 0.10 was used for determining which items to delete or retain. Each factor was assessed individually before evaluating the final scale on a single factor.

We used oblique rotation because it allows for naturally occurring correlation between the rotated factors [19]. Internal consistency of the full scale and each sub-scale was then assessed using Cronbach's alpha. Cronbach's alpha may range from 0 to 1, with a higher value implying greater reliability. An alpha of at least 0.70 is generally considered to suggest acceptable reliability [19].

Finally, criterion validity was assessed by examining whether the PCAC scale was related to other measures of perceived quality of care. This was done by first summing the responses across all items to obtain a total PCAC score, as well as total scores for each sub-scale, where higher scores indicate better PCAC. We then ran bivariate linear regression analyses to assess the association between PCAC score (for the full scale and each sub-scale) and receipt of adequate information regarding the abortion procedure as a measure of quality of care. Women were considered to have received adequate information regarding their abortion procedure if they reported to have been told the following: about the care to be received so they knew what to expect; what side effects to expect; the warning signs to look out for that would warrant returning to the clinic or nearest hospital; and that they would be able to get pregnant again quickly, even before their next menstruation. There was a measure of satisfaction of care; however, there was little heterogeneity in distribution of the responses, with more than 95% of women indicating that they were satisfied with their abortion care. We therefore used another measure of women's quality of care. All analyses were performed using StataSE version 15.

Results

Demographic characteristics

A total of 353 women completed all PCAC scale items; their demographic characteristics are presented in Table 2, stratified by abortion procedure type. Among both surgical abortion and medication abortion participants, most women were aged 20–29 years (63 and 68%, respectively) and not married, partnered, or cohabitating (73 and 81%, respectively). About half of women in both samples had a college or university degree (52% in the surgical abortion sample and 49% in the medication abortion sample). Slightly more women were employed for pay among the surgical abortion sample (62%) compared to the medication abortion sample (55%). A higher proportion of women among the medication abortion sample (54%) reported this to be their first pregnancy than in the surgical abortion sample (40%).

Exploratory factor analysis

Among the medication abortion sample, nearly 50% of women responded “not applicable” to the “pain medication given” item. Further, only one woman responded “yes” to the “bribe” item among the surgical abortion sample and all the women in the medication abortion sample responded “no, never.” As a result, the “bribe” item was removed from both samples and the “pain medication given” item was removed among the medication abortion sample. This left 25 scale items among the surgical abortion sample and 24 items among the medication abortion sample at the start of conducting factor analyses. All scale items among both samples had a KMO measure of sampling adequacy greater than 0.50, with an overall KMO value of 0.74 among the surgical abortion sample and 0.80 among the medication abortion sample, providing evidence that the items were sufficiently correlated for conducting principal components analysis.

Surgical abortion sample

Among the surgical abortion sample, the initial EFA yielded a 3-factor solution with eigenvalues of greater than 1, accounting for about 50% of the variance. All 25 items loaded onto at least one of the three factors at the 0.10 factor loading cutoff, with cross loading on 11 items. We therefore categorized the items that cross loaded to factors based on the factor they loaded higher on and theoretical reasoning. Of the 11 items that loaded positively onto more than one factor, eight were retained in the factor they loaded highest on. Two items, “explain exams” and “ask questions,” which had slightly higher factor loadings on Factor 1 than Factor 2, were categorized into Factor 2 for conceptual reasons. Similarly, the item “pain medication given” had a higher loading on Factor 2 but was ultimately categorized with Factor 1. Also, despite the item “language understand” having a positive loading of greater than 0.10, upon further review it was felt that the wording of the item was too ambiguous, and as a result, the item was removed. Thus, this process resulted in 12 items being categorized to Factor 1, 9 items being categorized to Factor 2, and only three items (“treated differently,” “verbal abuse,” and “physical abuse”) remaining in Factor 3. Upon further discussion, we decided three items were insufficient for a sub-scale, and given that these items were conceptually related to those included in Factor 1, they were categorized with the items in Factor 1.

We then performed EFA with oblique rotation again on the two factors, or sub-scales, separately (Table 3). For the first factor, a standardized alpha of 0.78 was obtained suggesting acceptable reliability. Eleven of the 15 items had factor loadings of at least 0.30 and two items had factor loadings of at least 0.10. The remaining two

Table 2 Demographic characteristics stratified by abortion type

| Characteristic | N (%) | |
|--|--------------------------------|-------------------------------------|
| | Surgical Abortion (N = 157) | Medication Abortion (N = 196) |
| Age, years | | |
| Less than 20 | 6 (3.8) | 25 (13.8) |
| 20–24 | 61 (38.9) | 76 (38.8) |
| 25–29 | 37 (23.6) | 57 (29.1) |
| 30–34 | 25 (15.9) | 25 (12.8) |
| 35 or older | 28 (17.8) | 13 (6.6) |
| Married, partnered, or cohabitating | | |
| Yes | 42 (26.8) | 37 (18.9) |
| No | 114 (72.6) | 158 (80.6) |
| Missing | 1 (0.6) | 1 (0.5) |
| Education | | |
| Primary or less | 20 (12.7) | 14 (7.1) |
| Secondary or vocational | 56 (35.7) | 87 (44.4) |
| College or University | 81 (51.6) | 95 (48.5) |
| Employed for pay | | |
| Yes | 97 (61.8) | 108 (55.1) |
| No | 60 (38.2) | 88 (44.9) |
| Religion | | |
| Catholic, Protestant, or other Christian | 148 (94.3) | 189 (96.4) |
| Muslim | 5 (3.2) | 5 (2.6) |
| None | 3 (1.9) | 2 (1.0) |
| Missing | 1 (0.6) | 0 (0.0) |
| Number of pregnancies | | |
| 1 | 63 (40.1) | 105 (53.6) |
| 2 | 41 (26.1) | 46 (23.5) |
| 3 | 25 (15.9) | 22 (11.1) |
| 4 | 11 (7.0) | 17 (8.7) |
| 5 or more | 17 (10.8) | 6 (3.1) |
| Number of births | | |
| 0 | 76 (48.4) | 120 (61.2) |
| 1 | 37 (23.6) | 40 (20.4) |
| 2 | 23 (14.7) | 25 (12.8) |
| 3 or more | 21 (13.4) | 11 (5.6) |
| Number of children | | |
| 0 | 78 (49.7) | 121 (61.7) |
| 1 | 35 (22.3) | 43 (21.9) |
| 2 | 24 (15.3) | 20 (10.2) |
| 3 or more | 20 (12.7) | 12 (6.1) |
| Received adequate information regarding abortion procedure | | |
| No | 88 (56.1) | 75 (38.3) |
| Yes | 69 (43.9) | 121 (61.7) |

items, “privacy” and “physical abuse,” had factor loadings less than 0.10 but were retained due to their theoretical importance. For the second factor, a standardized alpha of 0.69 was obtained. With the exception of “called by name” and “involvement in care,” the remaining seven items had factor loadings greater than 0.30. The standardized alpha for the nine items was 0.72 suggesting acceptable reliability.

A final factor analysis was conducted on the remaining 24 items restricted to a single factor (Table 4). Twenty-one of the 24 items loaded onto the single dominant factor at the 0.10 cutoff, and in fact, 17 of these had factor loadings greater than 0.30. The item “called by name” had a factor loading less than 0.10; however, it was ultimately retained because of its acceptable factor loading on the sub-scale (Factor 2). “Privacy” and “physical abuse” items also had factor loadings less than 0.10, as in the two-factor solution, but again, were retained due to their theoretical importance – lack of patient privacy and physical abuse are considered central components of poor person-centered care, and thus, would be important to measure. The standardized alpha of the 24-item scale was 0.82 (mean score = 61.84; SD = 7.69; Range = 25–71). A summary of standardized alphas and associated means, standard deviations (SD), and the range of scores for the full scale and each sub-scale are provided in Table 5.

Medication abortion sample

Among the medication abortion sample, the initial EFA using principal factors with oblique rotation also yielded a 3-factor solution with eigenvalues of greater than 1, accounting for about 53% of the variance. All 24 items loaded onto at least one of the three factors at the 0.10 factor loading cutoff, with cross loading on 12 items. We categorized the items that cross loaded to factors based on the factor they loaded higher on and theoretical reasoning. Ten of the 12 items that loaded positively onto more than one factor were categorized according to the higher factor loading. Two items, “friendly” and “privacy,” had similar loadings on two factors but were categorized with Factor 1 because these items were deemed to be more conceptually related to the items in that factor. Despite “explain medicines” loading to Factor 1, it was ultimately recategorized to Factor 3 for conceptual reasons. As was done in the surgical abortion sample, the “language understand” item was removed after it was concluded that the item was too ambiguous. This process resulted in 12 items being categorized to Factor 1, three items (“treated differently,” “verbal abuse,” and “physical abuse”) being categorized to Factor 2, and 9 items being categorized to Factor 3. For consistency, the three items in Factor 2 were ultimately categorized with Factor 1 to yield a final 2-factor solution, like that

Table 3 Rotated factor loadings of sub-scales stratified by abortion type

| Item | Rotated Factor Loadings | | | |
|---------------------------|--|---|--|---|
| | Surgical Abortion (24 items total) | | Medication Abortion (23 items total) | |
| | Respectful and Supportive Care (15 items) | Communication and Autonomy (9 items) | Respectful and Supportive Care (14 items) | Communication and Autonomy (9 items) |
| Time to care | 0.3208 | | 0.2312 | |
| Treated with respect | 0.5565 | | 0.6147 | |
| Friendly | 0.4865 | | 0.6244 | |
| Treated differently | 0.1974 | | 0.4978 | |
| Cared | 0.5379 | | 0.5557 | |
| Privacy | 0.0613 ^a | | 0.2580 | |
| Record confidentiality | 0.1548 | | 0.3769 | |
| Pain medication given | 0.5453 | | NA | |
| Paid attention | 0.8119 | | 0.6811 | |
| Verbal abuse | 0.4189 | | 0.6824 | |
| Physical abuse | 0.0539 ^a | | 0.4982 | |
| Enough staff | 0.6007 | | 0.2242 | |
| Took best care | 0.8304 | | 0.7268 | |
| Trust | 0.4935 | | 0.6571 | |
| Safe | 0.5398 | | 0.4339 | |
| Introduce self | | 0.4468 | | 0.4611 |
| Called by name | | 0.2496 | | 0.2876 |
| Involvement in care | | 0.2069 | | 0.4135 |
| Consent before procedures | | 0.6446 | | 0.3996 |
| Explain exams | | 0.4508 | | 0.3301 |
| Explain medicines | | 0.5950 | | 0.1952 |
| Ask about feeling | | 0.6432 | | 0.7358 |
| Ask questions | | 0.4473 | | 0.2669 |
| Ask about pain | | 0.6197 | | 0.6138 |

Notes: NA Not applicable as this item was deleted among medication abortion patients

^aFactor loading is less than 0.10 but item is being retained due to theoretical importance

proposed among the surgical abortion sample, with 23 items retained at this time.

Performing EFA with oblique rotation on the two factors, or sub-scales, separately, we found that the first factor had a standardized alpha of 0.82 suggesting acceptable reliability. All 14 items had factor loadings greater than 0.10, and 11 of the 14 items had factor loadings greater than 0.30 (Table 3). For the second factor, a standardized alpha of 0.65 was obtained. All items in Factor 2 had factor loadings greater than the 0.10 cutoff; six of the nine items had factor loadings greater than 0.30.

A final factor analysis was conducted on the remaining 23 items restricted to a single factor (Table 4), with all items loading onto the single factor at the 0.10 cutoff. The standardized alpha of the final 23 item scale was 0.82 (mean score = 56.98; SD = 7.44; Range = 13–69; Table 5).

To name the factors, we assessed the specific items and mapped the items out on the original domains that the authors conceptualized for person-centered care. For the first factor, items came from the domains of respectful and supportive care (RSC sub-scale). Items included in the second factor all related to communication and women's ability to be involved in care (CA sub-scale). Therefore, again, guided by the original conceptualized domains of person-centered care, we named the second factor "Communication and Autonomy."

Criterion validity

The results of the bivariate linear regressions assessing the association between the two PCAC sub-scales (RSC sub-scale and CA sub-scale) and the full PCAC scale and the receipt of adequate information regarding the abortion procedure, respectively, are provided in Table 6.

Table 4 Full PCAC scale rotated factor loadings stratified by abortion type

| Item | Rotated Factor Loading | |
|---------------------------|------------------------|---------------------|
| | Surgical Abortion | Medication Abortion |
| Time to care | 0.2977 | 0.2349 |
| Treated with respect | 0.5408 | 0.6111 |
| Friendly | 0.4508 | 0.6248 |
| Treated differently | 0.2419 | 0.4521 |
| Cared | 0.5051 | 0.5736 |
| Privacy | 0.0447 ^a | 0.2038 |
| Record confidentiality | 0.1710 | 0.4074 |
| Pain medication given | 0.6478 | NA |
| Paid attention | 0.7917 | 0.7092 |
| Verbal abuse | 0.3945 | 0.6321 |
| Physical abuse | 0.0035 ^a | 0.4484 |
| Enough staff | 0.5660 | 0.2288 |
| Took best care | 0.7686 | 0.7301 |
| Trust | 0.4592 | 0.6596 |
| Safe | 0.4708 | 0.4115 |
| Introduce self | 0.3128 | 0.2845 |
| Called by name | 0.0299 ^a | 0.2789 |
| Involvement in care | 0.1147 | 0.2810 |
| Consent before procedures | 0.5083 | 0.3673 |
| Explain exams | 0.5021 | 0.1683 |
| Explain medicines | 0.4877 | 0.2647 |
| Ask about feeling | 0.5783 | 0.2860 |
| Ask questions | 0.5160 | 0.4295 |
| Ask about pain | 0.5334 | 0.1539 |

Notes: NA Not applicable as this item was deleted among medication abortion patients

^aFactor loading is less than 0.10 but item is being retained due to theoretical importance

Among both samples, we found that women who reported receiving adequate information regarding their abortion procedure had significantly higher PCAC scores (for each sub-scale, as well as the full scale) than women who did not receive adequate information. These results confirm our hypothesis and suggest that higher PCAC scores are associated with receiving adequate information.

Discussion

This study utilized established scale development procedures to validate a new scale for PCAC. The resulting 24- and 23-item scale for surgical abortion and medication abortions, respectively, includes two sub-scales that measure: 1) Respectful and Supportive Care (i.e. whether women received care in a respectful manner, trusted providers, and felt safe in the facility); and 2) Communication and Autonomy (i.e. whether women were actively involved in making healthcare decisions and providers fully explained procedures, exams, treatment regimens, etc. in a way a woman could understand beforehand). The scale has high content, construct, and criterion validity, as well as strong reliability, based on the expert reviews, cognitive interviews, and psychometric analyses. The few existing related scales are either much more narrowly focused, for example measuring only one component of person-centered care related to stigma [20], or are much broader, including the quality of technical aspects of the abortion itself [13]. Our scale measures a diverse set of domains of person-centered care, while still being focused on the quality of women's experiences themselves.

This study examines the potential for customized scale items for medication abortion versus surgical abortion procedures, and we found significant overlap in items across the two methods. While surgical procedures can be completed in less than 15 min, the preparation and wait times at the health facility and, potentially, at home is typically longer for those receiving a medication abortion. Consequently, the interactions with the provider

Table 5 Standardized alphas and means for the PCAC scale and sub-domains stratified by abortion type

| | Alpha (standardized) | Mean Score | SD | Min | Max |
|---|----------------------|------------|------|-----|-----|
| Surgical Abortion | | | | | |
| Full PCAC Scale (24 items) | 0.82 | 61.84 | 7.69 | 25 | 71 |
| Respectful and Supportive Care (15 items) | 0.78 | 41.13 | 4.17 | 20 | 45 |
| Communication and Autonomy (9 items) | 0.72 | 20.71 | 4.79 | 4 | 27 |
| Medication Abortion | | | | | |
| Full PCAC Scale (23 items) | 0.82 | 56.98 | 7.44 | 13 | 69 |
| Respectful and Supportive Care (14 items) | 0.82 | 38.16 | 4.10 | 6 | 42 |
| Communication and Autonomy (9 items) | 0.65 | 18.82 | 4.98 | 7 | 27 |

Note: SD Standard deviation

are likely to be different. However, the one minor difference included a question regarding pain, which was not included in the person-centered scale for medication abortions. This minor difference suggests that a unified scale can be applicable for both medication abortion and surgical abortion clients, with an additional pain question asked for surgical clients only. Given that this same scale could potentially be used for all abortion clients regardless of abortion type, this scale is more generalizable and user-friendly for clinics and potentially scalable if integrated into other larger, on-going surveys. Additionally, many researchers and others consider the two types of abortion services to be vastly different; these findings suggest a convergence in this field that the same set of factors are equally important to women's person-centered abortion experiences, regardless of method.

The two identified sub-scales (RSC sub-scale and CA sub-scale) are in line with other measures of person-centered care for reproductive health, specifically PCMC [15] and PCFP [16], but also diverged in a couple areas. For example, similar to PCMC, autonomy and communication domains produced one sub-scale for PCAC. On the other hand, for PCFP, autonomy was more closely associated with items for respectful care. Additionally, in the respectful care domain, similar to PCMC, physical and verbal abuse were included in the abortion care scale, but not PCPF scale, although in both cases the item on physical abuse was retained based on theoretical rationale. Potential reasons for these similarities in maternity and abortion scales include that, similar to maternity care, provision of abortion procedures, particularly surgical abortions, requires greater attention and time from providers than providing contraceptive services. Therefore, the longer and more in-depth the

provider-patient interaction is may naturally lead to more opportunities for certain behaviors to occur during care. Nonetheless, the similarity of items across PCMC, PCFP, and PCAC scales suggest that the construction and manifestation of person-centered care may be similar across maternity care, family planning, and abortion care. Future studies may be able to use items found across the three scales to examine whether there are differences in experiences across reproductive health care services.

There were a few limitations to the PCAC scale validation for abortion. One limitation is that we recruited women who received an abortion-related service from a private clinic/hospital. The extent to which these measures hold up to other settings, including public hospitals where women may experience substandard care, is unknown. Recruiting in private clinics may have biased our sample. For example, the prevalence of women with a college or university degree is high and may be due to recruitment in private facilities, where the quality of care may be higher, and thus attract higher socioeconomic status women. Future studies may be conducted among women accessing abortion services from the public sector. Our study also includes women seeking abortions in legal settings and it is difficult to ascertain its applicability in settings where laws are even more restrictive than Kenya. Likely, these women would potentially face higher levels of poor person-centered care and experience additional issues captured by items that were dropped from our validation process, such as bribes. Another limitation is that due to constraints in the duration of the surveys, we were unable to include items related to the health facility environment. We anticipated that because private non-governmental organization clinics

Table 6 Bivariate linear regression of person-centered abortion care sub-scales and full-scale and receipt of adequate abortion procedure information stratified by abortion type

| | RSC Sub-scale | CA Sub-scale | PCAC Full Scale |
|--|-------------------------|-------------------------|-------------------------|
| Surgical Abortion | | | |
| Received adequate information regarding abortion procedure | | | |
| No | Ref | Ref | Ref |
| Yes | 2.01 (0.72, 3.30)** | 4.07 (2.68, 5.45)*** | 6.06 (3.82, 8.33)*** |
| Constant | 40.25 (39.39, 41.11)*** | 18.92 (18.00, 19.84)*** | 59.17 (57.68, 60.66)*** |
| R-squared | 0.0515 | 0.1730 | 0.1494 |
| Medication Abortion | | | |
| Received adequate information regarding abortion procedure | | | |
| No | Ref | Ref | Ref |
| Yes | 1.44 (0.27, 2.62)* | 1.68 (0.25, 3.10)* | 3.12 (1.00, 5.23)** |
| Constant | 37.27 (36.35, 38.19)*** | 17.79 (16.66, 18.91)*** | 55.05 (53.39, 56.72)*** |
| R-squared | 0.0245 | 0.0219 | 0.0368 |

Notes: Coefficient (95% confidence interval) provided. RSC Respectful and Supportive Care. CA Communication and Autonomy. PCAC=Person Centered Abortion Care

* $p < 0.05$; ** $p < 0.01$; *** $p < 0.001$

tend to have better health facility environments than public facilities, that this might play less of a role in our study sample, but is likely to be more important in public facilities. In addition, given that the surveys were conducted within the health facility, social desirability bias in responses is a limitation despite emphasis on confidentiality and privacy during the consent process.

Conclusion

We developed and validated a new scale for PCAC. This scale makes an important contribution to the literature given growing evidence of the importance of prioritizing person-centered care across reproductive health services and the lack of validated measurement tools. There are a number of potential uses of this scale. First, a validated scale for PCAC will facilitate measurement and research to better understand women's experiences when seeking care for abortion. Second, it will facilitate future research to identify the gaps in PCAC during women's abortion care experiences. This scale will also aid research into sources of disparities in women's abortion care experiences and help refine interventions aiming to improve abortion care, particularly for vulnerable groups. In addition, it provides a valuable tool for evaluating these types of interventions in other contexts. The scale will also facilitate standardized measurement across key places where person-centered care is most lacking during abortion care and simultaneously assess trends in the provision of abortion care. Future studies may assess how person-centered abortion care may influence or correlate with different outcomes, including abortion complications or post-abortion family planning counseling. Finally, given the similarities in items across the PCMC, PCFP, and PCAC scales, together these reproductive health scales could facilitate future research on person-centered care across the reproductive health continuum and improve women's reproductive health experiences and outcomes worldwide. This scale is potentially applicable in other settings beyond Kenya. Validation in other settings is however needed to assess its validity across settings.

Supplementary information

Supplementary information accompanies this paper at <https://doi.org/10.1186/s12905-020-01071-w>.

Additional file 1.

Additional file 2.

Abbreviations

CA: Communication and Autonomy; EFA: Exploratory factor analysis; KMO: Kaiser-Meyer-Olkin; PCAC: Person-centered abortion care; PCFP: Person-centered family planning care; PCMC: Person-centered maternity care; RSC: Respectful and Supportive Care

Acknowledgements

The authors would like to thank the research team including Ziporah Mugwanga, Edward Ikiugu, Ginger Golub, James Oput, Nancy Onditi, and Morine Odongo. We also thank Avery Seefeld for overseeing data collection activities.

Authors' contributions

MS conceptualized, oversaw data collection and analyses, and drafted the manuscript. AL led the analysis of the manuscript and drafted parts of the manuscript. PA and NDS contributed to data interpretation and reviewed and edited the manuscript. BP and SC contributed to data collection and reviewed and edited the manuscript. All authors read and approved the manuscript for publication.

Funding

The David and Lucile Packard Foundation funded this work (Grant Number A126422). They had no role in the research study design or research activities.

Availability of data and materials

The datasets analyzed during the current study are available from the corresponding author upon reasonable request.

Ethics approval and consent to participate

The full study protocol and all study materials were reviewed and approved by the Ethical Review Board at the University of California, San Francisco and the Kenya Medical Research Institute. All participants provided written informed consent prior to any study procedures as discussed in the methods section. Informed consent was not obtained from a parent or legal guardian of minors (under 16 years of age) who were eligible and interested to participate. This was done to maintain privacy and confidentiality given the sensitive nature of abortion care and was approved by the aforementioned Institutional Review Boards as described in our study protocol.

Consent for publication

Not applicable.

Competing interests

The authors declare that they have no competing interests.

Author details

¹Community Health Sciences, University of California, Los Angeles, Jonathan and Karin Fielding School of Public Health, 650 Charles E Young Dr. S, Los Angeles, CA, USA. ²Institute for Global Health Sciences, University of California, San Francisco, School of Medicine, 550 16th Street, San Francisco, CA, USA.

Received: 13 February 2020 Accepted: 8 September 2020

Published online: 19 September 2020

References

- Mutua MM, Manderson L, Musenge E, TNO A. Policy, law and post-abortion care services in Kenya. *PLoS ONE*. 2018;13 cited 2019 Feb 27]. Available from: <https://www.ncbi.nlm.nih.gov/pmc/articles/PMC6150499/>.
- Guttmacher Institute. Induced Abortion Worldwide. Guttmacher Inst. 2016; [cited 2019 Apr 5]. Available from: <https://www.guttmacher.org/fact-sheet/induced-abortion-worldwide>.
- The Government of Kenya (GOK). The Constitution of Kenya, 210 revised edition. Nairobi: National Council for Law Reporting (NCLR); 2010.
- Jayaweera RT, Nguu FM, Hall KS, Gerds C. Women's experiences with unplanned pregnancy and abortion in Kenya: a qualitative study. *PLoS One*. 2018;13:e0191412.
- Ganatra B, Gerds C, Rossier C, Johnson BR, Tunçalp Ö, Assifi A, et al. Global, regional, and subregional classification of abortions by safety, 2010–14: estimates from a Bayesian hierarchical model. *Lancet*. 2017;390:2372–81.
- Jewkes RK, Gumedde T, Westaway MS, Dickson K, Brown H, Rees H. Why are women still aborting outside designated facilities in metropolitan South Africa? *BJOG Int J Obstet Gynaecol*. 2005;112:1236–42.
- Altshuler AL, Whaley NS. The patient perspective: perceptions of the quality of the abortion experience. *Curr Opin Obstet Gynecol*. 2018;30:407–13.

8. Madeiro AP, Rufino AC. [Maltreatment and discrimination in induced abortion care: perception of women in Teresina, state of Piauí, Brazil]. *Cienc. Saude Coletiva*. 2017;22:2771–80.
9. Sudhinaraset M, Afulani P, Diamond-Smith N, Bhattacharyya S, Donnay F, Montagu D. Advancing a conceptual model to improve maternal health quality: the person-centered care framework for reproductive health equity. *Gates Open Res*. 2017;1:1.
10. Izugbara CO, Egesa C, Okelo R. 'High profile health facilities can add to your trouble': women, stigma and un/safe abortion in Kenya. *Soc Sci Med*. 2015; 141:9–18.
11. Izugbara CO, Egesa CP, Kabiru CW, Sidze EM. Providers, unmarried young women, and post-abortion Care in Kenya. *Stud Fam Plan*. 2017;48:343–58.
12. Izugbara CO, Otsola KJ, Ezeh AC. Men, women, and abortion in Central Kenya: a study of lay narratives. *Med Anthropol*. 2009;28:397–425.
13. Dennis A, Blanchard K, Bessenaar T. Identifying indicators for quality abortion care: a systematic literature review. *J Fam Plann Reprod Health Care*. 2017;43:7–15.
14. DeVellis RF. *Scale development: theory and applications*. 3rd ed. Thousand Oaks: SAGE Publications, Inc; 2011.
15. Afulani PA, Diamond-Smith N, Golub G, Sudhinaraset M. Development of a tool to measure person-centered maternity care in developing settings: validation in a rural and urban Kenyan population. *Reprod Health*. 2017;14:118.
16. Sudhinaraset M, Afulani PA, Diamond-Smith N, Golub G, Srivastava A. Development of a Person-Centered Family Planning Scale in India and Kenya. *Stud Fam Plann*. 2018;49 [cited 2018 Sep 13]. Available from: <https://onlinelibrary.wiley.com/doi/abs/10.1111/sifp.12069>.
17. Principal Component Analysis - Econometrics Academy [Internet]. [cited 2019 Apr 5]. Available from: <https://sites.google.com/site/econometricsacademy/econometrics-models/principal-component-analysis>.
18. Kaiser HF. The application of electronic computers to factor analysis. *Educ Psychol Meas*. 1960;20:141–51.
19. Hinkin T, Tracey J, Enz C. Scale Construction: Developing Reliable and Valid Measurement Instruments. *Artic Chapters*. 1997; Available from: <https://scholarship.sha.cornell.edu/articles/613>.
20. Cockrill K, Upadhyay UD, Turan J, Greene FD. The stigma of having an abortion: development of a scale and characteristics of women experiencing abortion stigma. *Perspect Sex Reprod Health*. 2013;45:79–88.

Publisher's Note

Springer Nature remains neutral with regard to jurisdictional claims in published maps and institutional affiliations.

Ready to submit your research? Choose BMC and benefit from:

- fast, convenient online submission
- thorough peer review by experienced researchers in your field
- rapid publication on acceptance
- support for research data, including large and complex data types
- gold Open Access which fosters wider collaboration and increased citations
- maximum visibility for your research: over 100M website views per year

At BMC, research is always in progress.

Learn more biomedcentral.com/submissions

