

UC Riverside

UC Riverside Previously Published Works

Title

AMPK mediates inhibition of electrolyte transport and NKCC1 activity by reactive oxygen species

Permalink

<https://escholarship.org/uc/item/8mm317rc>

Journal

AJP Gastrointestinal and Liver Physiology, 317(2)

ISSN

0193-1857

Authors

King, Stephanie J
Bunz, Michael
Chappell, Alfred
[et al.](#)

Publication Date

2019-08-01

DOI

10.1152/ajpgi.00317.2018

Peer reviewed

RESEARCH ARTICLE | *Epithelial Biology and Secretion*

AMPK mediates inhibition of electrolyte transport and NKCC1 activity by reactive oxygen species

Stephanie J. King,^{1*} Michael Bunz,^{2,3*} Alfred Chappell,² Michael Scharl,^{2,4} Michael Docherty,² Barbara Jung,² Christian Lytle,¹ and Declan F. McCole¹

¹Division of Biomedical Sciences, University of California, Riverside, Riverside, California; ²Division of Gastroenterology, School of Medicine, University of California, San Diego, La Jolla, California; ³Kreisklinik Woert an der Donau, Department of Anaesthesiology and Intensive Care Medicine, Woerth an der Donau, Germany; and ⁴Division of Gastroenterology and Hepatology, Zurich Center for Integrative Human Physiology, University Hospital and University of Zurich, Zurich, Switzerland

Submitted 20 September 2018; accepted in final form 3 May 2019

King SJ, Bunz M, Chappell A, Scharl M, Docherty M, Jung B, Lytle C, McCole DF. AMPK mediates inhibition of electrolyte transport and NKCC1 activity by reactive oxygen species. *Am J Physiol Gastrointest Liver Physiol* 317: G171–G181, 2019. First published May 9, 2019; doi:10.1152/ajpgi.00317.2018.—Reactive oxygen species such as H₂O₂ are believed to play a prominent role in the injury and loss of transport function that affect the intestinal epithelium in inflammatory conditions such as inflammatory bowel diseases. Defects in intestinal epithelial ion transport regulation contribute to dysbiosis and inflammatory phenotypes. We previously showed that H₂O₂ inhibits Ca²⁺-dependent Cl[−] secretion across intestinal epithelial cells (IECs) via a phosphatidylinositol 3-kinase (PI3K)- and extracellular signal-regulated kinase (ERK)-dependent mechanism that occurs, at least in part, through inhibition of the basolateral Na⁺-K⁺-2Cl[−] cotransporter NKCC1. NKCC1 governs Cl[−] entry into crypt IECs and thus plays a critical role in maintaining the driving force for Cl[−] secretion. Electrolyte transport consumes large amounts of cellular energy, and direct pharmacological activation of the cellular energy sensor AMP-activated protein kinase (AMPK) has been shown to inhibit a number of ion transport proteins. Here, we show that H₂O₂ activates AMPK in human IEC lines and ex vivo human colon. Moreover, we demonstrate that the inhibitory effect of H₂O₂ on Ca²⁺-dependent Cl[−] secretion and NKCC1 activity is AMPK-dependent. This inhibitory effect is associated with a physical interaction between AMPK and NKCC1, as well as increased phosphorylation (Thr^{212,217}) of NKCC1, without causing NKCC1 internalization. These data identify a key role for AMPK-NKCC1 interaction as a point of convergence for suppression of colonic epithelial ion transport by inflammatory reactive oxygen species.

NEW & NOTEWORTHY H₂O₂ inhibition of intestinal epithelial Ca²⁺-dependent Cl[−] secretion involves recruitment of AMP-activated protein kinase (AMPK) downstream of ERK and phosphatidylinositol 3-kinase signaling pathways, physical interaction of AMPK with the Na⁺-K⁺-2Cl[−] cotransporter NKCC1, and AMPK-dependent suppression of NKCC1-mediated electrolyte influx without causing NKCC1 internalization. It is intriguing that, in human intestinal epithelial cell lines and human colon, H₂O₂ activation of AMPK increased phosphorylation of NKCC1 residues required for promoting, not inhibiting, NKCC1 activity. These data identify an elevated

complexity of AMPK regulation of NKCC1 in the setting of an inflammatory stimulus.

carbachol; chloride; epithelial; HT-29; hydrogen peroxide; T84

INTRODUCTION

The overproduction of reactive oxygen species (ROS) during active episodes of Crohn's disease and ulcerative colitis is thought to play an important role in the pathophysiology of inflammatory bowel disease (IBD) (6, 51). A number of studies have shown that chronically inflamed colonic tissue is subject to significant oxidative stress due, in part, to decreased expression of endogenous antioxidants (12, 27, 33, 41). Therapeutic approaches using broad-spectrum antioxidants, such as vitamin E, or immunomodulatory agents that show antioxidant activity, such as 5-aminosalicylic acid, have proven effective to differing degrees in treating IBD and reducing markers of oxidative stress (52). In addition, antioxidants have proven effective in delaying the onset of, and enhancing recovery from, colitis including a protective effect against the onset of diarrhea in mouse models (11, 37, 50).

H₂O₂ is among the most important ROS believed to have a role in IBD, as it is a relatively stable and cell-permeable precursor of numerous more-toxic oxygen radicals such as the hydroxyl radical (38, 51, 69). Furthermore, the large influx of activated phagocytes in ulcerative colitis may secrete millimolar concentrations of H₂O₂ in close proximity to colonocytes (64). Although pathophysiological concentrations of H₂O₂ can cause damage, both direct and indirect, to cultured epithelial cells and cells isolated from IBD patients via generation of more ROS, H₂O₂ is also capable of more subtle influences on cell function. H₂O₂ has emerged as an important cell-signaling molecule involved in mediating responses to epidermal growth factor and other growth factors (51), as well as participating in MAPK and NF-κB signaling networks (20, 70). H₂O₂ has also been shown to exert discreet, and often opposing, effects on ion transport functions in intestinal tissue and colonic epithelial cell lines. High concentrations (i.e., 5.5 mmol/l) of H₂O₂ have been shown to induce a small increase in short-circuit current (*I*_{sc}), accounted for by Cl[−] secretion, across T84 colonic epithelial cells (48). Other studies have shown that lower

* S. J. King and M. Bunz contributed equally to this work.

Address for reprint requests and other correspondence: D. F. McCole, Div. of Biomedical Sciences, University of California, Riverside, 900 University Ave., 307 SOM Research Bldg., Riverside, CA 92521 (e-mail: declan.mccole@ucr.edu).

concentrations (500 μ M) of H₂O₂ inhibit cAMP-dependent Cl⁻ secretion through specific effects on transport proteins involved in Cl⁻ transport, without evidence of toxic or injurious effects on cell monolayers (20, 67). The overall effect of H₂O₂ on Cl⁻ secretion appears to be one of suppression based on studies of colonic epithelial cell lines and isolated mouse colonic mucosa, consistent with the reduced ion-transporting capability of intestinal tissues in the setting of inflammation (9, 35, 58). Indeed, we previously demonstrated that in vivo administration of the H₂O₂-degrading antioxidant catalase to mice alleviated ion transport dysfunction in dextran sulfate sodium-induced colitis and that this effect was associated with reduced H₂O₂ levels in the colonic mucosa (3).

In the intestinal tract, Ca²⁺-dependent Cl⁻ secretion is triggered by neuronal reflexes responding to changes in luminal composition, contact between luminal contents and the epithelium, and physical distension (13). Cl⁻ secretion serves useful "housekeeping" functions, such as hydration of the mucosal surface, dispersal of mucus and antimicrobial peptides from the crypt, and flushing of noxious substances and pathogenic bacteria. Thus, crypt fluid secretion is an important component of intestinal homeostasis and protection of the host epithelial barrier (47). Indeed, compromised secretory function likely contributes to a decrease in the overall effectiveness of the barrier, facilitating bacterial colonization of the intestinal crypt and subsequent excessive inflammatory responses that are a feature of diseases such as IBD. This hypothesis was supported by studies demonstrating that CFTR-knockout mice exhibit inflammation similar to that observed in IBD patients, as well as pronounced changes in richness and diversity of the intestinal microbiome (43, 49).

We previously demonstrated that H₂O₂ pretreatment is also capable of inhibiting Ca²⁺-dependent Cl⁻ secretion across colonic epithelial cells via Ca²⁺-dependent activation of MAPKs, phosphatidylinositol 3-kinase (PI3K), and functional inhibition of the Na⁺-K⁺-2Cl⁻ cotransporter NKCC1 (9). NKCC1 is located on the basolateral membrane of colonic epithelial cells at the crypt base and is the principal basolateral uptake mechanism for secreted Cl⁻ in crypt epithelium (2). Therefore, inhibition of NKCC1 compromises the normal entry of electrolytes into crypt epithelial cells and restricts secretory capability (21). Decreased electrogenic secretory capacity is associated with IBD and may be particularly relevant to loss of crypt flushing and dispersal of antimicrobial peptides associated with inflammatory conditions (1, 4, 26, 34, 43, 65).

In our earlier work we established that the inhibitory effect of H₂O₂ on Ca²⁺-dependent Cl⁻ secretion across colonic epithelial cells is mediated through activation of a number of signaling kinases previously shown to negatively regulate intestinal Cl⁻ secretion (9, 36) and subsequent inhibition of NKCC1 activity. However, the mechanism by which H₂O₂-activated kinases inhibit NKCC1 and the signaling effector(s) mediating this inhibition has not been identified. Therefore, in the present study we set out to identify the mechanism by which H₂O₂ inhibits NKCC1 by focusing on the cellular energy sensor AMP-activated protein kinase (AMPK). AMPK responds to metabolic stress, manifesting as depletion of ATP levels and elevation of the AMP-to-ATP ratio (10, 31, 32) by stimulating energy-producing pathways, such as glucose uptake and glycolysis, while inhibiting energy-consuming processes, such as protein and fatty acid synthesis (25, 28, 32). In

the intestinal epithelium, energy-consuming processes also include electrolyte transport and maintenance of the epithelial barrier. Indeed, previous studies have demonstrated that activation of AMPK suppresses intestinal ion transport and synthesis of various proteins (28, 53). In the Cl⁻-secreting epithelial cells within the colonic crypt, AMPK has been shown to physically interact with and inhibit the CFTR Cl⁻ channel (67). In absorptive intestinal cells, AMPK can promote removal of the epithelial Na⁺ channel from the apical membrane via increased activity of the ubiquitin ligase enzyme Nedd4-2 (5, 29, 30, 68). These effects would reduce the overall ion-transporting capability of the intestinal epithelium, as demonstrated in colonic tissues from the IL-10^{-/-} mouse model of colitis (67) and intestinal tissues from IBD patients (26, 54, 57).

Therefore, the goal of this study was to determine whether AMPK mediates H₂O₂-induced alterations in intestinal epithelial Ca²⁺-dependent Cl⁻ secretion and NKCC1 function. We found that AMPK is activated by H₂O₂ in human colon and intestinal epithelial cell (IEC) lines. H₂O₂ activation of AMPK leads to physical interaction of AMPK with NKCC1 and suppression of NKCC1 activity, and inhibition of AMPK partially protects NKCC1 from inhibition by H₂O₂. Our data indicate a novel role for the cellular energy sensor AMPK in the regulation of a critical epithelial transport protein in a cell model of inflammation. These findings may have implications for suppression of ion transport function in the setting of inflammatory conditions such as IBD.

MATERIALS AND METHODS

Materials. Carbachol (CCh), compound C (CC), and metformin hydrochloride were obtained from Sigma (St. Louis, MO); wortmannin, LY294002, PD98059, 5-aminoimidazole-4-carboxamide-1- β -D-ribofuranoside (AICAR), and H₂O₂ from Calbiochem (San Diego, CA); mouse anti-phosphotyrosine antibodies from Upstate Biotechnology (Lake Placid, NY); anti-phosphorylated ERK antibodies from New England Biolabs (Ipswich, MA); goat polyclonal anti-total NKCC1 (N-16) and mouse monoclonal anti-total human NKCC1 (T4) antibodies from Dr. C. Lytle (46); rabbit anti-lamin A/C antibody from Santa Cruz Biotechnology (Santa Cruz, CA); rabbit anti-phosphorylated (Thr¹⁷²) AMPK α antibody, rabbit anti-AMPK α antibody, rabbit anti-phosphorylated (Ser⁴⁷³) Akt antibody, and rabbit anti-Akt antibody from Cell Signaling Technology (Danvers, MA); and Tris-glycine electrophoresis gels from Bio-Rad. Rabbit polyclonal anti-human ERK1 (K-23; Santa Cruz Biotechnology) was used to measure total ERK. The R5 anti-phosphorylated (Thr^{212,217}) NKCC1 antibody was a kind gift from Dr. Biff Forbush (22). All other reagents were of analytical grade and obtained commercially.

Cell culture. Methods for maintenance of T84 cells (obtained from primary stocks of Dr. Kim Barrett, University of California, San Diego) in culture were described previously (9). Briefly, T84 cells were grown in DMEM/F-12 medium (Mediatech, Herndon, VA) supplemented with 5% newborn calf serum. For Ussing chamber/voltage-clamp experiments, 5×10^5 cells were seeded onto 12-mm Millicell polycarbonate filters. For Western blot experiments, 10^6 cells were seeded onto 30-mm filters. Cells were cultured for 10–15 days before use. When grown on polycarbonate filters, T84 cells are known to acquire the polarized phenotype of native colonic epithelia. In accordance with muscarinic M₃ receptor distribution on intestinal epithelia, CCh was added basolaterally in all experiments. HT-29.c119A cells (from Dr. Kim Barrett) were grown to confluence (~5 days) in DMEM supplemented with 10% FBS, 1% L-glutamine, and streptomycin (100 μ g/ml) in 75-cm² flasks. Thereafter, the cells were

seeded onto 12-mm Millicell polycarbonate filters for Ussing chamber experiments or onto 30-mm filters for Western blotting, as described above for T84 cells.

Physiological solutions. Composition of the Ringer solution used for Ussing chamber, Western blot, and ⁸⁶Rb⁺ uptake studies was as follows (in mM): 140 Na⁺, 120 Cl⁻, 5.2 K⁺, 25 HCO₃⁻, 0.4 H₂PO₄⁻, 2.4 HPO₄²⁻, 1.2 Ca²⁺, 1.2 Mg²⁺, and 10 D-glucose.

Electrophysiological studies. T84 cells grown to confluence on permeable 12-mm filters were mounted in Ussing chambers (0.6 cm² window area) and bathed in oxygenated (95% O₂-5% CO₂) Ringer solution at 37°C. Monolayers were voltage-clamped to zero potential difference by application of I_{sc}. Under these conditions, changes in I_{sc} in response to agonists are wholly reflective of electrogenic Cl⁻ secretion (8). Transepithelial resistance (TER) was measured using a "chopstick" voltohmmeter (World Precision Instruments, Sarasota, FL) (55, 56). The MEK inhibitor PD98059 was solubilized in DMSO. In control studies, the final concentration of DMSO (0.1%) had no effect on I_{sc} responses to CCh in the presence or absence of H₂O₂ (data not shown). This finding is consistent with our previous unpublished findings and with studies by other groups on MAPK phosphorylation using IEC lines (39).

Human subjects. Control patients [*n* = 3 (2 female and 1 male), 61 ± 5 (range 52–71) yr old] presenting for colon cancer screening were asymptomatic and showed no clinical or macroscopic signs of acute or chronic inflammation. Individuals with a history of IBD or intestinal inflammation at the time of the study were excluded. After informed consent was obtained from the patients, a reusable round-cup jumbo biopsy forceps (model no. FB-25 K-1, Olympus) was used to obtain mucosal punch biopsies (~2 mm² surface area) from macroscopically normal-appearing regions of the sigmoid colon. Tissue biopsies were placed in cold, oxygenated Ringer solution for transport from the endoscopy suite to the laboratory. Tissues were allowed to equilibrate at 37°C in Ringer solution for 20 min before incubation in test solutions. Biopsies were then washed three times in ice-cold Ringer solution before tissue homogenization in lysis buffer for Western blot analysis (see below). The protocol was evaluated and approved by the University of California, San Diego, Institutional Review Board.

Immunoprecipitation and Western blotting. T84 cell monolayers and tissue biopsies were washed three times with Ringer solution, allowed to equilibrate for 30 min at 37°C, and then stimulated with agonists (in the presence or absence of antagonists) as appropriate.




Fig. 1. H₂O₂ increases AMP-activated protein kinase (AMPK) phosphorylation in intestinal epithelial cells. T84 and HT-29.c119a cells grown as monolayers on semipermeable supports were treated bilaterally with H₂O₂ (500 μM) for 2–30 min or the AMPK activator 5-aminoimidazole-4-carboxamide-1-β-D-ribofuranoside (AICAR, 2.5 mM) for 30 min. **A:** Western blot analysis of phosphorylation of the activating Thr¹⁷² residue on the catalytic α-subunit of AMPK. Molecular mass (kDa) of each protein is shown at right. **B:** densitometric quantitation of Western blot results in **A** and normalization to total AMPKα levels. **C:** CCh-induced change in ion transport reported as change in short-circuit current (ΔI_{sc}). T84 monolayers mounted in Ussing chambers were not treated (control) or preincubated with compound C (1–100 μM, bilaterally) for 30 min before administration of H₂O₂ (500 μM) for 30 min followed by stimulation of electrogenic Cl⁻ secretion with carbachol (CCh, 100 μM, basolaterally, *n* = 4). **D:** change in ion transport reported as change in short-circuit current (ΔI_{sc}). HT-29.c119a monolayers mounted in Ussing chambers were untreated (control) or pretreated with compound C (50 μM, bilaterally) for 30 min before administration of H₂O₂ (500 μM, bilaterally) for 30 min and subsequent CCh stimulation of ion transport (*n* = 3). Values are means ± SE. **P* < 0.05, ***P* < 0.01 vs. control (**B**) and vs. H₂O₂ + CCh (**C** and **D**); #*P* < 0.05 vs. compound C (1 μM).




Fig. 2. H₂O₂ activation of AMP-activated protein kinase (AMPK) is downstream of ERK and phosphatidylinositol 3-kinase (PI3K) pathways. **A**: Western blot analysis of AMPK, ERK, and Akt phosphorylation. T84 monolayers were preincubated with the MEK inhibitor PD98059 (50 μ M) or the PI3K inhibitor wortmannin (50 nM) for 30 min before treatment with H₂O₂ (500 μ M) for 5 min. Molecular mass (kDa) of each protein is shown at right. **B**: densitometric quantitation of Western blot results in **A** and normalization to total AMPK α levels ($n = 3$). Values (arbitrary units) are means \pm SE. * $P < 0.05$ vs. control/untreated; # $P < 0.05$ vs. H₂O₂ alone.

The reaction was stopped by washing in ice-cold PBS, and the cells were lysed in ice-cold lysis buffer (1% Triton X-100, 1 μ g/ml leupeptin, 1 μ g/ml pepstatin, 1 μ g/ml antipain, 100 μ g/ml phenylmethylsulfonyl fluoride, 1 mM sodium vanadate, 1 mM sodium fluoride, and 1 mM EDTA in PBS) for 45 min. Cell lysate supernatants were assayed for protein content (Bio-Rad protein assay kit) and adjusted so that each sample contained an equal amount of protein. For immunoprecipitation studies, lysates were incubated with immunoprecipitating antibody according to the manufacturer's instructions for 1 h at 4°C and then with protein A-Sepharose for 1 h at 4°C. For NKCC1 immunoprecipitation, the goat polyclonal anti-total NKCC1 antibody (N-16) was used. Lysates were centrifuged for 3 min at 21,000 g , and the supernatant was discarded. The pellets were washed three times in ice-cold PBS and resuspended in 2 \times gel loading buffer [50 mM Tris (pH 6.8), 2% SDS, 200 mM dithiothreitol, 40% glycerol, and 0.2% bromophenol blue] and boiled before separation by SDS-polyacrylamide gel electrophoresis. Biopsy lysates were passed through a needle, and insoluble material was removed by centrifugation at 350 g for 5 min at 4°C. The supernatant was centrifuged at 43,000 g for 30 min at 4°C, and pellets were resuspended in lysate buffer. Lysate supernatants were assayed for protein content (Bio-Rad protein assay kit) and adjusted so that each sample contained an equal amount of protein. Aliquots of 20 μ g of protein were separated by polyacrylamide gel electrophoresis. Resolved proteins were transferred onto polyvinylidene difluoride membranes (NEN Life Science Products, Boston, MA). Thereafter, the membrane was preblocked with a 1% solution of blocking buffer (Upstate Biotechnology) for 30

min and then incubated for 1 h with the appropriate concentration of primary antibody in 1% blocking buffer. After five 10-min washes in Tris-buffered saline with 1% Tween, membranes were incubated for 30 min with a horseradish peroxidase-conjugated secondary antibody (anti-mouse or anti-rabbit IgG; BD Biosciences, San Jose, CA) in 1% blocking buffer. After five 10-min washes in Tris-buffered saline with 1% Tween, immunoreactive proteins were detected using an enhanced chemiluminescence detection kit (Roche Molecular Biochemicals, Indianapolis, IN) on film. National Institutes of Health Image software was used for densitometric analysis of bands.

Basolateral ⁸⁶Rb⁺ uptake studies. Basolateral K⁺ uptake was measured using ⁸⁶Rb⁺ as a tracer and an adaptation of a previously described method (9). Confluent monolayers of T84 cells grown on 12-mm Transwell inserts were rinsed three times with warm Ringer solution and incubated for 1 h at 37°C. After 1 h of preincubation, cells were exposed bilaterally to H₂O₂ (500 μ M) or vehicle control for 30 min on a warming plate at 37°C. Inserts were transferred to wells containing Ringer solution with or without 100 μ M CCh for 1 min and then to wells containing Ringer solution with 1 μ Ci/ml ⁸⁶Rb⁺ and CCh and maintained at 37°C for 3 min. Inserts were immersed several times in ice-cold 100 mM MgCl₂ containing 10 mM Tris-HCl (pH 7.5) to terminate ⁸⁶Rb⁺ uptake. Filters were immediately excised from the inserts and placed directly into 5 ml of scintillation fluid, and ⁸⁶Rb⁺ content was measured by standard scintillation methods. K⁺ influx (μ mol K⁺·g protein⁻¹·min⁻¹) was calculated as (cpm·g protein⁻¹·min⁻¹)/SA, where SA is the specific activity of the uptake buffer (cpm/ μ mol K⁺).

siRNA transfection. T84 cells (2 \times 10⁶) were seeded 3 days before transfection and grown to 50–70% confluence in T75 flasks. Three different annealed Silencer predesigned siRNA oligonucleotides targeting AMPK α 1 were obtained from Applied Biosystems (Foster City, CA). For transfection reactions, 100 pmol of each of the three gene-specific siRNA oligonucleotides were transfected into T84 cells using the Nucleofector system (Amaxa, Gaithersburg, MD) according to the manufacturer's instructions. After transfection, T84 cells were cultured on filter membranes for 48 h before further treatment. The




Fig. 3. H₂O₂ inhibits basolateral K⁺ influx and activity of the Na⁺-K⁺-2Cl⁻ cotransporter NKCC1. T84 monolayers were pretreated with the AMP-activated protein kinase (AMPK) inhibitor compound C (50 μ M) or Ringer solution (control) and then incubated with vehicle control or H₂O₂ (500 μ M) for 30 min. Cells were subsequently transferred to Ringer solution or Ringer solution + carbachol (CCh, 100 μ M, basolaterally) for 1 min and then to a basolateral ⁸⁶Rb⁺-containing buffer for 3 min ($n = 4$). Values are means \pm SE. ** $P < 0.01$ vs. control/untreated cells; # $P < 0.05$ vs. CCh alone.

nonspecific control siRNA SMARTpool (100 pmol; Upstate Biotechnology/Dharmacon, Chicago, IL) was used as a negative control.

NKCC1 biotinylation assay. NKCC1 surface expression was determined as previously described (7). Briefly, T84 monolayers were treated with 500 μ M H₂O₂ with or without 30 min of pretreatment with compound C (50 μ M), metformin (5 mM) alone, or capsaicin (20 μ M) alone. After treatment, monolayers were washed twice with ice-cold 1 \times PBS (pH 8.0). EZ-Link Sulfo-NHS-SS-biotin (10 μ M; Thermo Fisher Scientific) was added basolaterally, and monolayers were incubated for 30 min at 4°C. Fresh biotin was added, and monolayers were incubated for an additional 30 min. Monolayers were washed twice with ice-cold 1 \times PBS (pH 8.0), incubated in quenching buffer [100 mM glycine in 1 \times PBS (pH 8.0)] for 15 min, and then aspirated. Radioimmunoprecipitation lysis buffer (500 μ l) was added, and monolayers were incubated for 30 min at 4°C with gentle agitation. Cells were scraped and collected into 1.7-ml tubes and then subjected to sonication (QSonica) for 30 s (10 s on/off) at 30% amplitude. Lysates were centrifuged at 4°C for 10 min at 16,200 g, and the supernatant was aspirated into new tubes. Protein was quantified using a bicinchoninic acid assay (Pierce). Pierce streptavidin-agarose resin (50 μ l; Thermo Fisher Scientific) was placed in 1.7-ml tubes and washed according to the manufacturer's instructions. Equal concentrations of protein (~1 mg/ml) were added to respective streptavidin-bead aliquots, which were then incubated overnight at 4°C with rotation. On the following day, samples were centrifuged according to instructions of the manufacturer of the streptavidin-agarose beads. Supernatants were aspirated into new tubes for total NKCC1 samples. Beads were washed twice with a high-salt buffer [0.1% Triton X-100, 500 mM NaCl, 50 mM Tris, and 5 mM EDTA (pH 7.5)] for 10 min and once with a low-salt buffer [0.1% Triton X-100, 100 mM NaCl, 50 mM Tris, and 5 mM EDTA (pH 7.5)] for 10 min. Biotinylated proteins were eluted from streptavidin beads using 50 μ l of Laemmli sample buffer (Bio-Rad). Samples were run on 8% SDS-PAGE and transferred to polyvinylidene difluoride membranes.

Statistical analysis. All data are expressed as means \pm SE for a series of *n* experiments. Student's *t*-test or ANOVA with the Student-Newman-Keuls post test was used to compare mean values as appropriate. *P* < 0.05 was considered to represent significant difference.

RESULTS

H₂O₂ inhibits CCh-stimulated Cl⁻ secretion through activation of AMPK. The metabolic-sensing serine-threonine kinase AMPK has been shown to play an important role in restoration of cellular energy levels under conditions of metabolic stress, including after exposure to oxidants such as H₂O₂ (10, 31, 32). AMPK activation occurs in response to AMP binding or via activation by upstream kinases (14). Moreover, binding of AMP to AMPK makes AMPK a better substrate for kinase activation, resulting in phosphorylation of its catalytic α -subunit on the Thr¹⁷² residue, with a 50- to 100-fold increase in enzyme activity (14). Therefore, we measured phosphorylation (Thr¹⁷²) of AMPK α to investigate whether H₂O₂ increases activation of AMPK. H₂O₂ increased phosphorylation transiently, with a maximal effect between 5 and 15 min (*P* < 0.05–0.01, *n* = 7; Fig. 1, A and B). The AMPK activator AICAR, administered for 30 min, also significantly increased AMPK α phosphorylation (*P* < 0.05, *n* = 4; Fig. 1, A and B). While AMPK has been shown to inhibit cAMP-dependent Cl⁻ secretion through effects on CFTR, its effects on Ca²⁺-dependent epithelial Cl⁻ secretion have not been determined (29, 67). Consequently, we investigated whether AMPK activation was involved in the inhibitory effect of H₂O₂ on Ca²⁺-dependent Cl⁻ secretion stimulated by the acetylcholine analog CCh.

Pretreatment of T84 monolayers with the specific AMPK inhibitor compound C (1–100 μ M) blocked H₂O₂-mediated inhibition of CCh-stimulated Cl⁻ secretion, indicating that AMPK is involved in H₂O₂ inhibition of Ca²⁺-dependent Cl⁻ secretion (50 μ M compound C, *P* < 0.05, *n* = 4; Fig. 1C). This finding was also observed in HT-29.cl19a monolayers, where the inhibitory effect of H₂O₂ was significantly blocked by pretreatment with compound C (*P* < 0.05; Fig. 1D).




Fig. 4. H₂O₂ induces AMP-activated protein kinase (AMPK)-Na⁺-K⁺-2Cl⁻ cotransporter type 1 (NKCC1) association. T84 monolayers were treated with H₂O₂ (500 μ M) for 2–30 min or the AMPK activator 5-aminoimidazole-4-carboxamide-1- β -D-ribofuranoside (AICAR, 2.5 mM) for 30 min. Cell lysates were immunoprecipitated with anti-AMPK α (A) or anti-NKCC1 (N-16) antibody (B) and subsequently probed for NKCC1 and/or AMPK α binding by Western blotting. The catalytic α -subunit of AMPK was immunoprecipitated (IP) and immunoblotted (Blot) for NKCC1 (T4 antibody). Results from blots were quantified by densitometric analysis (*n* = 4). Molecular mass of each protein (kDa) is shown at right. Values (arbitrary units) are means \pm SE and were normalized to total levels of the primary immunoprecipitate (AMPK α or NKCC1). **P* < 0.05, ***P* < 0.01, ****P* < 0.001 vs. control/untreated cells.

H₂O₂-induced AMPK phosphorylation occurs downstream of PI3K and ERK activation. We previously demonstrated that the inhibitory effect of H₂O₂ on CCh-stimulated I_{sc} involves activation of two separate, but Ca²⁺-dependent, signaling pathways. One pathway involves activation of Src, Pyk-2, and ERK/p38 MAPK while the other involves p38-dependent ac-

tivation of PI3K (9). We therefore investigated whether these signaling pathways are also involved in activation of AMPK by H₂O₂. Preincubation of T84 monolayers with the MEK1 inhibitor PD98059 or the PI3K inhibitor wortmannin reduced phosphorylation of AMPK α in response to H₂O₂ ($P < 0.05$ – 0.001 , $n = 3$; Fig. 2). AMPK phosphorylation levels in monolayers treated with PD98059 or wortmannin alone were not statistically different from those in monolayers treated with either agent in the presence of H₂O₂. Consistent with our previous findings, MEK1 inhibition had no effect on H₂O₂-stimulated PI3K signaling measured by Akt phosphorylation; similarly, PI3K inhibition by wortmannin had no effect on ERK phosphorylation, whereas it successfully reduced Akt phosphorylation (Fig. 2A). These data indicate that the ERK and PI3K signaling pathways activated by H₂O₂ converge at the level of AMPK.

AMPK activity mediates H₂O₂ inhibition of NKCC1. Our prior work indicated that H₂O₂ pretreatment inhibits CCh-stimulated NKCC1 activity, as measured by bumetanide-sensitive ⁸⁶Rb⁺ influx, and that the inhibitory effect of H₂O₂ does not involve changes in intracellular Cl⁻ (a major regulator of NKCC1 activity) (9). Further experiments explored whether H₂O₂ inhibition of CCh-stimulated K⁺ influx also involves AMPK activation. Preincubation of T84 monolayers with the AMPK inhibitor compound C (50 μ M) for 30 min reversed the inhibitory effect of H₂O₂ on CCh-stimulated ⁸⁶Rb⁺ influx ($P < 0.01$ vs. Ringer solution, $n = 4$; Fig. 3), while there was no significant difference in ⁸⁶Rb⁺ influx between CCh alone and CCh in the presence of compound C. These data indicate that H₂O₂ inhibition of Ca²⁺-dependent K⁺ uptake, likely via NKCC1 based on our prior studies, is mediated, at least in part, through AMPK activation.

AMPK physically associates with NKCC1 in IECs. AMPK activators such as AICAR and A-769662 have been shown to have no effect or to inhibit NKCC1 activity, with the latter effect due, in part, to phosphorylation at the NH₂ terminus (23, 60). To determine whether AMPK inhibits NKCC1 through a physical association, we used lysates from T84 cells treated with H₂O₂ to carry out coimmunoprecipitation experiments (Fig. 4). We observed a significant level of coimmunoprecipitation of AMPK and NKCC1 (detected with anti-NKCC1 T4 antibody), regardless of whether AMPK or NKCC1 was initially immunoprecipitated [$P < 0.01$, $n = 3$ (Fig. 4A); $P < 0.05$, $n = 3$ (Fig. 4B)]. Thus, inhibition of NKCC1 by AMPK could involve a direct association between the two proteins.




Fig. 5. H₂O₂ increases Na⁺-K⁺-2Cl⁻ cotransporter type 1 (NKCC1) phosphorylation in human colon and in cell lines in an AMP-activated protein kinase (AMPK)-dependent manner. **A**: H₂O₂ treatment of T84 monolayers significantly increased phosphorylation (Thr^{212,217}) of NKCC1 (500 μ M, $P < 0.05$, $n = 4$). Treatment with 5-aminoimidazole-4-carboxamide-1- β -D-ribofuranoside (AICAR, 2.5 mM) for 30 min also increased NKCC1 (Thr^{212,217}) phosphorylation ($P < 0.05$, $n = 4$). **B**: experiment described in **A** was repeated in ex vivo human colonic biopsies from control subjects without active inflammatory intestinal disease ($n = 3$). H₂O₂ (1 mM) also increased AMPK α (Thr¹⁷²) phosphorylation in human colonic tissue. **C**: preincubation of T84 monolayers with compound C (50 μ M, bilaterally) for 30 min significantly reduced H₂O₂ induction of NKCC1 phosphorylation ($P < 0.01$, $n = 4$). Densitometric data (arbitrary units) are presented as means \pm SE for levels of phosphorylation and normalized to total NKCC1 levels. * $P < 0.05$, ** $P < 0.01$, *** $P < 0.001$ vs. vehicle control; & $P < 0.05$ vs. compound C control; ## $P < 0.01$ vs. vehicle H₂O₂.

H₂O₂ modifies the phosphorylation status of NKCC1 through AMPK. Full activity of human NKCC1 requires phosphorylation of Thr residues at positions 212 and 217 (16). We therefore explored whether H₂O₂ affects NKCC1 phosphorylation and whether this involves AMPK activation. We were surprised to find that treatment of T84 monolayers with H₂O₂ (500 μM) induced a rapid increase in phosphorylation (Thr^{212,217}) of NKCC1 that peaked at 15 min ($P < 0.05$, $n = 4$; Fig. 5A). The AMPK activator AICAR also significantly increased NKCC1 phosphorylation ($P < 0.05$, $n = 4$; Fig. 5A). These findings were reproduced using human colonic mucosal biopsies isolated from subjects who were undergoing routine colorectal cancer screening but showed no evidence of intestinal inflammation. Ex vivo treatment with H₂O₂ (1 mM) and subsequent Western blotting for phosphorylated and total forms of NKCC1 and AMPK showed that H₂O₂ increased NKCC1 (Thr^{212,217}), as well as AMPK (Thr¹⁷²), phosphorylation ($n = 3$; Fig. 5B). Follow-up studies in T84 cells showed that H₂O₂-stimulated phosphorylation of NKCC1 was significantly reduced by preincubation with the AMPK inhibitor compound C ($P < 0.01$, $n = 4$; Fig. 5C). These data indicate that H₂O₂ actually increases phosphorylation of NKCC1 residues required for NKCC1 activity.

H₂O₂-activated AMPK does not promote internalization of membrane-bound NKCC1. NKCC1 has also been shown to be regulated via internalization of membrane-bound NKCC1 in colonic epithelial cells (7). Therefore, we employed a surface biotinylation assay to determine if surface expression of NKCC1 is altered and if NKCC1 internalization is AMPK-dependent. Monolayers of T84 cells were treated with H₂O₂ with or without pretreatment with the AMPK inhibitor compound C (50 μM), metformin (5 mM) alone as a positive control for AMPK activation, or capsaicin (20 μM) alone as a

positive control for NKCC1 internalization (7, 67). After treatment, proteins expressed in the basolateral membrane were biotinylated and pulled down by immunoprecipitation with streptavidin-coated agarose beads. We observed no significant change in NKCC1 surface expression in response to H₂O₂ treatment, as well as no effect of compound C pretreatment, although there appeared to be an increasing trend of NKCC1 surface expression with H₂O₂ treatment (Fig. 6). Furthermore, AMPK activation by metformin (5 mM) did not alter NKCC1 internalization. Capsaicin (20 μM) also did not induce NKCC1 internalization, despite our use of a previously established incubation time and concentration of capsaicin that were shown to promote NKCC1 internalization in T84 cells (7). Our data suggest that H₂O₂ inhibition of Cl⁻ secretion is independent of NKCC1 internalization and that AMPK does not mediate distinct patterns of membrane turnover of NKCC1.

DISCUSSION

Regulation of epithelial transport is a complex and multifaceted physiological process that requires careful orchestration of acute vs. chronic and transient vs. sustained events. In the acute regulation of electrolyte transport in response to exposure by ROS, we investigated whether the cellular energy sensor AMPK, itself capable of acute and/or sustained regulation of many metabolic events, was involved in the H₂O₂ regulation of Ca²⁺-dependent ion transport. We previously showed that H₂O₂ inhibits CCh-stimulated Cl⁻ secretion (9). CCh is a more stable analog of the neurotransmitter acetylcholine that plays a prominent role in the minute-to-minute regulation of fluid secretion in intestinal crypts through activation of Ca²⁺-dependent signals following binding to muscarinic receptors on IECs (2). We further found that the inhibitory




Fig. 6. AMP-activated protein kinase (AMPK) inhibition of the Na⁺-K⁺-2Cl⁻ cotransporter NKCC1 is independent of internalization. NKCC1 surface expression was detected using a biotinylation assay, where biotinylation indicated membrane insertion. H₂O₂ treatment of T84 monolayers slightly, but not statistically significantly, increased surface expression of NKCC1 (500 μM; $n = 3$). Pretreatment with the AMPK inhibitor compound C (CC, 50 μM, bilaterally) for 30 min did not decrease NKCC1 internalization. Neither metformin (Metf, 5 mM, bilaterally; $n = 3$) nor capsaicin (Cap, 20 μM, bilaterally, $n = 3$) treatment significantly altered surface expression of NKCC1. Densitometric data (arbitrary units) are presented as means ± SE for levels of biotinylated NKCC1 and normalized to total NKCC1 levels.

effect of H₂O₂ occurred by virtue of its ability to reduce the activity of the basolateral NKCC1, which is responsible for basolateral Cl⁻ entry into IECs and is critically involved in overall epithelial homeostasis (9). This finding was supported by *in vivo* studies showing that administration of the antioxidant enzyme catalase, which degrades H₂O₂, restricted the suppression of colonic ion transport responsiveness and reduced levels of colonic epithelial NKCC1 caused by dextran sulfate sodium-induced colitis (3). In the present study we showed that AMPK activity is a critical mediator of the inhibitory effect of H₂O₂ on Cl⁻ secretion and NKCC1 activity. H₂O₂ increased phosphorylation (Thr¹⁷²) of AMPK α in human colonic epithelial cell lines and in *ex vivo* human colonic mucosal biopsies. While there was some variability between human samples from different patients with respect to the magnitude and kinetics of H₂O₂ induction of AMPK α and NKCC1 phosphorylation, the overall effect of H₂O₂ as a stimulus of phosphorylation of these proteins was consistent and, thus, broadly aligned with our cell line data. The AMPK inhibitor compound C dramatically reduced the inhibitory effect of H₂O₂ on overall ion transport and CCh-stimulated NKCC1 activity. The inhibitory effect of H₂O₂ on NKCC1 activity was associated with an increased physical association between AMPK and NKCC1 based on reciprocal coimmunoprecipitation studies. Since NKCC1 activity is dependent on phosphorylation of regulatory Thr²¹² and Thr²¹⁷ residues (16, 22, 44, 45), we investigated if H₂O₂ had an effect on NKCC1 phosphorylation. Using the R5 antibody, which recognizes phosphorylation of these residues, we were surprised to find that H₂O₂ actually increases NKCC1 phosphorylation in both IEC lines and human colonic biopsies, even while H₂O₂ exerted an inhibitory effect on overall NKCC1 activity. Phosphorylation of NKCC1 at its regulatory Thr²¹² and Thr²¹⁷ residues in response to H₂O₂ appears, ultimately, to require AMPK activity, since both AICAR and H₂O₂ elicited NKCC1 phosphorylation in a manner antagonized by compound C.

AMPK activity has been linked to inhibition of renal-specific NKCC2 and, recently, to NKCC1 activity in *in vitro* model systems. This was shown to occur in response to selective activators of AMPK, such as AICAR and A-769662, and also in response to osmotic shrinkage (23, 24, 60). However, in these studies, AMPK phosphorylation of NKCC1 at Ser³⁸ and Ser²¹⁴ (dogfish sequence), corresponding to Ser⁷⁷ and Ser²⁴² in human NKCC1, was, in fact, associated with reduced NKCC1 activity. The consensus amino acid sequence that is associated with AMPK binding and Ser/Thr phosphorylation involves the following recognition motif: $\varphi(X\beta)XXS/TXXX\varphi$, where φ is a hydrophobic residue (M, V, L, I, or F), β is a basic residue (R, K, or H), and the parentheses indicate that the order of residues at the P-4 and P-3 positions is not essential (15, 60). The sequence surrounding Ser²⁴² of human NKCC1 loosely fits the AMPK consensus recognition motif, with an Arg residue at the P-2, rather than P-3, position (60). Additional studies with cell-free assays showed increased phosphorylation of Ser⁷⁷ following AMPK activation by phenformin, while a NKCC1 construct with a point mutation at Ser⁷⁷ showed less phosphorylation and a 16% decrease in the inhibitory effect of phenformin on NKCC1 transport of ⁸⁶Rb⁺, thus indicating a partial role for this residue in AMPK regulation of NKCC1 (23). One caveat is that phenformin can exert AMPK-independent signaling events (66).

Further studies indicate that AMPK can inhibit activity of Ste-related proline-alanine-rich kinase (SPAK)/oxidative stress-responsive kinase (OSR1), which phosphorylates and activates NKCC1 in response to several stimuli (17, 60). However, there does appear to be a context- and cell type-dependent pattern of regulation of NKCC1 by AMPK that is not always consistent with NKCC1 phosphorylation levels or kinetics. Indeed, this is not an unusual feature of AMPK, as it has been shown to play context-specific roles in tumorigenesis (42). Interestingly, the cell type-specific effects of individual AMPK activators were emphasized by the use of A-769662: Sid et al. observed that A-769662, while increasing phosphorylation of Ser²⁴², had no effect on NKCC1 activity in erythrocytes (60), whereas Fraser et al. observed inhibition of NKCC1 activity by A-769662 in Madin-Darby canine kidney cells (23). There also appear to be important differences between the effects of basal and stimulated AMPK activity in the regulation of NKCC1: Davies et al. observed that in erythrocytes the constitutive absence or suppression of AMPK activity could regulate SPAK/OSR1 through mechanisms different from those invoked by acute




Fig. 7. Signaling pathways activated by H₂O₂ to inhibit Na⁺-K⁺-2Cl⁻ cotransporter NKCC1 converge at AMP-activated protein kinase (AMPK). H₂O₂ stimulates an increase in cytosolic Ca²⁺ concentration (1). Two separate Ca²⁺-stimulated signaling pathways [Src-Pyk2-ERK and p38-phosphatidylinositol 3-kinase (PI3K)] are activated (2). These pathways converge to increase AMPK activation (3). H₂O₂ also inhibits basolateral Ca²⁺-responsive K⁺ influx mediated by NKCC1 in an AMPK-dependent manner, and H₂O₂ increases physical association of AMPK with NKCC1 and increases NKCC1 phosphorylation via AMPK activity (4). Suppression of NKCC1 activity likely contributes to the overall inhibitory effect of H₂O₂ on Ca²⁺-dependent ion transport in colonic epithelial cells (5).

changes in AMPK activity (17). Furthermore, the nuanced regulation of NKCC1 activity was also emphasized by findings that the extent of NKCC1 phosphorylation did not directly correlate with NKCC1 activation by hyperosmolarity in mouse erythrocytes; thus, Sid et al. concluded that other factors are likely involved (60).

Even though we observed an increase in H₂O₂-stimulated phosphorylation of two key residues (Thr²¹² and Thr²¹⁷) that are required for increasing NKCC1 activity, increased AMPK activity in Madin-Darby canine kidney cells treated with the AMPK activator A-769662 correlated with reduced NKCC1 phosphorylation on Thr²¹² and Thr²¹⁷ (23). Our studies indicate a clear functional role for H₂O₂-activated AMPK in inhibiting Ca²⁺-dependent electrolyte transport and NKCC1 activity (cf. Figs. 1A, 2, and 4A). We also observed a partial, although not statistically significant, increase in NKCC1 phosphorylation in cells incubated with compound C alone (cf. Fig. 5), with no effect on NKCC1 activity (cf. Fig. 3), which is consistent with the findings of Fraser et al. (23). Our results suggest that AMPK may have a dual role with respect to NKCC1 phosphorylation in unstimulated/resting conditions vs. that in response to an inflammatory stimulus. Moreover, we did not see an increase in ⁸⁶Rb⁺ flux in colonic epithelial cells treated with compound C alone, which is also in agreement with the findings of Fraser et al. (23). AMPK activity is also stimulus-specific, as well-established activators such as AICAR and A-769662 act via different mechanisms. AICAR is phosphorylated by adenylate kinase to generate the AMP analog ZMP, which directly binds to the α -subunit of AMPK, while A-769662 allosterically activates AMPK through binding to its regulatory β 1-subunit (14, 59). Indeed, coadministration of these agents has been shown to have a synergistic effect with respect to AMPK activation in mouse hepatocytes (19). Another important consideration of select activators of AMPK is their potential for off-target effects. AICAR and metformin have been shown to inhibit proliferation by AMPK-independent mechanisms, while A-769662 can effectively inhibit proteasomal activity, which is relevant to the study of NKCC1 function, since NKCC1 can be regulated by proteasomal degradation (18, 42).

With respect to the stimulus used to activate AMPK in these studies, several groups have reported that oxidative stress can activate AMPK in a cell type-specific manner independently of alterations in the AMP-to-ATP ratio (reviewed in Refs. 40 and 61). We previously demonstrated that H₂O₂, at doses used in the present study and in the same cell lines, did not cause disruption of barrier function in T84 cells, thus indicating that cell integrity is maintained (9). Our previous work demonstrated that H₂O₂ caused a sustained increase in intracellular Ca²⁺ levels that triggered activation of two signaling pathways, leading to increased phosphorylation of MEK and Akt downstream of PI3K. Our data suggest that these pathways feed into AMPK phosphorylation, as phosphorylated AMPK levels were reduced by ERK or PI3K inhibition (cf. Fig. 3). Our biotinylation-based internalization assays did not detect a clear effect of H₂O₂ on membrane localization of NKCC1, although there was a counterintuitive trend toward increased membrane localization. This pattern was also observed in response to capsaicin treatment, which was included as a positive control to promote internalization based on a previous study (7). Given the lack of a significant effect of H₂O₂ on

membrane NKCC1 levels, we interpret our findings as indicating that H₂O₂-AMPK signaling primarily inhibits NKCC1 by altering its activation status, rather than by causing a substantial internalization of NKCC1 away from the basolateral membrane.

In conclusion, we have demonstrated that pathophysiologically relevant concentrations of H₂O₂ can inhibit Ca²⁺-dependent Cl⁻ secretion by activating kinase-signaling pathways, leading to activation of AMPK and inhibition of NKCC1 activity and, overall, ion transport responsiveness (Fig. 7). Our findings emphasize the context-dependent nature of the relationship between AMPK and NKCC1 but are consistent with an overall inhibitory effect of H₂O₂ and AMPK on the ion-transporting capability of IECs that aligns with the loss of transport capacity in intestinal inflammation (35). In agreement with the findings of others, it appears that regulation of NKCC1 activity is more complex than a linear and uniform series of protein kinase signals and may involve multiple protein kinase complexes acting in different cell types and resulting in varied outcomes determined by specific stimuli (62, 63). The findings in our study serve to further expand our understanding of how AMPK may act to negatively regulate energy-consuming cellular activities in epithelia in the setting of inflammation.

ACKNOWLEDGMENTS

We are grateful to Merrie Mosedale (University of California, San Diego) and Kristian Lucas (University of California, Riverside) for technical assistance. We are grateful to Dr. Kim E. Barrett (University of California, San Diego) for providing T84 and HT-29.cl19a cell lines and to Dr. Biff Forbush (Yale University) for providing the R5 anti-phosphorylated (Thr²¹², Thr²¹⁷) NKCC1 antibody.

Current addresses: A. Chappell, Ionis Pharmaceuticals, 2855 Gazelle Ct., Carlsbad, CA 92010; B. Jung, Div. of Gastroenterology and Hepatology, University of Illinois at Chicago, Chicago, IL 60612.

GRANTS

This work was supported by a Pease-Burden Fellowship (to S. J. King), a Crohn's and Colitis Foundation Career Development Award (to D. F. McCole), and National Institute of Diabetes and Digestive and Kidney Diseases Grant 2R01 DK-091281 (to D. F. McCole).

DISCLOSURES

No conflicts of interest, financial or otherwise, are declared by the authors.

AUTHOR CONTRIBUTIONS

S.J.K., M.B., C.L., and D.F.M. conceived and designed research; S.J.K., M.B., A.C., M.S., M.D., B.J., and D.F.M. performed experiments; S.J.K., M.B., A.C., M.S., and D.F.M. analyzed data; S.J.K., M.B., C.L., and D.F.M. interpreted results of experiments; S.J.K., M.B., A.C., and D.F.M. prepared figures; S.J.K., M.B., B.J., C.L., and D.F.M. edited and revised manuscript; S.J.K., M.B., A.C., M.S., M.D., B.J., C.L., and D.F.M. approved final version of manuscript; D.F.M. drafted manuscript.

REFERENCES

1. Amasheh S, Barmeyer C, Koch CS, Tavalali S, Mankertz J, Epple HJ, Gehring MM, Florian P, Kroesen AJ, Zeitz M, Fromm M, Schulzke JD. Cytokine-dependent transcriptional down-regulation of epithelial sodium channel in ulcerative colitis. *Gastroenterology* 126: 1711–1720, 2004. doi:10.1053/j.gastro.2004.03.010.
2. Barrett KE, Keely SJ. Chloride secretion by the intestinal epithelium: molecular basis and regulatory aspects. *Annu Rev Physiol* 62: 535–572, 2000. doi:10.1146/annurev.physiol.62.1.535.
3. Barrett KE, McCole DF. Hydrogen peroxide scavenger, catalase, alleviates ion transport dysfunction in murine colitis. *Clin Exp Pharmacol Physiol* 43: 1097–1106, 2016. doi:10.1111/1440-1681.12646.

4. **Bevins CL, Salzman NH.** The potter's wheel: the host's role in sculpting its microbiota. *Cell Mol Life Sci* 68: 3675–3685, 2011. doi:10.1007/s00018-011-0830-3.
5. **Bhalla V, Oyster NM, Fitch AC, Wijngaarden MA, Neumann D, Schlattner U, Pearce D, Hallows KR.** AMP-activated kinase inhibits the epithelial Na⁺ channel through functional regulation of the ubiquitin ligase Nedd4-2. *J Biol Chem* 281: 26159–26169, 2006. doi:10.1074/jbc.M606045200.
6. **Biasi F, Leonarduzzi G, Oteiza PI, Poli G.** Inflammatory bowel disease: mechanisms, redox considerations, and therapeutic targets. *Antioxid Redox Signal* 19: 1711–1747, 2013. doi:10.1089/ars.2012.4530.
7. **Bouyer PG, Tang X, Weber CR, Shen L, Turner JR, Matthews JB.** Capsaicin induces NKCC1 internalization and inhibits chloride secretion in colonic epithelial cells independently of TRPV1. *Am J Physiol Gastrointest Liver Physiol* 304: G142–G156, 2013. doi:10.1152/ajpgi.00483.2011.
8. **Cartwright CA, McRoberts JA, Mandel KG, Dharmasathaphorn K.** Synergistic action of cyclic adenosine monophosphate- and calcium-mediated chloride secretion in a colonic epithelial cell line. *J Clin Invest* 76: 1837–1842, 1985. doi:10.1172/JCI112176.
9. **Chappell AE, Bunz M, Smoll E, Dong H, Lytle C, Barrett KE, McCole DF.** Hydrogen peroxide inhibits Ca²⁺-dependent chloride secretion across colonic epithelial cells via distinct kinase signaling pathways and ion transport proteins. *FASEB J* 22: 2023–2036, 2008. doi:10.1096/fj.07-099697.
10. **Choi SL, Kim SJ, Lee KT, Kim J, Mu J, Birnbaum MJ, Soo Kim S, Ha J.** The regulation of AMP-activated protein kinase by H₂O₂. *Biochem Biophys Res Commun* 287: 92–97, 2001. doi:10.1006/bbrc.2001.5544.
11. **Choudhary S, Keshavarzian A, Yong S, Wade M, Bocchino S, Day BJ, Banan A.** Novel antioxidants zolimid and AEOL11201 ameliorate colitis in rats. *Dig Dis Sci* 46: 2222–2230, 2001. doi:10.1023/A:1011975218006.
12. **Conner EM, Brand SJ, Davis JM, Kang DY, Grisham MB.** Role of reactive metabolites of oxygen and nitrogen in inflammatory bowel disease: toxins, mediators, and modulators of gene expression. *Inflamm Bowel Dis* 2: 133–147, 1996. doi:10.1097/00054725-199606000-00011.
13. **Cooke HJ.** "Enteric tears": chloride secretion and its neural regulation. *News Physiol Sci* 13: 269–274, 1998. doi:10.1152/physiologyonline.1998.13.6.269.
14. **Corton JM, Gillespie JG, Hardie DG.** Role of the AMP-activated protein kinase in the cellular stress response. *Curr Biol* 4: 315–324, 1994. doi:10.1016/S0960-9822(00)00070-1.
15. **Dale S, Wilson WA, Edelman AM, Hardie DG.** Similar substrate recognition motifs for mammalian AMP-activated protein kinase, higher plant HMG-CoA reductase kinase-A, yeast SNF1, and mammalian calmodulin-dependent protein kinase I. *FEBS Lett* 361: 191–195, 1995. doi:10.1016/0014-5793(95)00172-6.
16. **Darman RB, Forbush B.** A regulatory locus of phosphorylation in the N terminus of the Na-K-Cl cotransporter, NKCC1. *J Biol Chem* 277: 37542–37550, 2002. doi:10.1074/jbc.M206293200.
17. **Davies M, Fraser SA, Galic S, Choy SW, Katerelos M, Gleich K, Kemp BE, Mount PF, Power DA.** Novel mechanisms of Na⁺ retention in obesity: phosphorylation of NKCC2 and regulation of SPAK/OSR1 by AMPK. *Am J Physiol Renal Physiol* 307: F96–F106, 2014. doi:10.1152/ajprenal.00524.2013.
18. **Del Castillo IC, Fedor-Chaikin M, Song JC, Starlinger V, Yoo J, Matlin KS, Matthews JB.** Dynamic regulation of Na⁺-K⁺-2Cl⁻ cotransporter surface expression by PKC-ε in Cl⁻-secretory epithelia. *Am J Physiol Cell Physiol* 289: C1332–C1343, 2005. doi:10.1152/ajpcell.00580.2004.
19. **Ducommun S, Ford RJ, Bultot L, Deak M, Bertrand L, Kemp BE, Steinberg GR, Sakamoto K.** Enhanced activation of cellular AMPK by dual-small molecule treatment: AICAR and A769662. *Am J Physiol Endocrinol Metab* 306: E688–E696, 2014. doi:10.1152/ajpendo.00672.2013.
20. **DuVall MD, Guo Y, Matalon S.** Hydrogen peroxide inhibits cAMP-induced Cl⁻ secretion across colonic epithelial cells. *Am J Physiol Cell Physiol* 275: C1313–C1322, 1998. doi:10.1152/ajpcell.1998.275.5.C1313.
21. **Flagella M, Clarke LR, Miller ML, Erway LC, Giannella RA, Andringa A, Gawenis LR, Kramer J, Duffy JJ, Doetschman T, Lorenz JN, Yamoah EN, Cardell EL, Shull GE.** Mice lacking the basolateral Na-K-2Cl cotransporter have impaired epithelial chloride secretion and are profoundly deaf. *J Biol Chem* 274: 26946–26955, 1999. doi:10.1074/jbc.274.38.26946.
22. **Flemmer AW, Gimenez I, Dowd BF, Darman RB, Forbush B.** Activation of the Na-K-Cl cotransporter NKCC1 detected with a phospho-specific antibody. *J Biol Chem* 277: 37551–37558, 2002. doi:10.1074/jbc.M206294200.
23. **Fraser SA, Davies M, Katerelos M, Gleich K, Choy SW, Steel R, Galic S, Mount PF, Kemp BE, Power DA.** Activation of AMPK reduces the co-transporter activity of NKCC1. *Mol Membr Biol* 31: 95–102, 2014. doi:10.3109/09687688.2014.902128.
24. **Fraser SA, Gimenez I, Cook N, Jennings I, Katerelos M, Katsis F, Levidiotis V, Kemp BE, Power DA.** Regulation of the renal-specific Na⁺-K⁺-2Cl⁻ co-transporter NKCC2 by AMP-activated protein kinase (AMPK). *Biochem J* 405: 85–93, 2007. doi:10.1042/BJ20061850.
25. **Fryer LG, Hajdudch E, Rencurel F, Salt IP, Hundal HS, Hardie DG, Carling D.** Activation of glucose transport by AMP-activated protein kinase via stimulation of nitric oxide synthase. *Diabetes* 49: 1978–1985, 2000. doi:10.2337/diabetes.49.12.1978.
26. **Greig E, Sandle GI.** Diarrhea in ulcerative colitis. The role of altered colonic sodium transport. *Ann NY Acad Sci* 915: 327–332, 2000. doi:10.1111/j.1749-6632.2000.tb05260.x.
27. **Grisham MB.** Oxidants and free radicals in inflammatory bowel disease. *Lancet* 344: 859–861, 1994. doi:10.1016/S0140-6736(94)92831-2.
28. **Hallows KR.** Emerging role of AMP-activated protein kinase in coupling membrane transport to cellular metabolism. *Curr Opin Nephrol Hypertens* 14: 464–471, 2005. doi:10.1097/01.mnh.0000174145.14798.64.
29. **Hallows KR, Kobinger GP, Wilson JM, Witters LA, Foskett JK.** Physiological modulation of CFTR activity by AMP-activated protein kinase in polarized T84 cells. *Am J Physiol Cell Physiol* 284: C1297–C1308, 2003. doi:10.1152/ajpcell.00227.2002.
30. **Hallows KR, Raghuram V, Kemp BE, Witters LA, Foskett JK.** Inhibition of cystic fibrosis transmembrane conductance regulator by novel interaction with the metabolic sensor AMP-activated protein kinase. *J Clin Invest* 105: 1711–1721, 2000. doi:10.1172/JCI9622.
31. **Hardie DG, Carling D.** The AMP-activated protein kinase—fuel gauge of the mammalian cell? *Eur J Biochem* 246: 259–273, 1997. doi:10.1111/j.1432-1033.1997.00259.x.
32. **Hardie DG, Carling D, Carlson M.** The AMP-activated/SNF1 protein kinase subfamily: metabolic sensors of the eukaryotic cell? *Annu Rev Biochem* 67: 821–855, 1998. doi:10.1146/annurev.biochem.67.1.821.
33. **Harris ML, Schiller HJ, Reilly PM, Donowitz M, Grisham MB, Bulkeley GB.** Free radicals and other reactive oxygen metabolites in inflammatory bowel disease: cause, consequence or epiphenomenon? *Pharmacol Ther* 53: 375–408, 1992. doi:10.1016/0163-7258(92)90057-7.
34. **Hirota CL, McKay DM.** Loss of Ca-mediated ion transport during colitis correlates with reduced ion transport responses to a Ca-activated K channel opener. *Br J Pharmacol* 156: 1085–1097, 2009. doi:10.1111/j.1476-5381.2009.00122.x.
35. **Kachur JF, Keshavarzian A, Sundaresan R, Doria M, Walsh R, de las Alas MM, Gaginnella TS.** Colitis reduces short-circuit current response to inflammatory mediators in rat colonic mucosa. *Inflammation* 19: 245–259, 1995. doi:10.1007/BF01534465.
36. **Keely SJ, Calandrella SO, Barrett KE.** Carbachol-stimulated transactivation of epidermal growth factor receptor and mitogen-activated protein kinase in T₈₄ cells is mediated by intracellular Ca²⁺, PYK-2, and p60^{src}. *J Biol Chem* 275: 12619–12625, 2000. doi:10.1074/jbc.275.17.12619.
37. **Keshavarzian A, Haydek J, Zabihi R, Doria M, D'Astice M, Sorenson JR.** Agents capable of eliminating reactive oxygen species. Catalase, WR-2721, or Cu(II)2(3,5-DIPS)4 decrease experimental colitis. *Dig Dis Sci* 37: 1866–1873, 1992.
38. **Kruiderien L, Kuiper I, Lamers CB, Verspaget HW.** Intestinal oxidative damage in inflammatory bowel disease: semi-quantification, localization, and association with mucosal antioxidants. *J Pathol* 201: 28–36, 2003. doi:10.1002/path.1409.
39. **Li W, Duzgun A, Sumpio BE, Basson MD.** Integrin and FAK-mediated MAPK activation is required for cyclic strain mitogenic effects in Caco-2 cells. *Am J Physiol Gastrointest Liver Physiol* 280: G75–G87, 2001. doi:10.1152/ajpgi.2001.280.1.G75.
40. **Liangpunsakul S, Wou SE, Zeng Y, Ross RA, Jayaram HN, Crabb DW.** Effect of ethanol on hydrogen peroxide-induced AMPK phosphorylation. *Am J Physiol Gastrointest Liver Physiol* 295: G1173–G1181, 2008. doi:10.1152/ajpgi.90349.2008.
41. **Lih-Brody L, Powell SR, Collier KP, Reddy GM, Cerchia R, Kahn E, Weissman GS, Katz S, Floyd RA, McKinley MJ, Fisher SE, Mullin GE.** Increased oxidative stress and decreased antioxidant defenses in

- mucosa of inflammatory bowel disease. *Dig Dis Sci* 41: 2078–2086, 1996. doi:10.1007/BF02093613.
42. Liu X, Chhipa RR, Pooya S, Wortman M, Yachyshin S, Chow LM, Kumar A, Zhou X, Sun Y, Quinn B, McPherson C, Warnick RE, Kendler A, Giri S, Poels J, Norga K, Viollet B, Grabowski GA, Dasgupta B. Discrete mechanisms of mTOR and cell cycle regulation by AMPK agonists independent of AMPK. *Proc Natl Acad Sci USA* 111: E435–E444, 2014. doi:10.1073/pnas.1311121111.
 43. Lynch SV, Goldfarb KC, Wild YK, Kong W, De Lisle RC, Brodie EL. Cystic fibrosis transmembrane conductance regulator knockout mice exhibit aberrant gastrointestinal microbiota. *Gut Microbes* 4: 41–47, 2013. doi:10.4161/gmic.22430.
 44. Lytle C, Forbush B III. Na-K-Cl cotransport in the shark rectal gland. II. Regulation in isolated tubules. *Am J Physiol Cell Physiol* 262: C1009–C1017, 1992. doi:10.1152/ajpcell.1992.262.4.C1009.
 45. Lytle C, Forbush B III. The Na-K-Cl cotransport protein of shark rectal gland. II. Regulation by direct phosphorylation. *J Biol Chem* 267: 25438–25443, 1992.
 46. Lytle C, Xu JC, Biemesderfer D, Forbush B III. Distribution and diversity of Na-K-Cl cotransport proteins: a study with monoclonal antibodies. *Am J Physiol Cell Physiol* 269: C1496–C1505, 1995. doi:10.1152/ajpcell.1995.269.6.C1496.
 47. McCole DF, Barrett KE. Varied role of the gut epithelium in mucosal homeostasis. *Curr Opin Gastroenterol* 23: 647–654, 2007. doi:10.1097/MOG.0b013e3282f0153b.
 48. Nguyen TD, Canada AT. Modulation of human colonic T84 cell secretion by hydrogen peroxide. *Biochem Pharmacol* 47: 403–410, 1994. doi:10.1016/0006-2952(94)90032-9.
 49. Norkina O, Kaur S, Ziemer D, De Lisle RC. Inflammation of the cystic fibrosis mouse small intestine. *Am J Physiol Gastrointest Liver Physiol* 286: G1032–G1041, 2004. doi:10.1152/ajpgi.00473.2003.
 50. Oz HS, Chen TS, McClain CJ, de Villiers WJ. Antioxidants as novel therapy in a murine model of colitis. *J Nutr Biochem* 16: 297–304, 2005. doi:10.1016/j.jnutbio.2004.09.007.
 51. Pavlick KP, Laroux FS, Fuseler J, Wolf RE, Gray L, Hoffman J, Grisham MB. Role of reactive metabolites of oxygen and nitrogen in inflammatory bowel disease. *Free Radic Biol Med* 33: 311–322, 2002. doi:10.1016/S0891-5849(02)00853-5.
 52. Rezaie A, Parker RD, Abdollahi M. Oxidative stress and pathogenesis of inflammatory bowel disease: an epiphenomenon or the cause? *Dig Dis Sci* 52: 2015–2021, 2007. doi:10.1007/s10620-006-9622-2.
 53. Rogers AC, Huetter L, Hoekstra N, Collins D, Collaco A, Baird AW, Winter DC, Ameen N, Geibel JP, Kopic S. Activation of AMPK inhibits cholera toxin stimulated chloride secretion in human and murine intestine. *PLoS One* 8: e69050, 2013. doi:10.1371/journal.pone.0069050.
 54. Sandle GI, Higgs N, Crowe P, Marsh MN, Venkatesan S, Peters TJ. Cellular basis for defective electrolyte transport in inflamed human colon. *Gastroenterology* 99: 97–105, 1990. doi:10.1016/0016-5085(90)91235-X.
 55. Scharl M, Paul G, Barrett KE, McCole DF. AMP-activated protein kinase mediates the interferon- γ -induced decrease in intestinal epithelial barrier function. *J Biol Chem* 284: 27952–27963, 2009. doi:10.1074/jbc.M109.046292.
 56. Scharl M, Paul G, Weber A, Jung BC, Docherty MJ, Hausmann M, Rogler G, Barrett KE, McCole DF. Protection of epithelial barrier function by the Crohn's disease associated gene protein tyrosine phosphatase N2. *Gastroenterology* 137: 2030–2040.e5, 2009. doi:10.1053/j.gastro.2009.07.078.
 57. Schmitz H, Barmeyer C, Gitter AH, Wullstein F, Bentzel CJ, Fromm M, Riecken EO, Schulzke JD. Epithelial barrier and transport function of the colon in ulcerative colitis. *Ann NY Acad Sci* 915: 312–326, 2000. doi:10.1111/j.1749-6632.2000.tb05259.x.
 58. Schultheiss G, Hennig B, Diener M. Sites of action of hydrogen peroxide on ion transport across rat distal colon. *Br J Pharmacol* 154: 991–1000, 2008. doi:10.1038/bjp.2008.162.
 59. Scott JW, van Denderen BJ, Jorgensen SB, Honeyman JE, Steinberg GR, Oakhill JS, Iseli TJ, Koay A, Gooley PR, Stapleton D, Kemp BE. Thienopyridone drugs are selective activators of AMP-activated protein kinase beta1-containing complexes. *Chem Biol* 15: 1220–1230, 2008. doi:10.1016/j.chembiol.2008.10.005.
 60. Sid B, Miranda L, Vertommen D, Viollet B, Rider MH. Stimulation of human and mouse erythrocyte Na⁺-K⁺-2Cl⁻ cotransport by osmotic shrinkage does not involve AMP-activated protein kinase, but is associated with STE20/SPS1-related proline/alanine-rich kinase activation. *J Physiol* 588: 2315–2328, 2010. doi:10.1113/jphysiol.2009.185900.
 61. Sid B, Verrax J, Calderon PB. Role of AMPK activation in oxidative cell damage: implications for alcohol-induced liver disease. *Biochem Pharmacol* 86: 200–209, 2013. doi:10.1016/j.bcp.2013.05.007.
 62. Smith L, Litman P, Liedtke CM. COMMD1 interacts with the COOH terminus of NKCC1 in Calu-3 airway epithelial cells to modulate NKCC1 ubiquitination. *Am J Physiol Cell Physiol* 305: C133–C146, 2013. doi:10.1152/ajpcell.00394.2012.
 63. Smith L, Smallwood N, Altman A, Liedtke CM. PKC δ acts upstream of SPAK in the activation of NKCC1 by hyperosmotic stress in human airway epithelial cells. *J Biol Chem* 283: 22147–22156, 2008. doi:10.1074/jbc.M801752200.
 64. Test ST, Weiss SJ. Quantitative and temporal characterization of the extracellular H₂O₂ pool generated by human neutrophils. *J Biol Chem* 259: 399–405, 1984.
 65. Tomas J, Mulet C, Saffarian A, Cavin JB, Ducroc R, Regnault B, Kun Tan C, Duszka K, Burcelin R, Wahli W, Sansonetti PJ, Pédrón T. High-fat diet modifies the PPAR- γ pathway leading to disruption of microbial and physiological ecosystem in murine small intestine. *Proc Natl Acad Sci USA* 113: E5934–E5943, 2016. doi:10.1073/pnas.1612559113.
 66. Vincent EE, Coelho PP, Blagih J, Griss T, Viollet B, Jones RG. Differential effects of AMPK agonists on cell growth and metabolism. *Oncogene* 34: 3627–3639, 2015. doi:10.1038/ncr.2014.301.
 67. Walker J, Jijon HB, Churchill T, Kulka M, Madsen KL. Activation of AMP-activated protein kinase reduces cAMP-mediated epithelial chloride secretion. *Am J Physiol Gastrointest Liver Physiol* 285: G850–G860, 2003. doi:10.1152/ajpgi.00077.2003.
 68. Walker J, Jijon HB, Diaz H, Salehi P, Churchill T, Madsen KL. 5-Aminoimidazole-4-carboxamide riboside (AICAR) enhances GLUT2-dependent jejunal glucose transport: a possible role for AMPK. *Biochem J* 385: 485–491, 2005. doi:10.1042/BJ20040694.
 69. Watson AJ, Askew JN, Sandle GI. Characterisation of oxidative injury to an intestinal cell line (HT-29) by hydrogen peroxide. *Gut* 35: 1575–1581, 1994. doi:10.1136/gut.35.11.1575.
 70. Zheng Y, Shen X. H₂O₂ directly activates inositol 1,4,5-trisphosphate receptors in endothelial cells. *Redox Rep* 10: 29–36, 2005. doi:10.1179/135100005X21660.