

UC San Diego

UC San Diego Previously Published Works

Title

Chronic air pollution and social deprivation as modifiers of the association between high temperature and daily mortality

Permalink

<https://escholarship.org/uc/item/8mw282kp>

Journal

Environmental Health, 13(1)

ISSN

1476-069X

Authors

Benmarhnia, Tarik

Oulhote, Youssef

Petit, Claire

et al.

Publication Date

2014-12-01

DOI

10.1186/1476-069x-13-53

Copyright Information

This work is made available under the terms of a Creative Commons Attribution License, available at <https://creativecommons.org/licenses/by/4.0/>

Peer reviewed



RESEARCH

Open Access

Chronic air pollution and social deprivation as modifiers of the association between high temperature and daily mortality

Tarik Benmarhnia^{1,2*}, Youssef Oulhote², Claire Petit^{1,3}, Annabelle Lapostolle^{4,5}, Pierre Chauvin^{4,5}, Denis Zmirou-Navier^{1,3,6} and Séverine Deguen^{1,3}

Abstract

Background: Heat and air pollution are both associated with increases in mortality. However, the interactive effect of temperature and air pollution on mortality remains unsettled. Similarly, the relationship between air pollution, air temperature, and social deprivation has never been explored.

Methods: We used daily mortality data from 2004 to 2009, daily mean temperature variables and relative humidity, for Paris, France. Estimates of chronic exposure to air pollution and social deprivation at a small spatial scale were calculated and split into three strata. We developed a stratified Poisson regression models to assess daily temperature and mortality associations, and tested the heterogeneity of the regression coefficients of the different strata. Deaths due to ambient temperature were calculated from attributable fractions and mortality rates were estimated.

Results: We found that chronic air pollution exposure and social deprivation are effect modifiers of the association between daily temperature and mortality. We found a potential interactive effect between social deprivation and chronic exposure with regards to air pollution in the mortality-temperature relationship.

Conclusion: Our results may have implications in considering chronically polluted areas as vulnerable in heat action plans and in the long-term measures to reduce the burden of heat stress especially in the context of climate change.

Keywords: Heat, Population health, Time-series models, Air pollution, Socioeconomic status

Introduction

Among environmental determinants of health, climatic effects are of high concern, especially given the growing body of literature concerning the impacts of climate change. Ambient temperature has long been recognized as a physical hazard, and is associated with a wide range of adverse health effects [1]. Increases in summer ambient temperatures are associated with increases in mortalities over a city specific threshold. Consequently, future heat-related mortality is likely to increase in the context of climate change [2,3].

Identification of factors that increase vulnerability to hot temperature in human populations has become a growing public health concern [4,5]. Some studies have explored different aspects of heat-related mortality or morbidity heterogeneity across different levels. These heterogeneity factors or effect modifiers in relation to temperature and mortality represent the heat vulnerability factors which can be observed at the individual or at the community level. At the scale of the individual, many studies have reported increased vulnerability of elderly individuals to hot temperatures [6-14]. The effects of gender on temperature impacts have seen mixed evidence [12,13]. At the community level, vulnerability of populations has been shown to be influenced by factors such as low density of green spaces [15], poor urban design and planning [16], urban

* Correspondence: benmarhnia.tarik@ouranos.ca

¹EHESP School of Public Health, Sorbonne-Paris Cité, Rennes, France

²Université de Montréal, DSEST, Montréal, QC, Canada

Full list of author information is available at the end of the article

heat islands [17], and the modification effect by air pollutants [18,19]. Ozone in particular has been studied as an effect modifier in the mortality-temperature relationship [20,21], while the link between temperature and particulate matter (PM) [22], sulfur oxides (SOx) [23] or Nitrogen oxides (NOx) [24] has been less explored. Furthermore, urban air pollution has never been studied from a chronic exposure perspective but only using daily levels. Some studies have also focused on community-based social characteristics that may influence mortality and morbidity outcomes through effects of community organization [25], social cohesion or social networks [26], and social deprivation using an aggregated index [27].

The interaction between ambient temperature and air pollution has been investigated in air pollution mortality time series studies and, to a lesser extent, in temperature mortality time series analyses [28]. To our knowledge, no data are available on the relationship between temperature and mortality effect modified by chronic air pollution exposure and socio-economic factors (or ecological social deprivation) assessed jointly.

The objective of the study is to identify whether and how the magnitude of the effects of mean temperature on all-cause mortality were modified by chronic air pollution exposure (here nitrogen dioxide (NO₂) representing urban traffic), social deprivation, and a combination of these two factors. For this purpose, we studied the city of Paris (France) where NO₂ long-term average concentrations vary substantially across the city according to traffic density (i.e. the main source of NO₂ emissions), and where the different city neighbourhoods host populations with contrasted socio-economic profiles. We hypothesized that a better understanding of these vulnerability factors should provide relevant information for developing public health programs targeting the most vulnerable populations and territories.

Methods

Study setting and small-area level

Paris, the capital city of France, has a population of roughly 2.25 million inhabitants. The spatial scale used was the French census block (i.e. IRIS - a French acronym for "blocks for incorporating statistical information"), which constitute the smallest census unit areas in France, designed by the National Census Bureau (INSEE - Institut national de la statistique et des études économiques), for which aggregate data is available. The city of Paris is subdivided into 992 census blocks with a mean population of 2,199 inhabitants (range = 0-5,456 inhabitants) and a mean area of 0.11 km² (range = 0.009-5.4 km²).

Mortality and population data

We considered all deaths occurred in the city of Paris (excluding woods of Boulogne and Vincennes) for residents

older than 35 years old from May to August for the years 2004–2009 included. All-cause mortality data were provided by the death registry of the city of Paris. Individual information on age, sex, date of death, and census block of residence was available for each case of death. For confidentiality issues it was not possible to distinguish causes of mortality, thus external causes of deaths could not be excluded. The analysis included only subjects older than 35 years old at the time of death to minimize this bias because accidental causes of death are dominant in subjects under 35 years old [29]. We obtained the number of population in each stratum from the INSEE.

Ethical approval was obtained from the French commission on data privacy and public liberties (CNIL - Commission Nationale Informatique et Liberté).

Climate data

Daily mean outdoor temperatures and relative humidity were obtained from Météo-France, as measured at the Montsouris station in Paris and were computed using data of the corresponding period (summers from 2004 to 2009).

Air pollution data

Annual NO₂ concentrations were modeled at a grid scale of 25x25m throughout the period 2004–2009 (only summers) by the local association for the monitoring and the study of air quality (AirParif). We used a dispersion model (ESMERALDA) to produce annual NO₂ concentrations at a fine spatial resolution. A description of the methods used to produce NO₂ levels at the French census block is provided in Additional file 1. Chronic NO₂ exposure at the census block scale was defined as the average of annual NO₂ concentrations from 2004 to 2009. We did not include other air pollutants such as ozone due to the lack of data at this spatial scale.

Community based socio-economic characteristics: social deprivation

To characterize the socio-economic characteristics at a community level, we computed an aggregated, multidimensional deprivation index, fitted at the census block scale [30]. We provided a summary description of the deprivation index and its categorization in Additional file 1.

Statistical analyses

Social deprivation was defined and stratified according to the 3-classes deprivation index described above. NO₂ chronic exposure was categorized into three groups according to the terciles of its distribution. To create a double stratification with sufficient statistical power, we stratified the chronic NO₂ exposure into two strata according to the median of NO₂ chronic concentrations (i.e.

we obtained a total of 6 strata for this double stratification). We also assessed age-related vulnerability (i.e. number of daily deaths for the ≥ 65 years people compared to the number of daily deaths for the <65 years) and sex-related vulnerability (number of daily deaths for men versus the number of daily deaths women).

First, in a crude model, we developed stratified Poisson regression models to assess daily temperature-related mortality associations with daily mean temperatures and daily death counts [31]. We used cubic B-splines (5 knots) of time to control for secular trends in the mortality series [9]. Seasonal patterns of mortality were controlled by including a quadratic function represented by the day of the season (1 to 123) [32]. Sensitivity analyses were performed by using a cubic polynomial for seasonality. We used natural cubic splines (2 knots) to consider the non-linearity of the temperature-mortality relationship. Daily levels of humidity were incorporated into regression models as possible confounding variables based on evidence in the literature [1,33,34]. Then, we developed stratified models for each category [35]. The threshold for statistical significance was set at $p \leq 0.05$ and all the tests were two-sided. Model assumptions and validity were verified graphically (i.e. with quintile-quintile and partial autocorrelation function plots). A white noise test [32] was also used to ensure that no auto-correlation remained in the residuals. Estimates of exposure (daily temperature) – response (mortality) functions - were expressed as Relative Risks (RR) and their confidence intervals (CI) at 95% for each temperature degree. Relative Risks for the relation between mortality and temperature were estimated relative to the daily mean number of deaths for the entire period [34]. We assessed the heterogeneity of temperature–mortality associations across different strata using the method developed by Payton et al. [36] to compare the regressions' coefficients between different strata. This test was conducted on all regression coefficients across the strata.

Calculation of deaths attributable to temperature

We calculated deaths attributable to ambient temperatures for each stratum. Attributable deaths were calculated

from attributable fractions (AF), using heat-related mortality relationships (RR) described in the previous section [37,38]. We considered only days with daily mean temperature related to a RR strictly greater than 1 for any temperature value (corresponding to 280 days in total). AF were calculated for each temperature degree (T_i : represents a temperature unit) and using the equation $[(RR(T_i)-1)/RR(T_i)]$.

The Attributable Number of deaths (AN) for the period 2004–2009 (months of May, June, July, and August), for a given temperature variable, was then estimated from the following equation:

$$AN = \sum_{i=\min}^{\max} AF(T_i) \times MDC \times ND(T_i)$$

In this equation, i is the temperature degree Celsius, $AF(T_i)$ is attributable fraction for temperature degree (T_i), MDC is the mean observed daily death count, and $ND(T_i)$ is the number of days for which temperature was i (°C). We then divided the total number of attributable deaths for the period 2004–2009 by 6 to obtain an average attributable number of deaths by summer. We used the RR estimated within the strata of interest to estimate the AR in those strata. We calculated confidence intervals at 95% of attributable number of deaths. Then we compared the number of deaths attributable to temperature in each of the groups stratified by chronic air pollution exposure (3strata), social deprivation (3 strata) and the two simultaneously (6 strata). We made sure that there was no homogeneity in the temperature–mortality associations across different strata, as described in the last section. Finally we calculated attributable mortality rates, by dividing the total number of deaths attributable to temperature in each stratum (by summer) by the total population in each of these strata. The final results were mapped to visualize heat vulnerability according to social deprivation and chronic air pollution exposure.

Results

46,056 deaths were registered during the study period, and the mean age at death was 68 years (SD = 10.21).

Table 1 Summary statistics for health outcomes, climate and air pollutants data (Paris, 2004–2009)

Variable	Mean	Minimum	25th Percentile	Median	75th Percentile	Maximum	Standard deviation
Daily death count (n)	33.3	3	27	33	42	93	4.5
Minimum Temperature (°C)*	12.5	0	9	13	16	21	4.2
Mean Temperature (°C)*	17.3	7	14	17	20	26	4.4
Maximum Temperature (°C)*	20.9	8	17	21	24	35	5.2
Relative Humidity (%)*	68.4	37	61	69	76	94	10.7
Daily NO2 (µg/m ³)	48.9	16	38	47	58	121	14.3
Chronic NO2 (µg/m ³)	52.5	38.7	47.88	51.8	56.1	81.1	6.8

*Summary statistics for weather variables (minimum, mean and maximum temperatures as well as relative humidity) are based on daily data.

As presented in Table 1, daily death count ranged from 3 to 93 (mean, 33.3, IQR: 27 to 42). Mean temperatures ranged from 7°C to 26°C (mean: 17.3°C, IQR: 14°C to 20°C), while atmospheric pollution assessed by daily NO₂ ranged from 16 to 121 µg/m³ (mean: 48.9 µg/m³, IQR: 38 µg/m³ to 58 µg/m³) (Table 1). Chronic NO₂ exposure ranged from 39 µg/m³ to 81 µg/m³ (mean: 52.5 µg/m³, IQR: 48 µg/m³ to 56 µg/m³). From the 992 census blocks in our study, we had 54 census blocks with missing data concerning social deprivation which corresponded to 160 deaths. These census blocks are non-residential (activity and miscellaneous) and with few residents. Descriptive statistics of the social deprivation index and chronic air pollution are presented in (Additional file 1: Table S1). As shown in Additional file 1: Figure S1, chronic NO₂ concentrations were slightly higher in the most favoured census blocks. In the crude model (for the whole population), mortality was significantly associated with daily mean temperature (RR for each temperature unit and for all strata are shown in Additional file 1: Table S2 and Table S3). The association between temperature and mortality was U-shaped. Applying the central point estimate of the temperature-mortality relationships to the observed mean temperatures, an average number of 121 attributable deaths by summer (May to August) was estimated. Deaths attributable to mean temperature are presented with their 95% confidence intervals in Additional file 1: Table S4.

As presented in the Figure 1, we observed a gradient in the raw deaths attributable to temperature according to social deprivation: the higher the degree of social deprivation, the higher the number of deaths attributable to mean temperature. Likewise, higher chronic air pollution exposure as assessed by NO₂ levels was also identified as a vulnerability factor in the relationship between temperature and mortality. We observed a significant difference in regression coefficients (between strata)




Figure 1 Number of deaths attributable to mean temperature by social deprivation (3 strata) and chronic air pollution exposure (3 strata).

separately according to social deprivation and chronic air pollution ($p < 0.05$).

Finally, we calculated mortality rates (Table 2), including the population living in each strata to consider the differences in population density across strata. We found some heterogeneity ($p < 0.10$) between ambient temperature

Table 2 Mortality rates attributable to summer temperatures (per 100 000) by strata

Strata	Population	Mortality rates (per 100 000) ^a	p for heterogeneity ^b
Total	2234105	5.37 [5.01;5.73]	
Age			
Under 65 years	1921330	0.78 [0.62;0.88]	0.001
More than 65 years	312774	33.57 [31.65;35. 81]	
Sex			
Female	1161734	5.51 [4.99;5.85]	0.51
Male	1072370	5.22 [4.76;5.67]	
Social Deprivation			
Low Social Deprivation	692572	4.33 [3.31;5.23]	0.08
Medium Social Deprivation	781936	4.60 [3.59;5.41]	
High Social Deprivation	743956	7.26 [6.74;7.85]	
Chronic air pollution exposure			
Low Chronic NO ₂ Exposure ≤ 50.6 µg/m ³	737254	4.75 [4.13;5.22]	0.03
Medium Chronic NO ₂ Exposure 50.6-55.8 µg/m ³	795341	5.97 [5.36;6.32]	
High Chronic NO ₂ Exposure > 55.8 µg/m ³	670231	7.89 [7.28;8.14]	
Double stratification Low Chronic Exposure Group			
Low Social Deprivation	370567	3.78 [2.87;5.03]	0.14
Medium Social Deprivation	381190	4.19 [3.34;5.25]	
High Social Deprivation	375341	6.92 [5.11;8.12]	
Double stratification High Chronic Exposure Group			
Low Social Deprivation	389872	3.59 [2.29;5.07]	0.07
Medium Social Deprivation	391432	5.36 [4.22;6.41]	
High Social Deprivation	325701	9.82 [7.79;10.93]	

^aFor social deprivation and chronic air pollution strata, mortality rates were standardized for age and sex.

^bHomogeneity test of Chi-square Pearson, for central estimates.

and mortality according to age, social deprivation and chronic air pollution exposure. Sex was not an effect modifier in the relationship between heat and temperature ($p = 0.51$). For the double stratification, we found that, in the low chronic exposure group, social deprivation did not significantly modify the relation between heat and mortality ($p = 0.14$), while we found that social deprivation may have modified this relation in the low chronic exposure group ($p = 0.07$). We also conducted heterogeneity tests across the two chronic exposure groups. We found a heterogenic effect (data not shown) only between the strata of low chronic exposure/high social deprivation and the strata of high chronic exposure/high social deprivation.

Figure 2 presents the spatial distribution of social deprivation and chronic air pollution exposure. The spatial variations of both social characteristics are related to deaths attributable to temperature as presented in Table 2.

Discussion

In this study, we assessed the effects of mean temperature and chronic air pollution (with NO₂ representing urban traffic) on all-cause mortality with additional attention paid to the interactive effects of social deprivation. To do this we conducted stratified time-series analyses. We found that chronic air pollution exposure modifies the association between daily temperature and mortality. We also found that social deprivation is a heat-related vulnerability factor. Finally we found that there is a potential combined modification effect of social deprivation and chronic exposure to NO₂ with regards to heat-related mortality.

Our results for the temperature-mortality relationship are comparable to other studies conducted in Paris [8,32]. Concerning social vulnerability to heat-related mortality, our results are consistent with other studies, using time series analyses [5] or using different methods, such as a case crossover design [11].




Figure 2 Spatial distribution of social deprivation and chronic air pollution.

Ecological social vulnerability can be explained by an accumulation of different types of vulnerabilities for individuals within a census block (e.g., social isolation [39], material conditions [40], poor urban design including micro heat islands [17] or poorer health [14]). In the same way, social mobility may also partly explain the observed social gradient, similar to previous studies looking at socio-economic scales [41]. Thus, populations with low socio-economic levels will be inclined to stay in their neighbourhoods, with a low daily mobility, accumulating other heat vulnerability factors and with other consequences on certain determinants of health such as physical activity.

We found that air pollution modifies the relation between ambient temperature and daily mortality as found in previous studies using daily exposure [21-23,28,42]. It is possible that chronic NO₂ levels modify the effects of temperature on mortality. A range of studies have shown that NO₂ is consistently associated with many health outcomes such as cardiovascular effects [43,44], which can explain how the NO₂ chronic effects can make some populations more vulnerable to heat impacts.

Our results suggest that ambient temperature has a greater impact on daily mortality on chronically polluted areas, especially if these areas are socially deprived. However our results only suggest this double effect modifier and further studies may confirm this result.

Methods for estimating the heat related mortality relationships inevitably rely on some assumptions. In our study, we chose to not take into account lag effects to simplify the interpretation of deaths attributable to temperature. We considered that contribution of lag effects compared to the day of death was not differential according to the different strata characteristics. We did not take into account harvesting effect or mortality displacement. We also did not consider intra urban variations of temperatures. Future studies should consider these aspects. We considered the social deprivation at the ecological level, and we did not explore social inequalities in mortality at the individual level, because of the lack of data. The use of a synthetic index rather than using independent ecological socio-economic measures allows us to represent an accumulation of social and material disadvantages [45] and these kinds of indexes are more useful to lead population interventions [27]. The division of neighbourhoods into census blocks aims to maximize their homogeneity in terms of population size, socioeconomic characteristics, land use and zoning. In Paris, the area of a block census is quite small so the ecological bias is likely to be small. Finally, we did not consider the spatial auto-correlation in our analysis which is a result of the correlation between the spatial unit and the adjacent geographical areas. Variables observed at small area level are interdependent because proximity and

linkages between neighbouring areas. This issue could influence the effect of social deprivation and chronic air pollution [46], and further studies could consider it by conducting spatial analyses such as a hierarchical Bayesian modeling approach.

Conclusion

In this study, we showed that areas which are chronically polluted by air pollution can be characterized as vulnerable to heat in the sense that they have more deaths attributable to heat than less exposed areas. We also presented a potential combined vulnerability of social deprivation and chronic exposure to air pollution in the mortality-temperature relationship. Our results may have important implications considering chronically polluted areas as vulnerable in heat actions plans (especially including adapted surveillance and warning systems) and in the long-term measures to reduce the burden of heat stress (as building regulations, urban planning or land-use changes), especially in the context of climate change. However, further studies are necessary to determine whether similar results could be found in other settings.

Ethics approval

French commission on data privacy and public liberties (CNIL - Commission Nationale Informatique et Liberté).

Additional file

Additional file 1: Supplemental Material.

Abbreviations

AF: Attributable fraction; CI: Confidence interval; INSEE: Institut National de la Statistique et des Etudes Economiques (National Census Bureau); IQR: Inter quartile range; NO: Nitrogen oxides; PM: Particle matter; RR: Relative risk; SD: Standard deviation; SES: Socio economic status; SO: Sulfur oxides.

Competing interests

The authors declare that they have no competing interests.

Authors' contributions

TB, YO and SD conceived the study. TB and CP analyzed the data in consultation with SD and DZN. TB, CP and YO wrote the draft version and revisions of the manuscript. SD, DZN, AL and PC contributed to the writing of the manuscript. All authors agree with manuscript results and conclusions. All authors read and approved the final version of the manuscript.

Acknowledgments

The authors would like to acknowledge AirParif for providing air pollution data, Météo France for providing the temperature data, and Allan Brand from the INSPQ (Montréal, Canada) for his help in data management and English editing.

Funding

This work and the Equit'Area project are supported by the French National Research Agency (ANR, contract-2010-PRSP-002-01) and the EHESP School of Public Health.

Author details

¹EHESP School of Public Health, Sorbonne-Paris Cité, Rennes, France.

²Université de Montréal, DSEST, Montréal, QC, Canada. ³INSERM U1085

(IRSET), Rennes, France. ⁴INSERM UMRS707, DS3, Paris, France. ⁵UPMC Univ Paris 06, UMRS 707, Paris, France. ⁶Lorraine University Medical School, Vandœuvre-les, Nancy, France.

Received: 24 December 2013 Accepted: 5 June 2014
Published: 18 June 2014

References

- Basu R, Samet JM: Relation between elevated ambient temperature and mortality: a review of the epidemiologic evidence. *Epidemiol Rev* 2002, **24**:190–202.
- Armstrong B, Hajat S, Kovats S, Lloyd S, Scovronick N, Wilkinson P: Commentary: climate change: how can epidemiology best inform policy? *Epidemiology* 2012, **23**(6):780–784.
- McMichael AJ: Globalization, climate change, and human health. *N Engl J Med* 2013, **368**:1335–43.
- Basu R: High ambient temperature and mortality: a review of epidemiologic studies from 2001 to 2008. *Environ Health* 2009, **8**:40.
- Yu W, Vaneckova P, Mengersen K, Pan X, Tong S: Is the association between temperature and mortality modified by age, gender and socio-economic status? *Sci Total Environ* 2010, **408**(17):3513–3518.
- Anderson BG, Bell ML: Weather-related mortality: how heat, cold, and heat waves affect mortality in the United States. *Epidemiology* 2009, **20**:205.
- Astrom DO, Forsberg B, Rocklov J: Heat wave impact on morbidity and mortality in the elderly population: a review of recent studies. *Maturitas* 2011, **69**:99–105.
- Baccini M, Biggeri A, Accetta G, Kosatsky T, Katsouyanni K, Analitis A, Anderson HR, Bisanti L, D'Ippoliti D, Danova J: Heat effects on mortality in 15 European cities. *Epidemiology* 2008, **19**(5):711–719.
- Hajat S, Kovats RS, Lachowycz K: Heat-related and cold-related deaths in England and Wales: who is at risk? *Occup Environ Med* 2007, **64**:93–100.
- Ishigami A, Hajat S, Kovats RS, Bisanti L, Rognoni M, Russo A, Paldy A: An ecological time-series study of heat-related mortality in three European cities. *Environ Health* 2008, **7**(5):1–7.
- Stafoggia M, Forastiere F, Agostini D, Biggeri A, Bisanti L, Cadum E, Caranci N, de' donato F, De Lisio S, De Maria M: Vulnerability to heat-related mortality: a multicity, population-based, case-crossover analysis. *Epidemiology* 2006, **17**(3):315–323.
- Kysely J, Kim J: Mortality during heat waves in South Korea, 1991 to 2005: How exceptional was the 1994 heat wave? *Climate Res* 2009, **38**:105.
- Robine J-M, Michel J-P, Herrmann F: Excess male mortality and age-specific mortality trajectories under different mortality conditions: a lesson from the heat wave of summer 2003. *Mech Ageing Dev* 2012, **133**:378–83.
- Bell ML, O'Neill MS, Ranjit N, Borja-Aburto VH, Cifuentes LA, Gouveia NC: Vulnerability to heat-related mortality in Latin America: a case-crossover study in Sao Paulo, Brazil, Santiago, Chile and Mexico City, Mexico. *Int J Epidemiol* 2008, **37**(4):796–804.
- Reid CE, O'Neill MS, Gronlund CJ, Brines SJ, Brown DG, Diez-Roux AV, Schwartz J: Mapping community determinants of heat vulnerability. *Environ Health Perspect* 2009, **117**(11):1730.
- Stone B Jr, Rodgers MO: Urban form and thermal efficiency: how the design of cities influences the urban heat island effect. *J Am Plann Assoc* 2001, **67**:186–98.
- Smargiassi A, Goldberg MS, Plante C, Fournier M, Baudouin Y, Kosatsky T: Variation of daily warm season mortality as a function of micro-urban heat islands. *J Epidemiol Community Health* 2009, **63**(8):659–664.
- Ye X, Wolff R, Yu W, Vaneckova P, Pan X, Tong S: Ambient temperature and morbidity: a review of epidemiological evidence. *Environ Health Perspect* 2012, **120**(1):19.
- Analitis A, Michelozzi P, D'Ippoliti D, De' Donato F, Menne B, Matthies F, Atkinson RW, Iniguez C, Basagana X, Schneider A, Lefranc A, Paldy A, Bisanti L, Katsouyanni K: Effects of heat waves on mortality: effect modification and confounding by air pollutants. *Epidemiology* 2014, **25**(1):15–22.
- Filleul L, Cassadou S, Medina S, Fabres P, Lefranc A, Eilstein D, Le Tertre A, Pascal L, Chardon B, Blanchard M: The relation between temperature, ozone, and mortality in nine French cities during the heat wave of 2003. *Environ Health Perspect* 2006, **114**(9):1344.
- Ren C, Williams GM, Morawska L, Mengersen K, Tong S: Ozone modifies associations between temperature and cardiovascular mortality: analysis of the NMMAPS data. *Occup Environ Med* 2008, **65**(4):255–260.
- Ren C, Williams GM, Tong S: Does particulate matter modify the association between temperature and cardiorespiratory diseases? *Environ Health Perspect* 2006, **114**:1690.
- Katsouyanni K, Pantazopoulou A, Touloumi G, Tselepidaki I, Moustris K, Asimakopoulou D, Pouloupoulou G, Trichopoulos D: Evidence for interaction between air pollution and high temperature in the causation of excess mortality. *Arch Environ Health* 1993, **48**(4):235–242.
- Peel JL, Haeuber R, Garcia V, Russell AG, Neas L: Impact of nitrogen and climate change interactions on ambient air pollution and human health. *Biogeochemistry* 2013, **114**(1-3):121–134.
- Romero-Lankao P, Qin H, Dickinson K: Vulnerability to temperature-related hazards: a meta-analysis and meta-knowledge approach. *Glob Environ Chang* 2012, **22**:670–83.
- Klinenberg E: *Heat wave: A social autopsy of disaster in Chicago*. Chicago: University of Chicago Press; 2003.
- Rey G, Jouglé E, Fouillet A, Hémon D: Ecological association between a deprivation index and mortality in France over the period 1997-2001: variations with spatial scale, degree of urbanicity, age, gender and cause of death. *BMC Public Health* 2009, **9**(1):33.
- Roberts S: Interactions between particulate air pollution and temperature in air pollution mortality time series studies. *Environ Res* 2004, **96**:328–37.
- Meslé F, Vallin J: Reconstructing long-term series of causes of death: the case of France. *J Quant Interdiscip Hist* 1996, **29**:72–87.
- Lalloué B, Monnez J-M, Padilla C, Kihal W, Le Meur N, Zmirou-Navier D, Deguen S: A statistical procedure to create a neighborhood socioeconomic index for health inequalities analysis. *Int J Equity Health* 2013, **12**(1):21.
- Armstrong B: Models for the relationship between ambient temperature and daily mortality. *Epidemiology* 2006, **17**:624–31.
- Fouillet A, Rey G, Jouglé E, Frayssinet P, Bessemoulin P, Hémon D: A predictive model relating daily fluctuations in summer temperatures and mortality rates. *BMC Public Health* 2007, **7**(1):114.
- Buckley JP, Samet JM, Richardson DB: Commentary: does air pollution confound studies of temperature? *Epidemiology* 2014, **25**:242–245.
- Yu W, Mengersen K, Hu W, Guo Y, Pan X, Tong S: Assessing the relationship between global warming and mortality: lag effects of temperature fluctuations by age and mortality categories. *Environ Pollut* 2011, **159**(7):1789–1793.
- Chan EYY, Goggins WB, Kim JJ, Griffiths SM: A study of intracity variation of temperature-related mortality and socioeconomic status among the Chinese population in Hong Kong. *J Epidemiol Community Health* 2012, **66**(4):322–327.
- Payton ME, Greenstone MH, Schenker N: Overlapping confidence intervals or standard error intervals: what do they mean in terms of statistical significance? *J Insect Sci* 2003, **3**:34.
- Baccini M, Kosatsky T, Analitis A, Anderson HR, D'Ovidio M, Menne B, Michelozzi P, Biggeri A: Impact of heat on mortality in 15 European cities: attributable deaths under different weather scenarios. *J Epidemiol Community Health* 2011, **65**(1):64–70.
- Hajat S, Armstrong B, Baccini M, Biggeri A, Bisanti L, Russo A, Paldy A, Menne B, Kosatsky T: Impact of high temperatures on mortality: is there an added heat wave effect? *Epidemiology* 2006, **17**(6):632–638.
- Semenza JC, Rubin CH, Falter KH, Selanikio JD, Flanders WD, Howe HL, Wilhelm JL: Heat-related deaths during the July 1995 heat wave in Chicago. *N Engl J Med* 1996, **335**(2):84–90.
- Rogot E, Sorlie PD, Backlund E: Air-conditioning and mortality in hot weather. *Am J Epidemiol* 1992, **136**:106–16.
- Perchoux C, Chaix B, Cummins S, Kestens Y: Conceptualization and measurement of environmental exposure in epidemiology: Accounting for activity space related to daily mobility. *Health Place* 2013, **21**:86–93.
- Buckley JP, Samet JM, Richardson DB: Commentary: does air pollution confound studies of temperature? *Epidemiology* 2014, **25**(2):242–245.
- Latza U, Gerdes S, Baur X: Effects of nitrogen dioxide on human health: systematic review of experimental and epidemiological studies conducted between 2002 and 2006. *Int J Hyg Environ Health* 2009, **212**:271.

44. Hart JE, Garshick E, Dockery DW, Smith TJ, Ryan L, Laden F: **Long-term ambient multipollutant exposures and mortality.** *Am J Respir Crit Care Med* 2011, **183**(1):73.
45. Townsend P: **Deprivation.** *J Soc Policy* 1987, **16**:125–46.
46. Deguen S, Lalloue B, Bard D, Havard S, Arveiler D, Zmirou-Navier D: **A small-area ecologic study of myocardial infarction, neighborhood deprivation, and sex: a Bayesian modeling approach.** *Epidemiology* 2010, **21**(4):459–466.

doi:10.1186/1476-069X-13-53

Cite this article as: Benmarhnia et al.: Chronic air pollution and social deprivation as modifiers of the association between high temperature and daily mortality. *Environmental Health* 2014 **13**:53.

**Submit your next manuscript to BioMed Central
and take full advantage of:**

- Convenient online submission
- Thorough peer review
- No space constraints or color figure charges
- Immediate publication on acceptance
- Inclusion in PubMed, CAS, Scopus and Google Scholar
- Research which is freely available for redistribution

Submit your manuscript at
www.biomedcentral.com/submit

