

# Lawrence Berkeley National Laboratory

## Recent Work

### Title

INFRARED STUDY OF THE CRAB PULSAR: THE ""SHOULDER"" PULSE AND THE 3.45um PULSE PROFILE

### Permalink

<https://escholarship.org/uc/item/8nc2p0dx>

### Authors

Middleditch, J.  
Pennypacker, C.  
Burns, M.S.

### Publication Date

1982-12-01



# Lawrence Berkeley Laboratory

UNIVERSITY OF CALIFORNIA

## Physics, Computer Science & Mathematics Division

RECEIVED  
LAWRENCE  
BERKELEY LABORATORY

FEB 9 1983

LIBRARY AND  
DOCUMENTS SECTION

Submitted to the Astrophysical Journal


INFRARED STUDY OF THE CRAB PULSAR: THE "SHOULDER"  
PULSE AND THE 3.45 $\mu$ m PULSE PROFILE

J. Middleditch, C. Pennypacker, and M.S. Burns

December 1982

### TWO-WEEK LOAN COPY

*This is a Library Circulating Copy  
which may be borrowed for two weeks.  
For a personal retention copy, call  
Tech. Info. Division, Ext. 6782.*



LBL-15362  
c 2

## **DISCLAIMER**

This document was prepared as an account of work sponsored by the United States Government. While this document is believed to contain correct information, neither the United States Government nor any agency thereof, nor the Regents of the University of California, nor any of their employees, makes any warranty, express or implied, or assumes any legal responsibility for the accuracy, completeness, or usefulness of any information, apparatus, product, or process disclosed, or represents that its use would not infringe privately owned rights. Reference herein to any specific commercial product, process, or service by its trade name, trademark, manufacturer, or otherwise, does not necessarily constitute or imply its endorsement, recommendation, or favoring by the United States Government or any agency thereof, or the Regents of the University of California. The views and opinions of authors expressed herein do not necessarily state or reflect those of the United States Government or any agency thereof or the Regents of the University of California.

**Infrared Study of the Crab Pulsar:  
the "Shoulder" Pulse and the 3.45  $\mu\text{m}$  Pulse Profile**

J. Middleditch<sup>1,2,3</sup>, C. Pennypacker<sup>1,4</sup>, M.S. Burns<sup>1,4</sup>

Received 1982

**ABSTRACT**

Infrared measurements of the Crab Pulsar with the NASA IRTF 3.0 m telescope show that the spectrum of the main pulse turns downward for wavelengths longer than 3  $\mu\text{m}$ . The "shoulder" pulse discovered by Pennypacker is measured in the 0.9 to 2.4  $\mu\text{m}$  region, but disappears at 3.5  $\mu\text{m}$ . This pulse rises from 0 to 20% of the height of the main pulse within 1 to 2 ms after the main pulse peak, and decays with a 4 to 5 ms time constant. Excess infrared flux also appears after the interpulse. The main peak itself may be narrower at 3.45  $\mu\text{m}$  than in the optical to 2.2  $\mu\text{m}$  band.

**Subject Headings:** stars: neutron - infrared: spectra

1. Visiting Astronomer at the Infrared Telescope Facility which is operated by the University of Hawaii under contract with the National Aeronautics and Space Administration.
2. Visiting Astronomer at the University of Arizona Observatories.
3. Los Alamos National Laboratory, Los Alamos, New Mexico.
4. Space Sciences Laboratory and Lawrence Berkeley Laboratory, University of California, Berkeley.

## INTRODUCTION

We have measured the Crab Nebula Pulsar (NP 0532) in the near infrared. The pulsar had previously been measured at these wavelengths by Becklin, et al (1973) and the pulse had been shown to be similar to the optical measurements. More sensitive measurements by Pennypacker (1981) confirmed the previous work's flux measurements, but the main peak of the averaged infrared light curve was shown to have excess flux following the peak which is not found in the optical light curve. The "shoulder" pulse has been studied in more detail in this work and we have also measured the 3.5  $\mu\text{m}$  light curve for the first time, where the shoulder pulse is absent. In addition, we measure the spectrum of the main pulse which reveals a turndown at wavelengths longer than 3.5  $\mu\text{m}$ .

## OBSERVATIONS

NP 0532 was observed from the University of Hawaii/NASA IRTF 3.0 m telescope at Mauna Kea from the 18<sup>th</sup> to the 20<sup>th</sup> of December 1981. In addition to the usual J, H, K, L, and M bands, observations were made with a special broadband filter ("UC") (see Table 1) which transmitted between 0.9 and 2.4  $\mu\text{m}$ . A liquid helium-cooled InSb detector and pre-amplifier (RC2) was provided by the IRTF. This detector was optimized for high time resolution photometry yielding a 3 dB decrease in gain at 200 Hz. The detector time response was measured using an infrared LED with a signal generator. In contrast, the InSb detector in normal use (RC1) was measured to have a 3 dB point at 5 Hz.

The internal amplifier gains of the detectors were set to 10 while an external pre-amplifier gain was set between 10 and 50. The signals were then digitized at 0 to 1023 counts for -2.500 to +2.495 volts and the resulting bit patterns were translated into 16-bit words in the PDP 11/45 data acquisition computer. The computer was controlled by an IRTF program which continuously recorded data on magnetic tape at rate of from 1000 to 4000 words  $\text{s}^{-1}$ . Most of the observations were made through a circular aperture

of 7" diameter; a few were made through a 5" aperture.

Optical observations of NP 0532 were made with the University of Arizona/NASA 1.5 m telescope at Mt. Lemmon on the 17<sup>th</sup> January 1982, using a 16" circular aperture. The counts from an unfiltered EMI 9658 A photomultiplier tube were recorded onto magnetic tape at one ms intervals for later analysis in which the optical and infrared data were compared.

## ANALYSIS

The data from observations of NP 0532 were Fourier-transformed in order to establish the apparent frequency of the pulsar. In general, the apparent pulse frequency from the IRTF observations differed from the frequency predicted by an ephemeris generated from the data of Lohsen (1981). This inconsistency was traced to a drift in the clock of the data acquisition system by a few parts in  $10^7$  on the timescale of a few hours. This drift was monitored during the observations and found to be too small to adversely affect the data.

An integrated pulse profile was produced from the optical data by folding the time series at 30.0585424(4) Hz and then convolving the resulting light curve with the measured RC2 infrared detector response. The convolution may be expressed as:

$$(1) \quad G'(t) = (2\pi) 200 \int_0^{\infty} G(t-\tau) e^{-(2\pi)200\tau} d\tau$$

where  $G$  is the raw optical pulse profile and  $G'$  is the pulse profile to be compared with the raw infrared pulse profiles.

The pulse profiles for the infrared data were examined and the light curves obtained while using the J(1.25  $\mu$ m), H(1.65  $\mu$ m), K(2.20  $\mu$ m), and "UC"(0.9 - 2.4  $\mu$ m) filters all appeared to be equivalent to within statistical error. Accordingly, all of the runs (except those in L, M, or "UC" with the slower RC1 detector) were summed together

directly with equal detector voltages receiving equal weights. To sum the different runs, each was fitted to the pulse profile of run 1, so that the offsets in phase of the profiles could be standardized. The results of this analysis are plotted in Figure 1a, together with the optical G' light curve.

In order to correct the infrared light curves for the 5 ms detector response, the pulse profile of each individual run was Fourier-analyzed and the statistical significance of each harmonic was computed using the noise levels derived from the Fourier transforms of the time series. All harmonics, up to and including the 13th at approximately 390 Hz, which were more statistically significant than 2 standard deviations (power/local power greater than 3.0), were scaled according to the formula:

$$(2) \quad a'_n = [(a_n^2 - r)(1 + (nf_0/200)^2)]^{\frac{1}{2}}$$

and the phases of the harmonics were increased according to:

$$(3) \quad \phi'_n = \phi_n + \tan^{-1} (nf_0/200)$$

where  $f_0$  is the frequency of the fundamental (approx. 30.06 Hz) and  $r$  is the local power level at  $nf_0$ . The original harmonics were subtracted from the pulse profiles and the corrected harmonics were re-added. The G' light curve was similarly corrected; harmonics 1 through 13 were large enough that the noise level could be neglected (the power level in the 13th harmonic was 218 times the local level). A composite corrected infrared light curve was generated exactly as done for Figure 1a and is plotted against the best fit of the "corrected" optical light curve in Figure 1b. The ripples present in both curves of Figure 1b are due to the incomplete harmonic reconstruction of the corrected pulse profiles. The curves in Figure 1b are plotted at their corrected position, approximately 0.5 ms earlier than the uncorrected pulse profiles.

Figure 1c shows the difference of the two curves in Figure 1b. The curve rises from zero to its peak of 20% of the main peak height in 1 ms. The decay of this infrared excess is much slower, with an exceptional time constant of 4 to 5 ms. A similar excess is also evident for the interpulse; the excess is possibly slower to decay than the excess associated with the main pulse. Both of these postcursor pulses peak approximately 2 ms following the usual pulse and interpulse peaks. The good agreement of the fits between the optical and infrared pulse profiles for most of the main peak argues against any flaw in the analysis described above. In addition, in tests with an infrared LED, the system faithfully reproduced a pulse similar to the Crab pulsar optical pulse. The harmonic reconstruction technique was also tested by restoring the pulse profile taken on RC1 (run 8), which was significantly degraded by this detector's slow response. We can think of no reason for this difference other than that it represents a real physical difference between the optical and infrared pulse profiles. This flux is barely discernible as a break in the slope of the main pulse in the less statistically significant data of Becklin, et al. (1973; see their Figure 1), but was originally noted by Pennypacker (1981).

More evidence in favor of the reality of the postcursor pulse and against any possible unusual behavior in the detector comes from the pulse profile obtained with the L filter (3.45  $\mu\text{m}$ ) shown in Figure 2. Although this curve is noisier than the JHK-UC composite curve, the fortuitous statistical result of "negative" flux following the main peak makes the existence of a postcursor pulse much less probable. In addition, the main peak of this pulse profile appears to be significantly narrower than the optical main pulse—the optical pulse is too wide and consequently the fit falls short of the top of the infrared peak. In contrast, the fits to the J, H, and K profiles all satisfactorily match the widths of the main pulse until 1 ms past the peak, when the optical profile drops below the infrared profiles. There is also a suggestion that the pulse-interpulse separation may be less than in the optical pulse profile, although this difference may be due to statistical effects. Further study of the L pulse profile of NP 0532 is clearly



warranted.

Because of the difference between the optical and infrared pulse profiles, the infrared fluxes were estimated by using the height of the main pulse, rather than by integrating the pulse area as done by Neugebauer, et al. (1969) and Becklin, et al. (1973). The RC2 detector was calibrated by measurements of Zeta Tau, whose J, H, K, and L fluxes were taken to be 79.8 Jy, 62.4 Jy, 41.3 Jy, and 22 Jy respectively. An M flux of 13.6 Jy was estimated by extrapolating a power law fit from the K-L region. The power law index was in turn determined by an extrapolation of the indices determined by sections of the infrared data. This index fell between 1.5 and 2.0, so that if the flux from Zeta Tau is Rayleigh-Jeans beyond  $3.4 \mu\text{m}$ , this technique will overestimate the M flux.

The infrared data were corrected for interstellar absorption by assuming the net absorption to be  $0.817/\lambda$  ( $\mu\text{m}$ ) magnitudes. This is consistent with Miller's (1973) measurement of 1.6 magnitudes of absorption in the visual and the review of Annestad and Purcell (1973) which generally places the infrared absorption slightly below the extrapolation from the visible bands. The absorptions applied are listed in Table 2. It is assumed that the airmass extinction corrections for the runs on NP 0532 and the calibrations on Zeta Tau will cancel.

One other small systematic correction was applied to the data, the correction for the addition of noise. In effect, because of statistical fluctuations, the actual peak of the pulse profile as determined by rigorous fitting, will be higher than the real peak because of the addition of noise (and the center of the fit will be shifted toward the location of the positive fluctuation). Accordingly, the J, H, and K fluxes were each decreased by approximately 5% depending on the net amount of noise in the composite pulse profiles. Because of the poor match of the L pulse profile to the optical light curve, no correction was made for this band.

The optical data of Oke (1969) was corrected for the interstellar absorption

measured by Miller (1973). For this purpose, the optical pulse profile used for the data analysis was measured to estimate the height of the main pulse, given the integrated flux of the entire profile. If the integrated magnitude is 16.55 as seems likely (Kristian, et al., 1970; Oke, 1969), then the peak of the main pulse is 14.17 magnitudes. This calibration was used to plot the data of Oke together with the infrared data of this work in Figure 3. The upper limit for the M pulse profile was established by folding the data at an estimated value for the apparent frequency of NP 0532.

The flux of 30 - 35 mJy for the JHK bands is consistent with Becklin, et al.'s K filter measurement of  $4.5 (+ 0.4) \times 10^{-31} \text{ Jm}^{-2} \text{ Hz}^{-1} = 45 (+ 4) \text{ Jy} \times \text{s}$  for the time averaged energy in the main pulse. Using 1.75 ms as the FWHM of the main pulse and assuming an approximately triangular waveform, this time-averaged K flux corrects to  $26 (+ 2) \text{ mJy}$  which agrees reasonably with this work. However, the measurement of the L flux of Becklin, et al. relative to their K flux is about 2 sigma higher than our value. This inconsistency may be due to unfortunate statistics in the previously measured K and L fluxes or it may have been caused by the addition of noise to Becklin's data, by the process described above for the systematic correction to our infrared data. It is unlikely that this difference is caused by light loss outside of the 7" circular aperture used in the L observation, since our entire data set was taken through 7" or 5" circular apertures. Moreover, the J flux derived from a run in a 5" aperture (run 9) was at least 85% of the fluxes measured through the 7" aperture (runs 3 and 5). Furthermore, the turnover in the L and M bands cannot be due to the systematics of the detailed analysis procedure described above, which has possibly overestimated these fluxes.

In view of the uncertainties involved in the estimation of the infrared main pulse heights, the errors were set at 15% of the measured values. Estimates for the heights of the postcursors to the main pulse are also listed in Table 2.

## DISCUSSION

No one single spectral model could adequately fit the data shown in Figure 3. Blackbody fits with  $T$  approx.  $9,000^0\text{K}$  fit Oke's optical data but fell far below the infrared points. Similar fits to the M, L, and K points of the infrared data surpass the H and J fluxes before turning over to fall through the optical data. Optically thin synchrotron models (Rybicki and Lightman, 1979) can come close to matching the turnover toward the ultraviolet, but cannot match the fall of the infrared points at the longer wavelengths.

It is possible that the turnover of the spectrum at infrared wavelengths is due to synchrotron self-absorption. The infrared spectrum rises with a spectral index of about  $2/3$  in the infrared. Self-absorption produced in a non-uniform plasma and the non-uniform dipolar magnetic field of the pulsar can yield a spectral index of this value. However, self-absorption might tend to broaden the main pulse (Smith 1980) and this is not observed.

The origin of the postcursor pulse(s) is even more of an enigma. The long timescale decay is suggestive of radiative cooling, while the rapid rise suggests a geometrical change of the radiative direction—both effects might be produced by the neutron star. Finally the apparent change of shape of the L pulse profile merits further investigation.

## ACKNOWLEDGEMENTS

We would like to thank Dr. Rich Capps for modifying the RC2 detector and Dr. Eric Becklin for donating engineering time on the night of 18 December 1981 for our observations. We owe a great deal to Deanne Thompson for writing the data acquisition program, and to the entire staff of the IRTF for their support. Finally, we would like to thank Dr. David Daniels of MP Division of Los Alamos National Laboratory for the RSX 11-M system which enabled us to perform some online analysis of our data. This work was supported by the Director, Office of Energy Research, Office of High Energy and Nuclear Physics, Division of High Energy Physics of the U.S. Department of Energy under Contract No. DE-AC03-76SF00098.

TABLE 1  
 Journal of Observations  
 Infrared, NASA IRTF 3.0 m

UT Run	Date (1981)	Filter	Start (UT)	Sampling (Hz)	Duration (words)	Aperture (mm)	Detector
1	Dec 18	UC	1203	4000	2640000	4.0	RC2
2	"	K	1258	"	2400000	4.0	"
3	"	J	1320	"	2400000	"	"
4	Dec 19	UC	0902	"	1052000	"	"
5	"	J	0917	"	2400000	"	"
6	"	L	0934	2000	6800000	"	"
7	"	K	1038	4000	2400000	"	"
8	"	UC	1138	1000	528000	3.0	RC1
9	Dec 20	J	0812	2000	1052000	"	RC2
10	"	H	0833	2000	1052000	"	"
11	"	M	0903	1250	8372000	"	"

Optical -- University of Arizona/NASA 1.5 m

A	Jan 17(82) --	0434	1000	1060000	16"	EMI 9658 A
---	---------------	------	------	---------	-----	------------

TABLE 2  
Calibrations

<u>Band</u>	<u>Tau</u> <u>(Volts)</u>	<u>Magnitudes</u>	<u>Jy</u>	<u>0 Mag.</u> <u>(Jy)</u>
J	0.33	3.20	79.8	1520
H	0.235	2.99	62.4	980
K	0.195	2.94	41.3	620
L	0.050	2.76	22.0	280
M	0.005	2.62*	13.6*	153
UC	0.860			

Measurements

<u>Band</u>	<u>V</u>		<u>J</u>		<u>H</u>		<u>K</u>		<u>L</u>		<u>M</u>	
	<u>Peak</u>	<u>Mean</u>	<u>Peak</u>	<u>Mean</u>	<u>Peak</u>	<u>Mean</u>	<u>Peak</u>	<u>Mean</u>	<u>Peak</u>	<u>Mean</u>	<u>Peak</u>	<u>Mean</u>
Raw. Mag.	14.17	16.55	12.17	14.13	11.63	13.64	11.08	13.03	10.69	13.26	>10.85	>12.51
Interstellar												
Absorption	1.6	1.6	0.65	0.65	0.50	0.50	0.37	0.37	0.24	0.24	0.17	0.17
Corrected	12.57	14.95	11.52	13.48	11.13	13.14	10.71	12.66	10.45	13.02	>10.7	>12.4
Fv (mJy)	33.9	3.8	37.5	6.16	34.6	5.43	32.24	5.35	18.5	1.73	<8.0	<1.7
Fv (corr. for noise)			35.5		32.2		30.0		18.5			
Postcursor	<2.7		8.5		9.1		7.5		<1.0			
( $\mu$ m)	0.5435		1.25		1.65		2.20		3.45		4.80	

\*Values extrapolated from the shorter wavelengths.

FIGURE CAPTIONS

1. The 0.9 - 2.4  $\mu\text{m}$  infrared pulse profile of NP0532 (histogram) is plotted against the optical pulse profile (curve). (a) Shows the raw infrared data and the optical convolved with the  $4\text{ms}/2\pi$  infrared detector response. (b) Shows the two profiles after correction for this detector response through the 13th harmonic ( $\sim 390\text{Hz}$ ). The ripples present in both curves are due to the incomplete harmonic reconstruction of the corrected pulse profiles. (c) Shows the difference between the infrared and optical pulse profiles--the "postcursor" or "shoulder" pulse of Pennypacker (1981).

2. The pulse profile of NP0532 in the L filter (3.45  $\mu\text{m}$ ) is plotted against the best (but not very good) fit of the optical profile.

3. The infrared main pulse height of NP0532 (squares with error bars) of this work are plotted with the optical measurements of Oke (1969) (diamonds). No one simple model can fit this data (see the Discussion).




Figure 1






Figure 2

CRAB PULSAR MAIN PULSE INFRARED/OPTICAL SPECTRUM  
DKE'S + IRTF 3.0 M DATA ON 1B-22DEC81 WITH IN SB




Figure 3

## REFERENCES

- Aannestad, P. A. and Purcell, E. M. 1973, *Ann. Rev. As. and Ap.*, 11, 309.
- Becklin, E. E., Kristian, J., Matthews, K., and Neugebauer, G. 1973, *Ap. J. (Letters)*, 186, L137.
- Kristian, J., Visvanathan, N., Westphal, J. A. and Snellen, G. H. 1970, *Ap. J.*, 162, 475.
- Lohsen, E. H. G. 1981, *Astron. and Ap. Supp.*, 44, 1.
- Miller, J. S. 1973, *Ap. J. (Letters)*, 180, L83.
- Neugebauer, G., Becklin, E. E., Kristian, J., Leighton, R. B., Snellen, G. and Westphal, J. A. 1969, *Ap. J. (Letters)*, 156, L115.
- Oke, J. B. 1969, *Ap. J. (Letters)*, 156, L49.
- Pennypacker, C. R. 1981, *Ap. J.*, 244, 286.
- Rybicki, G. B. and Lightman, A. P. 1979, *Radiative Processes in Astrophysics* (N. Y.: Wiley).
- Smith, F. G. 1980, *Proc. I. A. U. Symp.*, 95, 221.

This report was done with support from the Department of Energy. Any conclusions or opinions expressed in this report represent solely those of the author(s) and not necessarily those of The Regents of the University of California, the Lawrence Berkeley Laboratory or the Department of Energy.

Reference to a company or product name does not imply approval or recommendation of the product by the University of California or the U.S. Department of Energy to the exclusion of others that may be suitable.

TECHNICAL INFORMATION DEPARTMENT  
LAWRENCE BERKELEY LABORATORY  
UNIVERSITY OF CALIFORNIA  
BERKELEY, CALIFORNIA 94720