

UC Santa Cruz

UC Santa Cruz Previously Published Works

Title

Paths and timings of the peopling of Polynesia inferred from genomic networks.

Permalink

<https://escholarship.org/uc/item/8r28n2r7>

Journal

Nature: New biology, 597(7877)

Authors

Ioannidis, Alexander
Blanco-Portillo, Javier
Sandoval, Karla
[et al.](#)

Publication Date

2021-09-01

DOI

10.1038/s41586-021-03902-8

Peer reviewed



HHS Public Access

Author manuscript

Nature. Author manuscript; available in PMC 2022 November 30.

Published in final edited form as:

Nature. 2021 September ; 597(7877): 522–526. doi:10.1038/s41586-021-03902-8.

Paths and timings of the peopling of Polynesia inferred from genomic networks

Alexander G. Ioannidis^{1,2,20,*}, Javier Blanco-Portillo^{2,20}, Karla Sandoval², Erika Hagelberg³, Carmina Barberena-Jonas², Adrian V. S. Hill^{4,5}, Juan Esteban Rodriguez-Rodriguez², Keolu Fox⁶, Kathryn Robson⁷, Sonia Haoa-Cardinali⁸, Consuelo D. Quinto-Cortés², Juan Francisco Miquel-Poblete⁹, Kathryn Auckland⁴, Tom Parks⁴, Abdul Salam M. Sofro¹⁰, María C. Avila-Arcos¹¹, Alexandra Sockell¹², Julian R. Homburger¹², Celeste Eng¹³, Scott Huntsman¹³, Esteban G. Burchard¹³, Christopher R. Gignoux¹⁴, Ricardo A. Verdugo^{15,16}, Mauricio Moraga^{15,17}, Carlos D. Bustamante^{12,18}, Alexander J. Mentzer^{4,19}, Andrés Moreno-Estrada^{2,*}

¹Institute for Computational and Mathematical Engineering, Stanford University, Stanford, CA 94305, USA

²National Laboratory of Genomics for Biodiversity (LANGE BIO) - UGA, CINVESTAV, Irapuato, Guanajuato 36821, Mexico

³Department of Biosciences, University of Oslo, PO Box 1066, 0316 Oslo, Norway

⁴Wellcome Centre for Human Genetics, University of Oxford, Roosevelt Drive, Oxford OX3 7BN, UK

⁵The Jenner Institute, Nuffield Department of Medicine, University of Oxford, Oxford, OX3 7DQ, UK.

⁶Department of Anthropology, University of California San Diego, 9500 Gilman Drive, La Jolla, CA 92093, USA

⁷MRC Weatherall Institute of Molecular Medicine, University of Oxford, Oxford OX3 9DS, UK.

⁸Mata Ki Te Rangi Foundation, Hanga Roa, Easter Island, Chile.

⁹Departamento de Gastroenterología, Facultad de Medicina, Pontificia Universidad Católica de Chile, Santiago, Chile

*Correspondence and requests for materials should be addressed to A.G.I and A.M.-E. ioannidis@stanford.edu; andres.moreno@cinvestav.mx.

²⁰These authors contributed equally to this work.

Author contributions A.M.-E. and A.G.I. conceived the study. A.G.I., A.J.M., C.D.B., and A.M.-E. provided overall project supervision and management. A.M.-E., A.G.I., E.H., K.S., A.J.M., and C.R.G. contributed to study design. A.S., C.E., S.H., E.B., C.D.B., and A.M.-E. carried out genotyping experiments and quality control. A.G.I., J.B.-P., C. B.-J., J.E.R.-R., C.D.Q.-C., J.R.H., and A.M.-E. analyzed the data. A.G.I. developed the analytical methods. A.G.I., J.B.-P., and A.M.-E. interpreted the results. A.M.-E., K.S., E.H., A.V.S.H., J.F.M.-P., K.A., T.P., K.R., M.C.A.-A., A.S., A.S.M.S., C.E., S.H., E.B., R.A.V., M.M., A.J.M., and C.D.B. contributed to acquisition of the data. A.G.I. wrote the manuscript, and A.M.-E., E.H., K.F., and S.H.-C. provided feedback on the manuscript.

Competing interests C.D.B. is a member of the scientific advisory boards for Liberty Biosecurity, Personalis, 23andMe Roots into the Future, [Ancestry.com](https://ancestry.com), IdentifyGenomics, Genomelink, and Etalon and is a founder of CDB Consulting. C.R.G. owns stock in 23andMe and is member of the scientific advisory board for Encompass Bioscience.

Supplementary information is available for this paper.

Code availability All new techniques described in Methods are available from <https://github.com/ai-sandbox> and all existing software packages and versions used are noted in Methods.

¹⁰Department of Biochemistry, Faculty of Medicine, Yayasan Rumah Sakit Islam (YARSI) University, Cempaka Putih, Jakarta, Indonesia.

¹¹International Laboratory for Human Genome Research (LIIGH), UNAM Juriquilla, Queretaro 76230, Mexico

¹²Center for Computational, Evolutionary and Human Genomics, Stanford University, Stanford, CA 94305, USA

¹³Program in Pharmaceutical Sciences and Pharmacogenomics, Department of Medicine, University of California San Francisco, San Francisco, CA, USA

¹⁴Division of Biomedical Informatics and Personalized Medicine, University of Colorado, Denver, CO, USA

¹⁵Human Genetics Program, Institute of Biomedical Sciences, Faculty of Medicine, University of Chile, Santiago 8380453, Chile

¹⁶Translational Oncology Department, Faculty of Medicine, University of Chile, Santiago 8380453, Chile

¹⁷Department of Anthropology, Faculty of Social Sciences, University of Chile, Santiago 8380453, Chile

¹⁸Department of Biomedical Data Science, Stanford University, Stanford, CA 94305, USA

¹⁹Big Data Institute, Li Ka Shing Centre for Health Information and Discovery, University of Oxford, Oxford, UK

Abstract

Polynesia was settled in a series of extraordinary voyages across an ocean spanning one third of the planet¹, but the sequences of islands settled remain unknown and their timings disputed. Currently, several centuries separate the dates suggested by different archaeological surveys²⁻⁴. Here, using genome-wide data from merely 430 contemporary individuals from 21 key Pacific island populations and novel ancestry-specific computational analyses, we unravel the detailed genetic history of this vast, dispersed island network. Our reconstruction of the branching Polynesian migration sequence reveals a serial founder expansion, characterized by directional loss of variants, that originated in Samoa and spread first through the Cook Islands (Rarotonga), then to the Society (Taiteti) Islands (11th century), the western Austral (Tuha'a Pae) Islands and Tu motu Archipelago (12th century), and finally to the widely separated, but genetically connected, megalithic statue-building cultures of the Marquesas (Te Henua 'Enana) Islands in the north, Raivavae in the south, and Easter Island (Rapa Nui), the easternmost of the Polynesian islands, settled in approximately 1200 CE via Mangareva.

Main

The history of the human settlement of Polynesia has long been examined by its residents⁵, and has been an open question worldwide, since at least the time of Captain James Cook^{6,7}. More recently, the prevalence of certain health conditions in these island founder populations has attracted the interest of medical geneticists⁸. However, although essential for

both medical research and historical understanding, little is known about the human genetic structure of this oceanic expanse, our planet's last habitable region to be settled.

Background

The history of the human settlement of Polynesia has long been examined by its residents⁵, and has been an open question worldwide, since at least the time of Captain James Cook^{6,7}. More recently, the prevalence of certain health conditions in these island founder populations has attracted the interest of medical geneticists⁸. However, although essential for both medical research and historical understanding, little is known about the human genetic structure of this oceanic expanse, our planet's last habitable region to be settled.

The settlement sequence of Polynesian islands remains particularly difficult to unravel using comparative linguistic or cultural approaches due to the rapidity of the initial expansion and the subsequent cultural exchanges between islands^{7,9–11}. Meanwhile, the archaeological estimates for settlement dates remain debated, and have recently been revised forward across eastern Polynesia by up to a millennium^{2–4,12,13}. Previous region-wide Polynesian genetics studies have considered only globin gene polymorphisms¹⁴, or have been restricted to near (western) Polynesia¹⁵ and the Society Islands¹⁶ and lacked an ancestry-specific approach. Meanwhile, ancient DNA studies have sequenced only four samples from one island in western Polynesia and three near-modern samples from one island in eastern Polynesia, all with low genotype density, still lower between-sample genotype overlap, and different time frames^{17,18}. Here we use a dataset of modern samples that is two orders of magnitude larger to examine detailed intra- and inter-island population substructure across all of Polynesia (Supplementary Tables 1–2). We leverage our sample size to perform directionality and network analyses, and leverage our high-density, overlapping genotypes from coexistent individuals to perform within-generation autosomal haplotype matching, allowing us to date and reconstruct the settlement paths of these islands for the first time. This study also allows us to demonstrate new ancestry-specific techniques for analyzing genomic data from underrepresented, admixed populations.

The Polynesians are predominantly descended from Austronesian-speaking voyagers¹⁷, who trace their linguistic origins to Taiwan⁹; their ancestral expansion is thought to have proceeded into Island Southeast Asia and eventually out into the Pacific¹⁹. The Austronesian-speaking settlers of the western Pacific (Fiji, Tonga, and Samoa) went on to people the widely dispersed islands in the vast ocean to their east through extraordinary voyages of exploration and settlement^{2,20}. Historians believe that family groups of 30–200 individuals sailed in double-hulled canoes across thousands of kilometers of open ocean to inhabit each new Polynesian island group^{21,22}. The first arrivals to these isolated island groups are thought to have experienced rapid initial growth, driven by the abundant resources of unfished reefs, huge seabird colonies and flightless birds (that soon became extinct) unhabituated to humans^{2,7,22–25}. These rapidly expanding island populations then initiated new voyages of exploration and colonization in search of—according to some theories—further untapped resources²⁶, a model supported by early oral histories²⁷. Geological analyses of Polynesian trade goods, particularly adzes, indicate that the remote Polynesian islands remained in trade contact with one another for several centuries^{25,28,29}.

However, these contacts were necessarily limited in frequency by the vast distances between island groups and limited in size by the capacities of the double-hulled sailing canoes²¹.

Under this historical model, we would expect the minor alleles on these isolated Pacific islands to be lost in a telescoping fashion following the order of the islands' colonization—a range expansion³⁰—due to the compounding succession of founding bottlenecks. We confirm this hypothesis below and then use its consequence—that the genetic composition of each remote island group is dominated by the contribution of its founders (see Extended Data Figure 3), whose descendants rapidly populated it—to reconstruct the Polynesian settlement sequence. We finally evaluate this model for self-consistency to test its validity.

Dimensionality of Polynesian genetics

In direct contrast to continental (and nearshore island) populations, where genetic substructure is shaped by large historical migrations, conquests, and diffusions occurring freely across the two dimensional landmass surface, thus producing two dimensional projections of genetic variance that mirror geography³¹, we find that Polynesian population structure exhibits high dimensionality (Supplementary Fig. 1) not at all reflective of geography (Fig. 1a), with islands diverging separately in a standard principal component analysis (PCA) (Supplementary Fig. 2–3). Indeed, the first two dimensions of major genomic variation—even in an ancestry-specific PCA of the Polynesian individuals (Fig. 1b)—do not separate islands geographically, as they do for within-continent populations^{32,33} (Extended Data Figs 1 – 2). Instead, each successive principal component captures the genetic drift of a particular island or island group (Fig. 1b, c and Supplementary Fig. 2), illustrating that genetic variance between these islands is dominated by their founder effects, not by diffusion clines or migration gradients. To further complicate such a standard variance-based approach (Supplementary Figs. 2 – 4) to genomic dimensionality reduction, the Polynesian islands differ widely in genetic diversity. Because the originating islands have much greater diversity (as discussed below), they dominate the first principal component when included in the PCA (Supplementary Fig. 3). Furthermore, many individuals, including all samples from certain islands, have some amount of non-Polynesian ancestry: European, Native American, and African³². The presence of large-scale post-colonial admixture from such divergent ancestry sources completely confounds Polynesian-focused interpretations of within-island and between-island variance when these admixed samples are included in the PCA (Supplementary Fig. 4).

To overcome these threefold obstacles to visualizing relationships between islands, we applied a novel ancestry-specific version of a nonlinear dimensionality reduction technique, t-Distributed Stochastic Neighbor Embedding (tSNE), applying it only to the genomic segments of Polynesian ancestry in our sampled individuals and employing a matrix completion step (Fig. 1d and Supplementary Fig. 5). In a plot of this ancestry-specific tSNE method (Fig. 1d), the islands of the ancestral west—Taiwan, Island Southeast Asia (Sumatra), Fiji, Tonga, and Samoa—are grouped on the left and the more recently settled eastern islands on the right. Islands in archipelagos, for instance the Cook Islands of Mauke, Atiu, and Rarotonga, form neighboring clusters. Rarotonga and Palliser appear at the center of the eastern Polynesian islands with the other eastern islands radiating out from them.

This pattern is consistent across alternate dimensionality reduction methods (see Methods), including our ancestry-specific formulations of uniform manifold approximation (UMAP) (Supplementary Figs 6, 7) and self-organizing map (SOM) (Supplementary Fig. 8), as well as our genetic drift projection method (Supplementary Fig. 9).

Tree building and path reconstruction

Because individuals from each of the islands form coherent, separate clusters in all of the non-linear, variant-based projections (tSNE, UMAP, and SOM), we can define a meaningful variant-frequency vector for each island by averaging the single nucleotide polymorphism (SNP) dosage vectors across all individuals on that island. Again, we consider only genomic segments of Polynesian origin (Supplementary Tables 3–4), since standard non-ancestry specific analyses are confounded by the recent introduction of highly differentiated colonial ancestry, such as European, even when that ancestry proportion is small (Supplementary Fig. 10). Averaging across all individuals reduces noise and produces composite Polynesian-specific frequency vectors with little to no remaining missingness from masking. Using these island-specific Polynesian-variant frequency vectors, we compute statistics for each pair of islands (Extended Data Figs. 4a–d, 7, and Supplementary Figs 11–19), including the average number of pairwise differences (π)³⁴, variant inner product (outgroup- F_3)³⁵, fixation index F_{st} , and directionality index^{36,37} (range expansion statistic) (ψ).

The directionality index ψ (Fig. 2a) measures the aggregate increase in frequencies of retained rare variants across the genome due to founder events, following the direction of a range expansion (see Fig. 2b and Supplementary Discussion ‘On psi’). The ψ -statistic gives crucial information that is not available from any genetic distance (π , F_2 , MixMapper³⁸) or inner product (F_3 , TreeMix³⁹) based methods; namely, a directionality arrow delineating a parental population from its child. Most human population studies have not required such directionality, as modern human populations are generally siblings, both having genetic drift from a no-longer extant, ancient parental population. That parental population, if available from ancient samples, is clearly indicated by the arrow of time (typically carbon dating). However, amongst the relatively recently settled Polynesian islands, genetic drift is created not so much by time, but by founder effects. Thus, the largely un-drifted (parental) populations for most of these islands are still extant: they are the populations of the originating islands. When constructing a population tree, this means that our dataset contains not only the terminal (leaf) nodes, but also the internal nodes, and we know their hierarchy from the ψ -statistic. This directional knowledge enables us to use a tree-building algorithm that, unlike population tree algorithms currently in use (Supplementary Figs. 20–21)^{35,38–40}, is guaranteed to find the optimal tree out of the space of all possible trees in the presence of perfect data (see Methods section ‘Migration network reconstruction’). Using this more robust directionality-based algorithm (see Supplementary Discussion ‘On tree-building’), the settlement path of Polynesia can be reconstructed (Fig. 2a).

Dating

To estimate dates for the settlement events that we infer, we use a method for detecting DNA segments that have been inherited from a common ancestor (identical-by-descent, IBD) for all pairs of individuals on different islands. Again, we consider only genomic

segments of Polynesian ancestry. For each pair of islands A and B, we pool all of the Polynesian IBD segments shared between individuals on A paired with individuals on B, and fit an exponential curve to the resulting segment length distribution (Fig. 2c and Extended Data Fig. 4d). From the decay constant of this exponential curve, we compute the number of generations elapsed since divergence of the island pair (Extended Data Fig. 6 and Supplementary Figs. 22–24). Fig. 2A shows the estimated divergence dates for all pairs of islands that are connected by a settlement path. Recent movement between islands, such as post-settlement trade contact, can introduce small numbers of longer, inter-island IBD segments, shifting the estimated divergence time towards the present, so we fit a truncated exponential. Nevertheless, these divergence dates should be seen as the terminus ante quem for the settlement of each child island (Fig. 2A and Extended Data Table 1 and Extended Data Fig. 6). In the case of the most remote islands such as Rapa Nui, which are believed to have had no large-scale population exchanges with other islands, the IBD-based date should coincide closely with the actual date of settlement.

The dates that we infer from our genome-wide network analyses support the radiocarbon-based ‘short chronology’ from the comprehensive re-analysis of Wilmshurst et al.¹² as corrected by Mulrooney et al.³ (Extended Data Table 1), as opposed to the prior nearly one thousand year older ‘long chronology’^{2,4}, and as opposed to the intermediate chronology suggested by Spriggs and Anderson¹³ (Marquesas 300–600 CE, remainder of eastern Polynesia 600–950 CE). Only in the settlement of the Marquesas Island group, dated by Mulrooney to the late 1100s CE, and the Southern Cook Islands, dated even later by Wilmshurst to the mid-1200s CE, do we find different (earlier) dates. However, as Mulrooney et al. explain, the small sample size of early-dated historical sites on each island mean that new archaeological discoveries could revise Wilmshurst’s chronology (backward). Our dates, from the full island-wide ancestral history coded within modern Polynesians themselves, do not have these sampling issues affecting ancient DNA and artifacts. Indeed, modern genomes complement ancient artifacts, since issues affecting the artifacts—finding the earliest human sites on each island, determining whether objects within them are anthropogenic, and determining whether those artifacts, often wood or charcoal, stem from young or old trees^{2,4,41} (inbuilt age), do not affect the former and vice versa.

Our date for the settlement of Rapa Nui is consistent with Wilmshurst and Mulrooney and also agrees closely with the date found by Hunt et al. (1200 CE) based on analyses of pollen in lake cores and soil erosion patterns⁴², as well as with recent radiocarbon dates of archaeological sites⁴³. Furthermore, unlike the long chronology estimates (200 BCE in the Marquesas), our settlement dates (1140 CE on Fatu Hiva in the Marquesas, or 28.4 generations before 1989) agree with the genealogical oral histories of many Pacific islanders themselves (1005 CE, or 29 generations before 1875, on Fatu Hiva)²⁷. In the Tuamotus our dates (1110 CE, or 29.3 generations before 1989) agree even more closely with island’s oral histories (1125 CE, or 28 generations before 1965)⁴⁴.

Our later divergence dates (1330–1360 CE) for some islands within archipelagos—North Marquesas (Nuku Hiva) in the Marquesas, Raivavae and Rimatara in the Australs—fall within the period of greatest inter-island trade contact in eastern Polynesia²⁵. Either the last islands were discovered during this period of long-distance trade voyaging, as suggested by

the dates of Schmid et. al⁴, or sufficient migration-to-inhabitant ratios still existed within archipelagos then to influence IBD distribution dates (see Supplementary Fig. 23). Note that our reconstruction of the settlement path is independent of these date estimates, which are overlaid upon it, and is more robust to later sporadic contact than IBD distributions are (see Methods sections ‘Polynesian ancestry-specific allele frequency analyses’ and ‘F₄’).

Discussion

Our analyses indicate the following scenario for the settlement of eastern Polynesia. From western Polynesia, Polynesian voyagers reached Rarotonga in the Cook Islands around 830 CE, having passed from Samoa along a route shared with the settlement of Fiji and Tonga. Rarotonga is the tallest and largest of the Cook Islands, with fertile volcanic soil watered by orographic rainfall²⁵, creating distinct clouds. These clouds on its prominent mountain make the island visible for long distances at sea and probably facilitated its discovery⁴⁵. From this base, we find that settlers continued south around 1190 CE to Rapa Iti (a branch recently hypothesized from linguistic evidence⁴⁶) and, separately, east to the smaller Cook Islands (Mauke and Atiu in our dataset).

Settlers also fanned out from Rarotonga northeast to the Society Islands (represented by Tahiti in our dataset but also containing the culturally significant island Ra‘i tea) around 1050 CE, thence northeast to the Tuamotu Archipelago (represented by Mataiva in the Palliser group in our dataset) by 1110 CE. At this time the widely scattered Tuamotu hub and other critical atolls in the expansion path (e.g. Nororotu in the Austral group), would have only recently emerged above falling sea levels (900 CE) and finished solidifying their topsoil and forests^{44,47} (Extended Data Fig. 5). Thus, our inferred dates and settlement path lend support to the idea that expansion into eastern Polynesia was mediated by the birth of those intermediary island clusters at the turn of the last millennium.

Stretching across central eastern Polynesia, the Tuamotu Archipelago was previously hypothesized to have served as a regional voyaging hub^{20,25,28}, and our analysis indicates that it was from this hub that settlers made their way north to the Marquesas Islands (Nuku Hiva and Fatu Hiva in our dataset) and south to the Gambier Islands (Mangareva in our dataset) beginning in the mid-1100s CE. From Mangareva, we find that the expansion reached the easternmost inhabited Polynesian island, Rapa Nui (Easter Island), around 1210 CE. This final leg had been suggested by some based on similarities between the Mangarevan and Rapanui languages⁴⁸, and by similarities in their traditional stone ceremonial platforms⁴⁹. This settlement sequence is also supported by our marker frequency-based genetic analyses, including ancestry-specific UMAP (Supplementary Fig. 6), drift projection (Supplementary Fig. 9), F-statistics (Supplementary Figs. 11–12), principal curve analyses (Extended Data Fig. 7 and Supplementary Fig. 17), diversity statistics (Supplementary Figs. 25–31), and ADMIXTURE clustering (Supplementary Figs. 32, 33).

Notably, we find that the population of Raivavae in the Australs arrived via the distant Tuamotus and Mangareva, rather than via the other Austral islands of Tubuai and Rimatara (Fig. 1a and Supplementary Figs. 6, 7). Together with even more distant North and South

Marquesas and Rapa Nui, each also with inferred settlement stemming from the Tuamotus, Raivavae had an ancient tradition of carving monumental anthropomorphic statues in stone. No other Austral island had these⁵⁰; indeed, such immense sculptures are found only on those far-flung islands that we now, however, show to had a common genetic source in the Tuamotu archipelago (Fig. 2a). It is also notable that it is only on islands that we infer were settled via the Tuamotus that pre-colonial Native American genetic contact has been identified, and its timing corresponds closely with our voyaging dates for this region³². This supports the theory that that contact occurred while the Polynesians were embarking on their easternmost, and longest, voyages of discovery.

The modern peoples of Polynesia carry strong genetic evidence for a range expansion beginning in Samoa and propagating out across eastern Polynesia through a series of telescoping founder events from the 11th through 12th centuries. Since this telescoping series of bottlenecks increased (via genetic surfing) the frequency of retained rare variants along the settlement path (see ψ -statistic, Fig. 2b), and since some of these variants are likely deleterious, future studies characterizing the individual frequencies and effects of these rare variants are desirable. We suggest that such large-scale sequencing and phenotyping studies should focus on the terminal islands in the settlement sequences that we have described, where compounded bottlenecking created the largest increase in frequencies (Fig 2b). We have shown that these particular islands also have high levels of homozygosity (Supplementary Figs. 25–27), which should increase the power to detect trait associations, and significant IBD, enabling IBD mapping, another useful approach⁵¹. Of note, two large modern Polynesian populations lie at the geographic termini of these serial bottleneck chains—Hawaii in the north and New Zealand in the south—and are thus notable candidates for such future large-scale association studies. We have introduced ancestry-specific computational methods for detailed characterization of Polynesian variant frequencies within admixed, modern samples, so potential admixture within future cohorts from such diverse populations should not be considered a barrier to designing these studies. Continued partnerships with these communities will be crucial⁵², since such studies will benefit both the personalized health understandings of these populations, as well as the global genetic understandings of all of us.

Methods

Data reporting

No statistical methods were used to predetermine sample size. The experiments were not randomized and the investigators were not blinded to allocation during experiments and outcome assessment.

Sample collection and approvals

This work combines publicly available sequence data and newly generated SNP array data from samples collected over different time periods by the participating institutions (see Supplementary Tables 1, 2). Written informed consent was obtained from all participants and research/ethics approval and permits were obtained from the following institutions: Stanford University Institutional Review Board (IRB approval No. 20839),

Oxford University Tropical Research Ethics Committee (reference No. 537–14), and the Scientific Ethics Committee of the Catholic University of Chile (reference No. 1971092). This study was also approved by the Council of Polynesian Elders for the community of Rapa Nui, along with local educational institutions, including the Lyceum Hoŋa'a o te Mana and the Lorenzo Baeza school for adults. Community engagement, including pre-participation presentations and post-participation return of results, were conducted throughout the project. Local approvals for engagement with the Rapa Nui community were obtained from the mayor (Pedro Pablo Edmunds Paoa) of the municipality of Easter Island, and the study was registered with the National Corporation for Indigenous Development (CONADI), in accordance with the indigenous law No. 19.253. The guidelines of the UNESCO International Declaration on Human Genetic Data and the Declaration of Helsinki were followed throughout the study.

Genotyping

Sampled populations and genotyping platforms are detailed in Supplementary Tables 1, 2. A total of 26 populations were genotyped at the University of California, San Francisco (UCSF) using Affymetrix (Mountain View, CA) Axiom LAT-1 arrays. Genotype calling was performed following default parameters using Affymetrix's Genotyping Console software. The average call rate was 98.5% for all newly genotyped samples. Before filtering and merging, the total number of SNPs called was 813,036. The resulting SNP density after merging with different reference panels varied across working datasets for downstream analyses, as detailed throughout the methods below.

Data curation

Quality control filters were applied across all sampled individuals using the Plink 1.9 package⁵³, removing individuals with >1% of genotyped sites missing (mind .01), removing genotyped sites missing in >1% of individuals (geno .01), and removing sites (18 SNPs) with extreme deviations from Hardy-Weinberg equilibrium (p-value less than 10e-110). The independence of drift between these separated, small island populations leads us to expect some deviation from Hardy-Weinberg in Polynesia, so we do not apply a typical threshold here. All samples were analyzed on the GRCh37 (hg19) genome build⁵⁴. REAP⁵⁵ was used to determine kinship coefficients using the ADMIXTURE clustering results discussed below; individuals with a kinship coefficient of >.2 (first degree relatives) were iteratively removed. Total numbers of individuals from each population after all filters were applied are given in Supplementary Table 1. After merging reference sequence data with sample genotyped data, strand inconsistencies were flipped when unambiguous, while ambiguous SNPs were removed, leaving 689,899 SNP sites. The recombination map from the 1000 Genomes project was used to assign genetic positions in centimorgans⁵⁶.

Admixture analyses

Principal component analysis (PCA)—EIGENSOFT 7.2.1⁵⁷ was used for all principal component analyses (PCA). LD-pruning was used across sliding 50 SNP windows with 10 SNP steps to remove variants with >0.5 squared correlation (--indep-pairwise 50 10 .5), leaving 461,571 SNPs for PCA. Plots were made with ggplot2 3.1.0⁵⁸ using R 3.5.2⁵⁹.

Global ancestry clustering analysis—Unsupervised ancestry clustering was performed using ADMIXTURE 1.3.0⁶⁰ on the LD-pruned dataset described above for PCA combining samples from all Pacific island populations together with continental references from Africa (Yoruba), Europe (Britain and Spain), East Asia (Japan and China), and the Americas (Aymara, Mapuche, Huilliche, Pehuenche) for a total of 686 samples. The numbers of samples from each population are given in Supplementary Tables 1–2. An elbow⁶¹ was found in the cross-validation error plot at $K = 7$ clusters, with larger numbers of clusters delivering little improvement (Supplementary Fig. 32).

Local ancestry analysis—Semi-supervised local ancestry inference was performed for all filtered Pacific island samples (430 samples, 689,899 SNP sites) using RFMix v1.5.4⁶² with two EM iterations and references from the five ancestry clusters, viz. African (60 West African Yoruba individuals), European (30 Spanish and 30 British individuals), Native American (60 Native American individuals from Puno, Peru), Ni-Vanuatuan (all 19 individuals from Vanuatu) and Remote Oceanian (60 individuals with <1% ancestry from outside the Pacific islands as identified by ADMIXTURE). The existence of these five ancestries within the Pacific island samples had been indicated by the $K=7$ unsupervised global ADMIXTURE clustering run discussed above (Supplementary Fig. 32). The recommended RFMix settings (two EM iterations and a .2 centimorgan window size) were used, and unphased samples were first phased by SHAPEITv2.837 with default settings⁶³. A few Pacific island individuals, particularly in the Marquesas, were found to also have >5% East Asian ancestry in the ADMIXTURE results. (This is likely due to the post-colonial movement to those islands of Hakka immigrants from China for work in the 19th century⁶⁴.) Those individuals were removed, so that this sixth ancestry did not need to be separately resolved by local ancestry analysis.

Masking—As discussed above, modern Pacific islanders are often admixed, possessing European and occasionally Native American and African ancestries (Supplementary Fig. 32). European ancestries entered Polynesia during the colonial period with the first European explorer (Magellan) arriving in the 16th century and significant immigration commencing in the early 19th century⁶⁴. Native American ancestry, particularly from emigration of admixed Hispanic individuals from Chile, which annexed Rapa Nui (Easter Island), and African ancestry also entered³². Because ancestries fully (or partially) introduced via colonial settlement did not necessarily follow the same island settlement process (or founder sizes and dates) as the original Polynesian settlement, such ancestries need to be distinguished, necessitating an ancestry-specific approach⁶⁵, cf. Supplementary Figs. 10, 20. For this reason we removed European chromosomal segments, as well as African and Native American, from the Pacific island samples. This step is called masking^{66,67}, since variants located in certain ancestry segments (identified above by RFMix via haplotype sequence pattern matching) are masked (removed) from the analysis. We refer to the remaining (unmasked) chromosomal segments as Polynesian ancestry chromosomal segments (Supplementary Table 3), and we refer to analyses that use only these segments as Polynesian ancestry-specific analyses. (Such analyses may still include as references non-Polynesian populations, such as from Europe or Taiwan. These reference populations will of course have non-Polynesian ancestry and are not masked.) A description

of which analyses were performed masked and which unmasked, and when references were used, is given in Supplementary Table 4.

Polynesian ancestry-specific allele frequency analyses

Treemix analysis—Treemix³⁹ was run on the combined set of Pacific island and reference populations (Supplementary Tables 1–2) using raw marker counts for each population. It was also run on the Pacific island populations using only the counts of markers found in Polynesian-ancestry chromosomal segments for each population, as described above.

Creation of Polynesian ancestry-specific genotype frequency vectors and matrix—For each of an individual's two haplotypes, variants located in non-Polynesian ancestry segments were masked, as described above. The two haplotypes for each individual were then averaged to create a genotype frequency vector having, for each site, 0 when no alternate allele was present, 0.5 when one alternate allele and one reference allele were present, and 1 when no reference allele was present. Some sites, where an individual had no Polynesian variant on either haplotype, remained missing. These missing values were accounted for in the following manner. The genotype frequency vector for each individual from the dataset was placed into the row of an N (individuals) by p (genotyped markers) matrix and the nuclear norm regularized matrix completion algorithm of Mazumder et al. was applied to create a reduced rank approximation to the original, incomplete 689,899-dimensional masked genotype matrix^{32,68}. Unlike earlier methods⁶⁹, this method permits the use of all samples, rather than only a panel of reference samples, for the completion step; thus, far more data is used allowing for more accurate completion. In addition, instead of using haplotypes (haploid genomes) as the unit of analysis, this method uses genotype frequency vectors (frequency vectors for the diploid genome). Since there is no linkage present in the genome across chromosome boundaries (due to independent assortment of chromosomes), population phasing cannot resolve parental haplotypes across these boundaries. Thus, a genome-wide haplotype vector constructed by assembling all chromosomes sequentially into a single row vector will switch phase arbitrarily across chromosome boundaries and so is already a mixture of an individual's two true parental haplotypes. By explicitly averaging an individual's two haplotype vectors to form a single genotype frequency vector for that individual, we are able to fill in much of the masked data that is missing from either of the two haplotypes.

Ancestry-specific drift projection—Each Pacific island individual's Polynesian ancestry-specific genotype frequency vector, described above, was projected onto the axis (drift axis), defined as the axis between the centroid of the indigenous Taiwan (Atayal and Paiwan) genotype frequency vectors and the centroid of the Rapa Nui (Easter Island) genotype frequency vectors. Each Pacific island individual's genotype frequency vector was also projected onto the first principal component of the subspace orthogonal to this axis to provide a second coordinate for two-dimensional visualization. (The first principal component of this orthogonal subspace is computed by finding the residual of each data point after subtracting off its component parallel to the drift axis and then determining the direction of greatest variation for these residuals.) The percent variance explained by

each dimension was computed directly by finding the variance of the projections on that dimension.

Ancestry-specific t-SNE—The number of significant ($p > 0.05$) dimensions for the genotype frequency matrix, described above, was determined ($n = 14$) using a Tracy-Widom distribution⁵⁷ and verified via a scree plot⁷⁰. To ensure that all population structure was captured, the genotype frequency matrix was projected onto its first twenty principal component axes. A tSNE was generated by applying the Barnes-Hut tSNE implementation to this projected matrix using: theta = 0, perplexity = 15, exaggeration factor = 10, max iter = 10000, and lying iter = 1000 parameters^{71,72}. Both a two-dimensional and three-dimensional embedding were created. (Projections onto fewer dimensions yielded similar results, with some clusters beginning to disappear in the range 12–15 dimensions, as predicted by the Tracy-Widom analysis.)

Ancestry-specific UMAP—The left singular vectors of the completed genotype frequency matrix were used as input for computing a two-dimensional uniform manifold approximation with a Manhattan distance metric and 80 nearest neighbors^{73,74}.

Ancestry-specific SOM—A two-dimensional self-organizing map (SOM) of the genotype frequency matrix was produced on a 100×100 rectangular grid using a Gaussian neighborhood⁷⁵. The package Somoclu, a massively parallel implementation of SOM, was used for optimization with parameters: 10 epochs, stdcoef .5, and linear cooling⁷⁶.

Principal coordinate analysis (PCoA) and principal curves—PCoA and principal curves were constructed from the relevant distance matrices (either π or F3, described below) using R 3.5.2 together with the package buds⁷⁷.

Population statistics—All population statistics described below (ψ , π , F3, F4, Fst, heterozygosity) were computed on population variant frequency vectors \tilde{p}_i created by computing, for each site, $\tilde{p}_i = \frac{\tilde{a}_i}{\tilde{n}_i}$, where \tilde{a}_i is the minor allele count at the site aggregated across all individuals' haplotypes (two haploid genomes per individual) in a population having that site located in a Polynesian chromosomal segment, and \tilde{n}_i is the total count of minor and major alleles. (A tilde is used throughout to denote counts from only Polynesian-specific chromosomal haplotypes.) Any sites not located in a Polynesian segment for any of the individuals within a population (or located in only one haplotype within the entire population) were removed from the dataset for all populations, so as to have no populations with one or fewer total allele observations at any site. This filtering resulted in the loss of 60,377 SNPs (8.75% of the total 689,899 SNPs), leaving 629,522 SNPs across all populations for computation of population allele frequency statistics.

Psi (ψ)—The range expansion statistic (ψ) of Peter et. al³⁶ (see Supplementary Discussion 'On directionality' and 'On psi') was computed first by polarizing all markers (identifying the minor allele) using the indigenous Taiwanese samples (Atayal and Paiwan) as an outgroup. To investigate the effect of using a different outgroup in a separate analysis a repolarization was performed using the western islands (Tonga, Samoa, Fiji) as an outgroup.

The latter calculation reduced the standard errors for the range expansion statistic on islands settled subsequent to western Polynesia, i.e. the eastern Polynesian islands; nevertheless, the general ordering of islands in the range expansion was the same for both calculations (see comparison Supplementary Fig. 14). Because allele frequencies drifted during the Pacific island settlement process, some minor alleles in Taiwan would have become major alleles by the time the settlers reached western Polynesia (see Supplementary Discussion ‘On psi’), so the intermediate repolarization using Tonga, Samoa, and Fiji as an outgroup increased the resolution of the range expansion statistic (reduced standard errors) for downstream islands. The larger number of samples from Tonga, Samoa, and Fiji (51), as opposed to Taiwan (22), also contributed, as it allowed us to set a more permissive bound for confirming that an allele observed minor in the outgroup samples was also minor in the outgroup population. This in turn increased the number of markers present in the latter analysis. (A 0.1 or lower minor allele frequency was required in the merged Tonga, Samoa, Fiji outgroup samples, yielding 228,262 SNPs, as opposed to the more stringent requirement of minor alleles being fixed in the Taiwan outgroup samples, following the procedure of Zhan et. al³⁷, which yielded only 137,383 SNPs.) Psi was calculated using the formula of Peter et. al,

$$\psi(A, B) = \frac{1}{\text{No. of shared SNPs}} \sum_{j \in \text{shared SNPs}} (\bar{p}_{A, j} - \bar{p}_{B, j}),$$

where the sum is taken only over SNPs shared polymorphic in both the population A sample and the population B sample⁷⁸. When using Taiwan as an outgroup, we masked the small Ni-Vanuatuan segments seen within Polynesia, since these segments trace their predominant ancestral origin back to a Papuan outgroup in New Guinea, rather than to the Austronesian outgroup, Taiwan. (Admixture between populations stemming from these two sources occurred on Vanuatu and other Melanesian islands in the thousand years before the settlement of Polynesia and was carried into Polynesia during its settlement^{17,79}.) However, both masking methods (both the Taiwan and the Tonga, Samoa, Fiji outgroup polarizations) gave the same ordering of islands settled.

Pi (π)—This quantity is the average number of pairwise differences per pair of haplotypes (haploid genomes) selected at random, one from each population, normalized by the number of sites^{34,80,81}. Also known as the nucleotide diversity⁸², it can be computed by first taking the ratio of the total number of mismatch combinations at a site to the total number of combinations, i.e. where a_1 is the number of alleles of one type in population 1 at a biallelic marker, b_1 is the number of the other type, $n_1 = a_1 + b_1$ is the total number of haplotypes in population 1, and thus $p_1 = a_1/n_1$ is the allele frequency in population 1, then at this site,

$$\begin{aligned} \pi_{12} &= \frac{a_1 \cdot b_2 + b_1 \cdot a_2}{(a_1 + b_1)(a_2 + b_2)} \\ &= \frac{a_1(n_2 - a_2) + a_2(n_1 - a_1)}{n_1 n_2} \end{aligned}$$

$$= p_1 - p_1 p_2 + p_2 - p_2 p_1$$

$$= p_1(1 - 2p_2) + p_2$$

This is an unbiased estimator that can be averaged over all sites to find the average number of pairwise differences per haplotype pair per site^{34,82}. Using this frequency-based formulation, this estimator can be generalized to Polynesian-specific allele frequencies for each island \tilde{p}_i .

F₃—The F3 shared drift statistic of Patterson et al.⁸³ was computed using the formula,

$$\hat{F}_3(C; A, B) = (\tilde{p}_C - \tilde{p}_A)(\tilde{p}_C - \tilde{p}_B) - \hat{h}_C/\tilde{n}_C,$$

where $\tilde{p}_A = \tilde{a}_A/\tilde{n}_A$ is the sample allele frequency in the ancestry of interest in population A (\tilde{n}_A total observations and \tilde{a}_A observations of the allele a in segments with the ancestry of interest) and $\hat{h}_A = \frac{\tilde{a}_A(\tilde{n}_A - \tilde{a}_A)}{\tilde{n}_A(\tilde{n}_A - 1)}$, similarly for B and C. For multiple sites these values are computed for each site and then averaged across all sites³⁵.

F₄—To detect departures from the reconstructed settlement tree (inter-island admixture), the F4 statistic was computed for each site using the formula of Patterson et al.³⁵

$$\hat{F}_4(A, B; C, D) = (\tilde{p}_A - \tilde{p}_C)(\tilde{p}_B - \tilde{p}_C),$$

and was then averaged across all sites. The F4 statistic is expected to be zero unless groups A, B do not form a separate clade from C, D within the actual population tree. Thus, when computing statistics of the form $F_4(\text{parental_island}, \text{child_island}; \text{Samoa}, X)$, where X varies across all islands that are not descended from *parental_island* in our model, a zero value of F4 is expected if the data completely support our settlement model. This is because all non-descendant islands (X) must lie in a common clade with outgroup *Samoa*; that is, external to the *parental_island*, *child_island* subclade. We look for significant evidence ($p < .001$) of departure from this model for each *parental_island*, *child_island* pair in our settlement sequence, and across all possible non-descendant islands X , while accounting for the multiple tests ($n = 52$) with a Bonferroni correction. We find deviations from our settlement tree only for 3 of its branches: Mangareva – Raivavae (migration from Tahiti), Mangareva – Palliser (migration from Tahiti), and North Marquesas – Palliser (migration from Tahiti and also from the Cooks). The Tahitian migrations go only to French Polynesian islands and likely reflect modern (see Supplementary Fig. 23) introgression to those islands from Tahiti, the modern capital, source of teachers, ministers, and civil servants, and center of employment, transportation, and residential education for French Polynesia. The migration from the Cooks directly to North Marquesas (bypassing the Palliser group) is intriguing, especially in light of our late dated Palliser-North Marquesas connection (1330

CE). It could be that North Marquesas (Nuku Hiva) was settled earlier more directly from the Cooks, whereas South Marquesas (Fatu Hiva) was, we have found, settled early (1140 CE) from Palliser. Later within-island-group migration between these neighboring islands may have led North Marquesas to exhibit these two origin signals, one from Palliser and one from the Cooks. If so, North and South Marquesas would be an unusual case, where two neighboring islands were settled from different parental islands, then, because they were not separated by large oceanic distances, were able to exchange enough subsequent migrants to leave a notable genetic trace within their post-growth population base.

F_{st}—The Hudson estimator for F_{st} is

$$\hat{F}_{st}^{\text{Hudson}} = \frac{(\tilde{p}_A - \tilde{p}_B)^2 - \frac{\tilde{p}_A(1 - \tilde{p}_A)}{\tilde{n}_A - 1} - \frac{\tilde{p}_B(1 - \tilde{p}_B)}{\tilde{n}_B - 1}}{\tilde{p}_A(1 - \tilde{p}_B) + \tilde{p}_B(1 - \tilde{p}_A)},$$

for a given SNP. For multiple sites, the numerator and the denominator (unbiased estimators of the variance between populations and the variance in the ancestral population respectively) are averaged across all SNPs separately before taking the ratio to create a consistent estimator⁸⁴.

Heterozygosity—The unbiased estimator for heterozygosity, first given by Nei⁸⁵, is, for a specific site

$$\hat{h} = 1 - \frac{N \sum \tilde{p}_\ell^2 - 1}{N - 1},$$

where \tilde{p}_ℓ is the frequency of the ℓ^{th} allele at the site, and N is the total number of alleles at that site (two for each of our SNPs). This estimator was aggregated across each SNP locus k using

$$\hat{H} = \sum_{k=1}^r \frac{\hat{h}_k}{r},$$

for all r of our SNP loci^{34,85}.

Standard errors—Standard errors for all allele frequency based statistics were computed using the block-bootstrap using 100 replicates and a block size of 1000 markers⁸⁶. This gives better variance estimates than the jackknife for these pairwise allele frequency comparisons³⁴. The markers are bootstrapped together as long contiguous blocks to preserve the effects of linkage on the variance of the estimates³⁵.

Migration network reconstruction

The various population measures of distance and directionality (ψ , π , F_3) between all pairs of islands define together tensors that annotate the complete graph of island connectivity. It remains to prune this graph judiciously to arrive at the tree representing the branching

settlement process of the serially founded Pacific islands; that is, a tree describing which islands were settled from which other islands (see Supplementary Discussion ‘On differences between range expansion trees and typical population trees’ and ‘on tree building’).

In brief, we use the range expansion statistic ψ (Fig. 2B) to determine the upstream islands along the range expansion; that is, the set of potential parent islands for each island. Beginning with the island with the largest ψ (measured against Samoa), we work backward in order of decreasing ψ (Fig. 2B and Extended Data Fig. 4A), joining each still orphaned island (j) to its closest related potential parent island (i) as defined by ψ . To measure genetic distance (closeness), we use the average number of pairwise differences π_{ij} (Extended Data Fig. 4B and Supplementary Fig. 17), since π_{ij} has been shown to have higher correlation with the divergence time between two populations (i and j) than the outgroup- F_3 statistic⁸¹ (see Supplementary Discussion ‘On Different Drift Distance Metrics’), although the same settlement sequence is also returned when using the latter metric instead (Supplementary Fig. 12).

Begin with the island with the most potential parents (at the end of the range expansion) (Fig. 2B) or in other words the largest ψ , Rapa Nui. Consistent with its terminal position in the range expansion, Rapa Nui also has the lowest heterozygosity (Supplementary Fig. 31) and the highest intra-island IBD (Supplementary Figs. 25–26). Starting with this terminal island, we consider all potential parent islands as indicated by the ψ directionality index, and connect Rapa Nui to the most closely related potential parent as indicated by the smallest average number of pairwise differences (π). We then proceed to the island with the second most potential parents according the ψ -statistic (here Raivavae, see Fig. 2b) and repeat. For Samoa, Fiji, Tonga and upstream islands, we use the directionality index (ψ) polarized using the Taiwan outgroup. For islands downstream of Samoa, as indicated by the Taiwan-polarization ψ , we use a ψ -statistic repolarized using the more proximal Samoa, Fiji, and Tonga outgroup, since it has smaller standard errors (see ψ discussion above).

This recursive algorithm for building the branching settlement path of the Pacific islands is a form of the Chu-Liu-Edmonds algorithm, which is guaranteed to produce the minimum spanning tree of a directed acyclic graph (see Supplementary Discussion ‘On tree-building’)^{87,88}. That our graph is acyclic can be shown (proof derived in Supplementary Discussion ‘On the acyclicity of ψ ’) from the formal definition of ψ , which defines our edge directionality. The lack of significant internal cross-migration edges was determined by our F_4 analysis above.

In the case of parental islands with multiple child islands, we can now use an inner product measure, the F_3 statistic (Extended Data Fig. 4C), which measures shared genetic drift, to determine whether any of those child islands share additional drift with each other beyond what they share with their common parent (Extended Data Fig. 7). Such additional shared drift is indicated in Fig. 2A by branching arrows; that is, arrows from a parent island that share an initial path before later branching to each child island. The order of arrow divergence indicates the ordering of shared drift amongst the child populations. These shared paths may suggest that intermediate islands in the settlement sequence are missing from our

dataset (Extended Data Fig. 5), since the founding bottleneck of an intermediate island could account for the additional shared drift. To further verify our settlement sequence, and to look for signs of post-settlement inter-island admixture, we compute F_4 statistics of the form $F_4(\textit{parental island}, \textit{child island}; \textit{Samoa}, X)$ with X ranging over all Polynesian islands not stemming from *parental island* in the settlement tree (described above). These F_4 statistics indicate whether there is statistically significant evidence for deviations from our settlement model; that is, later migrations across the ocean of sufficient size to significantly alter the genetic base of the post-growth island populations. Only three branches in our settlement sequence show any significant deviations, and each of these indicate a migration from Tahiti to an outlying French Polynesian island, consistent with Tahiti's recent role as the capital of French Polynesia.

Principal curve analysis

To independently verify our settlement sequence map, we compute unsupervised principal curves^{75,77} between the islands using genetic distances defined by both the outgroup- F_3 and π metrics (Extended Data Fig. 7 and Supplementary Fig. 17, respectively).

Identity-by-descent (IBD) analyses

In highly related populations, such as populations that have passed through a population size bottleneck in the recent past, individuals will share many ancestors, and thus many identical-by-descent (IBD) genetic fragments⁸⁹. In such cases, for example serially founded small island populations, IBD-based analyses become a powerful tool for reconstructing migrations.

Germline—GERMLINE 1.5.3 was run on the phased Pacific islander samples to find all identity-by-descent (IBD) shared segments of 5cM or greater using the `-min_m` flag. Fragments shorter than this length are prone to false positives due to insufficient SNPs^{90–92}. Up to four homozygous marker mismatches were permitted per IBD slice (`-err_hom`), and one heterozygous marker mismatch was permitted per IBD slice (`-err_het`).

Polynesian ancestry-specific filtering—To deconvolve Polynesian ancestral history from later (colonial and post-colonial) ancestry histories (European etc.) we used an ancestry-specific approach to IBD⁶⁵. Inter-island IBD segments lying wholly within post-colonial ancestries, or spanning post-colonial and pre-colonial ancestries, are necessarily the result of post-colonial inter-island contact events and were discarded. IBD segments lying wholly within chromosomal regions of known pre-colonial ancestry sources, that is Polynesian ancestry, were identified and analyzed together.

Runs of homozygosity (ROH)—Polynesian runs of homozygosity (ROH) were computed by summing together only Polynesian-specific IBD segments found shared between an individual's two haploid genomes, then normalizing by the effective fraction of homozygous Polynesian ancestry segments found in that individual. (These are the only segments of the diploid genome that could have shared a Polynesian ancestry ROH.) Population Polynesian-specific ROH values were computed by averaging these values for

all individuals within each island population. Standard errors were calculated by using the jackknife over individuals in a population⁹³.

Ancestry-specific sum of IBD segment lengths—When analyzing IBD segments, it has been typical to sum the total length (W_{ab}) of segments shared between a pair of individuals (a and b), one from each of a pair of populations (A and B), and then sum over all such pairs to arrive at a total sum of IBD sharing between each pair of populations⁹⁴. This sum can be normalized, dividing by the total number of possible cross-population pairs of individuals, one from each of the populations ($n_A n_B$), to give the average total IBD length shared (W_{AB}) per cross-population individual pair^{91,94,95}

$$W_{AB} = \frac{\sum_{a \in A} \sum_{b \in B} W_{ab}}{n_A n_B}$$

This normalization can also be performed over the total number of cross-population haplotype (haploid genome) pairs ($2n_A \cdot 2n_B$), rather than all individual pairs ($n_A n_B$)⁶⁵.

When considering only IBD segments found in those portions of both individuals' genomes that belong to a particular ancestry, the normalization must be modified to reflect the reduced fraction of the pairs' genomes that were considered. Thus, we replace the number of cross-population pair comparisons by an effective number of pair comparisons. If f_a is the fraction of the genome of a particular ancestry in individual a , and similarly for f_b , then the expected fraction of pairwise overlap between the two individuals is $f_a f_b$, rather than 1 as it is for non-admixed individuals. The denominator of the normalization above is now modified by the factor $\bar{f}_A \bar{f}_B$, where \bar{f}_A is the average fraction of the ancestry of interest in population A

$$\sum_{a \in A} \sum_{b \in B} f_a f_b = \left(\sum_{a \in A} f_a \right) \left(\sum_{b \in B} f_b \right) = (n_A \bar{f}_A) (n_B \bar{f}_B)$$

Within a single non-admixed population, the normalized intra-population IBD length sharing per haplotype pair is

$$W_{AA} = \frac{\sum_{a \in A} \sum_{\alpha \in A} W_{a\alpha}}{2n_A(2n_A - 1)}$$

The ancestry-specific normalization factor for intra-population IBD in an admixed population can be derived by considering the sum of all possible same-ancestry haplotype pair comparisons within the population of interest

$$\sum_{i=1}^{n_A} (f_i (\sum_{j < i} f_j)) = \frac{1}{2} \left(\sum_{i=1}^{n_A} \sum_{j=1}^{n_A} f_i f_j - \sum_{i=1}^{n_A} f_i^2 \right)$$

$$= \frac{1}{2} \left(\sum_{i=1}^{n_A} f_i \sum_{j=1}^{n_A} f_j - \sum_{i=1}^{n_A} f_i^2 \right) = \frac{1}{2} \left(n_A^2 \bar{f}_A^2 - \sum_{i=1}^{n_A} f_i^2 \right)$$

These ancestry-specific normalization factors make clear that, although the normalized total length of IBD sharing between two populations gives a measure of the relatedness of the populations, it is quite sensitive to an accurate estimation of the average fraction in each population of the ancestry of interest.

A heat map showing the normalized Polynesian-specific IBD sum values for each pair of Pacific islands in our dataset is displayed in Supplementary Fig. 24. Trends of increasing IBD sharing along the course of the inferred settlement chain (see the map in Extended Data Fig. 5) are evident, but there is significant noise.

IBD segment length distributions—A better approach is to compute the distribution of lengths of IBD segments shared between pairs of individuals, one from each of the two populations being compared. Although the total count (integral) of this distribution will be influenced by the fraction in each population of the ancestry of interest, the shape of the distribution will not be. Such robustness to the estimate of each population's ancestry fraction, which can vary by a few percent between different ancestry inference methods, is of great benefit. In addition, the shape of the IBD length distribution (decay rate) changes steadily each generation. It does not depend, as genetic drift does, on the fluctuations, which are generally unknown, of the historical population sizes.

Assuming no interference, recombination can be modeled as a Poisson process occurring along the genome at a rate of one recombination break per generation per unit of genomic length (measured in Morgans)⁹⁶. Thus, the length of a recombination segment, that is the distance between recombination events, is the waiting time of a Poisson process of rate T , where T is measured in generations. Hence, the distribution of the length of fragments (x) from a particular ancestor T generations ago will be exponential with $\lambda = T$ decay rate⁹⁷, $f(x) = \lambda e^{-\lambda x}$.

If we are considering recombination segments shared between two present day individuals stemming from the same common ancestor, that is IBD segments, we must adjust the rate for the number of recombination events per unit length that have occurred down both sides of the pedigree from this common ancestor, which gives a λ of $2T$ total^{92,95,98}. Each of these $2T$ opportunities for recombination to occur along the genome is called a meiosis event. For our empirical calculations (and all plots), we use centimorgans, rather than Morgans, so the λ rate constant is divided by 100, yielding $T/50$.

The total distribution of tract lengths shared between all individuals can be viewed as independent samples from the same exponential distribution. Coop et. al have shown that the decay rate parameter λ of this distribution is a weighted average of the distribution of times to the most recent common ancestor across all genomic sites⁹⁹. This distribution of times can be a complicated function of the demography, when the latter is not simple, leading to an ill-conditioned inverse problem⁹⁹. However, for our problem—dating the

founding of an isolated island group—the demography is amenable. Consider a parent island whose Polynesian explorers crossed thousands of kilometers of Pacific waves to discover and colonize a child island during the Polynesian settlement process. A pair of present-day Pacific islanders, one from the child island and one from the parent island, cannot share a common ancestor at *any* site (in their Polynesian ancestry segments) more recently than the founding date of the child island. Moreover, because of the small size of the founding populations arriving on double-hulled sailing canoes^{2,7,21,22}, all individuals on the child island will share ancestors with one another dating to at least the time of this founding bottleneck (plus a population size constant) before which time they coalesce rapidly with the ancestral lineages (also reduced due to prior bottlenecks) of modern individuals on the parent island. Thus, to first order the decay rate parameter λ will measure the time ($T/50$ with T in generations) to the split of the parent and child islands plus a constant, (which we assume is common for these remote islands and founder events and calibrate empirically, see Extended Data Table 1).

Example IBD length distributions for all pairs of individuals in our dataset—with one individual from Rapa Nui (Easter Island) and one from Mangareva, Palliser, Rarotonga, or Samoa—are plotted in Supplementary Fig. 22. Note that altering the normalization factor based on the estimated fractions of Polynesian ancestry amounts to a rescaling of the y-axis in Supplementary Fig. 22A or a translation of the y-axis in Supplementary Fig. 22B. This alters the amplitude, but not the decay rate shape parameter λ , of the exponential in Supplementary Fig. 22A, or, equivalently, the intercept, but not the slope of the lines in Supplementary Fig. 22B; thus demonstrating graphically that λ is robust to noise (errors) in ancestry normalization. However, the sum of IBD lengths, which is the integral of the curves in Supplementary Fig. 22A, is clearly not robust to such errors in the normalization (rescaling the y-axis).

Our empirically observed IBD length distributions are left truncated at 5 cM, since fragments shorter than this length are prone to false positives due to insufficient SNPs^{90–92}. The distributions are also right truncated at 15 cM, because outlier segments longer than this are expected to stem from recent contact, dating to less than 10 generations ago (18th century or later) as computed from the expected generation time (g) based on a single fragment length ℓ in Morgans, $E[g|\ell] \approx \frac{3}{2(\ell/1M)}$, where $1M$ is one Morgan⁹¹.

Such occasional post-colonial, inter-island Polynesian contact is not the focus of our Polynesian settlement analysis, so we filter out these few outlier inter-island IBD segments. Not removing these outliers does not change our island settlement dates significantly, but, by distorting the tail of the exponential decay distributions, does increase our standard error (see Supplementary Discussion ‘On Quantification of Error in IBD Dating’).

To estimate each pairwise λ , we use the maximum likelihood estimator for a left and right truncated exponential distribution. Since the exponential distribution is memoryless, the left truncation is trivially handled by translation of the distribution. That is, the distribution of the length in excess of 5 cM for each fragment is also exponential with the same decay constant λ . For what follows we assume that the IBD lengths have been thus recentered by

subtracting 5 cM. The right truncation is less elegant to handle, yielding an equation for the maximum likelihood estimator ($\hat{\lambda}$) of λ given by¹⁰⁰

$$\frac{\partial \ln \mathcal{L}_n}{\partial \lambda} \Big|_{\lambda = \hat{\lambda}} = n \left[\hat{\lambda}^{-1} - x_0 e^{-\hat{\lambda} x_0} (1 - e^{-\hat{\lambda} x_0})^{-1} - \bar{x} \right] = 0,$$

where \mathcal{L}_n (the sample likelihood for the n IBD segments) is the product of their individual likelihoods \mathcal{L} , \bar{x} is the (recentered) mean IBD length, x_0 is the (recentered) right truncation point and n is sample size (number of IBD segments). This transcendental equation must be solved numerically. The standard error (SE) can be obtained directly from the observed Fisher Information $I(\hat{\lambda})$

$$SE \approx \frac{1}{\sqrt{n \cdot I(\hat{\lambda})}}$$

since

$$(\hat{\lambda} - \lambda) \sqrt{n \cdot I(\hat{\lambda})} \stackrel{n \rightarrow \infty}{\sim} \mathcal{N}(0,1)$$

where the observed Fisher Information is found by

$$I(\hat{\lambda}) = - \frac{\partial^2 \ln \mathcal{L}}{\partial \lambda^2} \Big|_{\lambda = \hat{\lambda}} = \frac{1}{\hat{\lambda}^2} - \frac{x_0^2}{4 \sinh^2\left(\frac{1}{2} \hat{\lambda} x_0\right)}.$$

Using this method, the estimated λ values for the exponential distributions of Polynesian-specific IBD segment lengths for pairs of individuals, one from Rapa Nui (Easter Island) and one from each of the other remote Pacific islands in our dataset, are shown in Extended Data Fig. 4d. The pattern confirms the results of our drift statistics; Mangareva is the island most recently connected to Rapa Nui, while Samoa, the root of the expansion into remote Polynesia, is the most archaic connection.

A few caveats remain. The model of a Poisson process of recombination events along a continuous genome holds for small IBD segment lengths, viz. $T > 5$, but for more recent relatedness, leading to very long IBD segments, one must consider the finite size of the chromosomes themselves when computing the IBD length distribution¹⁰¹. In addition, the model of IBD segment independence holds only for ($N \gg T$) with N the population size and T the number of generations⁹⁷. Fortunately, our dataset has T values of 25–30 generations and N values in the thousands²¹, so we do not fall into these problematic regimes. However, because of the founding bottlenecks for each island, there is some intra-island IBD shared between pairs of individuals on the island (Supplementary Figs. 25, 26), so some recombination events will be non-productive. That is, when a haplotype stemming from one islander recombines with a haplotype from another with a recombination event occurring in the midst of an IBD segment on one haplotype that is shared with a third

individual, the recombination break will occasionally not break up the IBD sequence, since the two different recombining haplotype segments might themselves be identical (IBD) at the recombination point and thus both in IBD with the third individual. This will happen with a frequency equal to the percentage of the genome shared IBD on average between pairs of individuals on the same island. Therefore, we can correct for the frequency of these non-productive recombination events at each meiosis event. The correction factor ρ (the proportion of the genome shared on average intra-island) is specific to each island (dependent on the intra-island IBD on each island accrued through preceding founding bottlenecks) and so must be applied separately to the two branches of the pedigree, one from the common ancestor on the parent island down to the present-day on the parent island (A) and from that common ancestor population down to the present-day on the child island (B). The number of effective meioses, which is equal to λ , can be expressed after correction as

$$\lambda = g(1 - \rho_A) + g(1 - \rho_B) = g(2 - \rho_A - \rho_B)$$

where g is the number of generations to a common ancestor, ρ_A is the average fraction of the genome in IBD between pairs of individuals on A , equivalent to the probability of non-productive recombination on A , and similarly for ρ_B .

The correction factor ρ can be found by dividing the average total sum of IBD segments (S) between pairs of individuals on an island by the length of the genome (35.3 M)^{90,102}. Since we empirically observe only the sum of IBD segments longer than 5 cM, we must extrapolate the total sum of all IBD segments (N total segments, number generally unknown) by integrating our fitted approximate exponential distribution of IBD segment lengths (e.g. Supplementary Fig. 22A). Thus, the total sum of IBD is given by

$$S = N \cdot \int_{0cM}^{\infty} x \lambda e^{-\lambda x} dx = \frac{N}{\lambda},$$

and the truncated sum of IBD by,

$$s = N \cdot \int_{5cM}^{\infty} x \lambda e^{-\lambda x} dx = N \frac{5\lambda + 1}{\lambda e^{5\lambda}}.$$

By inspection we can see that,

$$S = s \frac{e^{5\lambda}}{5\lambda + 1}.$$

The extrapolated sums of IBD in the Polynesian component on average between pairs of individuals on each island as a percent of the genome are plotted in Supplementary Fig. 26, showing that these correction factors represent an adjustment of only a few percent.

We can now construct the symmetric matrix of Polynesian-specific pairwise island λ values (shown in Supplementary Fig. 23), and convert it, using our ρ adjustment factors

for each island, to a generation count to common ancestor for each island pair. For a detailed discussion of the uncertainty in our estimates of these dates see the Supplementary Discussion ('On quantification of error in IBD dating') and Extended Data Fig. 6.


Some island pairs, particularly distantly related islands or island pairs each with small numbers of samples, have large standard errors. Removing all entries corresponding to λ values that have standard errors above .07 (representing errors larger than 15% of the average λ value), creates a matrix of more precise generation values, but with some missing entries. Because this is a distance matrix, entries must be consistent with the triangle inequality, so we can impute these missing entries using triangulation from the precisely known entries. In fact, we can use something stronger than the standard triangle inequality

$$d(i, j) \leq d(i, k) + d(j, k) \quad \forall i, j, k \in \{islands\},$$

Because for distances $d(i, j)$ between two islands i and j and a third island k ; we can instead use the tighter ultrametric inequality

$$d(i, j) \leq \max\{d(i, k); d(j, k)\} \quad \forall i, j, k \in \{islands\}.$$

This holds in our case because all samples were taken from contemporary populations, and so all are leaf nodes of the ancestry tree dating to the same period (the present). Thus, as long as we use a distance metric $d(i, j)$ that is uniform in time for each population, for instance the ρ -adjusted generation (or year) matrix, the distances back from each pair of populations to their common ancestor will be identical, yielding an ultrametric tree. This works because the per generation recombination rate is constant over time, so the distance from an ancestral population to any of its sampled contemporary descendants is the same when measured by segment length distributions. (Note that this matrix, measuring the total distance along the tree from one island to another, is twice the matrix measuring how many generations have passed since an island pair split, since the former sums down both tree branches descending from the split.) To complete the proof of the ultrametric inequality we notice that for any two contemporary populations i and j and a third population k , k must either coalesce in ancestry with i first (k_1), with j first (k_2), or after i and j have themselves first coalesced (k_3). In the last case, $d(i, j)$ is clearly less than both $d(i, k_3)$ and $d(j, k_3)$, so the inequality above holds. In the first case, where k coalesces first with i , i and k_1 have a shared common ancestor before coalescing with the branch from j , so $d(i, j) = d(k_1, j)$, similarly for k_2 , making the bound of the ultrametric inequality valid (tight) for both cases.




Using the ultrametric inequality, we can impute unknown distances $d(i,j)$ simply by searching across all intermediate populations k and finding the minimum¹⁰³,

$$\min_{k \in \text{pops}} \{ \max\{d(i,k), d(j,k)\} \}.$$


From this completed distance matrix of generations, we can apply dates to each of the migrations using the average human generation time (see Supplementary Discussion ‘On generation times and meiosis events’).

Extended Data




Extended Data Figure 1. Comparison of genetic and geographic coordinates for European vs. Polynesian samples

(a) A principal component analysis of samples from Europe (15 from each nation) is shown to closely fit the geography of Europe. (See Extended Data Figure 2 for a quantitative comparison.) (b) A principal component analysis of samples from Polynesia (with non-Polynesian ancestry masked) is shown not to match the vast geography of the Pacific (c), and instead splits out island groups one at a time, reflecting the founder effects that dominate the variance of these populations.




Extended Data Figure 2. Permutation test for fit between genetic and geographic coordinates 100,000 random permutations of the population labels were created for the European populations' genetic data (blue, left) vs. the Polynesian populations (orange, right). For the European populations, out of 100,000 random permutations of the population labels on the genetic PCA, none better fits the geography of Europe (after fitting using a Procrustes analysis), than the correct labels, showing that the genetic coordinates of Europeans fit the geographic coordinates of Europe better than chance every time. However, for the Polynesian data 5% of the random permutations of the labels on the genetic PCA fit the

geographic coordinates of the Pacific islands better (after fitting using Procrustes), showing that the genetic data in Polynesia does not fit Polynesia's geography better than random chance. In the box and whiskers plots the mean and upper and lower quartiles of the rms error of the fits of the random permutations of population labels are indicated by horizontal lines. The fits of the actual population labels are indicated by asterisks.



Extended Data Figure 3. Continuity between ancient and modern Polynesian island populations


F_3 statistics were computed between ancient Rapanui samples and the Polynesian component from modern samples from each island in our dataset (top)¹¹⁵. Indigenous Austronesian language speakers from Taiwan (the Atayal) were used as an outgroup. The ancient Rapanui were found to be the most similar genetically to the modern Rapanui, indicating genetic continuity. A similar comparison was performed between the only other ancient samples from an island in our study, Tonga (bottom)¹⁸. Again, the modern Tongans appear most similar genetically; however, all islands downstream from Tonga in our inferred settlement path also share the same amount of genetic drift with the ancient Tongan samples (to within one standard error), as they should, since they are all descendants of these ancient Tongan sample according to our settlement reconstruction.



Extended Data Figure 4. Statistics used for settlement path inference

All statistics are based on the Polynesian-specific aggregate SNP frequency vectors computed for each island from all sampled individuals. The number (n) of individuals used are given for each island in Supplementary Table 1. (A) Directionality index (ψ), used to define sets of potential parent islands, plotted for each island relative to Samoa (equivalent to the top row of the matrix in Fig. 2B). (B) Average number of pairwise differences (π), measuring genetic distance and used to select the closest of potential parents, plotted for each island relative to Rapa Nui. (C) F_3 statistic, used to find additional shared genetic drift,


plotted for each island relative to Rapa Nui, with Taiwan as an outgroup. Standard errors in A-C were determined by a block bootstrap analysis. (D) Exponential decay constant (λ) for the Polynesian-specific IBD fragment length distributions between all pairs of individuals from Rapa Nui and each plotted island. The λ values can be used to calculate the number of generations elapsed since each pair of island populations were joined. Error bars show 95% confidence intervals of the maximum likelihood estimates determined analytically from the Fisher Information.



Extended Data Figure 5. Settlement map with candidate intermediate islands added

A reproduction of the map of Fig. 2a showing intermediate islands that are in the settlement path but not in our dataset that are possible candidates for explaining the additional shared drift observed in the corresponding colored settlement branches, that is, genetic drift shared between the child islands but not shared with the parent island. The additional shared drift of the Austral islands (Rimatarara and Tubuai) with the Society islands (Tahiti) and Tuamotus (Palliser) beyond what they each share with their parental island (Rarotonga in the Cooks) could indicate that there exists a shared intermediate island in their settlement path that we do not have in our dataset, for instance Mangaia¹¹⁶. Geological analyses of ancient tools found on Mangaia (green) have shown that it served as a connection between the Cook islands and remote eastern Polynesia²⁸, now uninhabited Nororotu (Maria Atoll) is also believed to have played a role as an intermediary island¹¹⁶. Traditional histories give Raiatea (pink) and its surrounding islands a role in the settling of remote eastern Polynesia¹¹⁶. Finally, linguistic studies have found connections between Marquesic languages (Marquesas


and Mangareva) and the central Tuamotus (orange)¹¹⁷. North Marquesas, South Marquesas, and Mangareva share drift with one another beyond what they share with Palliser, the westernmost island group in the Tuamotus, which could indicate that these three populations shared a common settlement path eastward through some of the Tuamotu Archipelago before diverging. Another possible explanation for additional shared drift is the settlement of each child island from a common subpopulation within the parental island, such as from the same clan or village.



Extended Data Figure 6. Effect of Phasing Errors on IBD Dates

IBD segments on the island of Rapa Nui were identified between all male X chromosomes. The log of the number of IBD segments (y axis) of a given genetic length (x axis) is plotted (orange; bottom left). The expected exponential decay of IBD segment lengths (linear semilog plot) is seen. The slope of this line (-0.161) is the exponential (decay) constant λ . Since the X chromosome is perfectly phased in men, because it is haploid, the identification of these IBD segments is unaffected by errors introduced through phasing algorithms. To quantify the effect of such errors, synthetic-female individuals were constructed by combining two male X chromosomes to make a diploid pair and to erase the phase information by recording only the genotype. The unphased diploid genotypes so constructed were phased and IBD segments were again identified and plotted (green; bottom right). The difference between the exponential decay constant (-0.166) of these statistically phased genotypes and the previous one is seen to be minor (top panel), amounting to three percent (3.01%), which corresponds to a difference of around 25 years for dates approximately eight hundred years ago (as in Polynesia). Uncertainty in the slope of the lines (equivalent to the uncertainty in the estimate exponential decay constant) is shaded.

Polynesian ancestry-specific F3 Ordination plot

**Extended Data Figure 7. Polynesian ancestry-specific shared drift ordination plot with principal curve**

A principal coordinate analysis (PCoA) projection of the pairwise shared drift distances (the Polynesian ancestry-specific outgroup- F_3) between each Pacific island population using Taiwan as an outgroup (Supplementary Fig. 12). This PCoA projection uses only the pairwise distance matrix and is fully unsupervised; that is, it does not presuppose that Rapa Nui is a terminal island along some settlement path. Nevertheless, it shows the same ordering as in Supplementary Fig. 9, confirming that Rapa Nui is indeed the terminal island in our dataset along the longest drift path, and confirming the drift ordering along that path. For further confirmation, a principal curve was also fit to the full dimensional space (Supplementary Fig. 12) and then projected into the two-dimensional PCoA space for visualization. The orthogonal projections of each island onto the principal curve are shown as thinner grey lines. This fully unsupervised principal curve confirms the visually

apparent path from Island Southeast Asia (Sumatra, far right) through Samoa, Fiji, Tonga and ending in Raivavae, Mangareva, and Rapa Nui (far left) in that order (cf. migration map in Fig. 2a). This projection of the high dimensional principal curve does not double back on itself, showing that the apparent ordering in this projection is consistent with the original high dimensional ordering. Note that this principal curve is able to fit only one settlement path (the principal one, that is, the longest drift path), which ends in Rapa Nui. Other settlement paths that branch away from this principal (longest) path appear simply as clusters projected onto the principal curve, since islands on those paths share no further drift with the principal path. That is, islands settled along secondary branching paths appear as clusters lying very close to one another along the principal curve. For example, Rapa Iti, which branches off from Rarotonga separately from the main settlement path (Fig. 2a), appears here as coincident with Rarotonga along the principal curve. The eigenvalue for PC1 over the sum of eigenvalues is .997 and for PC 2 is .002 (all eigenvalues are non-negative).

Extended Data Table 1. Archaeological and genetic inferred dates for first settlement

Settlement dates for each island group from our genomic data analysis (shown for the earliest settled island in each island group), compared with settlement date estimates from archaeological studies with (‘short chronology’) and without (‘long chronology’) the use of chronometric hygiene¹³. The differences between our earliest dates for each island group and the Wilmshurst et. al radiocarbon dates are slight, with the exception of the Cooks. For the latter our dates are within those of Allen and Steadman¹⁰⁴, and, in contrast to the model described by Kirch², we have the Cooks preceding the Polynesian islands to their east in the settlement sequence. Such an order (Cook Islands first) had been suggested before based on the early establishment there of Polynesian rats¹¹⁴.

Island Groups (or single island if solitary)	Our genetic estimates (earliest date within each group)	Archaeological estimates	
		Short chronology	Long chronology ²
Cook Islands	830 CE (Rarotonga)	810–1170 CE (Allen 1990 ¹⁰⁴ , Spriggs 1993 ¹³) 850–1136 CE (Niespolo 2019 ¹⁰⁵ , Kirch 2017 ¹⁰⁶) 900 CE (Atiu, Sear 2020 ¹⁰⁷) mid-1200s CE (Wilmshurst 2011 ¹²) 1231–1290 CE (Schmid 2018 ⁴)	1 CE
Society Islands	1050 CE (Tahiti)	997–1079 CE (Schmid 2018 ⁴) 1000–1100 CE (Kahn 2017 ¹⁰⁸) 1025–1121 CE (Wilmshurst 2011 ¹² , Mulrooney 2011 ³) 1050–1160 CE (Anderson 2019 ⁴¹)	200 BCE
Marquesas Islands	1140 CE (Fatu Hiva)	850–900 CE (Conte 2014 ¹⁰⁹) 1166–1258 CE (Allen 2014 ¹¹⁰) late 1100s CE (Mulrooney 2011 ³) 1224–1265 CE (Schmid 2018 ⁴)	200 CE
Austral Islands	1150 CE (Tubuai)	1128–1228 CE (Prebble 2009 ¹¹¹) 1391–1517 CE (Schmid 2018 ⁴)	“
Rapa Iti	1190 CE	1100–1200 CE (Anderson 2012 ¹¹²)	“
Gambier Islands	1130 CE (Mangareva)	950 CE (Kirch 2010 ¹¹³) 1000–1250 CE (Anderson 2019 ⁴¹) 1099–1208 CE (Schmid 2018 ⁴) 1109–1275 CE (Wilmshurst 2011 ¹² , Mulrooney 2011 ³)	“

Island Groups (or single island if solitary)	Our genetic estimates (earliest date within each group)	Archaeological estimates	
		Short chronology	Long chronology ²
Rapa Nui (Easter Island)	1210 CE	1000–1100 CE (Kirch 2017 ²) 1200 CE (Hunt 2008 ⁴² , Mulrooney 2013 ⁴³) 1210–1253 CE (Wilmshurst 2011 ¹² , Mulrooney 2011 ³) 1221–1268 CE (Schmid 2018 ⁴)	300 CE

Supplementary Material

Refer to Web version on PubMed Central for supplementary material.

Acknowledgements

We thank the participants and volunteers who donated DNA samples for this study and the local community leaders and authorities who provided approval and logistics support in the various sampling sites, including Mr. T. Teariki (Cook Islands), Dr. S. Foliaki (Tonga), Dr. N. Tuuau (Samoa), Dr. H.-W. Peng (Taiwan), Professor J. Roux (French Polynesia), Mayor Edmunds Paoa (Rapa Nui), as well as Professors J. Martinson, D. Weatherall and J. Clegg for their pioneering fieldwork in the region that ultimately led to this work. We thank H. Kane and S. Kahanamoku for critiques and suggestions on the text, as well as M. Spriggs for his comments on the manuscript concerning archaeological dates in Polynesia. We thank the support from the Core Staff at the UCSF Institute for Human Genetics for contributing genotyping capacity, and the Stanford Center for Computational, Evolutionary and Human Genomics (CEHG) for supporting the initial stages of this project. We thank J. Cervantes for IT support and M. Ortega and G. Mireles for technical support at LANGEBIO's Genomics Core Facility at CINVESTAV, Mexico. This work was supported by the George Rosenkranz Prize for Health Care Research in Developing Countries, Mexico's CONACYT Basic Research Program (grant number CB-2015-01-251380), and the International Center for Genetic Engineering and Biotechnology (ICGEB, Italy) Grant CRP/MEX15-04_EC (each awarded to A.M.-E); the Chilean funding programs FONDEF, FONDECYT, and CONICYT (grants D10I1007, 1130303, and USA2013-0015, respectively); the National Institute for Health Research (NIHR) Oxford Biomedical Research Centre (BRC) and a Wellcome Trust Fellowship with reference 106289/Z/14/Z (A.J.M.); together with National Library of Medicine (NLM) training grant T15LM007033 and an American Society of Engineering Education NDSEG Fellowship (each awarded to A.G.I.). Views expressed are those of the author(s) and not necessarily those of the NIHR, the NHS, or the Department of Health.

Data availability

The samples for this project were collected by the University of Oxford, Stanford University, and the University of Chile as part of different studies. SNP data for all newly genotyped individuals is available through a data access agreement to respect the privacy of the participants for the transfer of genetic data from the European Genome-Phenome Archive under accession number EGAS00001005362.

References

1. Low S. *Hawaiki Rising: Hākeleʻa, Nainoa Thompson, and the Hawaiian Renaissance*. (University of Hawaii Press, 2019).
2. Kirch PV *On the Road of the Winds*. (University of California Press, 2017).
3. Mulrooney MA, Bickler SH, Allen MS & Ladefoged TN High-precision dating of colonization and settlement in East Polynesia. *Proc. Natl. Acad. Sci. U.S.A* 108, E192–E194 (2011). [PubMed: 21633014]
4. Schmid MME et al. How 14C dates on wood charcoal increase precision when dating colonization: The examples of Iceland and Polynesia. *Quaternary Geochronology* 48, 64–71 (2018).
5. Kahaloʻi Keauokalani K. *Kepelino's Traditions of Hawaii*. Bernice P. Bishop Museum Bulletin 206 (1932).

6. Cook J. *The Journals of Captain James Cook on his Voyages of Discovery*. (Cambridge University Press, 1955).
7. Kirch PV & Green RC *Hawaiki, Ancestral Polynesia*. (Cambridge University Press, 2001).
8. Minster RL et al. A thrifty variant in CREBRF strongly influences body mass index in Samoans. *Nat. Genet* 48, 1049–1054 (2016). [PubMed: 27455349]
9. Gray RD, Drummond AJ & Greenhill SJ Language phylogenies reveal expansion pulses and pauses in Pacific settlement. *Science* 323, 479–483 (2009). [PubMed: 19164742]
10. Walworth M. Eastern Polynesian: The linguistic evidence revisited. *Oceanic Linguistics* 53, 256–272 (2014).
11. Martinsson-Wallin H, Wallin P. & Anderson A. Chronogeographic variation in initial East Polynesian construction of monumental ceremonial sites. *The Journal of Island and Coastal Archaeology* 8, 405–421 (2013).
12. Wilmshurst JM, Hunt TL, Lipo CP & Anderson AJ High-precision radiocarbon dating shows recent and rapid initial human colonization of East Polynesia. *Proc. Natl. Acad. Sci. U.S.A* 108, 1815–1820 (2011). [PubMed: 21187404]
13. Spriggs M. & Anderson A. Late colonization of east Polynesia. *Antiquity* 67, 200–217 (1993).
14. Hill AVS et al. Polynesian origins and affinities: Globin gene variants in eastern Polynesia. *Am. J. Hum. Genet* 40, 453–463 (1987). [PubMed: 2883894]
15. Wollstein A. et al. Demographic history of Oceania inferred from genome-wide data. *Curr. Biol* 20, 1983–1992 (2010). [PubMed: 21074440]
16. Hudjashov G. et al. Investigating the origins of eastern Polynesians using genome-wide data from the Leeward Society Isles. *Sci. Rep* 8, 1823 (2018). [PubMed: 29379068]
17. Skoglund P. et al. Genomic insights into the peopling of the Southwest Pacific. *Nature* 538, 510–513 (2016). [PubMed: 27698418]
18. Posth C. et al. Language continuity despite population replacement in Remote Oceania. *Nat Ecol Evol* 2, 731–740 (2018). [PubMed: 29487365]
19. McColl H. et al. The prehistoric peopling of Southeast Asia. *Science* 361, 88–92 (2018). [PubMed: 29976827]
20. Emory KP *The Tuamotuan Creation Charts by Paiore*. *J. Polynesian Soc* 48, 1–29 (1939).
21. Hunt T. & Lipo C. *The Statues That Walked*. (Free Press, 2011).
22. Whyte ALH, Marshall SJ & Chambers GK Human evolution in Polynesia. *Human Biology* 77, 157–177 (2005). [PubMed: 16201134]
23. Duncan RP, Boyer AG & Blackburn TM Magnitude and variation of prehistoric bird extinctions in the Pacific. *Proc. Natl. Acad. Sci. U.S.A* 110, 6436–6441 (2013). [PubMed: 23530197]
24. Kirch PV et al. Human ecodynamics in the Mangareva Islands: a stratified sequence from Nenega-Iti Rock Shelter (site AGA-3, Agakautai Island). *Archaeology in Oceania* 50, 23–42 (2015).
25. Rolett BV *Voyaging and interaction in ancient East Polynesia*. *Asian Perspectives* 41, 182–194 (2002).
26. Steadman DW *Extinction and Biogeography of Tropical Pacific Birds*. (University of Chicago Press, 2006).
27. Handy ESC *The Native Culture in the Marquesas*. (The Bishop Museum, 1923).
28. Weisler MI et al. Cook Island artifact geochemistry demonstrates spatial and temporal extent of pre-European interarchipelago voyaging in East Polynesia. *Proc. Natl. Acad. Sci. U.S.A* 113, 8150–8155 (2016). [PubMed: 27382159]
29. Collerson KD & Weisler MI Stone adze compositions and the extent of ancient Polynesian voyaging and trade. *Science* 317, 1907–1911 (2007). [PubMed: 17901331]
30. Slatkin M. & Excoffier L. Serial Founder Effects During Range Expansion: A Spatial Analog of Genetic Drift. *Genetics* 191, 171–181 (2012). [PubMed: 22367031]
31. Stephens M. & Novembre J. Interpreting principal component analyses of spatial population genetic variation. *Nat. Genet* 40, 646–649 (2008). [PubMed: 18425127]
32. Ioannidis AG et al. Native American gene flow into Polynesia predating Easter Island settlement. *Nature* 583, 572–577 (2020). [PubMed: 32641827]

33. Novembre J. et al. Genes mirror geography within Europe. *Nature* 456, 98–101 (2008). [PubMed: 18758442]
34. Nei M. & Kumar S. *Molecular Evolution and Phylogenetics*. (Oxford University Press, USA, 2000).
35. Patterson N. et al. Ancient admixture in human history. *Genetics* 192, 1065–1093 (2012). [PubMed: 22960212]
36. Peter BM & Slatkin M. Detecting range expansions from genetic data. *Evolution* 67, 3274–3289 (2013). [PubMed: 24152007]
37. Zhan S. et al. The genetics of monarch butterfly migration and warning colouration. *Nature* 514, 317–321 (2014). [PubMed: 25274300]
38. Lipson M. et al. Efficient moment-based inference of admixture parameters and sources of gene flow. *Mol. Biol. Evol* 30, 1788–1802 (2013). [PubMed: 23709261]
39. Pickrell JK & Pritchard JK Inference of population splits and mixtures from genome-wide allele frequency data. *PLoS Genet.* 8, e1002967 (2012).
40. Leppälä K, Nielsen SV & Mailund T. admixturegraph: An R package for admixture graph manipulation and fitting. *Bioinformatics* 33, 1738–1740 (2017). [PubMed: 28158333]
41. Anderson AJ, Conte E, Smith I. & Szabo K. New excavations at Fa’ahia (Huahine, Society Islands) and chronologies of central East Polynesian colonization. *J. Pac. Arch* 10, 1–14 (2019).
42. Hunt TL & Lipo CP Evidence for a shorter chronology on Rapa Nui (Easter Island). *The Journal of Island and Coastal Archaeology* (2008).
43. Mulrooney MA An island-wide assessment of the chronology of settlement and land use on Rapa Nui (Easter Island) based on radiocarbon data. *J. Archaeological Science* 40, 4377–4399 (2013).
44. Pirazzoli PA & Montagnoni LF Late Holocene sea-level changes in the northwest Tuamotu islands, French Polynesia. *Quaternary Research* 25, 350–368 (1986).
45. Di Piazza A, Di Piazza P. & Pearthree E. Sailing virtual canoes across Oceania: revisiting island accessibility. *J. Archaeological Science* 34, 1219–1225 (2007).
46. Walworth M. *The Language of Rapa Iti*. (University of Hawaii, 2015).
47. Dickinson W. Pacific atoll living: how long already and until when. *Geological Society of America Today* 19, 4–10 (2009).
48. Fischer SR Mangarevan doublets: Preliminary evidence for Proto-Southeastern Polynesian. *Oceanic Linguistics* 40, 112–124 (2001).
49. Flenley J. & Bahn P. *The Enigmas of Easter Island*. (Oxford University Press, 2003).
50. Buck Te Rangi H roa PH *Vikings of the Sunrise*. (Lippincott JB, 1938).
51. Belbin GM et al. Toward a fine-scale population health monitoring system. *Cell* 184, 2068–2083.e11 (2021). [PubMed: 33861964]
52. Claw KG et al. A framework for enhancing ethical genomic research with Indigenous communities. *Nat. Comms* 9, 1–7 (2018).
53. Chang CC et al. Second-generation PLINK: rising to the challenge of larger and richer datasets. *GigaScience* 4, 7 (2015). [PubMed: 25722852]
54. Tynes C. et al. The UCSC Genome Browser database: 2017 update. *Nucleic Acids Res* 45, D626–D634 (2017). [PubMed: 27899642]
55. Thornton T. et al. Estimating kinship in admixed populations. *Am. J. Hum. Genet* 91, 122–138 (2012). [PubMed: 22748210]
56. The 1000 Genomes Project Consortium, The 1000 Genomes Project. A map of human genome variation from population-scale sequencing. *Nature* 467, 1061–1073 (2010). [PubMed: 20981092]
57. Patterson N, Price AL & Reich D. Population structure and eigenanalysis. *PLoS Genet.* 2, e190 (2006). [PubMed: 17194218]
58. Wickham H. *ggplot2*. (Springer, 2016).
59. R Core Team. *R: A Language and Environment for Statistical Computing*. (2017).
60. Alexander DH & Lange K. Enhancements to the ADMIXTURE algorithm for individual ancestry estimation. *BMC Bioinformatics* 12, 1–6 (2011). [PubMed: 21199577]

61. Holmes S. & Huber W. *Modern Statistics for Modern Biology*. (Cambridge University Press, 2019).
62. Maples BK, Gravel S, Kenny EE & Bustamante CD RFMix: A discriminative modeling approach for rapid and robust local-ancestry inference. *Am. J. Hum. Genet* 93, 278–288 (2013). [PubMed: 23910464]
63. O’Connell J. et al. A general approach for haplotype phasing across the full spectrum of relatedness. *PLoS Genet.* 10, e1004234 (2014).
64. D’Arcy P. The Chinese Pacifics: A brief historical review. *The Journal of Pacific History* 49, 396–420 (2014).
65. Browning SR et al. Ancestry-specific recent effective population size in the Americas. *PLoS Genet.* 14, e1007385 (2018).
66. Schroeder H. et al. Origins and genetic legacies of the Caribbean Taino. *Proc. Natl. Acad. Sci. U.S.A* 115, 2341–2346 (2018). [PubMed: 29463742]
67. Moreno-Estrada A. et al. The genetics of Mexico recapitulates Native American substructure and affects biomedical traits. *Science* 344, 1280–1285 (2014). [PubMed: 24926019]
68. Mazumder R, Hastie T. & Tibshirani R. Spectral regularization algorithms for learning large incomplete matrices. *J. Machine Learning Research* 11, 2287–2322 (2010).
69. Moreno-Estrada A. et al. Reconstructing the population genetic history of the Caribbean. *PLoS Genet.* 9, e1003925 (2013).
70. Nguyen LH & Holmes S. Ten quick tips for effective dimensionality reduction. *PLoS Comp. Biol* 15, e1006907 (2019).
71. Maaten LVD & Hinton G. Visualizing data using t-SNE. *J. Machine Learning Research* 9, 2579–2605 (2008).
72. Van Der Maaten L2014. Accelerating t-SNE using tree-based algorithms. *The Journal of Machine Learning Research* (2014).
73. McInnes L, Healy J. & Melville J. UMAP: Uniform manifold approximation and projection for dimension reduction. [arXiv.org](https://arxiv.org/abs/1802.03426) (2018).
74. Diaz-Papkovich A, Anderson-Trocmé L, Ben-Eghan C. & Gravel S. UMAP reveals cryptic population structure and phenotype heterogeneity in large genomic cohorts. *PLoS Genet.* 15, e1008432 (2019).
75. Hastie T, Tibshirani R. & Friedman J. *The Elements of Statistical Learning*. (Springer Science & Business Media, 2009).
76. Wittek P, Gao SC, Lim IS & Zhao L. Somoclu: An efficient parallel library for self-organizing maps. *J. Statistical Software* 78, (2017).
77. Nguyen LH & Holmes S. Bayesian Unidimensional Scaling for visualizing uncertainty in high dimensional datasets with latent ordering of observations. *BMC Bioinformatics* 18, 65–79 (2017). [PubMed: 28251866]
78. Peter BM & Slatkin M. The effective founder effect in a spatially expanding population. *Evolution* 69, 721–734 (2015). [PubMed: 25656983]
79. Pugach I. et al. The Complex Admixture History and Recent Southern Origins of Siberian Populations. 33, 1777–1795 (2016).
80. Takahata N. & Nei M. Gene genealogy and variance of interpopulational nucleotide differences. *Genetics* 110, 325–344 (1985). [PubMed: 4007484]
81. Peter BM Admixture, population structure, and F-statistics. *Genetics* 202, 1485–1501 (2016). [PubMed: 26857625]
82. Nei M. *Molecular Evolutionary Genetics*. (Columbia University Press, 1987).
83. Patterson N. et al. Reconstructing Indian population history. *Nature* 461, 489–494 (2009). [PubMed: 19779445]
84. Bhatia G, Patterson N, Sankararaman S. & Price AL Estimating and interpreting FST: The impact of rare variants. *Genome Research* 23, 1514–1521 (2013). [PubMed: 23861382]
85. Nei M. & Roychoudhury AK Sampling variances of heterozygosity and genetic distance. *Genetics* 76, 379–390 (1974). [PubMed: 4822472]

86. Davison AC & Hinkley DV *Bootstrap Methods and Their Application*. (Cambridge University Press, 1997).
87. Chu YJ & Lui TH On the shortest arborescence of a directed graph. *Science Sinica* 1396–1400 (1965).
88. Edmonds J. Optimum branchings. *J. Res. Nat. Bureau Standards* 71B, 233–240 (1967).
89. Ceballos FC, Joshi PK, Clark DW, Ramsay M. & Wilson JF Runs of homozygosity: windows into population history and trait architecture. *Nat Rev Genet* 19, 220–234 (2018). [PubMed: 29335644]
90. Huff CD et al. Maximum-likelihood estimation of recent shared ancestry (ERSA). *Genome Research* 21, 768–774 (2011). [PubMed: 21324875]
91. Baharian S. et al. The Great Migration and African-American genomic diversity. *PLoS Genet.* 12, e1006059 (2016).
92. Gusev A. et al. Whole population, genome-wide mapping of hidden relatedness. *Genome Research* 19, 318–326 (2009). [PubMed: 18971310]
93. Efron B. Bootstrap methods: Another look at the jackknife. *The Annals of Statistics* 7, 1–26 (1979).
94. Botigué LR et al. Gene flow from North Africa contributes to differential human genetic diversity in southern Europe. *Proc. Natl. Acad. Sci. U.S.A* 110, 11791–11796 (2013). [PubMed: 23733930]
95. Atzmon G. et al. Abraham’s children in the genome era: Major Jewish diaspora populations comprise distinct genetic clusters with shared Middle Eastern ancestry. *Am. J. Hum. Genet* 86, 850–859 (2010). [PubMed: 20560205]
96. Jobling M, Hurles M. & Tyler-Smith C. *Human Evolutionary Genetics*. (Garland Science, 2013).
97. Liang M. & Nielsen R. The lengths of admixture tracts. *Genetics* 197, 953–967 (2014). [PubMed: 24770332]
98. Palamara PF, Lencz T, Darvasi A. & Pe’er I. Length distributions of identity by descent reveal fine-scale demographic history. *Am. J. Hum. Genet* 91, 809–822 (2012). [PubMed: 23103233]
99. Ralph P. & Coop G. The geography of recent genetic ancestry across Europe. *PLoS Biol* 11, e1001555 (2013).
100. Deemer WL Jr, 1955 & Votaw DVJTAOM Jr. Estimation of parameters of truncated or censored exponential distributions. *The Annals of Mathematical Statistics* 498–504 (1955).
101. Hill WG & White IMS Identification of pedigree relationship from genome sharing. *G3: Genes, Genomes, Genetics* 3, 1553–1571 (2013). [PubMed: 23893739]
102. McVean GAT et al. The Fine-Scale Structure of Recombination Rate Variation in the Human Genome. *Science* 304, 581–584 (2004). [PubMed: 15105499]
103. Makarenkov V. & Lapointe F-J A weighted least-squares approach for inferring phylogenies from incomplete distance matrices. *Bioinformatics* 20, 2113–2121 (2004). [PubMed: 15059836]
104. Allen MS & Steadman DW Excavations at the Ureia Site, Aitutaki, Cook Islands: Preliminary Results. *Archaeology in Oceania* 25, 24–37 (1990).
105. Niespolo EM, Sharp WD & Kirch PV 230Th dating of coral abraders from stratified deposits at Tangatatau Rockshelter, Mangaia, Cook Islands: Implications for building precise chronologies in Polynesia. *J. Archaeological Science* 101, 21–33 (2019).
106. Kirch PV Tangatatau Rockshelter: the evolution of an eastern Polynesian socio-ecosystem. (Cotsen Institute of Archaeology Press, 2017).
107. Sear DA et al. Human settlement of East Polynesia earlier, incremental, and coincident with prolonged South Pacific drought. *Proc. Natl. Acad. Sci. U.S.A* 117, 8813–8819 (2020). [PubMed: 32253300]
108. Kahn JG & Sinoto Y. Refining the Society Islands cultural sequence: Colonisation phase and developmental phase coastal occupation on Mo’orea Island. *J. Polynesian Society* 126, 33 (2017).
109. Conte E. & Molle G. Reinvestigating a key site for Polynesian prehistory: new results from the Hane dune site, Ua Huka (Marquesas). *Archaeology in Oceania* 49, 121–136 (2014).
110. Allen MS Marquesan Colonisation Chronologies and Postcolonisation Interaction: Implications for Hawaiian origins and the ‘Marquesan Homeland’ hypothesis. *J. Pac. Arch* 5, 1–17 (2014).

111. Prebble M. & Wilmshurst JM Detecting the initial impact of humans and introduced species on island environments in Remote Oceania using palaeoecology. *Biol Invasions* 11, 1529–1556 (2009).
112. Anderson A, Kennett DJ, Culleton BJ & Southon J. in *Taking the High Ground* (eds. Anderson A. & Kennett DJ) 288 (ANU E Press, 2012). doi:10.2307/j.ctt24hd92.13
113. Kirch PV, Conte E, Sharp W. & Nickelsen C. The Onemea Site (Taravai Island, Mangareva) and the human colonization of Southeastern Polynesia. *Archaeology in Oceania* 45, 66–79 (2010).
114. Matisoo-Smith E. et al. Patterns of prehistoric human mobility in Polynesia indicated by mtDNA from the Pacific rat. *Proc. Natl. Acad. Sci. U.S.A* 95, 15145–15150 (1998). [PubMed: 9844030]
115. Fehren-Schmitz L. et al. Genetic ancestry of Rapanui before and after European contact. *Current Biology* 27, 3209–3215 (2017). [PubMed: 29033334]
116. Crowe A. *Pathway of the Birds*. (University of Hawai'i Press, 2018).
117. Marck JC *Topics in Polynesian Language and Culture History*. (Pacific Linguistics, Research School of Pacific and Asian Studies, The Australian National University, 2000). doi:10.15144/PL-504




Fig. 1. Dimensionality reduction of genetic variation in Pacific islanders.

(a-c) Ancestry-specific PCA of islanders (with non-Asian derived ancestries, such as post-colonial European ancestry, masked) shows islands (a) diverging separately along each component (b, c), due to the independence of genetic drift from each island's founder effect. Neither geography, nor settlement sequence can be discerned. The westernmost islands are omitted, as their greater diversity would otherwise dominate the first principal component (see Supplementary Fig. 2). The percent variance explained by each of the first four principal component dimensions is listed along each axis. Dots represent individuals, and colors represent islands. (d) Ancestry specific tSNE plot of all sampled islanders, providing superior separation of each island group. The ancestral western Pacific islands are on the left and the easternmost Polynesian island (Rapa Nui) on the right. Important patterns are now evident; for instance, Rarotonga and the Palliser group appear at the center of the eastern Polynesian islands while the other eastern islands radiate out from them, consistent with the settlement patterns we infer below. tSNE preserves local relationships, but not global relationships (between widely separated clusters).




Fig. 2. Serial bottlenecks and relatedness define the settlement sequence and timings for the Polynesian islands.

(a) Inferred genetic-based map of Polynesian origins for the islands sampled in our study (not to scale). The direction, line width, and date for each arrow are based on inter-island statistics as described in the key and the text. For example, the widths of the arrows are inversely proportional to the value of the range expansion statistic (ψ) relative to Samoa. The order of arrow divergences indicates the order of shared drift amongst the child populations. Where they occur, these shared paths may indicate that one or more intermediate islands in the settlement sequence are missing from our dataset (Extended Data Fig. 5). This settlement sequence is consistent with a principal curve analysis (Extended Data Fig. 7). A sex-averaged generation time of 30 years was used, as found in several studies of pre-industrial populations (see Supplementary Discussion ‘On generation times and meiosis events’). Locations with prehistoric remains of megalithic statue building are also indicated (red asterisk). (b) The range expansion statistic (ψ) shows a steady increase in retained rare variant frequencies (genetic surfing) along paths of settlement as a result of each successive founder effect. Note that each matrix element is computed on a different SNP set (rare variants found in some samples from both islands), so the matrix need not have a similar ordering across all rows or all columns—that it does is a confirmation of the range expansion process. Rapa Nui (Easter Island) is the easternmost island in our dataset with the most compounded series of founder effects. (c) Example IBD segment length distributions for all pairs of individuals, one on Rapa Nui and the other on Mangareva (green), Palliser (blue), Rarotonga (red), and Samoa (black), used to fit the respective exponential decay constants (λ).