

# Lawrence Berkeley National Laboratory

## Recent Work

### Title

DEHYDROPEPTIDE PRODUCTION IN THE RADIOLYSIS OF SOLID PEPTIDES: A RADIOLYTIC SYNTHESIS OF PYRAZINE DERIVATIVES FROM CYCLIC DIPEPTIDES

### Permalink

<https://escholarship.org/uc/item/8r32m0rm>

### Authors

Kland-English, Mathilde  
Garrison, Warren M.

### Publication Date

1962-09-21

**University of California**  
**Ernest O. Lawrence**  
**Radiation Laboratory**

**TWO-WEEK LOAN COPY**

*This is a Library Circulating Copy  
which may be borrowed for two weeks.  
For a personal retention copy, call  
Tech. Info. Division, Ext. 5545*

**Berkeley, California**

## **DISCLAIMER**

This document was prepared as an account of work sponsored by the United States Government. While this document is believed to contain correct information, neither the United States Government nor any agency thereof, nor the Regents of the University of California, nor any of their employees, makes any warranty, express or implied, or assumes any legal responsibility for the accuracy, completeness, or usefulness of any information, apparatus, product, or process disclosed, or represents that its use would not infringe privately owned rights. Reference herein to any specific commercial product, process, or service by its trade name, trademark, manufacturer, or otherwise, does not necessarily constitute or imply its endorsement, recommendation, or favoring by the United States Government or any agency thereof, or the Regents of the University of California. The views and opinions of authors expressed herein do not necessarily state or reflect those of the United States Government or any agency thereof or the Regents of the University of California.

UNIVERSITY OF CALIFORNIA  
Lawrence Radiation Laboratory  
Berkeley, California  
Contract No. W-7405-eng-48

DEHYDROPEPTIDE PRODUCTION IN THE RADIOLYSIS OF SOLID PEPTIDES:  
A RADIOLYTIC SYNTHESIS OF PYRAZINE DERIVATIVES  
FROM CYCLIC DIPEPTIDES

Mathilde Kland-English and Warren M. Garrison

September 21, 1962

DEHYDROPEPTIDE PRODUCTION IN THE RADIOLYSIS OF SOLID PEPTIDES:  
A RADIOLYTIC SYNTHESIS OF PYRAZINE DERIVATIVES  
FROM CYCLIC DIPEPTIDES.

Mathilde Kland-English and Warren M. Garrison

Lawrence Radiation Laboratory  
University of California, Berkeley, California

September 21, 1962


Among the chemical changes observed on radiolysis of compounds containing the peptide bond are those attributed<sup>1-3</sup> to the production of unsaturated linkages which may be represented in terms of the tautomeric structures,  $\sim\text{CON}=\text{C}(\text{CH}_2\text{R})\sim \rightleftharpoons \sim\text{CONH}-\text{C}(=\text{CHR})\sim$ . Compounds containing this type of chemical function have been designated dehydropeptides; such compounds are stable in water yet are readily hydrolyzed to yield ammonia and a carbonyl product on heating in dilute mineral acid,  $\sim\text{CON}=\text{C}(\text{CH}_2\text{R}) + 2\text{H}_2\text{O} \rightarrow \sim\text{COOH} + \text{NH}_3 + \text{RCH}_2\text{CO}\sim$ .<sup>4</sup> The experimental evidence for production of dehydropeptides in the radiolysis of proteins and simpler peptide systems has recently been reviewed.<sup>5</sup> Identification has been based primarily on studies of the formation of ammonia and carbonyl function during hydrolysis of the irradiated peptide system. For example, the hydrolytic production of ammonia and pyruvic acid from  $\gamma$ -irradiated acetylalanine (solid, evacuated) has been found to parallel very closely the hydrolytic degradation of authentic acetyldehydroalanine.

The cyclic dipeptides, such as glycine anhydride and alanine anhydride also show an analogous post-irradiation production of ammonia and  $\alpha$ -keto acid during acid hydrolysis.<sup>6</sup> We note too that the dehydropeptide derivatives of these cyclic compounds represent isomeric forms of the 2,5-dihydroxypyrazines. Quite recently, Sorkin<sup>7</sup> has given evidence that the dehydro derivative of glycylalanine anhydride, that is, 3-methylene-2,5-diketopiperazine (prepared

by the classical chemical method of Bergman and Miekely<sup>8</sup>), undergoes isomerization in alkaline solution to give 3-methyl-2,5-dihydroxypyrazine. We report here analogous evidence for the formation of pyrazine derivatives through  $\gamma$ -radiolysis of the amino acid anhydrides followed by dissolution of the irradiated solids in alkaline solution.

Glycylalanine anhydride was synthesized after the method of Fischer and Scheibler.<sup>9</sup> Glycine anhydride and alanine anhydride were obtained from the Nutritional Biochemicals Corp. The anhydrides were recrystallized three times from water, dried, and ground to a microcrystalline powder. Samples were evacuated on the vacuum line in pyrex ampoules for a minimum of 48 hours. After irradiation ( $\text{Co}^{60}$   $\gamma$ -rays,  $1.7 \times 10^{18}$  ev/gm/min), the ampoules were opened to the atmosphere and the contents were dissolved to a concentration of 0.1 M in 0.02 N sodium hydroxide under nitrogen aeration. Complete dissolution required two to three minutes. Changes in the absorption spectra of the oxygen-free solutions were followed differentially (against a 0.1 M solution of unirradiated anhydride) over the range 300 m $\mu$  to 400 m $\mu$ .

Solutions of irradiated glycylalanine anhydride (3-methyl-2,5-diketopiperazine) show a rapid increase in optical density with time over the 300-400 m $\mu$  range with  $\lambda_{\text{max}} = 370$  m $\mu$ . The spectrum is completely developed after about two hours. Authentic 3-methylene-2,5-diketopiperazine<sup>8</sup> was found to exhibit almost identical spectrophotometric properties. The fully developed spectra obtained with irradiated glycylalanine anhydride and authentic 3-methylene-2,5-diketopiperazine in oxygen-free .02 N sodium hydroxide solution are shown in Fig. 1. Appearance of the absorption maximum at 370 m $\mu$  is attributed to the formation of the dihydroxypyrazine nucleus (II) via the isomerization reaction:



The absorption spectrum of an authentic dihydroxypyrazine (3,6-dimethyl) prepared after Karmas and Spoerri<sup>10</sup> is included in Fig. 1 for purposes of comparison. It is noted that the dihydroxypyrazines in alkaline solution are rapidly destroyed on exposure to air.

Isolation of 3-methylene-2,5-diketopiperazine (Ia, Ib) from irradiated glycylalanine anhydride has been accomplished. A methanolic extract of the irradiated solid was transferred to filter paper and chromatographed in parallel with authentic 3-methylene-2,5-diketopiperazine; butanol saturated with water was used as the developing liquid. The authentic methylene derivative is readily detected at  $R_f \approx 0.4$  under ultraviolet illumination; the irradiated glycylalanine anhydride also gives a strongly absorbing component at this  $R_f$  value. The two areas were separately eluted and examined spectrophotometrically in oxygen-free 0.02 N sodium hydroxide. The characteristic growth in the 300 to 400  $\mu$  region was observed in both cases and the final curves were essentially identical to those given in Fig. 1. The yield of 3-methylene-2,5-diketopiperazine from glycylalanine anhydride at the lowest dosage studied ( $5 \times 10^{20}$  ev/gm) corresponds to a G value of  $\approx 0.55$ .


The results obtained with alanine anhydride appear to be somewhat more complicated. In this case the development of the absorption curve is accompanied by a gradual shift in  $\lambda_{\max}$  from 370 m $\mu$  to 360 m $\mu$ . However, the spectra obtained within 15 minutes of the time of dissolution coincide with that given by authentic 2,5-dihydroxy-3,6-dimethylpyrazine.

Solutions prepared from irradiated glycine anhydride show the characteristic dihydroxypyrazine spectrum and on standing there is no detectable shift in  $\lambda_{\max}$  from 370 m $\mu$  to shorter wavelengths. However, glycine anhydride differs from both glycylalanine and alanine anhydrides in that the absorption spectrum shown in Fig. 1 is fully developed within the two to three minute period required for dissolution. This behavior may be interpreted as evidence for rapid isomerization of dehydroglycine anhydride which must necessarily have only the endocyclic form as given by structure Ib.

Finally we should like to comment on the fact that all three solid anhydrides develop a yellow coloration which is visually detectable at radiation dosages as low as  $\sim 5 \times 10^{19}$  ev/gm. The effect is most pronounced in the case of glycine anhydride. Coloration is not appreciably diminished by exposure to oxygen and remains visible in water extracts of the irradiated solids. This coloration then, cannot be ascribed to an optical absorption by free radicals or ionic transients. It is noted that the authentic dihydroxypyrazines are yellow in the solid state and, although we have made only preliminary spectrophotometric studies of the irradiated solid anhydrides, the available evidence suggests that isomerization of dehydropeptides occurs to some extent in the solid state. A mechanism for the radiation-induced isomerization of the  $\sim\text{CO-NH}_2$  linkage in solid peptide systems has recently been described.<sup>5</sup> A detailed report of the various aspects of the radiation chemistry of the cyclic dipeptides will be forthcoming.

REFERENCES

1. M. E. Jayko and W. M. Garrison, J. Chem. Phys., 25, 1084 (1956).
2. A. Caputo and K. Dose, Z. Naturforschung. 12b, 172 (1957).
3. W. M. Garrison, M. E. Jayko and W. Bennett, Radiation Research 16, 483 (1962).
4. J. P. Greenstein and M. Winitz, Chemistry of the Amino Acids, John Wiley and Sons Inc., 1961.
5. W. M. Garrison and B. M. Weeks, Radiation Research 17, 000 (1962).
6. M. Kland-English and W. M. Garrison, Lawrence Radiation Laboratory Report UCRL-8988 (Dec. 1959).
7. H. Sorkin, Isomerization of Methylene Diketopiperazines, Ph.D. Dissertation, Cornell University (1960).
8. M. Bergmann and A. Miekely, Z. Physiol. Chem., 140, 128 (1924).
9. E. Fischer and H. Scheibler, Ann., 363, 136 (1908).
10. G. Karmas and R. E. Spoerri, J. Am. Chem. Soc., 79, 680 (1957).



MU-28088

Fig. 1. Absorption spectra of irradiated cyclic dipeptides and authentic dehydropeptides in .02N sodium hydroxide 0, irradiated glycine anhydride; □ authentic 3-methylene-2,5-diketopiperazine; ● authentic 2,5-dihydroxy-3,6-dimethylpyrazine.

This report was prepared as an account of Government sponsored work. Neither the United States, nor the Commission, nor any person acting on behalf of the Commission:

- A. Makes any warranty or representation, expressed or implied, with respect to the accuracy, completeness, or usefulness of the information contained in this report, or that the use of any information, apparatus, method, or process disclosed in this report may not infringe privately owned rights; or
- B. Assumes any liabilities with respect to the use of, or for damages resulting from the use of any information, apparatus, method, or process disclosed in this report.

As used in the above, "person acting on behalf of the Commission" includes any employee or contractor of the Commission, or employee of such contractor, to the extent that such employee or contractor of the Commission, or employee of such contractor prepares, disseminates, or provides access to, any information pursuant to his employment or contract with the Commission, or his employment with such contractor.

