

UC Berkeley

UC Berkeley Previously Published Works

Title

Observation of Decays $B_0 \rightarrow D_s(^*) + \pi^-$ and $B_0 \rightarrow D_s(^*) - K^+$

Permalink

<https://escholarship.org/uc/item/8r63g6j5>

Journal

Physical Review Letters, 98(8)

ISSN

0031-9007

Authors

Aubert, B
Barate, R
Bona, M
[et al.](#)

Publication Date

2007-02-23

DOI

10.1103/physrevlett.98.081801

Copyright Information

This work is made available under the terms of a Creative Commons Attribution License, available at <https://creativecommons.org/licenses/by/4.0/>

Peer reviewed

Observation of Decays $B^0 \rightarrow D_s^{(*)+} \pi^-$ and $B^0 \rightarrow D_s^{(*)-} K^+$

B. Aubert,¹ R. Barate,¹ M. Bona,¹ D. Boutigny,¹ F. Couderc,¹ Y. Karyotakis,¹ J. P. Lees,¹ V. Poireau,¹ V. Tisserand,¹ A. Zghiche,¹ E. Grauges,² A. Palano,³ J. C. Chen,⁴ N. D. Qi,⁴ G. Rong,⁴ P. Wang,⁴ Y. S. Zhu,⁴ G. Eigen,⁵ I. Ofte,⁵ B. Stugu,⁵ G. S. Abrams,⁶ M. Battaglia,⁶ D. N. Brown,⁶ J. Button-Shafer,⁶ R. N. Cahn,⁶ E. Charles,⁶ M. S. Gill,⁶ Y. Groysman,⁶ R. G. Jacobsen,⁶ J. A. Kadyk,⁶ L. T. Kerth,⁶ Yu. G. Kolomensky,⁶ G. Kukartsev,⁶ G. Lynch,⁶ L. M. Mir,⁶ P. J. Oddone,⁶ T. J. Orimoto,⁶ M. Pripstein,⁶ N. A. Roe,⁶ M. T. Ronan,⁶ A. Suzuki,⁶ D. Troost,⁶ W. A. Wenzel,⁶ M. Barrett,⁷ K. E. Ford,⁷ T. J. Harrison,⁷ A. J. Hart,⁷ C. M. Hawkes,⁷ S. E. Morgan,⁷ A. T. Watson,⁷ K. Goetzen,⁸ T. Held,⁸ H. Koch,⁸ B. Lewandowski,⁸ M. Pelizaeus,⁸ K. Peters,⁸ T. Schroeder,⁸ M. Steinke,⁸ J. T. Boyd,⁹ J. P. Burke,⁹ W. N. Cottingham,⁹ D. Walker,⁹ T. Cuhadar-Donszelmann,¹⁰ B. G. Fulson,¹⁰ C. Hearty,¹⁰ N. S. Knecht,¹⁰ T. S. Mattison,¹⁰ J. A. McKenna,¹⁰ A. Khan,¹¹ P. Kyberd,¹¹ M. Saleem,¹¹ L. Teodorescu,¹¹ V. E. Blinov,¹² A. D. Bukin,¹² V. P. Druzhinin,¹² V. B. Golubev,¹² A. P. Onuchin,¹² S. I. Serednyakov,¹² Yu. I. Skovpen,¹² E. P. Solodov,¹² K. Yu. Todyshev,¹² D. S. Best,¹³ M. Bondioli,¹³ M. Bruinsma,¹³ M. Chao,¹³ S. Curry,¹³ I. Eschrich,¹³ D. Kirkby,¹³ A. J. Lankford,¹³ P. Lund,¹³ M. Mandelkern,¹³ R. K. Mommsen,¹³ W. Roethel,¹³ D. P. Stoker,¹³ S. Abachi,¹⁴ C. Buchanan,¹⁴ S. D. Foulkes,¹⁵ J. W. Gary,¹⁵ O. Long,¹⁵ B. C. Shen,¹⁵ K. Wang,¹⁵ L. Zhang,¹⁵ H. K. Hadavand,¹⁶ E. J. Hill,¹⁶ H. P. Paar,¹⁶ S. Rahatlou,¹⁶ V. Sharma,¹⁶ J. W. Berryhill,¹⁷ C. Campagnari,¹⁷ A. Cunha,¹⁷ B. Dahmes,¹⁷ T. M. Hong,¹⁷ D. Kovalskyi,¹⁷ J. D. Richman,¹⁷ T. W. Beck,¹⁸ A. M. Eisner,¹⁸ C. J. Flacco,¹⁸ C. A. Heusch,¹⁸ J. Kroseberg,¹⁸ W. S. Lockman,¹⁸ G. Nesom,¹⁸ T. Schalk,¹⁸ B. A. Schumm,¹⁸ A. Seiden,¹⁸ P. Spradlin,¹⁸ D. C. Williams,¹⁸ M. G. Wilson,¹⁸ J. Albert,¹⁹ E. Chen,¹⁹ A. Dvoretiskii,¹⁹ D. G. Hitlin,¹⁹ I. Narsky,¹⁹ T. Piatenko,¹⁹ F. C. Porter,¹⁹ A. Ryd,¹⁹ A. Samuel,¹⁹ R. Andreassen,²⁰ G. Mancinelli,²⁰ B. T. Meadows,²⁰ M. D. Sokoloff,²⁰ F. Blanc,²¹ P. C. Bloom,²¹ S. Chen,²¹ W. T. Ford,²¹ J. F. Hirschauer,²¹ A. Kreisel,²¹ U. Nauenberg,²¹ A. Olivas,²¹ W. O. Ruddick,²¹ J. G. Smith,²¹ K. A. Ulmer,²¹ S. R. Wagner,²¹ J. Zhang,²¹ A. Chen,²² E. A. Eckhart,²² A. Soffer,²² W. H. Toki,²² R. J. Wilson,²² F. Winklmeier,²² Q. Zeng,²² D. D. Altenburg,²³ E. Feltresi,²³ A. Hauke,²³ H. Jasper,²³ B. Spaan,²³ T. Brandt,²⁴ V. Klose,²⁴ H. M. Lacker,²⁴ W. F. Mader,²⁴ R. Nogowski,²⁴ A. Petzold,²⁴ J. Schubert,²⁴ K. R. Schubert,²⁴ R. Schwierz,²⁴ J. E. Sundermann,²⁴ A. Volk,²⁴ D. Bernard,²⁵ G. R. Bonneaud,²⁵ P. Grenier,^{25,*} E. Latour,²⁵ Ch. Thiebaux,²⁵ M. Verderi,²⁵ D. J. Bard,²⁶ P. J. Clark,²⁶ W. Gradl,²⁶ F. Muheim,²⁶ S. Playfer,²⁶ A. I. Robertson,²⁶ Y. Xie,²⁶ M. Andreotti,²⁷ D. Bettoni,²⁷ C. Bozzi,²⁷ R. Calabrese,²⁷ G. Cibinetto,²⁷ E. Lippi,²⁷ M. Negrini,²⁷ A. Petrella,²⁷ L. Piemontese,²⁷ E. Prencipe,²⁷ F. Anulli,²⁸ R. Baldini-Ferrolli,²⁸ A. Calcaterra,²⁸ R. de Sangro,²⁸ G. Finocchiaro,²⁸ S. Pacetti,²⁸ P. Patteri,²⁸ I. M. Peruzzi,^{28,†} M. Piccolo,²⁸ M. Rama,²⁸ A. Zallo,²⁸ A. Buzzo,²⁹ R. Capra,²⁹ R. Contri,²⁹ M. Lo Vetere,²⁹ M. M. Macri,²⁹ M. R. Monge,²⁹ S. Passaggio,²⁹ C. Patrignani,²⁹ E. Robutti,²⁹ A. Santroni,²⁹ S. Tosi,²⁹ G. Brandenburg,³⁰ K. S. Chaisanguanthum,³⁰ M. Morii,³⁰ J. Wu,³⁰ R. S. Dubitzky,³¹ J. Marks,³¹ S. Schenk,³¹ U. Uwer,³¹ W. Bhimji,³² D. A. Bowerman,³² P. D. Dauncey,³² U. Egede,³² R. L. Flack,³² J. R. Gaillard,³² J. A. Nash,³² M. B. Nikolich,³² W. Panduro Vazquez,³² X. Chai,³³ M. J. Charles,³³ U. Mallik,³³ N. T. Meyer,³³ V. Ziegler,³³ J. Cochran,³⁴ H. B. Crawley,³⁴ L. Dong,³⁴ V. Eyges,³⁴ W. T. Meyer,³⁴ S. Prell,³⁴ E. I. Rosenberg,³⁴ A. E. Rubin,³⁴ A. V. Gritsan,³⁵ M. Fritsch,³⁶ G. Schott,³⁶ N. Arnaud,³⁷ M. Davier,³⁷ G. Grosdidier,³⁷ A. Höcker,³⁷ F. Le Diberder,³⁷ V. Lepeltier,³⁷ A. M. Lutz,³⁷ A. Oyanguren,³⁷ S. Pruvot,³⁷ S. Rodier,³⁷ P. Roudeau,³⁷ M. H. Schune,³⁷ A. Stocchi,³⁷ W. F. Wang,³⁷ G. Wormser,³⁷ C. H. Cheng,³⁸ D. J. Lange,³⁸ D. M. Wright,³⁸ C. A. Chavez,³⁹ I. J. Forster,³⁹ J. R. Fry,³⁹ E. Gabathuler,³⁹ R. Gamet,³⁹ K. A. George,³⁹ D. E. Hutchcroft,³⁹ D. J. Payne,³⁹ K. C. Schofield,³⁹ C. Touramanis,³⁹ A. J. Bevan,⁴⁰ F. Di Lodovico,⁴⁰ W. Menges,⁴⁰ R. Sacco,⁴⁰ C. L. Brown,⁴¹ G. Cowan,⁴¹ H. U. Flaecher,⁴¹ D. A. Hopkins,⁴¹ P. S. Jackson,⁴¹ T. R. McMahon,⁴¹ S. Ricciardi,⁴¹ F. Salvatore,⁴¹ D. N. Brown,⁴² C. L. Davis,⁴² J. Allison,⁴³ N. R. Barlow,⁴³ R. J. Barlow,⁴³ Y. M. Chia,⁴³ C. L. Edgar,⁴³ M. P. Kelly,⁴³ G. D. Lafferty,⁴³ M. T. Naisbit,⁴³ J. C. Williams,⁴³ J. I. Yi,⁴³ C. Chen,⁴⁴ W. D. Hulsbergen,⁴⁴ A. Jawahery,⁴⁴ C. K. Lae,⁴⁴ D. A. Roberts,⁴⁴ G. Simi,⁴⁴ G. Blaylock,⁴⁵ C. Dallapiccola,⁴⁵ S. S. Hertzbach,⁴⁵ X. Li,⁴⁵ T. B. Moore,⁴⁵ S. Saremi,⁴⁵ H. Staengle,⁴⁵ S. Y. Willocq,⁴⁵ R. Cowan,⁴⁶ K. Koeneke,⁴⁶ G. Sciolla,⁴⁶ S. J. Sekula,⁴⁶ M. Spitznagel,⁴⁶ F. Taylor,⁴⁶ R. K. Yamamoto,⁴⁶ H. Kim,⁴⁷ P. M. Patel,⁴⁷ S. H. Robertson,⁴⁷ A. Lazzaro,⁴⁸ V. Lombardo,⁴⁸ F. Palombo,⁴⁸ J. M. Bauer,⁴⁹ L. Cremaldi,⁴⁹ V. Eschenburg,⁴⁹ R. Godang,⁴⁹ R. Kroeger,⁴⁹ J. Reidy,⁴⁹ D. A. Sanders,⁴⁹ D. J. Summers,⁴⁹ H. W. Zhao,⁴⁹ S. Brunet,⁵⁰ D. Côté,⁵⁰ P. Taras,⁵⁰ F. B. Viaud,⁵⁰ H. Nicholson,⁵¹ N. Cavallo,^{52,‡} G. De Nardo,⁵² D. del Re,⁵² F. Fabozzi,^{52,‡} C. Gatto,⁵² L. Lista,⁵² D. Monorchio,⁵² P. Paolucci,⁵² D. Piccolo,⁵² C. Sciacca,⁵² M. Baak,⁵³ H. Bulten,⁵³ G. Raven,⁵³ H. L. Snoek,⁵³ C. P. Jessop,⁵⁴ J. M. LoSecco,⁵⁴ T. Allmendinger,⁵⁵ G. Benelli,⁵⁵ K. K. Gan,⁵⁵ K. Honscheid,⁵⁵ D. Hufnagel,⁵⁵ P. D. Jackson,⁵⁵ H. Kagan,⁵⁵ R. Kass,⁵⁵ T. Pulliam,⁵⁵ A. M. Rahimi,⁵⁵ R. Ter-Antonyan,⁵⁵ Q. K. Wong,⁵⁵ N. L. Blount,⁵⁶ J. Brau,⁵⁶ R. Frey,⁵⁶ O. Igonkina,⁵⁶ M. Lu,⁵⁶ C. T. Potter,⁵⁶ R. Rahmat,⁵⁶ N. B. Sinev,⁵⁶ D. Strom,⁵⁶ J. Strube,⁵⁶

E. Torrence,⁵⁶ F. Galeazzi,⁵⁷ A. Gaz,⁵⁷ M. Margoni,⁵⁷ M. Morandin,⁵⁷ A. Pompili,⁵⁷ M. Posocco,⁵⁷ M. Rotondo,⁵⁷ F. Simonetto,⁵⁷ R. Stroili,⁵⁷ C. Voci,⁵⁷ M. Benayoun,⁵⁸ J. Chauveau,⁵⁸ P. David,⁵⁸ L. Del Buono,⁵⁸ Ch. de la Vaissière,⁵⁸ O. Hamon,⁵⁸ B. L. Hartfiel,⁵⁸ M. J. J. John,⁵⁸ J. Malclès,⁵⁸ J. Ocariz,⁵⁸ L. Roos,⁵⁸ G. Therin,⁵⁸ P. K. Behera,⁵⁹ L. Gladney,⁵⁹ J. Panetta,⁵⁹ M. Biasini,⁶⁰ R. Covarelli,⁶⁰ M. Pioppi,⁶⁰ C. Angelini,⁶¹ G. Batignani,⁶¹ S. Bettarini,⁶¹ F. Bucci,⁶¹ G. Calderini,⁶¹ M. Carpinelli,⁶¹ R. Cenci,⁶¹ F. Forti,⁶¹ M. A. Giorgi,⁶¹ A. Lusiani,⁶¹ G. Marchiori,⁶¹ M. A. Mazur,⁶¹ M. Morganti,⁶¹ N. Neri,⁶¹ E. Paoloni,⁶¹ G. Rizzo,⁶¹ J. Walsh,⁶¹ M. Haire,⁶² D. Judd,⁶² D. E. Wagoner,⁶² J. Biesiada,⁶³ N. Danielson,⁶³ P. Elmer,⁶³ Y. P. Lau,⁶³ C. Lu,⁶³ J. Olsen,⁶³ A. J. S. Smith,⁶³ A. V. Telnov,⁶³ F. Bellini,⁶⁴ G. Cavoto,⁶⁴ A. D'Orazio,⁶⁴ E. Di Marco,⁶⁴ R. Faccini,⁶⁴ F. Ferrarotto,⁶⁴ F. Ferroni,⁶⁴ M. Gaspero,⁶⁴ L. Li Gioi,⁶⁴ M. A. Mazzoni,⁶⁴ S. Morganti,⁶⁴ G. Piredda,⁶⁴ F. Polci,⁶⁴ F. Safai Tehrani,⁶⁴ C. Voena,⁶⁴ M. Ebert,⁶⁵ H. Schröder,⁶⁵ R. Waldi,⁶⁵ T. Adye,⁶⁶ N. De Groot,⁶⁶ B. Franek,⁶⁶ E. O. Olaiya,⁶⁶ F. F. Wilson,⁶⁶ S. Emery,⁶⁷ A. Gaidot,⁶⁷ S. F. Ganzhur,⁶⁷ G. Hamel de Monchenault,⁶⁷ W. Kozanecki,⁶⁷ M. Legendre,⁶⁷ G. Vasseur,⁶⁷ Ch. Yèche,⁶⁷ M. Zito,⁶⁷ W. Park,⁶⁸ M. V. Purohit,⁶⁸ J. R. Wilson,⁶⁸ M. T. Allen,⁶⁹ D. Aston,⁶⁹ R. Bartoldus,⁶⁹ P. Bechtler,⁶⁹ N. Berger,⁶⁹ A. M. Boyarski,⁶⁹ R. Claus,⁶⁹ J. P. Coleman,⁶⁹ M. R. Convery,⁶⁹ M. Cristinziani,⁶⁹ J. C. Dingfelder,⁶⁹ D. Dong,⁶⁹ J. Dorfan,⁶⁹ G. P. Dubois-Felsmann,⁶⁹ D. Dujmic,⁶⁹ W. Dunwoodie,⁶⁹ R. C. Field,⁶⁹ T. Glanzman,⁶⁹ S. J. Gowdy,⁶⁹ M. T. Graham,⁶⁹ V. Halyo,⁶⁹ C. Hast,⁶⁹ T. Hryn'ova,⁶⁹ W. R. Innes,⁶⁹ M. H. Kelsey,⁶⁹ P. Kim,⁶⁹ M. L. Kocian,⁶⁹ D. W. G. S. Leith,⁶⁹ S. Li,⁶⁹ J. Libby,⁶⁹ S. Luitz,⁶⁹ V. Luth,⁶⁹ H. L. Lynch,⁶⁹ D. B. MacFarlane,⁶⁹ H. Marsiske,⁶⁹ R. Messner,⁶⁹ D. R. Muller,⁶⁹ C. P. O'Grady,⁶⁹ V. E. Ozcan,⁶⁹ M. Perl,⁶⁹ A. Perazzo,⁶⁹ B. N. Ratcliff,⁶⁹ A. Roodman,⁶⁹ A. A. Salnikov,⁶⁹ R. H. Schindler,⁶⁹ J. Schwiening,⁶⁹ A. Snyder,⁶⁹ J. Stelzer,⁶⁹ D. Su,⁶⁹ M. K. Sullivan,⁶⁹ K. Suzuki,⁶⁹ S. K. Swain,⁶⁹ J. M. Thompson,⁶⁹ J. Va'vra,⁶⁹ N. van Bakel,⁶⁹ M. Weaver,⁶⁹ A. J. R. Weinstein,⁶⁹ W. J. Wisniewski,⁶⁹ M. Wittgen,⁶⁹ D. H. Wright,⁶⁹ A. K. Yarritu,⁶⁹ K. Yi,⁶⁹ C. C. Young,⁶⁹ P. R. Burchat,⁷⁰ A. J. Edwards,⁷⁰ S. A. Majewski,⁷⁰ B. A. Petersen,⁷⁰ C. Roat,⁷⁰ L. Wilden,⁷⁰ S. Ahmed,⁷¹ M. S. Alam,⁷¹ R. Bula,⁷¹ J. A. Ernst,⁷¹ V. Jain,⁷¹ B. Pan,⁷¹ M. A. Saeed,⁷¹ F. R. Wappler,⁷¹ S. B. Zain,⁷¹ W. Bugg,⁷² M. Krishnamurthy,⁷² S. M. Spanier,⁷² R. Eckmann,⁷³ J. L. Ritchie,⁷³ A. Satpathy,⁷³ C. J. Schilling,⁷³ R. F. Schwitters,⁷³ J. M. Izen,⁷⁴ I. Kitayama,⁷⁴ X. C. Lou,⁷⁴ S. Ye,⁷⁴ F. Bianchi,⁷⁵ F. Gallo,⁷⁵ D. Gamba,⁷⁵ M. Bomben,⁷⁶ L. Bosisio,⁷⁶ C. Cartaro,⁷⁶ F. Cossutti,⁷⁶ G. Della Ricca,⁷⁶ S. Dittongo,⁷⁶ S. Grancagnolo,⁷⁶ L. Lanceri,⁷⁶ L. Vitale,⁷⁶ V. Azzolini,⁷⁷ F. Martinez-Vidal,⁷⁷ Sw. Banerjee,⁷⁸ B. Bhuyan,⁷⁸ C. M. Brown,⁷⁸ D. Fortin,⁷⁸ K. Hamano,⁷⁸ R. Kowalewski,⁷⁸ I. M. Nugent,⁷⁸ J. M. Roney,⁷⁸ R. J. Sobie,⁷⁸ J. J. Back,⁷⁹ P. F. Harrison,⁷⁹ T. E. Latham,⁷⁹ G. B. Mohanty,⁷⁹ M. Pappagallo,⁷⁹ H. R. Band,⁸⁰ X. Chen,⁸⁰ B. Cheng,⁸⁰ S. Dasu,⁸⁰ M. Datta,⁸⁰ A. M. Eichenbaum,⁸⁰ K. T. Flood,⁸⁰ J. J. Hollar,⁸⁰ P. E. Kutter,⁸⁰ H. Li,⁸⁰ R. Liu,⁸⁰ B. Mellado,⁸⁰ A. Mihalji,⁸⁰ A. K. Mohapatra,⁸⁰ Y. Pan,⁸⁰ M. Pierini,⁸⁰ R. Prepost,⁸⁰ P. Tan,⁸⁰ S. L. Wu,⁸⁰ Z. Yu,⁸⁰ and H. Neal⁸¹

(The BABAR Collaboration)

¹Laboratoire de Physique des Particules, F-74941 Annecy-le-Vieux, France

²Universitat de Barcelona, Facultat de Fisica Dept. ECM, E-08028 Barcelona, Spain

³Università di Bari, Dipartimento di Fisica and INFN, I-70126 Bari, Italy

⁴Institute of High Energy Physics, Beijing 100039, China

⁵University of Bergen, Institute of Physics, N-5007 Bergen, Norway

⁶Lawrence Berkeley National Laboratory and University of California, Berkeley, California 94720, USA

⁷University of Birmingham, Birmingham, B15 2TT, United Kingdom

⁸Ruhr Universität Bochum, Institut für Experimentalphysik I, D-44780 Bochum, Germany

⁹University of Bristol, Bristol BS8 1TL, United Kingdom

¹⁰University of British Columbia, Vancouver, British Columbia, Canada V6T 1Z1

¹¹Brunel University, Uxbridge, Middlesex UB8 3PH, United Kingdom

¹²Budker Institute of Nuclear Physics, Novosibirsk 630090, Russia

¹³University of California at Irvine, Irvine, California 92697, USA

¹⁴University of California at Los Angeles, Los Angeles, California 90024, USA

¹⁵University of California at Riverside, Riverside, California 92521, USA

¹⁶University of California at San Diego, La Jolla, California 92093, USA

¹⁷University of California at Santa Barbara, Santa Barbara, California 93106, USA

¹⁸University of California at Santa Cruz, Institute for Particle Physics, Santa Cruz, California 95064, USA

¹⁹California Institute of Technology, Pasadena, California 91125, USA

²⁰University of Cincinnati, Cincinnati, Ohio 45221, USA

²¹University of Colorado, Boulder, Colorado 80309, USA

²²Colorado State University, Fort Collins, Colorado 80523, USA

²³Universität Dortmund, Institut für Physik, D-44221 Dortmund, Germany

- ²⁴*Technische Universität Dresden, Institut für Kern- und Teilchenphysik, D-01062 Dresden, Germany*
- ²⁵*Ecole Polytechnique, LLR, F-91128 Palaiseau, France*
- ²⁶*University of Edinburgh, Edinburgh EH9 3JZ, United Kingdom*
- ²⁷*Università di Ferrara, Dipartimento di Fisica and INFN, I-44100 Ferrara, Italy*
- ²⁸*Laboratori Nazionali di Frascati dell'INFN, I-00044 Frascati, Italy*
- ²⁹*Università di Genova, Dipartimento di Fisica and INFN, I-16146 Genova, Italy*
- ³⁰*Harvard University, Cambridge, Massachusetts 02138, USA*
- ³¹*Universität Heidelberg, Physikalisches Institut, Philosophenweg 12, D-69120 Heidelberg, Germany*
- ³²*Imperial College London, London, SW7 2AZ, United Kingdom*
- ³³*University of Iowa, Iowa City, Iowa 52242, USA*
- ³⁴*Iowa State University, Ames, Iowa 50011-3160, USA*
- ³⁵*Johns Hopkins University, Baltimore, Maryland 21218, USA*
- ³⁶*Universität Karlsruhe, Institut für Experimentelle Kernphysik, D-76021 Karlsruhe, Germany*
- ³⁷*Laboratoire de l'Accélérateur Linéaire, IN2P3-CNRS et Université Paris-Sud 11, Centre Scientifique d'Orsay, B.P. 34, F-91898 ORSAY Cedex, France*
- ³⁸*Lawrence Livermore National Laboratory, Livermore, California 94550, USA*
- ³⁹*University of Liverpool, Liverpool L69 7ZE, United Kingdom*
- ⁴⁰*Queen Mary, University of London, E1 4NS, United Kingdom*
- ⁴¹*University of London, Royal Holloway and Bedford New College, Egham, Surrey TW20 0EX, United Kingdom*
- ⁴²*University of Louisville, Louisville, Kentucky 40292, USA*
- ⁴³*University of Manchester, Manchester M13 9PL, United Kingdom*
- ⁴⁴*University of Maryland, College Park, Maryland 20742, USA*
- ⁴⁵*University of Massachusetts, Amherst, Massachusetts 01003, USA*
- ⁴⁶*Massachusetts Institute of Technology, Laboratory for Nuclear Science, Cambridge, Massachusetts 02139, USA*
- ⁴⁷*McGill University, Montréal, Québec, Canada H3A 2T8*
- ⁴⁸*Università di Milano, Dipartimento di Fisica and INFN, I-20133 Milano, Italy*
- ⁴⁹*University of Mississippi, University, Mississippi 38677, USA*
- ⁵⁰*Université de Montréal, Physique des Particules, Montréal, Québec, Canada H3C 3J7*
- ⁵¹*Mount Holyoke College, South Hadley, Massachusetts 01075, USA*
- ⁵²*Università di Napoli Federico II, Dipartimento di Scienze Fisiche and INFN, I-80126, Napoli, Italy*
- ⁵³*NIKHEF, National Institute for Nuclear Physics and High Energy Physics, NL-1009 DB Amsterdam, The Netherlands*
- ⁵⁴*University of Notre Dame, Notre Dame, Indiana 46556, USA*
- ⁵⁵*Ohio State University, Columbus, Ohio 43210, USA*
- ⁵⁶*University of Oregon, Eugene, Oregon 97403, USA*
- ⁵⁷*Università di Padova, Dipartimento di Fisica and INFN, I-35131 Padova, Italy*
- ⁵⁸*Universités Paris VI et VII, Laboratoire de Physique Nucléaire et de Hautes Energies, F-75252 Paris, France*
- ⁵⁹*University of Pennsylvania, Philadelphia, Pennsylvania 19104, USA*
- ⁶⁰*Università di Perugia, Dipartimento di Fisica and INFN, I-06100 Perugia, Italy*
- ⁶¹*Università di Pisa, Dipartimento di Fisica, Scuola Normale Superiore and INFN, I-56127 Pisa, Italy*
- ⁶²*Prairie View A&M University, Prairie View, Texas 77446, USA*
- ⁶³*Princeton University, Princeton, New Jersey 08544, USA*
- ⁶⁴*Università di Roma La Sapienza, Dipartimento di Fisica and INFN, I-00185 Roma, Italy*
- ⁶⁵*Universität Rostock, D-18051 Rostock, Germany*
- ⁶⁶*Rutherford Appleton Laboratory, Chilton, Didcot, Oxon, OX11 0QX, United Kingdom*
- ⁶⁷*DSM/Dapnia, CEA/Saclay, F-91191 Gif-sur-Yvette, France*
- ⁶⁸*University of South Carolina, Columbia, South Carolina 29208, USA*
- ⁶⁹*Stanford Linear Accelerator Center, Stanford, California 94309, USA*
- ⁷⁰*Stanford University, Stanford, California 94305-4060, USA*
- ⁷¹*State University of New York, Albany, New York 12222, USA*
- ⁷²*University of Tennessee, Knoxville, Tennessee 37996, USA*
- ⁷³*University of Texas at Austin, Austin, Texas 78712, USA*
- ⁷⁴*University of Texas at Dallas, Richardson, Texas 75083, USA*
- ⁷⁵*Università di Torino, Dipartimento di Fisica Sperimentale and INFN, I-10125 Torino, Italy*
- ⁷⁶*Università di Trieste, Dipartimento di Fisica and INFN, I-34127 Trieste, Italy*
- ⁷⁷*IFIC, Universitat de Valencia-CSIC, E-46071 Valencia, Spain*
- ⁷⁸*University of Victoria, Victoria, British Columbia, Canada V8W 3P6*
- ⁷⁹*Department of Physics, University of Warwick, Coventry CV4 7AL, United Kingdom*
- ⁸⁰*University of Wisconsin, Madison, Wisconsin 53706, USA*
- ⁸¹*Yale University, New Haven, Connecticut 06511, USA*

(Received 5 April 2006; published 20 February 2007)

We report the observation of decays $B^0 \rightarrow D_s^{(*)+} \pi^-$ and $B^0 \rightarrow D_s^{(*)-} K^+$ in a sample of 230×10^6 $\Upsilon(4S) \rightarrow B\bar{B}$ events recorded with the *BABAR* detector at the SLAC PEP-II asymmetric-energy e^+e^- storage ring. We measure the branching fractions $\mathcal{B}(B^0 \rightarrow D_s^+ \pi^-) = (1.3 \pm 0.3(\text{stat}) \pm 0.2(\text{syst})) \times 10^{-5}$, $\mathcal{B}(B^0 \rightarrow D_s^- K^+) = (2.5 \pm 0.4(\text{stat}) \pm 0.4(\text{syst})) \times 10^{-5}$, $\mathcal{B}(B^0 \rightarrow D_s^{*+} \pi^-) = (2.8 \pm 0.6(\text{stat}) \pm 0.5(\text{syst})) \times 10^{-5}$, and $\mathcal{B}(B^0 \rightarrow D_s^{*-} K^+) = (2.0 \pm 0.5(\text{stat}) \pm 0.4(\text{syst})) \times 10^{-5}$. The significances of the measurements to differ from zero are 5, 9, 6, and 5 standard deviations, respectively. This is the first observation of $B^0 \rightarrow D_s^+ \pi^-$, $B^0 \rightarrow D_s^{*+} \pi^-$, and $B^0 \rightarrow D_s^{*-} K^+$ decays.

DOI: 10.1103/PhysRevLett.98.081801

PACS numbers: 13.25.Hw, 11.30.Er, 12.15.Hh

Within the Cabibbo-Kobayashi-Maskawa (CKM) model of quark-flavor mixing [1], CP violation manifests itself as a nonzero area of the unitarity triangle [2]. One of the important experimental tests of the model is the determination of the angle $\gamma = \arg(-V_{ud}V_{ub}^*/V_{cd}V_{cb}^*)$ of the unitarity triangle. A measurement of $\sin(2\beta + \gamma)$ can be obtained from the study of the time dependence of the B^0 , $\bar{B}^0 \rightarrow D^{(*)-} \pi^+$ [3] decay rates, and specifically of the interference between the CKM-favored B^0 decay amplitude and CKM-suppressed \bar{B}^0 amplitude [4]. The first measurements of the CP asymmetry in decays $B^0 \rightarrow D^{(*)\mp} \pi^\pm$ have recently been published [5].

The measurement of $\sin(2\beta + \gamma)$ in $B^0 \rightarrow D^{(*)\mp} \pi^\pm$ decays requires knowledge of the ratios of the decay amplitudes, $r(D^{(*)}\pi) = |A(B^0 \rightarrow D^{(*)+} \pi^-)/A(B^0 \rightarrow D^{(*)-} \pi^+)|$. The CP -violating observables in $B^0 \rightarrow D^{(*)\mp} \pi^\pm$ decays are proportional to $r(D^{(*)}\pi)$ [4,5]. However, direct measurement of the branching fractions $\mathcal{B}(B^0 \rightarrow D^{(*)+} \pi^-)$ is not possible with the currently available data sample due to the presence of the overwhelming background from $\bar{B}^0 \rightarrow D^{(*)+} \pi^-$. However, assuming $SU(3)$ flavor symmetry, $r(D^{(*)}\pi)$ can be related to the branching fraction (BF) of the decay $B^0 \rightarrow D_s^{(*)+} \pi^-$ [4]:

$$r(D^{(*)}\pi) = \tan\theta_c \frac{f_{D^{(*)}}}{f_{D_s^{(*)}}} \sqrt{\frac{\mathcal{B}(B^0 \rightarrow D_s^{(*)+} \pi^-)}{\mathcal{B}(B^0 \rightarrow D^{(*)-} \pi^+)}} \quad (1)$$

where θ_c is the Cabibbo angle, $f_{D^{(*)}}$ and $f_{D_s^{(*)}}$ are $D^{(*)}$ and $D_s^{(*)}$ decay constants [6]. Other $SU(3)$ -breaking effects are believed to affect $r(D^{(*)}\pi)$ by less than 30% [5].

Since $B^0 \rightarrow D_s^{(*)+} \pi^-$ has four different quark flavors in the final state, only a single amplitude contributes to the decay [Fig. 1(c)]. On the other hand, there are two diagrams contributing to $B^0 \rightarrow D^{(*)-} \pi^+$ and $B^0 \rightarrow D^{(*)+} \pi^-$: tree amplitudes [Figs. 1(a) and 1(b)] and color-suppressed direct W -exchange amplitudes [Figs. 1(d) and 1(e)]. The latter are assumed to be negligibly small in Eq. (1). The decays $B^0 \rightarrow D_s^{(*)-} K^+$ [Fig. 1(f)] probe the size of the W -exchange amplitudes relative to the dominant processes $B^0 \rightarrow D^{(*)-} \pi^+$. The rate of $B^0 \rightarrow D_s^{*-} K^+$ decays could be enhanced by final state rescattering [7], in addition to the W -exchange amplitude. The relative rates of $B^0 \rightarrow D_s^{(*)-} K^{(*)+}$ decays could shed light on the decay dynamics,

including relative contributions of short- and long-distance effects [8].

The branching fractions $\mathcal{B}(B^0 \rightarrow D_s^+ \pi^-)$ and $\mathcal{B}(B^0 \rightarrow D_s^- K^+)$ have been measured previously by the *BABAR* [9] and Belle [10] collaborations, but the decays $B^0 \rightarrow D_s^{*+} \pi^-$ and $B^0 \rightarrow D_s^{*-} K^+$ have never been observed. In this Letter we present new measurements of the decays $B^0 \rightarrow D_s^{(*)+} \pi^-$ and $B^0 \rightarrow D_s^{(*)-} K^+$. The analysis uses a sample of 230×10^6 $\Upsilon(4S)$ decays into $B\bar{B}$ pairs collected with the *BABAR* detector at the SLAC PEP-II asymmetric-energy B factory [11].

Since the *BABAR* detector is described in detail elsewhere [12], only the components that are crucial to this analysis are summarized here. Charged-particle tracking is provided by a five-layer double-sided silicon vertex tracker (SVT) and a 40-layer drift chamber (DCH). Ionization energy loss (dE/dx) in the DCH and SVT and Cherenkov radiation detected in a ring-imaging device are used for charged-particle identification. Photons are identified and measured using the electromagnetic calorimeter (EMC), which is comprised of 6580 thallium-doped CsI crystals. These systems are mounted inside a 1.5 T solenoidal superconducting magnet. We use the GEANT4 [13] software to simulate interactions of particles traversing the *BABAR* detector, taking into account the varying detector conditions and beam backgrounds.




FIG. 1. Dominant Feynman diagrams for (a) CKM-favored decay $B^0 \rightarrow D^{(*)-} \pi^+$, (b) doubly CKM-suppressed decay $B^0 \rightarrow D^{(*)+} \pi^-$, and (c) the $SU(3)$ flavor symmetry related decays $B^0 \rightarrow D_s^{(*)+} \pi^-$; (d) the color-suppressed W -exchange contributions to $B^0 \rightarrow D^{(*)-} \pi^+$, (e) $B^0 \rightarrow D^{(*)+} \pi^-$, and (f) decay $B^0 \rightarrow D_s^{(*)-} K^+$.

Candidates for D_s^+ mesons are reconstructed in the modes $D_s^+ \rightarrow \phi \pi^+$, $K_S^0 K^+$, and $\bar{K}^{*0} K^+$, with $\phi \rightarrow K^+ K^-$, $K_S^0 \rightarrow \pi^+ \pi^-$, and $\bar{K}^{*0} \rightarrow K^- \pi^+$. The K_S^0 candidates are formed from two oppositely charged tracks, and their momenta are required to make an angle $|\theta_{\text{flight}}| < 11^\circ$ with the line connecting their vertex and $e^+ e^-$ interaction point (IP). All other tracks are required to originate from the IP, whose average position and size are determined hourly using two-prong and hadronic events. In order to reject background from $D^+ \rightarrow K_S^0 \pi^+$ or $\bar{K}^{*0} \pi^+$, the K^+ candidate in the reconstruction of $D_s^+ \rightarrow K_S^0 K^+$ or $\bar{K}^{*0} K^+$ is required to satisfy positive kaon identification criteria with an efficiency of 85% and 5% pion misidentification probability. The same selection is used to identify kaon daughters of the B mesons in decays $B^0 \rightarrow D_s^{(*)-} K^+$. In all other cases, kaons are not positively identified, but instead candidates passing pion selection are rejected. Such ‘‘pion veto’’ has an efficiency of 95% for kaons and 20% for pions. Pion daughters of B mesons in the decays $B^0 \rightarrow D_s^{(*)+} \pi^-$ are required to be positively identified. Decay products of ϕ , \bar{K}^{*0} , K_S^0 , D_s^+ , and B^0 candidates are constrained to originate from a single vertex.

We reconstruct D_s^{*+} candidates in the mode $D_s^{*+} \rightarrow D_s^+ \gamma$ by combining D_s^+ and photon candidates. Photon candidates are required to be consistent with an electromagnetic shower in the EMC, and have an energy greater than 100 MeV in the laboratory frame. When forming a D_s^{*+} , the D_s^+ candidate is required to have invariant mass within 10 MeV/ c^2 of the nominal value [14].

After an initial preselection, we identify signal candidates using a likelihood ratio $R_L = \mathcal{L}_{\text{sig}} / (\mathcal{L}_{\text{sig}} + \mathcal{L}_{\text{bkg}})$, where $\mathcal{L}_{\text{sig}} = \prod_i \mathcal{P}_{\text{sig}}(x_i)$ is the multivariate likelihood for signal events and $\mathcal{L}_{\text{bkg}} = \prod_i \mathcal{P}_{\text{bkg}}(x_i)$ is the likelihood for background events. The ratio R_L has a maximum at $R_L = 1$ for signal events, and at $R_L = 0$ for background originating from continuum events. It also discriminates well against generic B decays without a real D_s^+ meson in the final state. The likelihoods \mathcal{L}_{sig} and \mathcal{L}_{bkg} are computed as products of the probability density functions (PDFs) $\mathcal{P}_{\text{sig}}(x_i)$ and $\mathcal{P}_{\text{bkg}}(x_i)$ for a number of selection variables x_i : invariant masses of the ϕ , \bar{K}^{*0} and K_S^0 candidates, χ^2 confidence level of the vertex fit for the B^0 and D_s^+ mesons, the helicity angles of the ϕ , \bar{K}^{*0} , and D_s^{*+} meson decays, the mass difference $\Delta m(D_s^{*+}) = m(D_s^{*+}) - m(D_s^+)$, the polar angle θ_B of the B candidate momentum vector with respect to the beam axis in the $e^+ e^-$ center-of-mass (c.m.) frame, the angle θ_T between the thrust axis of the B candidate and the thrust axis of all other particles in the event in c.m. frame, and event topology variable \mathcal{F} , discussed below. We have determined the correlations among these variables to be negligibly small. The helicity angle θ_H is defined as the angle between one of the decay products of a vector meson and the flight direction of its parent particle, in the meson’s rest frame. Polarization of

the vector mesons in the signal decays causes $\cos^2 \theta_H$ (ϕ and \bar{K}^{*0}) or $\sin^2 \theta_H$ (D_s^{*+}) distributions, while the random background combinations tend to produce a more uniform distribution in $\cos \theta_H$.

Variables $\cos \theta_B$, $\cos \theta_T$, and \mathcal{F} discriminate between spherically-symmetric $B\bar{B}$ events and jetty continuum background. $B\bar{B}$ pairs form a nearly uniform $|\cos \theta_T|$ distribution, while $|\cos \theta_T|$ distribution for the continuum peaks at 1. A linear (Fisher) discriminant \mathcal{F} is derived from the values of sphericity and thrust for the event, and the two Legendre moments L_0 and L_2 of the energy flow around the B -candidate thrust axis [15]. Finally, the polar angle θ_B is distributed as $\sin^2 \theta_B$ for real B decays, while being nearly flat in $\cos \theta_B$ for the continuum.

We select $B^0 \rightarrow D_s^+ \pi^-$ and $B^0 \rightarrow D_s^- K^+$ candidates that satisfy $R_L > 0.75$, and accept $B^0 \rightarrow D_s^{*+} \pi^-$ and $B^0 \rightarrow D_s^{*-} K^+$ candidates with $R_L > 0.8$. We measure the relative efficiency ε_{R_L} of the R_L selection in a copious data sample of decays $B^0 \rightarrow D^- \pi^+$ ($D^- \rightarrow K^+ \pi^- \pi^-$, $K_S^0 \pi^-$) and $B^+ \rightarrow \bar{D}^{*0} \pi^+$ ($\bar{D}^{*0} \rightarrow \bar{D}^0 \gamma$, $D^0 \rightarrow K^- \pi^+$) in which the kinematics is similar to that of our signal events, and find that it is consistent with Monte Carlo estimates $\varepsilon_{R_L} \approx 70\%$. The fraction of continuum background events passing the selection varies between 2% and 15%, depending on the mode.

We identify the signal using the invariant mass $m(D_s)$ of D_s candidates and two kinematic variables m_{ES} and ΔE . The first is the beam-energy-substituted mass $m_{\text{ES}} = \sqrt{(s/2 + \mathbf{p}_i \cdot \mathbf{p}_B)^2 / E_i^2 - \mathbf{p}_B^2}$, where \sqrt{s} is the total c.m. energy, (E_i, \mathbf{p}_i) is the four-momentum of the initial $e^+ e^-$ system and \mathbf{p}_B is the B^0 candidate momentum, both measured in the laboratory frame. The second variable is $\Delta E = E_B^* - \sqrt{s}/2$, where E_B^* is the B^0 candidate energy in the c.m. frame. For signal events, the m_{ES} distribution is Gaussian centered at the B meson mass with a resolution of about 2.5 MeV/ c^2 , and the ΔE distribution has a maximum near zero with a resolution of about 17 MeV. The invariant mass $m(D_s)$ has a resolution of (5–6) MeV/ c^2 , depending on the D_s^+ decay mode. We define a fit region $5.2 < m_{\text{ES}} < 5.3$ GeV/ c^2 , $|\Delta E| < 36$ MeV, and $|m(D_s) - m(D_s)_{\text{PDG}}| < 50$ MeV/ c^2 for $B^0 \rightarrow D_s^+ \pi^-$ and $B^0 \rightarrow D_s^- K^+$ candidates, where $m(D_s)_{\text{PDG}}$ is the world average D_s mass [14]. For $B^0 \rightarrow D_s^{*+} \pi^-$ and $B^0 \rightarrow D_s^{*-} K^+$, we require $|m(D_s) - m(D_s)_{\text{PDG}}| < 10$ MeV/ c^2 .

Less than 20% of the selected events in the $B^0 \rightarrow D_s^{*+} \pi^-$ and $B^0 \rightarrow D_s^{*-} K^+$ channels and less than 4% in $B^0 \rightarrow D_s^+ \pi^-$ and $B^0 \rightarrow D_s^- K^+$ channels contain two or more candidates that satisfy the criteria listed above. In such events we select a single B^0 candidate based on an event χ^2 formed with $m(D_s)$ (both D_s^+ and D_s^{*+} modes) and $\Delta m(D_s^{*+})$ (D_s^{*+} modes) and their average uncertainties, and the ΔE variable. Such selection does not bias background distributions significantly.

Four classes of background contribute to the fit region. First is the *combinatorial background*, in which a true or

fake $D_s^{(*)}$ candidate is combined with a randomly-selected pion or kaon. Second, B meson decays such as $\bar{B}^0 \rightarrow D^{(*)+} \pi^-$, ρ^- with $D^+ \rightarrow K_S^0 \pi^+$ or $\bar{K}^{*0} \pi^+$ can constitute a background for the $B^0 \rightarrow D_s^{(*)+} \pi^-$ modes if the pion in the D decay is misidentified as a kaon (*reflection background*). The reflection background has nearly the same m_{ES} distribution as the signal but different distributions in ΔE and $m(D_s)$. The corresponding backgrounds for the $B^0 \rightarrow D_s^- K^+$ mode ($B^0 \rightarrow D^- K^{(*)+}$) are negligible. Third, rare B decays into the same final state, such as $B^0 \rightarrow \bar{K}^{(*)0} K^+ \pi^-$ or $\bar{K}^{(*)0} K^+ K^-$ (*charmless background*), have the same m_{ES} and ΔE distributions as the $B^0 \rightarrow D_s^+ \pi^-$ or $B^0 \rightarrow D_s^- K^+$ signal, but are nearly flat in $m(D_s)$. The charmless background is significant in $B^0 \rightarrow D_s^+ \pi^-$ and $B^0 \rightarrow D_s^- K^+$ decays, but is negligible for $B^0 \rightarrow D_s^{*+} \pi^-$ and $B^0 \rightarrow D_s^{*-} K^+$. Finally, *crossfeed background* from misidentification of $\bar{B}^0 \rightarrow D_s^{(*)-} \pi^+$ events as $B^0 \rightarrow D_s^{(*)-} K^+$ signal, and vice versa, needs to be taken into account.

We perform a two-dimensional unbinned extended maximum-likelihood fit to the m_{ES} and $m(D_s)$ distributions to extract $\mathcal{B}(B^0 \rightarrow D_s^+ \pi^-)$ and $\mathcal{B}(B^0 \rightarrow D_s^- K^+)$ and constrain the contributions from charmless background modes. Charmless backgrounds are negligible for $B^0 \rightarrow D_s^{*+} \pi^-$ and $B^0 \rightarrow D_s^{*-} K^+$, and we determine the BFs of these decays with a one-dimensional fit to the m_{ES} distribution. For each B decay, we simultaneously fit distributions in three D_s^+ decay modes, constraining the signal BFs to a common value. The likelihood function contains the contributions of the signal and the four background components discussed above. The combinatorial background is described in m_{ES} by a threshold function [16], $dN/dx \propto x \sqrt{1 - 2x^2/s} \exp[-\xi(1 - 2x^2/s)]$. In $m(D_s)$, the combinatorial background is well described by a combination of a first-order polynomial (fake D_s^+ candidates) and a Gaussian with (5–6) MeV/ c^2 resolution (true D_s^+ candidates). The charmless background is parameterized by the signal Gaussian shape in m_{ES} and a first-order polynomial in $m(D_s)$.

For $B^0 \rightarrow D_s^+ \pi^-$ and $B^0 \rightarrow D_s^- K^+$ decays, the fit determines 14 free parameters: the shape of the combinatorial background ξ (1 parameter for all D_s^+ modes), the slope of the combinatorial and charmless backgrounds in $m(D_s)$ (3 parameters), the fraction of true D_s^+ candidates in combinatorial background (3), the number of combinatorial background events (3), the number of charmless events (3), and the BF of the signal mode (1). The signal yields for each D_s^+ mode are expressed as $N_{sigi} = N_{B\bar{B}} \mathcal{B}_{sig} \mathcal{B}_i \varepsilon_i$, where $N_{B\bar{B}} = 230 \times 10^6$, \mathcal{B}_i is the D_s^+ BF for the mode, ε_i is the reconstruction efficiency, and \mathcal{B}_{sig} is the BF (fit parameter) for the decay. For the $B^0 \rightarrow D_s^{*+} \pi^-$ and $B^0 \rightarrow D_s^{*-} K^+$ decays, 5 free parameters are determined by the fit: ξ (1 parameter for all D_s^+ modes), the number of combinatorial background events (3), and the BF of the signal




FIG. 2 (color online). (a),(c),(e),(f) m_{ES} projection of the fit with $|m(D_s^+) - m(D_s^+)_{PDG}| < 10$ MeV/ c^2 and (b),(d) $m(D_s)$ projection with $5.275 < m_{ES} < 5.285$ GeV for (a),(b) $B^0 \rightarrow D_s^+ \pi^-$, (c),(d) $B^0 \rightarrow D_s^- K^+$, (e) $B^0 \rightarrow D_s^{*+} \pi^-$, and (f) $B^0 \rightarrow D_s^{*-} K^+$. Bins with zero events are omitted for clarity. The black solid curve corresponds to the full PDF from the combined fit to all D_s^+ decay modes. Individual contributions are shown as solid red lines (signal PDF), green dashed lines (combinatorial background), and blue dotted lines (sum of reflection, charmless, and crossfeed backgrounds) curves.

mode (1). The signal efficiency ε_i varies between 6.7% and 29.3%, depending on the mode. The BFs of the channels contributing to the reflection background are fixed in the fit to the current world average values [14], and the BFs of the crossfeed backgrounds are determined by iterating the fits over each B decay mode. The fit samples contain 1305 events for $B^0 \rightarrow D_s^+ \pi^-$, 132 for $B^0 \rightarrow D_s^- \pi^-$, 539 for $B^0 \rightarrow D_s^- K^+$, and 41 events for $B^0 \rightarrow D_s^{*-} K^+$ mode. The results of the fits are shown in Fig. 2 and summarized in Table I.

Systematic errors are dominated by 13% relative uncertainty for $\mathcal{B}(D_s^+ \rightarrow \phi \pi^+)$ [17]. The relative BF uncertainties for $\mathcal{B}(D_s^+ \rightarrow \bar{K}^{*0} K^+)/\mathcal{B}(D_s^+ \rightarrow \phi \pi^+)$ and $\mathcal{B}(D_s^+ \rightarrow K_S^0 K^+)/\mathcal{B}(D_s^+ \rightarrow \phi \pi^+)$ contribute 5%–7%, depending on the decay channel. Uncertainties in the selection efficiency are estimated to be 3% for D_s^+ modes and 7% for D_s^{*+} modes. The uncertainties in the reflection and crossfeed backgrounds are below 1% for all decay channels. Other systematic errors include the uncertainties in tracking (5%), photon (3%), and K_S^0 reconstruction (0.2%–0.5%), charged-kaon identification (1%) efficiencies, and variations of the PDF shapes between data and

TABLE I. The results of the fit for the branching ratios. Shown are the probability (P_{bkg}) of the data being consistent with the background in the absence of signal, and the measured branching fraction \mathcal{B} . The first uncertainty is statistical, and the second is systematic.

B mode	P_{bkg}	$\mathcal{B}(10^{-5})$	$\mathcal{B} \times \mathcal{B}(D_s^+ \rightarrow \phi \pi^+)(10^{-6})$
$B^0 \rightarrow D_s^+ \pi^-$	3×10^{-6}	$1.3 \pm 0.3 \pm 0.2$	$0.63 \pm 0.15 \pm 0.05$
$B^0 \rightarrow D_s^{*+} \pi^-$	3×10^{-8}	$2.8 \pm 0.6 \pm 0.5$	$1.32 \pm 0.27 \pm 0.15$
$B^0 \rightarrow D_s^- K^+$	3×10^{-19}	$2.5 \pm 0.4 \pm 0.4$	$1.21 \pm 0.17 \pm 0.11$
$B^0 \rightarrow D_s^{*-} K^+$	2×10^{-5}	$2.0 \pm 0.5 \pm 0.4$	$0.97 \pm 0.24 \pm 0.12$

Monte Carlo calculations ([1%–5%], depending on decay mode).

The ratio $P_{\text{bkg}} = \mathcal{L}_0 / \mathcal{L}_{\text{max}}$, where \mathcal{L}_{max} is the maximum-likelihood value, and \mathcal{L}_0 is the likelihood for a fit with the signal contribution set to zero, describes the probability of the background to fluctuate to the observed number of events. The values P_{bkg} in Table I include all systematic uncertainties, which are assumed to be Gaussian-distributed. They correspond to the significance of signal observation of 5 ($B^0 \rightarrow D_s^+ \pi^-$), 6 ($B^0 \rightarrow D_s^{*+} \pi^-$), 9 ($B^0 \rightarrow D_s^- K^+$), and 5 ($B^0 \rightarrow D_s^{*-} K^+$) standard deviations. This is the first observation of $B^0 \rightarrow D_s^+ \pi^-$, $B^0 \rightarrow D_s^{*+} \pi^-$, and $B^0 \rightarrow D_s^{*-} K^+$ decays.

Assuming the SU(3) relation, Eq. (1), we determine $r(D\pi) = (1.29 \pm 0.15(\text{stat}) \pm 0.13(\text{syst})) \times 10^{-2}$, and $r(D^*\pi) = (1.87 \pm 0.19(\text{stat}) \pm 0.19(\text{syst})) \times 10^{-2}$, which implies small CP asymmetries in $B^0 \rightarrow D^{(*)\mp} \pi^\pm$ decays. The branching fractions for $B^0 \rightarrow D_s^{(*)-} K^+$ are small compared to the dominant decays $B^0 \rightarrow D^{(*)-} \pi^+$, implying relatively insignificant contributions from the color-suppressed W -exchange diagrams. These results supersede our previously published measurements [9].

We are grateful for the excellent luminosity and machine conditions provided by our PEP-II colleagues, and for the substantial dedicated effort from the computing organizations that support *BABAR*. The collaborating institutions wish to thank SLAC for its support and kind hospitality. This work is supported by DOE and NSF (USA), NSERC (Canada), IHEP (China), CEA and CNRS-IN2P3 (France), BMBF and DFG (Germany), INFN (Italy), FOM (The Netherlands), NFR (Norway), MIST (Russia), and PPARC (United Kingdom). Individuals have received support from CONACyT (Mexico), Marie Curie EIF (European Union), the A. P. Sloan Foundation, the Research Corporation, and the Alexander von Humboldt Foundation.

*Also at Laboratoire de Physique Corpusculaire, Clermont-Ferrand, France.

†Also at Università di Perugia, Dipartimento di Fisica, Perugia, Italy.

‡Also at Università della Basilicata, Potenza, Italy.

- [1] N. Cabibbo, Phys. Rev. Lett. **10**, 531 (1963); M. Kobayashi and T. Maskawa, Prog. Theor. Phys. **49**, 652 (1973).
- [2] C. Jarlskog, in *CP Violation*, edited by C. Jarlskog (World Scientific, Singapore, 1988).
- [3] Charge conjugation is implied throughout this Letter, unless explicitly stated.
- [4] I. Duniety, Phys. Lett. B **427**, 179 (1998).
- [5] B. Aubert *et al.* (*BABAR* Collaboration), Phys. Rev. Lett. **92**, 251801 (2004); Phys. Rev. D **71**, 112003 (2005); T. Sarangi *et al.* (Belle Collaboration), Phys. Rev. Lett. **93**, 031802 (2004).
- [6] C. Aubin *et al.*, Phys. Rev. Lett. **95**, 122002 (2005); D. Becirevic, hep-ph/0310072.
- [7] B. Blok *et al.*, Phys. Rev. Lett. **78**, 3999 (1997).
- [8] S. Mantry *et al.*, Phys. Rev. D **68**, 114009 (2003).
- [9] B. Aubert *et al.* (*BABAR* Collaboration), Phys. Rev. Lett. **90**, 181803 (2003).
- [10] P. Krokovny *et al.* (Belle Collaboration), Phys. Rev. Lett. **89**, 231804 (2002).
- [11] PEP-II Conceptual Design Report, No. SLAC-0418, 1993.
- [12] B. Aubert *et al.* (*BABAR* Collaboration), Nucl. Instrum. Methods Phys. Res., Sect. A **479**, 1 (2002).
- [13] S. Agostinelli *et al.* (GEANT4 Collaboration), Nucl. Instrum. Methods Phys. Res., Sect. A **479**, 1 (2002).
- [14] S. Eidelman *et al.* (Particle Data Group), Phys. Lett. B **592**, 1 (2004).
- [15] B. Aubert *et al.* (*BABAR* Collaboration), Phys. Rev. Lett. **89**, 281802 (2002); Phys. Rev. D **70**, 032006 (2004).
- [16] H. Albrecht *et al.* (ARGUS Collaboration), Z. Phys. C **48**, 543 (1990).
- [17] B. Aubert *et al.* (*BABAR* Collaboration), Phys. Rev. D **71**, 091104 (2005).