

# UC San Diego

## UC San Diego Previously Published Works

### Title

Sex differences in dementia with Lewy bodies: Focused review of available evidence and future directions.

### Permalink

<https://escholarship.org/uc/item/8s98b0nv>

### Authors

Chiu, Shannon

Wyman-Chick, Kathryn

Ferman, Tanis

et al.

### Publication Date

2023-02-01

### DOI

10.1016/j.parkreldis.2023.105285

Peer reviewed



Published in final edited form as:

*Parkinsonism Relat Disord.* 2023 February ; 107: 105285. doi:10.1016/j.parkreldis.2023.105285.

## Sex differences in dementia with Lewy bodies: focused review of available evidence and future directions

Shannon Y. Chiu, M.D.<sup>\*1,2</sup>, Kathryn A. Wyman-Chick, Psy.D.<sup>3</sup>, Tanis J. Ferman, Ph.D.<sup>4</sup>, Ece Bayram, M.D., Ph.D.<sup>5</sup>, Samantha K. Holden, M.D., M.S.<sup>6</sup>, Parichita Choudhury, M.D.<sup>7</sup>, Melissa J. Armstrong, M.D.<sup>1,2</sup>

<sup>1</sup>Department of Neurology, University of Florida College of Medicine, Gainesville, FL

<sup>2</sup>Norman Fixel Institute for Neurologic Diseases, University of Florida, Gainesville, FL

<sup>3</sup>Center for Memory and Aging, Department of Neurology, HealthPartners, Saint Paul, MN

<sup>4</sup>Department of Psychiatry & Psychology, Mayo Clinic, Jacksonville, FL

<sup>5</sup>Department of Neurosciences, University of California San Diego, La Jolla, CA

<sup>6</sup>Department of Neurology, University of Colorado School of Medicine, Aurora, CO

<sup>7</sup>Cleo Roberts Center, Banner Sun Health Research Institute, Sun City, AZ

### Abstract

In this review, we summarize the current knowledge on sex differences in dementia with Lewy bodies (DLB) relating to epidemiology, clinical features, neuropathology, biomarkers, disease progression, and caregiving. While many studies show a higher DLB prevalence in men, this finding is inconsistent and varies by study approach. Visual hallucinations may be more common and occur earlier in women with DLB, whereas REM sleep behavior disorder may be more common and occur earlier in men. Several studies report a higher frequency of parkinsonism in men with DLB, while the frequency of fluctuations appears similar between sexes. Women tend to be older, have greater cognitive impairment at their initial visit, and are delayed in meeting DLB criteria compared to men. Women are also more likely to have Lewy body disease with co-existing AD-related pathology than so-called “pure” Lewy body disease, while men may present with either. Research is mixed regarding the impact of sex on DLB progression. Biomarker and treatment research assessing for sex differences is lacking. Women provide the majority of caregiving in DLB but how this affects the caregiving experience is uncertain. Gaining a better understanding of sex differences will be instrumental in aiding future development of sex-specific strategies in DLB for early diagnosis, care, and drug development.

\*Corresponding author: Shannon Y. Chiu, MD, Department of Neurology, College of Medicine, University of Florida, P.O. Box 100268, Gainesville, FL 32611 (Shannon.Chiu@neurology.ufl.edu).

**Publisher's Disclaimer:** This is a PDF file of an unedited manuscript that has been accepted for publication. As a service to our customers we are providing this early version of the manuscript. The manuscript will undergo copyediting, typesetting, and review of the resulting proof before it is published in its final form. Please note that during the production process errors may be discovered which could affect the content, and all legal disclaimers that apply to the journal pertain.

Human and animal rights

This work did not involve the use of animal or human subjects. Ethical approval/consent for experimentation with human subjects was not applicable for this review paper.

## Keywords

dementia with Lewy bodies; dementia; Lewy body disease; sex

---

## 1. Introduction

Neurodegenerative diseases have documented sex differences in clinical phenotype, time to diagnosis, pathology, and treatment [1, 2]. For example, Alzheimer disease (AD) is more common in women [3-5]. This can partly relate to longevity differences, but women may have greater neurofibrillary tangle (NFT) pathology [6] and higher CSF total tau and A $\beta$ 42 levels [7] than men. Parkinson disease (PD) is more common in men [8-10]. Dementia with Lewy bodies (DLB) is more commonly diagnosed in men [11-13], resulting in fewer women in DLB cohorts. This has historically precluded the examination of sex differences in DLB. However, longstanding DLB programs and recent multicenter collaborations now have large cohorts available for examining potential sex differences [14-20]. A better understanding of sex differences in DLB is important for early disease detection, symptomatic management, caregiver support, and eventual implementation of personalized therapies.

In this review, we highlight current research on sex differences in DLB including epidemiology, clinical presentation, pathology, biomarkers, progression, management, and caregiving. Search criteria included peer-reviewed DLB research studies identified through EMBASE and PubMed published in English prior to July 2022. We excluded research solely on PD, AD, or other dementias. DLB studies were included if results had sex- or gender-specific analyses. While “sex” (biology) and “gender” (a socially-based construct) are related terms, they have different meanings but many studies used the terms interchangeably. Thus, the review focuses on identified sex differences in DLB with further comment on the potential role of gender in the discussion.

## 2. Sex differences in Clinical and Population Cohorts

DLB accounts for approximately 3.5% to 9.6% of dementia cases in epidemiologic studies [12, 21-24]. Clinical DLB samples typically have more men [11-13, 25], a pattern observed in some population studies [12] but not others [16, 21, 22] (Table 1). Some variability may relate to differences in community versus clinical populations. Specialty clinic DLB cohorts typically include more individuals with parkinsonism and/or REM sleep behavior disorder (RBD), which are more commonly diagnosed in men [12, 26-30]. Community samples often include older individuals with a greater proportion of women. In older patients with DLB, the sex ratio is more likely to be equal or higher for women. For those <75, a DLB diagnosis is more common in men [12, 16, 31]. Geographic location may also be relevant, although the basis for this is unclear. For example, Japanese multicenter DLB studies [32-34], French cohorts [16] and British samples [23, 35] showed more balanced sex ratio than U.S. cohorts.

It is unknown whether race or ethnicity affects cohort sex differences. An autopsy study of 81 Black and 154 White community participants from longitudinal aging studies revealed no differences in sex or in the frequency and distribution of Lewy body pathology [36]. In a National Alzheimer’s Coordinating Center (NACC) study comprised of 1,782 White, 130

Black, and 122 Hispanic participants, women represented only 26% of White participants with cognitive impairment due to Lewy body pathology, compared to 60% identifying as Black and 41% identifying as Hispanic [37]. More work is needed to determine whether this reflects disease-specific differences, geographic variation, and/or recruitment patterns.

### 3. Clinical Features

#### 3.1 Cognitive Function and Dementia:

DLB is characterized by impairments in attention and visual perceptual abilities, while memory and naming tend to be relatively preserved compared to patients with AD [38-40]. Lower memory and naming scores occur in DLB with AD-related pathology compared to DLB without AD-related pathology, and co-pathology is common in women with DLB [41, 42].

In a study of individuals with clinically probable DLB [43], men (n=101) had higher Mini Mental State Examination (MMSE) scores than women (n=133) ( $19.7 \pm 4.6$  vs.  $18.0 \pm 5.7$ ,  $p=0.029$ ), suggesting less severe cognitive impairment despite similar clinical presentation, disease duration, and age at onset [17]. In a European DLB Consortium sub-analysis (n=375), DLB participants with an AD-CSF profile were more likely to be female, older and with lower MMSE scores than those without an AD-CSF profile [44]. In the Amsterdam Dementia Cohort, women with DLB (n=39/223, 17%) had lower CSF A $\beta$ 42, as well as worse MMSE and animal fluency scores, but did not differ from men on other cognitive tasks [18]. This pattern was also shown in a NACC DLB study [20], wherein women had a higher Braak NFT stage, neuritic plaque frequency and Clinical Dementia Rating Scale Sum of Boxes scores at the last visit. Overall, data on differences in cognitive function between sexes in DLB are limited, but suggest that women may have worse MMSE scores, possibly relating to higher degrees of AD-related pathology.

#### 3.2 Cognitive Fluctuations.

Men and women with DLB have similar reported rates of cognitive fluctuations [14, 15, 17, 18, 20] (Table 2) based on the Mayo Fluctuation Scale [15], the Clinician Assessment of Fluctuation and One Day Fluctuation Assessment Scale scores [45], or history-taking [14, 17, 18, 20]. Reliable biomarkers of fluctuations in DLB are lacking [46]. In a NACC study of pathologically-defined DLB, there was also no sex effect on the presence of cognitive fluctuations during follow-up, but AD co-pathology reduced the likelihood of cognitive fluctuations more in men than women [20]. There is no convincing evidence of sex differences in the point prevalence or longitudinal progression of fluctuations in DLB, but how this is affected by co-pathology requires further study.

#### 3.3 REM Sleep Behavior Disorder (RBD).

RBD can occur decades before the onset of cognitive or motor features of DLB [15, 47]. In clinical sleep center samples, isolated RBD is more common in men [48-50]. A population study found a prevalence of 1% with no sex differences [48], however, suggesting RBD may be under-detected in women. Men with DLB are more likely to have a history of RBD (46-84%) than women (21-55%) [14, 15, 17, 18, 20] (Table 2). This may be due to greater

muscle phasic activity in men during REM sleep [50] and more vigorous dream-related movements that can cause injury [51-53]. In both men and women with DLB, RBD occurs less frequently in the presence of AD co-pathology [20]. In a longitudinal study at the Mayo Clinic, RBD emerged at a younger age in men than women (median 64 vs. 68 years) [15]. Although RBD developed before the other core DLB features in both sexes, it typically preceded cognitive symptom onset in men and was more likely to develop concurrently with cognitive symptoms in women [15]. The absence of RBD was associated with delays in diagnosis of probable DLB in both sexes [15]. These findings suggest that men with DLB and RBD may have a longer identifiable preclinical phase and window of opportunity for intervention.

### 3.4 Visual hallucinations (VH).

In DLB, VH may be more common in women (38-82%) than men (38-76%), with significant differences in 4 of 7 studies (Table 2) [15, 18, 54, 55]. However, this is not found in some autopsy [14, 20] or clinical samples [17]. While men were more likely to have VH compared to women in pathologically-defined DLB in the NACC, this sex difference disappeared after correcting for demographics and pathology staging [20]. In the presence of AD co-pathology, VH likelihood reduced further for men than women [20]. In both the clinical and pathologic subsets of a Mayo Clinic longitudinal study of clinically probable DLB [15], the time from cognitive symptom onset to developing VH was shorter in women than men. RBD typically preceded the development of VH for both men and women. However, women developed VH within the same time frame as fluctuations and parkinsonism, while men were more likely to develop VH after the other three core features of RBD, parkinsonism, and fluctuations [15]. Overall, research is mixed regarding sex differences in VH, with a suggestion that VH may occur more frequently and earlier in women with DLB. More research is needed regarding potential differences, such as use of dopaminergic treatments and follow-up duration (as VH may appear later in men).

### 3.5 Parkinsonism.

Several studies report a higher prevalence of parkinsonism in men than women with DLB (Table 2) [14, 15, 17, 18, 54]. In one study, sex differences for parkinsonism were statistically significant at diagnosis (59.0% men vs. 44.4% women,  $p=0.027$ ), but not symptom onset (5.0% men vs. 4.5% women) [17]. In another clinically-defined DLB cohort, parkinsonism was more common in men (91% men vs. 83% women,  $p=0.025$ ), but the difference was not significant in the autopsied subset (89% men vs. 88% women,  $p=0.83$ ) [15]. In a NACC pathologically-defined cohort, parkinsonism was less common in women with DLB [20]. In the setting of AD co-pathology, parkinsonism was less common in both sexes, but particularly in men [20]. Two DLB studies reported that women had lower rates of treatment for parkinsonism [15, 20], but milder symptom severity may be a confounder [15]. While women with DLB appear more likely to develop VH and parkinsonism concurrently, it remains unclear whether this timing makes women with DLB at greater risk for adverse effects from dopaminergic therapy (e.g., worsening hallucinations) and consequently, under-treatment of motor symptoms.

### 3.6 Supportive features.

In patients with DLB, Capgras syndrome appears equally likely in men and women [56], but depression and auditory hallucinations may be more common in women [17]. There were no sex differences in severe neuroleptic sensitivity reactions in a small DLB sample [57]. In another study, men more commonly had hyposmia and syncope in early DLB, but there were no sex-based differences in constipation or orthostatic hypotension [17].

### 3.7 Clinical subtypes.

Recent studies use data-driven approaches to identify clinical DLB subtypes (as opposed to pathological or imaging subtypes), with varying findings related to sex. In a study of 81 individuals with probable DLB, cluster analysis yielded three subtypes: cognitive predominant, neuropsychiatric-predominant, and parkinsonism-predominant [58]. Male:female representation was not statistically different across groups [58]. In a factorial analysis using 107 participants in the European DLB consortium [59], hierarchical clustering identified four subtypes: Cluster 1: A $\beta$  and cerebrovascular pathologies, medial temporal atrophy, and cognitive fluctuations; Cluster 2: posterior atrophy, poor cognitive performance, and parkinsonism; Cluster 3: tau pathology, posterior atrophy, and low parkinsonism frequency; and Cluster 4: cognitive fluctuations, higher MMSE scores, virtually normal AD biomarkers, and the least regional brain atrophy and cerebrovascular pathology. While the overall cohort was predominantly men (n=77/107, 72%), Cluster 4 was more common in women (n=12/19, 63%), though most patients in Cluster 4 came from one center [59]. Currently it remains uncertain whether these subtypes are clinically meaningful and whether there are true sex differences between groups.

## 4. Diagnosis of DLB

Individuals with DLB frequently experience diagnostic delays [60]. Sex differences in emergence of DLB core features [15, 42] may explain missed or delayed diagnoses in women. In a NACC study examining pathologically-defined DLB cohorts (excluding individuals with more severe AD co-pathology), women were older than men but manifested fewer core DLB features and were less likely to be diagnosed with probable DLB during life [14]. Delayed diagnosis in women may relate to milder parkinsonism and lower rates of RBD in women compared to men [15]. Diagnosis timing, however, may relate more to the presence or absence of RBD than sex per se: individuals without RBD met criteria for probable DLB later, irrespective of sex [15]. Compared to men, women with DLB tend to present at an older age, show greater cognitive impairment at their initial visit, and have a later emergence of two or more core features needed for a clinical diagnosis of probable DLB [14, 15, 18].

How underlying neuropathology intersects with sex to affect diagnostic timing is uncertain. In one study, DLB was diagnosed earlier in a neuropathologic subgroup of Lewy body disease without neocortical tangles, where individuals were disproportionately male and more likely to have RBD, greater parkinsonism severity, and relatively preserved memory and naming [42]. A NACC study found that sex did not affect the age of dementia onset in “pure” Lewy body disease (with Braak stage 0-II Alzheimer pathology). Male sex,

depression, and apolipoprotein E4 (ApoE-e4) status predicted earlier dementia onset in Lewy body disease with Braak stage III-VI pathology [61]. When evaluating pathologically-defined DLB in another NACC study, more severe AD co-pathology was associated with lower likelihood of DLB core clinical features and clinical DLB diagnoses, and a higher likelihood of dementia and clinical AD diagnoses, in both sexes. Sex modified the degree of these associations [20].

## 5. Neuropathology

DLB neuropathology includes Lewy-related pathology in brainstem-limbic regions or diffusely extending to the neocortex [43, 62]. At least 50% of people with DLB have co-existing AD-related pathology [11, 41, 62]. When A $\beta$  and tau pathology are present in DLB, the burden may still be lower than that of AD [62-64]. Lewy body disease without neocortical AD-related pathology often shows a male predominance, while men and women are similarly represented in pathologic subgroups of Lewy body disease with neocortical AD-related pathology [15, 42, 61, 65, 66]. In a NACC study, women and men had similar Lewy body and amyloid staging, but women had higher Braak Tau staging and less nigral neuronal loss than men [20].

Whether differential hormonal influences on  $\alpha$ -synuclein may contribute to sex differences in neocortical LB pathology is an area of active study. Most studies of hormonal influences on  $\alpha$ -synuclein have been conducted in PD [9, 67]. Sex hormones, especially estriol, contain antioxidant properties that harbor anti-aggregation and fibril-destabilizing effects on  $\alpha$ -synuclein in vitro [68]. Women may be protected against  $\alpha$ -synucleinopathies pre-menopause, after which  $\alpha$ -synuclein aggregation accelerates with decreasing estradiol levels, especially during the second half of menopause [4]. Whether the older age of onset of DLB in women [12, 16, 31] relates to these hormonal differences remains unknown.

## 6. Biomarkers

### 6.1 DLB Biomarkers.

While skin biopsy and CSF RT-QuIC are promising synuclein biomarkers [69], limited research evaluates sex differences in these assays. A small 2012 study reported lower levels of CSF  $\alpha$ -synuclein in women with DLB versus controls, but no difference in men with DLB versus controls [70]. A recent study found lower CSF  $\alpha$ -synuclein levels in women compared to men [18]. Notably, both studies used ELISA techniques rather than newer approaches assessing  $\alpha$ -synuclein seeding activity (e.g., RT-QuIC).

Apart from polysomnography to investigate RBD, there is minimal research on sex differences for indicative (dopamine transporter imaging,  $^{123}\text{I}$ -metaiodobenzylguanidine cardiac scintigraphy) or supportive (MRI, electroencephalography, fluorodeoxyglucose-positron emission tomography) biomarkers [43]. In a study examining DAT-SPECT in patients with parkinsonian disorders including DLB, correcting for age and sex did not improve sensitivity or specificity [71]. Studies of MRI atrophy patterns in DLB often control for sex rather than investigating it as a variable of interest [72, 73]. In individuals with probable DLB from the European DLB cohort, male sex was associated with

abnormal frontal cortical atrophy but sex did not predict atrophy in medial temporal or posterior cortices [74]. In DLB, female sex has been associated with higher white matter hyperintensity volumes but not with infarct frequency [66, 75, 76]. These findings highlight the need to evaluate for potential biomarker differences between sexes in DLB, not just control for sex as a confounding variable.

## 6.2 AD-related Biomarkers.

Age-adjusted AD-CSF biomarker analysis from 223 patients with DLB within the Amsterdam Dementia Cohort revealed women had lower levels of A $\beta$ 42 compared to men, but there were no differences in total or phosphorylated tau concentrations [18]. In a multicohort DLB study, women were more likely to have positive imaging biomarkers using Pittsburgh compound B and Tau-PET ([<sup>18</sup>F]AV-1451) or CSF A $\beta$ 42 and phosphorylated tau, including dual amyloid and tau positivity. The presence of A $\beta$  pathology was associated with greater cognitive impairment and participants with positive tau biomarkers demonstrated fewer core clinical DLB features [54]. Additional Tau-PET [77] and Amyloid-PET [78] studies included 77-86% male participants and did not specifically report sex differences. There is clearly a need for research into biomarkers of AD pathology in DLB.

## 6.3 Genetics.

ApoE-e4 is associated with an increased dementia risk in the synucleinopathies [79, 80]. In an autopsy-confirmed cohort of 652 patients with Lewy body disease (31% with dementia and parkinsonism, 43% with dementia without parkinsonism, 26% with parkinsonism without dementia), ApoE-e4 status was more common in those with a Thal phase of III-V and with neocortical tangle pathology (Braak NFT stage IV-VI). ApoE-e4 has also been associated with more severe Lewy-related pathology, even in cases with low AD pathology [81]. In one study of clinically probable DLB, women carried the ApoE-e4 allele more frequently than men [18], but another study showed no sex differences in the proportion of ApoE-e4 carriers [14]. GBA mutations are associated with an increased susceptibility for Lewy body disease and dementia [82, 83]. Some studies suggest that GBA mutations are associated with earlier symptom onset in men with DLB [84, 85]; however, a recent meta-analysis found no sex differences in GBA variants in DLB [86]. More work is needed to better understand the clinical heterogeneity of DLB and how it relates to genetic factors in men and women.

# 7. Progression

## 7.1 Clinical Progression.

Studies examining sex differences in cognitive progression have mixed findings, even within the same cohort. A study including 835 participants with probable DLB from the European DLB Consortium (54% male) did not find sex significant differences in the rate of decline on the MMSE over two years [87]. In a subgroup analysis (n=100), however, men had a more rapid MMSE decline over two years [87, 88]. Other studies have not found an effect of sex on cognitive decline [89-92]. It is possible that decline relates more to pathology than sex given that faster cognitive decline is associated with an AD-CSF profile [88] and the synergistic effects of  $\alpha$ -synuclein, amyloid- $\beta$ , and tau pathologies [11, 42, 93].



## 7.2 Survival.

Whether sex affects survival in DLB remains unclear. In an autopsy-confirmed DLB sample, women had shorter median survival following dementia onset than men (6.6 versus 8.1 years), though women died at an older age [94]. In a clinical cohort, however, sex did not affect median survival after first presentation with cognitive impairment in DLB (men: 3.57 years, women: 3.81 years) [95]. Similarly, sex was not a significant predictor of survival in a clinically-defined NACC DLB cohort [96] and duration of cognitive decline was similar between sexes in a pathological NACC study [20]. Discrepancies may relate to differences in cohort identification. In another NACC study with neuropathological data, male sex was associated with shorter survival in individuals with transitional or diffuse Lewy body disease pathology, but there was no difference in survival by sex in the subset of individuals with an antemortem diagnosis of probable DLB. Most of this sample had a Braak NFT stage of IV or greater [97].

Sex differences in clinical progression and survival are likely influenced by the extent of Lewy-related and AD-related pathology [62, 97]. Increased CSF phosphorylated tau is linked to shorter survival and faster rates of cognitive decline among individuals with DLB [88, 98, 99]. Additionally, while available studies focus on potential biological sex differences, survival is also influenced by gender roles, relating to the presence, identity, and involvement of caregivers and whether the person with DLB requires residential/institutional care.

## 8. Management

Under-representation of women in DLB clinical trials limits assessment of potential sex differences in treatment response. Completed studies on [Clinicaltrials.gov](https://clinicaltrials.gov) (accessed December 2, 2022) report only ~15-37% female participants in DLB drug trials. Additionally, although 15% of the sample in a trial of nelotanserin for RBD were female, none of the female participants were randomized to the treatment group [100]. Similarly, studies examining nucleus basalis of Meynert deep brain stimulation for DLB included 0-17% female participants [101, 102]. Reasons for low female participation are likely complex, relating to prevalence of DLB diagnoses in countries where trials are performed, the fact that women meet DLB diagnostic criteria less often and later than men, and potential differences in caregiving structures for men versus women.

## 9. Caregiving

While population-based studies of caregivers of individuals with DLB are lacking, 67-90% of caregivers in identified DLB studies were women [103-107]. Most of these caregivers were spouses, but a substantial minority were daughters and daughters-in-law [103-105, 108]. Men also serve as caregivers, but little is known about their unique experiences.

Higher caregiver burden in DLB is associated with dementia severity [103], neuropsychiatric symptoms (including sleep disturbance) [104-106, 108, 109], need for assistance with activities of daily living [105, 108, 109], caregiver anxiety [104], and social isolation [105]. Whether sex affects caregiving in DLB is uncertain. In one study combining caregivers of

individuals with DLB and AD, female caregivers reported higher relative stress than male caregivers [108]. In a study combining caregivers of individuals with DLB, PD dementia, and AD, sex was significantly correlated with caregiver burden, but not after adjusting for multiple comparisons [103]. Caregiver sex was not associated with pre-loss grief in a recent study in DLB [107]. Given that sex and gender likely align for many study participants, these results may reflect gender roles as well as potential biological differences affecting caregiving. Biological sex differences (e.g. physical stature, potentially affective symptoms) have caregiving implications, but so does gender (e.g. relating to societal and familial roles).

## 10. Limitations

This is a focused review of existing literature and while it is unlikely that publications specifically targeting sex differences in DLB were overlooked, studies in which sex was examined in secondary analyses may have been missed. The review also focused on sex rather than gender, as studies specifically evaluating gender were scarce.

## 11. Future Directions

This review focused on sex-related differences in DLB because research on gender in this context is lacking. While sex refers to biological distinctions (sex chromosomes, X and Y), gender is based on psychosocial, environmental, and cultural influences [110]. In the fields of dementia and AD, examples of gender-related differences in risk factors include education, income, marital status, smoking, depression, diet/exercise, and cerebrovascular comorbidities [111, 112]. Gender may also affect clinical diagnosis of dementia, as gender differences in functional roles can affect timing of recognizing functional decline [113]. While sex and gender intersect, more studies are needed to distinguish sex and gender effects in DLB, especially pertaining to modifiable risk factors and patient and caregiving experiences.

Clinical awareness of possible sex differences in clinical presentation and biomarker status of DLB will support more accurate and timely diagnoses, including at prodromal stages. Identification of endophenotypes of DLB, based on unique combinations of demographic and clinical features, along with pathological biomarkers, would inform diagnostic criteria revision, understanding of expected disease progression, and prediction of treatment response. Consideration of differential effects of sex and gender on individuals living with DLB and their caregivers [114-116] supports a trend towards individualized care and fulfillment of unmet needs (Table 3). Moving forward, DLB clinical trials should stratify randomization by sex so that women participants are not solely allocated to the placebo arm [100]. Excluding individuals with DLB and AD co-pathology also risks exacerbating under-representation of women in DLB trials [114, 115]. Clinical trial results should include sex/gender-specific data to identify whether treatment differences exist and allow for future meta-analyses.

## 12. Conclusions

DLB impacts women living with and caring for others living with the disease (Figure 1). Current literature supports sex differences in clinical presentations, pathology, and clinico-pathological correlations, informing efforts for early detection, appropriate counseling, and patient and caregiver support. Additional research is needed to address the presented gaps in the current understanding of potential sex and gender differences in biomarkers and treatment responses in DLB.

## Funding

Co-authors on this review have Lewy body dementia grants from the National Institutes of Health (K23AG073525-01A1, R21AG074368, K99AG073453, R01AG068128, P30AG047266, U01NS100620), which in part supported this work. The content is solely the responsibility of the authors and does not necessarily represent the official views of the National Institutes of Health.

## Declaration of competing interest

Dr. Chiu receives research support from the NIH (K23AG073525-01A1). Dr. Wyman receives research support from the NIH (R21AG074368). Dr. Ferman is supported by NIH U01NS100620, U19AG71754, P30AG62677 and from the Dorothy and Harry T. Mangurian Jr. Lewy Body Dementia Program. Dr. Bayram receives research support from the NIH (K99AG073453). Dr. Holden receives research support from the NIH (R21AG072153) and as the local PI of a Lewy Body Dementia Association Research Center of Excellence. Dr. Choudhury receives research support from Lewy Body Dementia Association and Arizona Alzheimer's Research Consortium. Dr. Armstrong receives research support from the NIH (R01AG068128, P30AG047266, R01NS121099, R44AG062072), the Florida Department of Health (grant 20A08), and as the local PI of a Lewy Body Dementia Association Research Center of Excellence. She serves on the DSMBs for the Alzheimer's Therapeutic Research Institute/Alzheimer's Clinical Trial Consortium and the Alzheimer's Disease Cooperative Study.

## REFERENCE:

- [1]. Gottgens I, van Halteren AD, de Vries NM, Meinders MJ, Ben-Shlomo Y, Bloem BR, Darweesh SKL, Oertelt-Prigione S, The Impact of Sex and Gender on the Multidisciplinary Management of Care for Persons With Parkinson's Disease, *Front Neurol* 11 (2020) 576121. [PubMed: 33071952]
- [2]. Ullah MF, Ahmad A, Bhat SH, Abu-Duhier FM, Barreto GE, Ashraf GM, Impact of sex differences and gender specificity on behavioral characteristics and pathophysiology of neurodegenerative disorders, *Neurosci Biobehav Rev* 102 (2019) 95–105. [PubMed: 30959072]
- [3]. Podcasy JL, Epperson CN, Considering sex and gender in Alzheimer disease and other dementias, *Dialogues Clin Neurosci* 18(4) (2016) 437–446. [PubMed: 28179815]
- [4]. Mielke MM, Vemuri P, Rocca WA, Clinical epidemiology of Alzheimer's disease: assessing sex and gender differences, *Clin Epidemiol* 6 (2014) 37–48. [PubMed: 24470773]
- [5]. Ferretti MT, Martinkova J, Biskup E, Benke T, Gialdini G, Nedelska Z, Rauen K, Mantua V, Religa D, Hort J, Santuccione Chadha A, Schmidt R, Sex and gender differences in Alzheimer's disease: current challenges and implications for clinical practice: Position paper of the Dementia and Cognitive Disorders Panel of the European Academy of Neurology, *Eur J Neurol* 27(6) (2020) 928–943. [PubMed: 32056347]
- [6]. Barnes LL, Wilson RS, Bienias JL, Schneider JA, Evans DA, Bennett DA, Sex differences in the clinical manifestations of Alzheimer disease pathology, *Arch Gen Psychiatry* 62(6) (2005) 685–91. [PubMed: 15939846]
- [7]. Koran MEI, Wagener M, Hohman TJ, Sex differences in the association between AD biomarkers and cognitive decline, *Brain Imaging Behav* 11(1) (2017) 205–213. [PubMed: 26843008]
- [8]. Moisan F, Kab S, Mohamed F, Canonico M, Le Guern M, Quintin C, Carcaillon L, Nicolau J, Dupont N, Singh-Manoux A, Boussac-Zarebska M, Elbaz A, Parkinson disease male-to-female ratios increase with age: French nationwide study and meta-analysis, *J Neurol Neurosurg Psychiatry* 87(9) (2016) 952–7. [PubMed: 26701996]

- [9]. Vaidya B, Dhamija K, Guru P, Sharma SS, Parkinson's disease in women: Mechanisms underlying sex differences, *Eur J Pharmacol* 895 (2021) 173862. [PubMed: 33450279]
- [10]. Jurado-Coronel JC, Cabezas R, Avila Rodriguez MF, Echeverria V, Garcia-Segura LM, Barreto GE, Sex differences in Parkinson's disease: Features on clinical symptoms, treatment outcome, sexual hormones and genetics, *Front Neuroendocrinol* 50 (2018) 18–30. [PubMed: 28974386]
- [11]. Irwin DJ, Grossman M, Weintraub D, Hurtig HI, Duda JE, Xie SX, Lee EB, Van Deerlin VM, Lopez OL, Kofler JK, Nelson PT, Jicha GA, Woltjer R, Quinn JF, Kaye J, Leverenz JB, Tsuang D, Longfellow K, Yearout D, Kukull W, Keene CD, Montine TJ, Zabetian CP, Trojanowski JQ, Neuropathological and genetic correlates of survival and dementia onset in synucleinopathies: a retrospective analysis, *Lancet Neurol* 16(1) (2017) 55–65. [PubMed: 27979356]
- [12]. Savica R, Grossardt BR, Bower JH, Boeve BF, Ahlskog JE, Rocca WA, Incidence of dementia with Lewy bodies and Parkinson disease dementia, *JAMA Neurol* 70(11) (2013) 1396–402. [PubMed: 24042491]
- [13]. Lemstra AW, de Beer MH, Teunissen CE, Schreuder C, Scheltens P, van der Flier WM, Sikkes SA, Concomitant AD pathology affects clinical manifestation and survival in dementia with Lewy bodies, *J Neurol Neurosurg Psychiatry* 88(2) (2017) 113–118. [PubMed: 27794030]
- [14]. Bayram E, Coughlin DG, Banks SJ, Litvan I, Sex differences for phenotype in pathologically defined dementia with Lewy bodies, *J Neurol Neurosurg Psychiatry* 92(7) (2021) 745–750. [PubMed: 33563809]
- [15]. Choudhury P, Graff-Radford J, Aakre JA, Wurtz L, Knopman DS, Graff-Radford NR, Kantarci K, Forsberg LK, Fields JA, Pedraza O, Chen Q, Miyagawa T, Day GS, Tipton P, Savica R, Botha H, Lachner C, Dredla B, Reichard RR, Petersen RC, Dickson DW, Boeve BF, Ferman TJ, The temporal onset of the core features in dementia with Lewy bodies, *Alzheimers Dement* (2021).
- [16]. Mouton A, Blanc F, Gros A, Manera V, Fabre R, Sauleau E, Gomez-Luporsi I, Tifratene K, Friedman L, Thummler S, Pradier C, Robert PH, David R, Sex ratio in dementia with Lewy bodies balanced between Alzheimer's disease and Parkinson's disease dementia: a cross-sectional study, *Alzheimers Res Ther* 10(1) (2018) 92. [PubMed: 30208961]
- [17]. Utsumi K, Fukatsu R, Yamada R, Takamaru Y, Hara Y, Yasumura S, Characteristics of initial symptoms and symptoms at diagnosis in probable dementia with Lewy body disease: incidence of symptoms and gender differences, *Psychogeriatrics* 20(5) (2020) 737–745. [PubMed: 32743894]
- [18]. van de Beek M, Babapour Mofrad R, van Steenoven I, Vanderstichele H, Scheltens P, Teunissen CE, Lemstra AW, van der Flier WM, Sex-specific associations with cerebrospinal fluid biomarkers in dementia with Lewy bodies, *Alzheimers Res Ther* 12(1) (2020) 44. [PubMed: 32303272]
- [19]. van de Beek M, van Steenoven I, van der Zande JJ, Porcelijn I, Barkhof F, Stam CJ, Raijmakers P, Scheltens P, Teunissen CE, van der Flier WM, Lemstra AW, Characterization of symptoms and determinants of disease burden in dementia with Lewy bodies: DEVELOP design and baseline results, *Alzheimers Res Ther* 13(1) (2021) 53. [PubMed: 33637117]
- [20]. Bayram E, Coughlin DG, Litvan I, Sex Differences for Clinical Correlates of Alzheimer's Pathology in People with Lewy Body Pathology, *Mov Disord* 37(7) (2022) 1505–1515. [PubMed: 35531707]
- [21]. Vann Jones SA, O'Brien JT, The prevalence and incidence of dementia with Lewy bodies: a systematic review of population and clinical studies, *Psychol Med* 44(4) (2014) 673–83. [PubMed: 23521899]
- [22]. Hogan DB, Fiest KM, Roberts JI, Maxwell CJ, Dykeman J, Pringsheim T, Steeves T, Smith EE, Pearson D, Jette N, The Prevalence and Incidence of Dementia with Lewy Bodies: a Systematic Review, *Can J Neurol Sci* 43 Suppl 1 (2016) S83–95. [PubMed: 27307129]
- [23]. Kane JPM, Surendranathan A, Bentley A, Barker SAH, Taylor JP, Thomas AJ, Allan LM, McNally RJ, James PW, McKeith IG, Burn DJ, O'Brien JT, Clinical prevalence of Lewy body dementia, *Alzheimers Res Ther* 10(1) (2018) 19. [PubMed: 29448953]
- [24]. Chin KS, Yassi N, Churilov L, Masters CL, Watson R, Prevalence and clinical associations of tau in Lewy body dementias: A systematic review and meta-analysis, *Parkinsonism Relat Disord* 80 (2020) 184–193. [PubMed: 33260030]

- [25]. Ferman TJ, Boeve BF, Smith GE, Lin SC, Silber MH, Pedraza O, Wszolek Z, Graff-Radford NR, Uitti R, Van Gerpen J, Pao W, Knopman D, Pankratz VS, Kantarci K, Boot B, Parisi JE, Dugger BN, Fujishiro H, Petersen RC, Dickson DW, Inclusion of RBD improves the diagnostic classification of dementia with Lewy bodies, *Neurology* 77(9) (2011) 875–82. [PubMed: 21849645]
- [26]. Fereshtehnejad SM, Religa D, Westman E, Aarsland D, Løkke J, Eriksdotter M, Demography, diagnostics, and medication in dementia with Lewy bodies and Parkinson's disease with dementia: data from the Swedish Dementia Quality Registry (SveDem), *Neuropsychiatr Dis Treat* 9 (2013) 927–35. [PubMed: 23847419]
- [27]. Cerri S, Mus L, Blandini F, Parkinson's Disease in Women and Men: What's the Difference?, *J Parkinsons Dis* 9(3) (2019) 501–515. [PubMed: 31282427]
- [28]. Boeve BF, Silber MH, Ferman TJ, Lin SC, Benarroch EE, Schmeichel AM, Ahlskog JE, Caselli RJ, Jacobson S, Sabbagh M, Adler C, Woodruff B, Beach TG, Iranzo A, Gelpi E, Santamaria J, Tolosa E, Singer C, Mash DC, Luca C, Arnulf I, Duyckaerts C, Schenck CH, Mahowald MW, Dauvilliers Y, Graff-Radford NR, Wszolek ZK, Parisi JE, Dugger B, Murray ME, Dickson DW, Clinicopathologic correlations in 172 cases of rapid eye movement sleep behavior disorder with or without a coexisting neurologic disorder, *Sleep Med* 14(8) (2013) 754–62. [PubMed: 23474058]
- [29]. Postuma RB, Iranzo A, Hu M, Hög B, Boeve BF, Manni R, Oertel WH, Arnulf I, Ferini-Strambi L, Puligheddu M, Antelmi E, Cochen De Cock V, Arnaldi D, Mollenhauer B, Videnovic A, Sonka K, Jung KY, Kunz D, Dauvilliers Y, Provini F, Lewis SJ, Buskova J, Pavlova M, Heidebreder A, Montplaisir JY, Santamaria J, Barber TR, Stefani A, St Louis EK, Terzaghi M, Janzen A, Leu-Semenescu S, Plazzi G, Nobili F, Sixel-Doering F, Dusek P, Bes F, Cortelli P, Ehgoetz Martens K, Gagnon JF, Gaig C, Zucconi M, Trenkwalder C, Gan-Or Z, Lo C, Rolinski M, Mahlkecht P, Holzkecht E, Boeve AR, Teigen LN, Toscano G, Mayer G, Morbelli S, Dawson B, Pelletier A, Risk and predictors of dementia and parkinsonism in idiopathic REM sleep behaviour disorder: a multicentre study, *Brain* 142(3) (2019) 744–759. [PubMed: 30789229]
- [30]. Yao C, Fereshtehnejad SM, Keezer MR, Wolfson C, Pelletier A, Postuma RB, Risk factors for possible REM sleep behavior disorder: A CLSA population-based cohort study, *Neurology* (2018).
- [31]. Zahirovic I, Wattmo C, Torisson G, Minthon L, Londos E, Prevalence of Dementia With Lewy Body Symptoms: A Cross-Sectional Study in 40 Swedish Nursing Homes, *J Am Med Dir Assoc* 17(8) (2016) 706–11. [PubMed: 27168051]
- [32]. Fujishiro H, Torii Y, Iritani S, Gender differences in early presentation and pathological subtypes in dementia with Lewy bodies, *Psychogeriatrics* 21(1) (2021) 142–143. [PubMed: 33107106]
- [33]. Mori E, Ikeda M, Nagai R, Matsuo K, Nakagawa M, Kosaka K, Long-term donepezil use for dementia with Lewy bodies: results from an open-label extension of Phase III trial, *Alzheimers Res Ther* 7(1) (2015) 5. [PubMed: 25713600]
- [34]. Komatsu J, Samuraki M, Nakajima K, Arai H, Arai H, Arai T, Asada T, Fujishiro H, Hanyu H, Iizuka O, Iseki E, Kashihara K, Kosaka K, Maruno H, Mizukami K, Mizuno Y, Mori E, Nakamura H, Nakano S, Nakashima K, Nishio Y, Orimo S, Takahashi A, Taki J, Tokuda T, Urakami K, Utsumi K, Wada K, Washimi Y, Yamashina S, Yamasaki J, Yoshita M, Yamada M, (123)I-MIBG myocardial scintigraphy for the diagnosis of DLB: a multicentre 3-year follow-up study, *J Neurol Neurosurg Psychiatry* 89(11) (2018) 1167–1173. [PubMed: 29853532]
- [35]. Surendranathan A, Kane JPM, Bentley A, Barker SAH, Taylor JP, Thomas AJ, Allan LM, McNally RJ, James PW, McKeith IG, Burn DJ, O'Brien JT, Clinical diagnosis of Lewy body dementia, *BJPsych Open* 6(4) (2020) e61.
- [36]. Nag S, Barnes LL, Yu L, Buchman AS, Bennett DA, Schneider JA, Wilson RS, Association of Lewy Bodies With Age-Related Clinical Characteristics in Black and White Decedents, *Neurology* 97(8) (2021) e825–e835. [PubMed: 34088871]
- [37]. Kurasz AM, Smith GE, McFarland MG, Armstrong MJ, Ethnoracial Differences in Lewy Body Diseases with Cognitive Impairment, *J Alzheimers Dis* 77(1) (2020) 165–174. [PubMed: 32804137]

- [38]. Ciafone J, Little B, Thomas AJ, Gallagher P, The Neuropsychological Profile of Mild Cognitive Impairment in Lewy Body Dementias, *J Int Neuropsychol Soc* 26(2) (2020) 210–225. [PubMed: 31826772]
- [39]. Azar M, Chapman S, Gu Y, Leverenz JB, Stern Y, Cosentino S, Cognitive tests aid in clinical differentiation of Alzheimer's disease versus Alzheimer's disease with Lewy body disease: Evidence from a pathological study, *Alzheimers Dement* 16(8) (2020) 1173–1181. [PubMed: 32558217]
- [40]. Ferman TJ, Smith GE, Boeve BF, Graff-Radford NR, Lucas JA, Knopman DS, Petersen RC, Ivnik RJ, Wszolek Z, Uitti R, Dickson DW, Neuropsychological differentiation of dementia with Lewy bodies from normal aging and Alzheimer's disease, *Clin Neuropsychol* 20(4) (2006) 623–36. [PubMed: 16980250]
- [41]. Peavy GM, Edland SD, Toole BM, Hansen LA, Galasko DR, Mayo AM, Phenotypic differences based on staging of Alzheimer's neuropathology in autopsy-confirmed dementia with Lewy bodies, *Parkinsonism Relat Disord* 31 (2016) 72–78. [PubMed: 27475955]
- [42]. Ferman TJ, Aoki N, Boeve BF, Aakre JA, Kantarci K, Graff-Radford J, Parisi JE, Van Gerpen JA, Graff-Radford NR, Uitti RJ, Pedraza O, Murray ME, Wszolek ZK, Reichard RR, Fields JA, Ross OA, Knopman DS, Petersen RC, Dickson DW, Subtypes of dementia with Lewy bodies are associated with  $\alpha$ -synuclein and tau distribution, *Neurology* 95(2) (2020) e155–e165. [PubMed: 32561678]
- [43]. McKeith IG, Boeve BF, Dickson DW, Halliday G, Taylor JP, Weintraub D, Aarsland D, Galvin J, Attems J, Ballard CG, Bayston A, Beach TG, Blanc F, Bohnen N, Bonanni L, Bras J, Brundin P, Burn D, Chen-Plotkin A, Duda JE, El-Agnaf O, Feldman H, Ferman TJ, Ffytche D, Fujishiro H, Galasko D, Goldman JG, Gomperts SN, Graff-Radford NR, Honig LS, Iranzo A, Kantarci K, Kaufer D, Kukull W, Lee VMY, Leverenz JB, Lewis S, Lippa C, Lunde A, Masellis M, Masliah E, McLean P, Mollenhauer B, Montine TJ, Moreno E, Mori E, Murray M, O'Brien JT, Orimo S, Postuma RB, Ramaswamy S, Ross OA, Salmon DP, Singleton A, Taylor A, Thomas A, Tiraboschi P, Toledo JB, Trojanowski JQ, Tsuang D, Walker Z, Yamada M, Kosaka K, Diagnosis and management of dementia with Lewy bodies: Fourth consensus report of the DLB Consortium, *Neurology* 89(1) (2017) 88–100. [PubMed: 28592453]
- [44]. van Steenoven I, Aarsland D, Weintraub D, Londos E, Blanc F, van der Flier WM, Teunissen CE, Mollenhauer B, Fladby T, Kramberger MG, Bonanni L, Lemstra AW, Cerebrospinal Fluid Alzheimer's Disease Biomarkers Across the Spectrum of Lewy Body Diseases: Results from a Large Multicenter Cohort, *J Alzheimers Dis* 54(1) (2016) 287–95. [PubMed: 27567832]
- [45]. Bradshaw J, Saling M, Hopwood M, Anderson V, Brodtmann A, Fluctuating cognition in dementia with Lewy bodies and Alzheimer's disease is qualitatively distinct, *J Neurol Neurosurg Psychiatry* 75(3) (2004) 382–7. [PubMed: 14966152]
- [46]. Matar E, Shine JM, Halliday GM, Lewis SJG, Cognitive fluctuations in Lewy body dementia: towards a pathophysiological framework, *Brain* 143(1) (2020) 31–46. [PubMed: 31612904]
- [47]. Claassen DO, Josephs KA, Ahlskog JE, Silber MH, Tippmann-Peikert M, Boeve BF, REM sleep behavior disorder preceding other aspects of synucleinopathies by up to half a century, *Neurology* 75(6) (2010) 494–9. [PubMed: 20668263]
- [48]. Haba-Rubio J, Frauscher B, Marques-Vidal P, Toriel J, Tობback N, Andries D, Preisig M, Vollenweider P, Postuma R, Heinzer R, Prevalence and determinants of rapid eye movement sleep behavior disorder in the general population, *Sleep* 41(2) (2018).
- [49]. Fernandez-Arcos A, Iranzo A, Serradell M, Gaig C, Santamaria J, The Clinical Phenotype of Idiopathic Rapid Eye Movement Sleep Behavior Disorder at Presentation: A Study in 203 Consecutive Patients, *Sleep* 39(1) (2016) 121–32. [PubMed: 26940460]
- [50]. Bugalho P, Salavisa M, Factors Influencing the Presentation of REM Sleep Behavior Disorder: The Relative Importance of Sex, Associated Neurological Disorder, and Context of Referral to Polysomnography, *J Clin Sleep Med* 15(12) (2019) 1789–1798. [PubMed: 31855164]
- [51]. Bjornara KA, Dietrichs E, Toft M, REM sleep behavior disorder in Parkinson's disease--is there a gender difference?, *Parkinsonism Relat Disord* 19(1) (2013) 120–2. [PubMed: 22726815]
- [52]. Zhou J, Zhang J, Li Y, Du L, Li Z, Lei F, Wing YK, Kushida CA, Zhou D, Tang X, Gender differences in REM sleep behavior disorder: a clinical and polysomnographic study in China, *Sleep Med* 16(3) (2015) 414–8. [PubMed: 25660814]

- [53]. Fernandez-Arcos A, Morenas-Rodriguez E, Santamaria J, Sanchez-Valle R, Llado A, Gaig C, Lleo A, Iranzo A, Clinical and video-polysomnographic analysis of rapid eye movement sleep behavior disorder and other sleep disturbances in dementia with Lewy bodies, *Sleep* 42(7) (2019).
- [54]. Ferreira D, Przybelski SA, Lesnick TG, Lemstra AW, Londres E, Blanc F, Nedelska Z, Schwarz CG, Graff-Radford J, Senjem ML, Fields JA, Knopman DS, Savica R, Ferman TJ, Graff-Radford NR, Lowe VJ, Jack CR Jr., Petersen RC, Mollenhauer B, Garcia-Ptacek S, Abdelnour C, Hort J, Bonanni L, Oppedal K, Kramberger MG, Boeve BF, Aarsland D, Westman E, Kantarci K, beta-Amyloid and tau biomarkers and clinical phenotype in dementia with Lewy bodies, *Neurology* 95(24) (2020) e3257–e3268. [PubMed: 32989106]
- [55]. Chiu PY, Teng PR, Wei CY, Wang CW, Tsai CT, Gender difference in the association and presentation of visual hallucinations in dementia with Lewy bodies: a cross-sectional study, *Int J Geriatr Psychiatry* 33(1) (2018) 193–199. [PubMed: 28295599]
- [56]. Thaipisuttikul P, Lobach I, Zweig Y, Gurnani A, Galvin JE, Capgras syndrome in Dementia with Lewy Bodies, *Int Psychogeriatr* 25(5) (2013) 843–9. [PubMed: 23211760]
- [57]. Aarsland D, Perry R, Larsen JP, McKeith IG, O'Brien JT, Perry EK, Burn D, Ballard CG, Neuroleptic sensitivity in Parkinson's disease and parkinsonian dementias, *J Clin Psychiatry* 66(5) (2005) 633–7. [PubMed: 15889951]
- [58]. Morenas-Rodriguez E, Sala I, Subirana A, Pascual-Goni E, Sanchez-Saudinos MB, Alcolea D, Illan-Gala I, Carmona-Iragui M, Ribosa-Nogue R, Camacho V, Blesa R, Fortea J, Lleo A, Clinical Subtypes of Dementia with Lewy Bodies Based on the Initial Clinical Presentation, *J Alzheimers Dis* 64(2) (2018) 505–513. [PubMed: 29889064]
- [59]. Abdelnour C, Ferreira D, van de Beek M, Cedres N, Oppedal K, Cavallin L, Blanc F, Bousiges O, Wahlund LO, Pilotto A, Padovani A, Boada M, Pagonabarraga J, Kulisevsky J, Aarsland D, Lemstra AW, Westman E, Parsing heterogeneity within dementia with Lewy bodies using clustering of biological, clinical, and demographic data, *Alzheimers Res Ther* 14(1) (2022) 14. [PubMed: 35063023]
- [60]. Zweig YR, Galvin JE, Lewy body dementia: the impact on patients and caregivers, *Alzheimers Res Ther* 6(2) (2014) 21. [PubMed: 25031635]
- [61]. Schaffert J, LoBue C, White CL 3rd, Wilmoth K, Didehbani N, Lacritz L, Nguyen T, Peters ME, Fields L, Li C, Cullum CM, Risk factors for earlier dementia onset in autopsy-confirmed Alzheimer's disease, mixed Alzheimer's with Lewy bodies, and pure Lewy body disease, *Alzheimers Dement* 16(3) (2020) 524–530. [PubMed: 32043803]
- [62]. Ferman TJ, Aoki N, Crook JE, Murray ME, Graff-Radford NR, van Gerpen JA, Uitti RJ, Wszolek ZK, Graff-Radford J, Pedraza O, Kantarci K, Boeve BF, Dickson DW, The limbic and neocortical contribution of  $\alpha$ -synuclein, tau, and amyloid  $\beta$  to disease duration in dementia with Lewy bodies, *Alzheimers Dement* 14(3) (2018) 330–339. [PubMed: 29100980]
- [63]. Walker L, McAleese KE, Thomas AJ, Johnson M, Martin-Ruiz C, Parker C, Colloby SJ, Jellinger K, Attems J, Neuropathologically mixed Alzheimer's and Lewy body disease: burden of pathological protein aggregates differs between clinical phenotypes, *Acta Neuropathol* 129(5) (2015) 729–48. [PubMed: 25758940]
- [64]. Kantarci K, Lowe VJ, Chen Q, Przybelski SA, Lesnick TG, Schwarz CG, Senjem ML, Gunter JL, Jack CR Jr., Graff-Radford J, Jones DT, Knopman DS, Graff-Radford N, Ferman TJ, Parisi JE, Dickson DW, Petersen RC, Boeve BF, Murray ME, beta-Amyloid PET and neuropathology in dementia with Lewy bodies, *Neurology* 94(3) (2020) e282–e291. [PubMed: 31862783]
- [65]. Nelson PT, Schmitt FA, Jicha GA, Kryscio RJ, Abner EL, Smith CD, Van Eldik LJ, Markesbery WR, Association between male gender and cortical Lewy body pathology in large autopsy series, *J Neurol* 257(11) (2010) 1875–81. [PubMed: 20563821]
- [66]. Barnes LL, Lamar M, Schneider JA, Sex differences in mixed neuropathologies in community-dwelling older adults, *Brain Res* 1719 (2019) 11–16. [PubMed: 31128096]
- [67]. Liu B, Dluzen DE, Oestrogen and nigrostriatal dopaminergic neurodegeneration: animal models and clinical reports of Parkinson's disease, *Clin Exp Pharmacol Physiol* 34(7) (2007) 555–65. [PubMed: 17581209]

- [68]. Hirohata M, Ono K, Morinaga A, Ikeda T, Yamada M, Anti-aggregation and fibril-destabilizing effects of sex hormones on alpha-synuclein fibrils in vitro, *Exp Neurol* 217(2) (2009) 434–9. [PubMed: 19289119]
- [69]. Scott GD, Arnold MR, Beach TG, Gibbons CH, Kanthasamy AG, Lebovitz RM, Lemstra AW, Shaw LM, Teunissen CE, Zetterberg H, Taylor AS, Graham TC, Boeve BF, Gomperts SN, Graff-Radford NR, Moussa C, Poston KL, Rosenthal LS, Sabbagh MN, Walsh RR, Weber MT, Armstrong MJ, Bang JA, Bozoki AC, Domoto-Reilly K, Duda JE, Fleisher JE, Galasko DR, Galvin JE, Goldman JG, Holden SK, Honig LS, Huddlestone DE, Leverenz JB, Litvan I, Manning CA, Marder KS, Pantelyat AY, Pelak VS, Scharre DW, Sha SJ, Shill HA, Mari Z, Quinn JF, Irwin DJ, Fluid and Tissue Biomarkers of Lewy Body Dementia: Report of an LBDA Symposium, *Front Neurol* 12 (2021) 805135. [PubMed: 35173668]
- [70]. Wennstrom M, Londos E, Minthon L, Nielsen HM, Altered CSF orexin and alpha-synuclein levels in dementia patients, *J Alzheimers Dis* 29(1) (2012) 125–32. [PubMed: 22207004]
- [71]. Schmitz-Steinkruger H, Lange C, Apostolova I, Mathies FL, Frings L, Klutmann S, Hellwig S, Meyer PT, Buchert R, Impact of age and sex correction on the diagnostic performance of dopamine transporter SPECT, *Eur J Nucl Med Mol Imaging* 48(5) (2021) 1445–1459. [PubMed: 33130960]
- [72]. Whitwell JL, Weigand SD, Shiung MM, Boeve BF, Ferman TJ, Smith GE, Knopman DS, Petersen RC, Benarroch EE, Josephs KA, Jack CR Jr., Focal atrophy in dementia with Lewy bodies on MRI: a distinct pattern from Alzheimer's disease, *Brain* 130(Pt 3) (2007) 708–19. [PubMed: 17267521]
- [73]. Nemoto K, Sakaguchi H, Kasai W, Hotta M, Kamei R, Noguchi T, Minamimoto R, Arai T, Asada T, Differentiating Dementia with Lewy Bodies and Alzheimer's Disease by Deep Learning to Structural MRI, *J Neuroimaging* 31(3) (2021) 579–587. [PubMed: 33476487]
- [74]. Abdelnour C, Ferreira D, Oppedal K, Cavallin L, Bousiges O, Wahlund LO, Hort J, Nedelska Z, Padovani A, Pilotto A, Bonanni L, Kramberger MG, Boada M, Westman E, Pagonabarraga J, Kulisevsky J, Blanc F, Aarsland D, The combined effect of amyloid-beta and tau biomarkers on brain atrophy in dementia with Lewy bodies, *Neuroimage Clin* 27 (2020) 102333. [PubMed: 32674011]
- [75]. Sarro L, Tosakulwong N, Schwarz CG, Graff-Radford J, Przybelski SA, Lesnick TG, Zuk SM, Reid RI, Raman MR, Boeve BF, Ferman TJ, Knopman DS, Comi G, Filippi M, Murray ME, Parisi JE, Dickson DW, Petersen RC, Jack CR Jr., Kantarci K, An investigation of cerebrovascular lesions in dementia with Lewy bodies compared to Alzheimer's disease, *Alzheimers Dement* 13(3) (2017) 257–266. [PubMed: 27521790]
- [76]. Ferreira D, Nedelska Z, Graff-Radford J, Przybelski SA, Lesnick TG, Schwarz CG, Botha H, Senjem ML, Fields JA, Knopman DS, Savica R, Ferman TJ, Graff-Radford NR, Lowe VJ, Jack CR, Petersen RC, Lemstra AW, van de Beek M, Barkhof F, Blanc F, Loureiro de Sousa P, Filippi N, Cretin B, Demuyneck C, Hort J, Oppedal K, Boeve BF, Aarsland D, Westman E, Kantarci K, Cerebrovascular disease, neurodegeneration, and clinical phenotype in dementia with Lewy bodies, *Neurobiol Aging* 105 (2021) 252–261. [PubMed: 34130107]
- [77]. Gomperts SN, Locascio JJ, Makaretz SJ, Schultz A, Caso C, Vasdev N, Sperling R, Growdon JH, Dickerson BC, Johnson K, Tau Positron Emission Tomographic Imaging in the Lewy Body Diseases, *JAMA Neurol* 73(11) (2016) 1334–1341. [PubMed: 27654968]
- [78]. Edison P, Rowe CC, Rinne JO, Ng S, Ahmed I, Kempainen N, Villemagne VL, O'Keefe G, Någren K, Chaudhury KR, Masters CL, Brooks DJ, Amyloid load in Parkinson's disease dementia and Lewy body dementia measured with [11C]PIB positron emission tomography, *J Neurol Neurosurg Psychiatry* 79(12) (2008) 1331–8. [PubMed: 18653550]
- [79]. Tsuang D, Leverenz JB, Lopez OL, Hamilton RL, Bennett DA, Schneider JA, Buchman AS, Larson EB, Crane PK, Kaye JA, Kramer P, Woltjer R, Trojanowski JQ, Weintraub D, Chen-Plotkin AS, Irwin DJ, Rick J, Schellenberg GD, Watson GS, Kukull W, Nelson PT, Jicha GA, Neltner JH, Galasko D, Masliah E, Quinn JF, Chung KA, Yearout D, Mata IF, Wan JY, Edwards KL, Montine TJ, Zabetian CP, APOE ε4 increases risk for dementia in pure synucleinopathies, *JAMA Neurol* 70(2) (2013) 223–8. [PubMed: 23407718]



- [80]. Pillai JA, Bena J, Bonner-Jackson A, Leverenz JB, Impact of APOE  $\epsilon 4$  genotype on initial cognitive symptoms differs for Alzheimer's and Lewy body neuropathology, *Alzheimers Res Ther* 13(1) (2021) 31. [PubMed: 33485373]
- [81]. Dickson DW, Heckman MG, Murray ME, Soto AI, Walton RL, Diehl NN, van Gerpen JA, Uitti RJ, Wszolek ZK, Ertekin-Taner N, Knopman DS, Petersen RC, Graff-Radford NR, Boeve BF, Bu G, Ferman TJ, Ross OA, APOE  $\epsilon 4$  is associated with severity of Lewy body pathology independent of Alzheimer pathology, *Neurology* 91(12) (2018) e1182–e1195. [PubMed: 30143564]
- [82]. Tsuang D, Leverenz JB, Lopez OL, Hamilton RL, Bennett DA, Schneider JA, Buchman AS, Larson EB, Crane PK, Kaye JA, Kramer P, Woltjer R, Kukull W, Nelson PT, Jicha GA, Neltner JH, Galasko D, Masliah E, Trojanowski JQ, Schellenberg GD, Yearout D, Huston H, Fritts-Penniman A, Mata IF, Wan JY, Edwards KL, Montine TJ, Zabetian CP, GBA mutations increase risk for Lewy body disease with and without Alzheimer disease pathology, *Neurology* 79(19) (2012) 1944–50. [PubMed: 23035075]
- [83]. Chia R, Sabir MS, Bandres-Ciga S, Saez-Atienzar S, Reynolds RH, Gustavsson E, Walton RL, Ahmed S, Viollet C, Ding J, Makarious MB, Diez-Fairen M, Portley MK, Shah Z, Abramzon Y, Hernandez DG, Blauwendraat C, Stone DJ, Eicher J, Parkkinen L, Ansorge O, Clark L, Honig LS, Marder K, Lemstra A, St George-Hyslop P, Londos E, Morgan K, Lashley T, Warner TT, Jaunmuktane Z, Galasko D, Santana I, Tienari PJ, Myllykangas L, Oinas M, Cairns NJ, Morris JC, Halliday GM, Van Deerlin VM, Trojanowski JQ, Grassano M, Calvo A, Mora G, Canosa A, Floris G, Bohannon RC, Brett F, Gan-Or Z, Geiger JT, Moore A, May P, Krüger R, Goldstein DS, Lopez G, Tayebi N, Sidransky E, Norcliffe-Kaufmann L, Palma JA, Kaufmann H, Shakkottai VG, Perkins M, Newell KL, Gasser T, Schulte C, Landi F, Salvi E, Cusi D, Masliah E, Kim RC, Caraway CA, Monuki ES, Brunetti M, Dawson TM, Rosenthal LS, Albert MS, Pletnikova O, Troncoso JC, Flanagan ME, Mao Q, Bigio EH, Rodríguez-Rodríguez E, Infante J, Lage C, González-Aramburu I, Sanchez-Juan P, Ghetti B, Keith J, Black SE, Masellis M, Rogaeva E, Duyckaerts C, Brice A, Lesage S, Xiromerisiou G, Barrett MJ, Tilley BS, Gentleman S, Logroscino G, Serrano GE, Beach TG, McKeith IG, Thomas AJ, Attems J, Morris CM, Palmer L, Love S, Troakes C, Al-Sarraj S, Hodges AK, Aarsland D, Klein G, Kaiser SM, Woltjer R, Pastor P, Bekris LM, Leverenz JB, Besser LM, Kuzma A, Renton AE, Goate A, Bennett DA, Scherzer CR, Morris HR, Ferrari R, Albani D, Pickering-Brown S, Faber K, Kukull WA, Morenas-Rodriguez E, Lleó A, Fortea J, Alcolea D, Clarimon J, Nalls MA, Ferrucci L, Resnick SM, Tanaka T, Foroud TM, Graff-Radford NR, Wszolek ZK, Ferman T, Boeve BF, Hardy JA, Topol EJ, Torkamani A, Singleton AB, Ryten M, Dickson DW, Chiò A, Ross OA, Gibbs JR, Dalgard CL, Traynor BJ, Scholz SW, Genome sequencing analysis identifies new loci associated with Lewy body dementia and provides insights into its genetic architecture, *Nat Genet* 53(3) (2021) 294–303. [PubMed: 33589841]
- [84]. Nalls MA, Duran R, Lopez G, Kurzawa-Akanbi M, McKeith IG, Chinnery PF, Morris CM, Theuns J, Crosiers D, Cras P, Engelborghs S, De Deyn PP, Van Broeckhoven C, Mann DM, Snowden J, Pickering-Brown S, Halliwell N, Davidson Y, Gibbons L, Harris J, Sheerin UM, Bras J, Hardy J, Clark L, Marder K, Honig LS, Berg D, Maetzler W, Brockmann K, Gasser T, Novellino F, Quattrone A, Annesi G, De Marco EV, Rogaeva E, Masellis M, Black SE, Bilbao JM, Foroud T, Ghetti B, Nichols WC, Pankratz N, Halliday G, Lesage S, Klebe S, Durr A, Duyckaerts C, Brice A, Giasson BI, Trojanowski JQ, Hurtig HI, Tayebi N, Landazabal C, Knight MA, Keller M, Singleton AB, Wolfsberg TG, Sidransky E, A multicenter study of glucocerebrosidase mutations in dementia with Lewy bodies, *JAMA Neurol* 70(6) (2013) 727–35. [PubMed: 23588557]
- [85]. Gamez-Valero A, Prada-Dacasa P, Santos C, Adame-Castillo C, Campdelacreu J, Rene R, Gascon-Bayarri J, Ispuerto L, Alvarez R, Ariza A, Beyer K, GBA Mutations Are Associated With Earlier Onset and Male Sex in Dementia With Lewy Bodies, *Mov Disord* 31(7) (2016) 1066–70. [PubMed: 27027900]
- [86]. Liu L, Li J, Quan W, Qin Y, Zhang Q, Pei X, Su H, Xu J, Chen J, Effect of GBA gene variants on clinical characteristics of dementia with Lewy bodies: a review and meta-analyses, *Neurol Sci* 43(6) (2022) 3541–3550. [PubMed: 35332438]
- [87]. Kramberger MG, Auestad B, Garcia-Ptacek S, Abdelnour C, Olmo JG, Walker Z, Lemstra AW, Londos E, Blanc F, Bonanni L, McKeith I, Winblad B, de Jong FJ, Nobili F, Stefanova E, Petrova

- M, Falup-Pecurariu C, Rektorova I, Bostantjopoulou S, Biundo R, Weintraub D, Aarsland D, E DLB, Long-Term Cognitive Decline in Dementia with Lewy Bodies in a Large Multicenter, International Cohort, *J Alzheimers Dis* 57(3) (2017) 787–795. [PubMed: 28304294]
- [88]. Abdelnour C, van Steenoven I, Londos E, Blanc F, Auestad B, Kramberger MG, Zetterberg H, Mollenhauer B, Boada M, Aarsland D, Alzheimer's disease cerebrospinal fluid biomarkers predict cognitive decline in lewy body dementia, *Mov Disord* 31(8) (2016) 1203–8. [PubMed: 27296778]
- [89]. Breivite MH, Hynninen MJ, Bronnick K, Chwiszczuk LJ, Auestad BH, Aarsland D, Rongve A, A longitudinal study of anxiety and cognitive decline in dementia with Lewy bodies and Alzheimer's disease, *Alzheimers Res Ther* 8 (2016) 3. [PubMed: 26812908]
- [90]. Olichney JM, Galasko D, Salmon DP, Hofstetter CR, Hansen LA, Katzman R, Thal LJ, Cognitive decline is faster in Lewy body variant than in Alzheimer's disease, *Neurology* 51(2) (1998) 351–7. [PubMed: 9710002]
- [91]. Hamilton JM, Salmon DP, Galasko D, Raman R, Emond J, Hansen LA, Masliah E, Thal LJ, Visuospatial deficits predict rate of cognitive decline in autopsy-verified dementia with Lewy bodies, *Neuropsychology* 22(6) (2008) 729–37. [PubMed: 18999346]
- [92]. Walker Z, McKeith I, Rodda J, Qassem T, Tatsch K, Booij J, Darcourt J, O'Brien J, Comparison of cognitive decline between dementia with Lewy bodies and Alzheimer's disease: a cohort study, *BMJ Open* 2 (2012) e000380.
- [93]. Compta Y, Parkkinen L, O'Sullivan SS, Vandrovcova J, Holton JL, Collins C, Lashley T, Kallis C, Williams DR, de Silva R, Lees AJ, Revesz T, Lewy- and Alzheimer-type pathologies in Parkinson's disease dementia: which is more important?, *Brain* 134(Pt 5) (2011) 1493–1505. [PubMed: 21596773]
- [94]. Williams MM, Xiong C, Morris JC, Galvin JE, Survival and mortality differences between dementia with Lewy bodies vs Alzheimer disease, *Neurology* 67(11) (2006) 1935–41. [PubMed: 17159097]
- [95]. Price A, Farooq R, Yuan JM, Menon VB, Cardinal RN, O'Brien JT, Mortality in dementia with Lewy bodies compared with Alzheimer's dementia: a retrospective naturalistic cohort study, *BMJ Open* 7(11) (2017) e017504.
- [96]. Armstrong MJ, Song S, Kurasz AM, Li Z, Predictors of Mortality in Individuals with Dementia in the National Alzheimer's Coordinating Center, *J Alzheimers Dis* 86(4) (2022) 1935–1946. [PubMed: 35253760]
- [97]. Graff-Radford J, Aakre J, Savica R, Boeve B, Kremers WK, Ferman TJ, Jones DT, Kantarci K, Knopman DS, Dickson DW, Kukull WA, Petersen RC, Duration and Pathologic Correlates of Lewy Body Disease, *JAMA Neurol* 74(3) (2017) 310–315. [PubMed: 28114455]
- [98]. Boström F, Hansson O, Blennow K, Gerhardsson L, Lundh T, Minthon L, Zetterberg H, Londos E, Cerebrospinal fluid total tau is associated with shorter survival in dementia with Lewy bodies, *Dement Geriatr Cogn Disord* 28(4) (2009) 314–9. [PubMed: 19844105]
- [99]. van Steenoven I, van der Flier WM, Scheltens P, Teunissen CE, Lemstra AW, Amyloid- $\beta$  peptides in cerebrospinal fluid of patients with dementia with Lewy bodies, *Alzheimers Res Ther* 11(1) (2019) 83. [PubMed: 31601267]
- [100]. Stefani A, Santamaria J, Iranzo A, Hackner H, Schenck CH, Högl B, Nelotanserin as symptomatic treatment for rapid eye movement sleep behavior disorder: a double-blind randomized study using video analysis in patients with dementia with Lewy bodies or Parkinson's disease dementia, *Sleep Med* 81 (2021) 180–187. [PubMed: 33714847]
- [101]. Gratwicke J, Zrinzo L, Kahan J, Peters A, Brechany U, McNichol A, Beigi M, Akram H, Hyam J, Oswal A, Day B, Mancini L, Thornton J, Yousry T, Crutch SJ, Taylor JP, McKeith I, Rochester L, Schott JM, Limousin P, Burn D, Rossor MN, Hariz M, Jahanshahi M, Foltynie T, Bilateral nucleus basalis of Meynert deep brain stimulation for dementia with Lewy bodies: A randomised clinical trial, *Brain Stimul* 13(4) (2020) 1031–1039. [PubMed: 32334074]
- [102]. Maltête D, Wallon D, Bourilhon J, Lefaucheur R, Danaila T, Thobois S, Defebvre L, Dujardin K, Houeto JL, Godefroy O, Krystkowiak P, Martinaud O, Gillibert A, Chastan M, Vera P, Hannequin D, Welter ML, Derrey S, Nucleus Basalis of Meynert Stimulation for Lewy Body Dementia: A Phase I Randomized Clinical Trial, *Neurology* 96(5) (2021) e684–e697. [PubMed: 33199437]

- [103]. Rigby T, Johnson DK, Taylor A, Galvin JE, Comparison of the Caregiving Experience of Grief, Burden, and Quality of Life in Dementia with Lewy Bodies, Alzheimer's Disease, and Parkinson's Disease Dementia, *J Alzheimers Dis* 80(1) (2021) 421–432. [PubMed: 33554912]
- [104]. Liu S, Liu J, Wang XD, Shi Z, Zhou Y, Li J, Yu T, Ji Y, Caregiver burden, sleep quality, depression, and anxiety in dementia caregivers: a comparison of frontotemporal lobar degeneration, dementia with Lewy bodies, and Alzheimer's disease, *Int Psychogeriatr* 30(8) (2018) 1131–1138. [PubMed: 29223171]
- [105]. Leggett AN, Zarit S, Taylor A, Galvin JE, Stress and burden among caregivers of patients with Lewy body dementia, *Gerontologist* 51(1) (2011) 76–85. [PubMed: 20667944]
- [106]. Galvin JE, Duda JE, Kaufer DI, Lippa CF, Taylor A, Zarit SH, Lewy body dementia: caregiver burden and unmet needs, *Alzheimer Dis Assoc Disord* 24(2) (2010) 177–181. [PubMed: 20505434]
- [107]. Park J, Galvin JE, Pre-Loss Grief in Caregivers of Older Adults with Dementia with Lewy Bodies, *J Alzheimers Dis* 82(4) (2021) 1847–1859. [PubMed: 34219734]
- [108]. Svendsboe E, Terum T, Testad I, Aarsland D, Ulstein I, Corbett A, Rongve A, Caregiver burden in family carers of people with dementia with Lewy bodies and Alzheimer's disease, *Int J Geriatr Psychiatry* 31(9) (2016) 1075–1083. [PubMed: 26765199]
- [109]. Kanemoto H, Sato S, Satake Y, Koizumi F, Taomoto D, Kanda A, Wada T, Yoshiyama K, Ikeda M, Impact of Behavioral and Psychological Symptoms on Caregiver Burden in Patients With Dementia With Lewy Bodies, *Front Psychiatry* 12 (2021) 753864. [PubMed: 34777057]
- [110]. Miller VM, Why are sex and gender important to basic physiology and translational and individualized medicine?, *Am J Physiol Heart Circ Physiol* 306(6) (2014) H781–8. [PubMed: 24414073]
- [111]. Mielke MM, Sex and Gender Differences in Alzheimer's Disease Dementia, *Psychiatr Times* 35(11) (2018) 14–17. [PubMed: 30820070]
- [112]. Azad NA, Al Bugami M, Loy-English I, Gender differences in dementia risk factors, *Gend Med* 4(2) (2007) 120–9. [PubMed: 17707846]
- [113]. Andrew MK, Tierney MC, The puzzle of sex, gender and Alzheimer's disease: Why are women more often affected than men?, *Womens Health* 14 (2018).
- [114]. Goldman JG, Forsberg LK, Boeve BF, Armstrong MJ, Irwin DJ, Ferman TJ, Galasko D, Galvin JE, Kaufer D, Leverenz J, Lippa CF, Marder K, Abler V, Biglan K, Irizarry M, Keller B, Munsie L, Nakagawa M, Taylor A, Graham T, Challenges and opportunities for improving the landscape for Lewy body dementia clinical trials, *Alzheimers Res Ther* 12(1) (2020) 137. [PubMed: 33121510]
- [115]. Chiu SY, Bowers D, Armstrong MJ, Lewy Body Dementias: Controversies and Drug Development, *Neurotherapeutics* (2021).
- [116]. Armstrong MJ, Gamez N, Alliance S, Majid T, Taylor A, Kurasz AM, Patel B, Smith G, Research priorities of caregivers and individuals with dementia with Lewy bodies: An interview study, *PLoS ONE* 15(10) (2020) e0239279. [PubMed: 33027276]

### Highlights

- Better understanding sex differences in DLB will aid diagnosis and management.
- RBD and parkinsonism may be more common in men.
- Visual hallucinations appear more common in women.
- Delayed DLB diagnosis may be more frequent in women.
- Women are more likely to have mixed Lewy body and Alzheimer-related pathologies.



**Figure 1. Sex differences in DLB**

Women with dementia with Lewy bodies (DLB) tend to be older, have greater cognitive impairment at their initial visit, and meet DLB criteria later when compared to men. Women with DLB may also be more likely to experience visual hallucinations than men, with visual hallucinations starting concurrently with parkinsonism and cognitive fluctuations. While the illustration includes a shadowed cat to visually represent the hallucination, hallucinations appear real to the individuals experiencing them. Parkinsonism can be milder in women with DLB than in men, one reason women might be diagnosed later. Caregivers in DLB are commonly women, reflecting either spouses or daughters and daughters-in-law. Caregiver burden can be affected by dementia severity, neuropsychiatric symptoms, and need to assist with activities of daily living, but whether sex affects caregiver burden is uncertain. Little is known about the experiences of DLB in individuals from diverse backgrounds. The illustration of two women of non-white race does not imply that DLB disproportionately affects Black women or that women from these backgrounds have a different experience.

**Table 1.**

Sex demographic profiles in sample DLB studies

| Reference                                      | Country | Referral base  | Study type  | Diagnostic criteria used  | Epidemiology (if reported)  | Men/women ratio  | Term Used in Publication (Sex or Gender) <sup>a</sup> |
|--|---------|--|---|---|---|--|---|
| <i>Population-based studies</i>                |         |  |   |   |   |  |   |
| Yue et al., 2016 [117]                         | China   | Door-to-door in rural China  | Cross-sectional, door-to-door, population-based study across 56 villages in Ji County | DSM-IV criteria   | Prevalence/total population, % (95% CI)<br>Men: 1.22 (0.78–1.65)<br>Women: 0.91 (0.57–1.25)<br><br>Prevalence/dementia population, % (95% CI)<br>Men: 15.15 (10.16–20.15)<br>Women: 7.45 (4.79–10.10) | <b>Men: n=30 (51.7%)</b><br><b>Women: n=28 (48.3%)</b>         | <b>Sex</b>  |
| Savica et al., 2013 [12]                       | U.S.    | Medical records-linkage system of the Rochester Epidemiology Project (REP)   | Retrospective study   | DLB Consortium 3rd Consensus Criteria                                 | Incidence/all ages, Rate per 100,000 person-years<br>Men: 4.8<br>Women: 2.2<br><br>Incidence/over 65 years old, Rate per 100,000 person-years<br>Men: 54.2<br>Women: 16.2                             | <b>Men: n=43 (67%)</b><br><b>Women: n=21 (33%)</b>             | <b>Sex</b>  |
| Williams et al., 2006 [94]                     | U.S.    | Recruited via word of mouth, public service announcements, and referrals from physicians in the St. Louis area   | Prospective longitudinal study  | DLB Consortium 1st Consensus Criteria and neuropathology confirmation | N/A   | <b>Men: n=38 (60%)</b><br><b>Women: n=25 (40%)</b>             | <b>Gender</b>   |
| Goodman et al., 2017 [118]                     | U.S.    | Centers for Medicare & Medicaid (CMS) administrative enrollment and claims data for 100% of Medicare beneficiaries enrolled in the fee-for-service (FFS) program | Retrospective study   | ICD-9 codes with 331.82 or 332.0 + 331.0 <sup>a</sup>                 | N/A   | <b>Women: n=85,234 (50.5%)</b><br><b>Men: n=83,395 (49.5%)</b> | <b>Sex</b>  |
| Rahkonen et al., 2003 [119]                    | Finland | Random sample drawn from "Kuopio 75+" population health survey study in the city of Kuopio   | Population healthy survey study   | DLB Consortium 1st Consensus Criteria                                 | N/A   | <b>Women: n=26 (87%)</b><br><b>Men: n=4 (13%)</b>              | <b>Sex</b>  |
| Gascón-Bayarri et al., 2007 [120]              | Spain   | Population-based in El Prat del Llobregat, in the Barcelona's Metropolitan Area, in Catalonia, Spain   | Population-based survey study   | DLB Consortium 1st Consensus Criteria                                 | Overall DLB prevalence: 9.09 (5.18–14.55)<br>Men: 0.4<br>Women: 1.2   | <b>Women: n=13 (87%)</b><br><b>Men: n=2 (13%)</b>              | <b>Sex</b>  |
| <i>Secondary care / large database studies</i> |         |  |   |   |   |  |   |

| Reference                   | Country | Referral base  | Study type                                | Diagnostic criteria used   | Epidemiology (if reported)  | Men/women ratio                              | Term Used in Publication (Sex or Gender) <sup>b</sup> |
|-----------------------------|---------|--|---|--|---|--|---|
| Gan et al., 2021 [121]      | China   | Nine memory clinics in the China Lewy Body Disease Collaborative Alliance  | Retrospective study                       | DSM-IV criteria  | N/A   | Male: n=177 (50.9%)<br>Female: n=171 (49.1%) | Sex/Gender <sup>b</sup>                               |
| Yang et al., 2018 [122]     | Taiwan  | DLB patient data obtained between 2000-2013 from the Taiwan National Health Insurance Research Database (NHIRD)                        | Retrospective study                       | ICD-9-CM code 331.82, or following 1-year rule of having parkinsonism and dementia (ICD-9-CM does 332, 333, 290.0, 290.1, 290.2, 290.3, 294.1, 3331.0) | Incidence/all ages, Rate per 100,000 person-years<br>Men 7.06<br>Women 7.14 | Men: n=438 (50.2%)<br>Women: n=434 (49.8)    | Sex   |
| Utsumi et al., 2020 [17]    | Japan   | Outpatient psychiatry clinic in Sunagawa City Medical Center   | Retrospective study                       | DLB Consortium 4th Consensus Criteria  | N/A   | Women: n=133 (57%)<br>Men: n=101 (43%)       | Gender  |
| Breivte et al., 2016 [89]   | Norway  | Outpatient clinics of old age psychiatry and geriatric medicine in western Norway  | Longitudinal study                        | DLB Consortium 3rd Consensus Criteria  | N/A   | Men: n=40 (55.6%)<br>Women: n=32 (44.4%)     | Sex   |
| Rongve et al., 2016 [123]   | Norway  | Outpatient geriatric, psychiatric and neurology clinics in western Norway  | Five-year prospective cohort study        | DLB Consortium 3rd Consensus Criteria  | N/A   | Men: n=35 (52.2%)<br>Women: n=32 (47.8%)     | Sex   |
| Aarsland et al., 2008 [124] | Norway  | Outpatient clinics in neurology, geriatric medicine and old age psychiatry in the counties of Rogaland and Hordaland in Western Norway | Retrospective cohort study                | DLB Consortium 3rd Consensus Criteria  | N/A   | Women: n=20 (51.3%)<br>Men: n=19 (48.7%)     | Gender  |
| Farina et al., 2009 [125]   | Italy   | Eight Italian memory clinics   | Mixed retrospective and prospective study | DLB Consortium 1st Consensus Criteria  | N/A   | Men: n=52 (51%)<br>Women: n=50 (49%)         | Sex   |
| Choudhury et al., 2021 [15] | U.S.    | Mayo Clinic Alzheimer's Disease Research Center (ADRC)   | Prospective longitudinal cohort study     | DLB Consortium 4th Consensus Criteria  | N/A   | Men: n=370 (76%)<br>Women: n=118 (24%)       | Sex   |
| Bayram et al., 2021 [14]    | U.S.    | Neuropathologically confirmed cases from the National Alzheimer's Coordinating Center (NACC) database                                  | Retrospective study                       | DLB Consortium 4th Consensus Criteria associated with a high likelihood of exhibiting a DLB phenotype  | N/A   | Men: n=156 (74%)<br>Women: n=55 (26%)        | Sex   |
| Kane et al., 2018 [23]      | U.K.    | Nine participating psychiatry of old age/memory clinic services in the UK  | Retrospective case series                 |  | Overall DLB prevalence 4.6 (4.0-5.2)  | Men: n=113 (54.6%)                           | Sex   |

| Reference                     | Country     | Referral base  | Study type                              | Diagnostic criteria used              | Epidemiology (if reported)   | Men/women ratio                                | Term Used in Publication (Sex or Gender) <sup>b</sup> |
|-------------------------------|-------------|--|---|---------------------------------------|--|--|---|
| Price et al., 2017 [95]       |             | across four NHS hospital trusts in East Anglia and North-East England  |   |                                       | <ul style="list-style-type: none"> <li>• 113 M/94 F (54.6% M)</li> <li>Overall DLB incidence 4.8 (4.0–5.7)</li> <li>• 62 M/61 F (50.4% M)</li> </ul> | Women: n=94 (45.4%)                            |   |
| Price et al., 2017 [95]       | U.K.        | Cambridge and Peterborough NHS Foundation Trust (CPFT), which provides secondary mental healthcare   | Retrospective naturalistic cohort study | DLB Consortium 3rd Consensus Criteria | N/A  | Women: n=129 (51.4%)<br>Men: n=122 (48.6%)     | Sex   |
| Mouton et al., 2018 [16]      | France      | French National Alzheimer database (Banque Nationale Alzheimer [BNA]), which stores information from French memory centers and private practice neurologists | Repeated cross-sectional study          | DLB Consortium 3rd Consensus Criteria | N/A  | Women: n=5,635 (54.7%)<br>Men: n=4,674 (45.3%) | Sex(Gender) <sup>b</sup>                              |
| van de Beek et al., 2020 [18] | Netherlands | Amsterdam Dementia Cohort (patient selection based on CSF availability)  | Retrospective study                     | DLB Consortium 4th Consensus Criteria | N/A  | Men: n=184 (83%)<br>Women: n=39 (17%)          | Sex   |

DLB=dementia with Lewy bodies; CI=Confidence Interval; ICD=International Classification of Diseases; DSM=Diagnostic and Statistical Manual of Mental Disorders

<sup>a</sup>Method of defining dementia allowed beneficiaries to be classified as having more than one dementia subtype.

<sup>b</sup>Sex and gender terms were used interchangeably, though authors' interpretations seemed to suggest biological definition of "sex".



**Table 2.**

DLB core features in men and women

| Reference                             | Cohort<br>(Total<br>N, %<br>men) | RBD (N, %) |           | p-value | Parkinsonism<br>(N, %) |           | p-value | Visual<br>hallucinations<br>(N, %) |           | p-value | Fluctuations<br>(N, %) |           | p-value |
|---------------------------------------|----------------------------------|------------|-----------|---------|------------------------|-----------|---------|------------------------------------|-----------|---------|------------------------|-----------|---------|
|                                       |                                  | Men        | Women     |         | Men                    | Women     |         | Men                                | Women     |         | Men                    | Women     |         |
| Choudhury et al., 2021 [15]           | 488 (76)                         | 311 (84)   | 61 (52)   | <0.001  | 335 (91)               | 98 (83)   | 0.025   | 244 (66)                           | 97 (82)   | 0.001   | 286 (77)               | 87 (74)   | 0.43    |
| Bayram et al., 2021 <sup>a</sup> [14] | 110 (50)                         | 28 (51)    | 16 (29)   | >0.29   | 47 (86)                | 42 (76)   | >0.29   | 35 (64)                            | 22 (40)   | <0.05   | 31 (56)                | 27 (49)   | >0.29   |
| Bayram et al., 2022 <sup>b</sup> [20] | 691 (68)                         | 165 (45.8) | 35 (20.8) | <0.001  | 278 (71.5)             | 90 (50.6) | <0.001  | 255 (54.4)                         | 85 (38.3) | <0.001  | 201 (55.2)             | 70 (40.9) | <0.005  |
| Utsumi et al., 2020 [17]              | 234 (43)                         | 76 (76)    | 66 (50)   | <0.001  | 60 (59)                | 59 (44)   | 0.027   | 77 (76)                            | 109 (82)  | NS      | 54 (54)                | 78 (58)   | NS      |
| Van de Beek et al., 2020 [18]         | 223 (83)                         | 99 (73)    | 11 (55)   | 0.17    | 130 (73)               | 23 (62)   | 0.28    | 69 (38)                            | 23 (59)   | 0.02    | 119 (84)               | 29 (94)   | 0.26    |
| Ferreira et al., 2020 [54]            | 416 (69)                         | 210 (73)   | 79 (61)   | 0.48    | 253 (88)               | 107 (83)  | 0.023   | 152 (53)                           | 93 (72)   | 0.04    | 230 (80)               | 112 (87)  | 0.53    |
| Chiu et al., 2018 <sup>c</sup> [55]   | 152 (57)                         | N/A        | N/A       | N/A     | N/A                    | N/A       | N/A     | 39 (45)                            | 39 (60)   | 0.028   | N/A                    | N/A       | N/A     |

DLB=dementia with Lewy bodies; NS=not significant (p-value not provided)

<sup>a</sup>Showing data from the study's matched subsamples (by age, education, tau burden; n=55 women, n=55 men); in the overall cohort, additional statistically significant sex differences were found for RBD (men 56% vs. women 29%, p<0.01), and parkinsonism (men 92% vs. women 76%, p<0.01) (Bayram et al., 2021).

<sup>b</sup>Unlike their prior study [14] that excluded individuals with Braak tau staging > III, this study [20] included pathologically-confirmed DLB sample with all Braak tau stages 0-VI.

<sup>c</sup>As this study focused on sex differences in DLB in visual hallucinations only, there are no available data on sex differences in other DLB core features.

**Table 3.**

## Gaps and Research Opportunities Relating to Sex and Gender in DLB

| <b>Knowledge or Care Gap</b>   | <b>Potential Research Opportunities</b>   |
|--|---|
| Differences in presenting phenotype and clinicopathological correlations | Large-scale clinicopathologic studies with assessment of both sex and gender  |
| Risk Factors   | <ul style="list-style-type: none"> <li>- Studies examining genetic risk factors (e.g. APOE, GBA)</li> <li>- Studies investigating hormonal influences (pregnancy, menopause, estrogen/hormone replacement therapy, androgen-deprivation treatment, transgender health, etc.)</li> <li>- Evaluation of environmental exposures, comorbidities (cerebrovascular risk factors, etc.), demographics (education, ethnicity/race, age, etc.) and how they may differ by sex/gender</li> </ul> |
| Under- or misdiagnosis   | Studies of pragmatic approaches to increase recognition of DLB generally and with sex- and gender-specific considerations   |
| Biomarker differences by sex/gender                                      | Targeted recruitment of diverse populations for biomarker studies and subanalyses by sex and gender   |
| Progression  | Natural history studies with diverse recruitment (including by sex/gender) and associated subanalyses   |
| Treatment  | <ul style="list-style-type: none"> <li>- Stratifying by sex/gender during randomization</li> <li>- Inclusion of biomarkers in DLB treatment trials (as this may be different by sex and affect treatment response)</li> <li>- Assessment of treatment response by sex/gender</li> </ul>   |
| Caregiving   | <ul style="list-style-type: none"> <li>- Research on differential caregiver experiences by sex and gender, access to healthcare/resources, and potential value of specific interventions</li> </ul>   |

Author Manuscript

Author Manuscript

Author Manuscript

Author Manuscript