

UC Irvine

UC Irvine Previously Published Works

Title

Melatonin signaling in mitochondria extends beyond neurons and neuroprotection: Implications for angiogenesis and cardio/gastroprotection

Permalink

<https://escholarship.org/uc/item/8sg5q9sf>

Journal

Proceedings of the National Academy of Sciences of the United States of America, 115(9)

ISSN

0027-8424

Authors

Ahluwalia, Amrita

Brzozowska, Iwona M

Hoa, Neil

et al.

Publication Date

2018-02-27

DOI

10.1073/pnas.1722131115

Copyright Information

This work is made available under the terms of a Creative Commons Attribution License, available at <https://creativecommons.org/licenses/by/4.0/>

Peer reviewed



Melatonin signaling in mitochondria extends beyond neurons and neuroprotection: Implications for angiogenesis and cardio/gastroprotection

Amrita Ahluwalia^{a,1}, Iwona M. Brzozowska^{b,1}, Neil Hoa^a, Michael K. Jones^{a,c}, and Andrzej S. Tarnawski^{a,c,2}

We read with interest the paper of Suofu et al. (1), demonstrating that neuronal mitochondria produce melatonin, which upon binding to its melatonin type 1 (MT₁) receptor on the mitochondrial membrane (MM) inhibits cytochrome c release, caspase activation, and apoptosis. We commend the authors on their thorough investigation but wish to point out some major questions: (i) Is the presence of the MT₁ receptor in mitochondria limited to the neuronal cells or may it also apply to other cells, for example, endothelial cells that are components of blood vessels and are critical for delivery of oxygen and nutrients to all tissues, vessels' regeneration, and angiogenesis? (ii) Is the MT₂ receptor expressed on mitochondria and what is its role?

Melatonin is abundantly expressed beyond the neural system; for example, in gastrointestinal tissues, including the stomach, where it has been shown to protect gastric mucosa against stress, ischemia, and nonsteroidal antiinflammatory drug-induced injury, and to accelerate ulcer healing (2–6). We recently demonstrated that both MT₁ and MT₂ are expressed on the MM of gastric endothelial cells (GECs); MT₁ was expressed in MM while MT₂ was expressed in the nucleus and MM (Fig. 1). However, in contrast to the study of Suofu et al. (1), we found that treatment of GECs with exogenous melatonin (10 μM) increased by 1.7-fold ($P < 0.001$) MM potential, which drives ATP synthesis (7). Moreover, treatment of GECs with exogenous melatonin increased the MM expression of MT₁ and MT₂ by 2.5- and 1.6-fold, respectively, and increased in vitro angiogenesis (new blood vessel formation) by 1.4-fold ($P < 0.001$). It should be noted that the angiogenic response to melatonin was independent of either cell proliferation or apoptosis, leading us to speculate that, at least in GECs, melatonin

signaling via mitochondria extends beyond cellular protection. Although, we have not yet determined the relative contributions of MT₁ vs. MT₂ in the angiogenic response of GECs to melatonin signaling, our finding that exogenous melatonin increased the MM expression of both MT₂ and MT₁ is at least indicative that both receptors are involved. Suofu et al. (1) did not assess the relative MM expression levels of MT₁ vs. MT₂, but rather used the MT₂ selective inhibitor, 4P-PDOT, as evidence that melatonin signaling in neuronal mitochondria is mediated exclusively through MM-expressed MT₁. Nevertheless, luzindole, which did prevent melatonin from blocking Ca²⁺-mediated cytochrome c release, has a much greater affinity for MT₂ vs. MT₁. Moreover, the possibility that MT₂ plays a “decoy” role, akin to Flt1 in VEGF signaling, cannot be excluded. In such a scenario, MT₂ may play a more subtle role in regulating (e.g., dampening) melatonin signaling. On the other hand, the role of MT₂ in mitochondrial signaling by melatonin may be restricted to certain cell types (e.g., endothelial cells but not to brain neurons). The work of Suofu et al. (1) demonstrates novel melatonin signaling in neuronal mitochondria. Our work uncovered that this mechanism operates in endothelial cells that are major components of all blood vessels. We further suggest that melatonin signaling in mitochondria may also be applicable to other physiological processes, such as angiogenesis, gastroprotection, cardioprotection, and aging (3–6, 8, 9).

Acknowledgments

This work was supported by Merit Review Award I01 BX000626-05A2 from the US Department of Veterans Affairs Biomedical Laboratory Research and Development Service (to A.S.T.).

^aMedical and Research Services, Veterans Affairs Long Beach Healthcare System, Long Beach, CA 90822; ^bDepartment of Anatomy, Jagiellonian University Medical College, 31-008 Cracow, Poland; and ^cDepartment of Medicine, University of California, Irvine, CA 92617

Author contributions: A.A. and A.S.T. designed research; A.A. and A.S.T. performed research; A.A., I.M.B., N.H., M.K.J., and A.S.T. analyzed data; and A.A., I.M.B., N.H., M.K.J., and A.S.T. wrote the paper.

The authors declare no conflict of interest.

Published under the [PNAS license](#).

¹A.A. and I.M.B. contributed equally to this work.

²To whom correspondence should be addressed. Email: atarnawski@yahoo.com.

Published online February 9, 2018.




Fig. 1. Localization of melatonin receptors MT_1 and MT_2 in MM of gastric endothelial cells (GECs). (A) Staining of mitochondria using MitoTracker, a fluorescent dye that stains mitochondria (red) in a manner dependent on mitochondrial membrane potential. (B) Immunofluorescence staining for MT_1 in GECs (green). (C) Overlay of MitoTracker and MT_1 immunostaining images showing localization of MT_1 in MM as yellow/orange staining (arrows). (D) Staining of mitochondria using MitoTracker (red). (E) Immunofluorescence staining for MT_2 in GECs (green); please note strong MT_2 expression in the nucleus (N). (F) Overlay of MitoTracker and MT_2 immunostaining images showing localization of MT_2 in MM as yellow/orange staining (arrows).

- 1 Suofu Y, et al. (2017) Dual role of mitochondria in producing melatonin and driving GPCR signaling to block cytochrome c release. *Proc Natl Acad Sci USA* 114:E7997–E8006.
- 2 Bubenik GA (2008) Thirty four years since the discovery of gastrointestinal melatonin. *J Physiol Pharmacol* 59:33–51.
- 3 Karasek M (2004) Melatonin, human aging, and age-related diseases. *Exp Gerontol* 39:1723–1729.
- 4 Ma Z, et al. (2017) Melatonin and mitochondrial function during ischemia/reperfusion injury. *Cell Mol Life Sci* 74:3989–3998.
- 5 Brzozowska I, et al. (2002) Role of prostaglandins, nitric oxide, sensory nerves and gastrin in acceleration of ulcer healing by melatonin and its precursor, L-tryptophan. *J Pineal Res* 32:149–162.
- 6 Brzozowska I, Strzalka M, Drozdowicz D, Konturek SJ, Brzozowski T (2014) Mechanisms of esophageal protection, gastroprotection and ulcer healing by melatonin. Implications for the therapeutic use of melatonin in gastroesophageal reflux disease (GERD) and peptic ulcer disease. *Curr Pharm Des* 20:4807–4815.
- 7 Klusch N, Murphy BJ, Mills DJ, Yildiz Ö, Kühlbrandt W (2017) Structural basis of proton translocation and force generation in mitochondrial ATP synthase. *eLife* 6:6.
- 8 Tengattini S, et al. (2008) Cardiovascular diseases: Protective effects of melatonin. *J Pineal Res* 44:16–25.
- 9 Lochner A, Huisamen B, Nduhirabandi F (2013) Cardioprotective effect of melatonin against ischaemia/reperfusion damage. *Front Biosci (Elite Ed)* 5:305–315.