

# UC Berkeley

## UC Berkeley Previously Published Works

### Title

Quantifying the Effect Size of Management Actions on Aboveground Carbon Stocks in Forest Plantations

### Permalink

<https://escholarship.org/uc/item/8t42v5q2>

### Journal

Current Forestry Reports, 9(3)

### ISSN

2198-6436

### Authors

Melikov, Cyril H

Bukoski, Jacob J

Cook-Patton, Susan C

et al.

### Publication Date

2023

### DOI

10.1007/s40725-023-00182-5

### Copyright Information

This work is made available under the terms of a Creative Commons Attribution License, available at <https://creativecommons.org/licenses/by/4.0/>

Peer reviewed



# Quantifying the Effect Size of Management Actions on Aboveground Carbon Stocks in Forest Plantations

Cyril H. Melikov<sup>1,2</sup> · Jacob J. Bukoski<sup>3,4</sup> · Susan C. Cook-Patton<sup>5</sup> · Hongyi Ban<sup>2</sup> · Jessica L. Chen<sup>2</sup> · Matthew D. Potts<sup>2,6</sup>

Accepted: 22 February 2023 / Published online: 11 April 2023  
© The Author(s) 2023

## Abstract

**Purpose of the Review** Improved forest management is a promising avenue for climate change mitigation. However, we lack synthetic understanding of how different management actions impact aboveground carbon stocks, particularly at scales relevant for designing and implementing forest-based climate solutions. Here, we quantitatively assess and review the impacts of three common practices—application of inorganic NPK fertilizer, interplanting with N-fixing species, and thinning—on aboveground carbon stocks in plantation forests.

**Recent Findings** Site-level empirical studies show both positive and negative effects of inorganic fertilization, interplanting, and thinning on aboveground carbon stocks in plantation forests. Recent findings and the results of our analysis suggest that these effects are heavily moderated by factors such as species selection, precipitation, time since practice, soil moisture regime, and previous land use. Interplanting of N-fixing crops initially has no effect on carbon storage in main tree crops, but the effect becomes positive in older stands. Conversely, the application of NPK fertilizers increases aboveground carbon stocks, though the effect lessens with time. Moreover, increases in aboveground carbon stocks may be partially or completely offset by emissions from the application of inorganic fertilizer. Thinning results in a strong reduction of aboveground carbon stocks, though the effect lessens with time.

**Summary** Management practices tend to have strong directional effects on aboveground carbon stocks in plantation forests but are moderated by site-specific management, climatic, and edaphic factors. The effect sizes quantified in our meta-analysis can serve as benchmarks for the design and scoping of improved forest management projects as forest-based climate solutions. Overall, management actions can enhance the climate mitigation potential of plantation forests, if performed with sufficient attention to the nuances of local conditions.

**Keywords** Natural climate solutions · Improved forest management · Carbon · Fertilization · Thinning · Forest plantation

---

✉ Cyril H. Melikov  
cyril\_melikov@berkeley.edu

Jacob J. Bukoski  
jbukoski@berkeley.edu

Susan C. Cook-Patton  
susan.cook-patton@tnc.org

Hongyi Ban  
stellaban@berkeley.edu

Jessica L. Chen  
jessicaliuchen@berkeley.edu

Matthew D. Potts  
mdpotts@berkeley.edu

<sup>1</sup> Environmental Defense Fund, New York, NY, USA

<sup>2</sup> Department of Environmental Science, Policy and Management, University of California, Berkeley, CA, USA

<sup>3</sup> Moore Center for Science, Conservation International, Arlington, VA, USA

<sup>4</sup> Department of Forest Ecosystems and Society, Oregon State University, Corvallis, OR, USA

<sup>5</sup> The Nature Conservancy, Arlington, VA, USA

<sup>6</sup> Carbon Direct Inc, New York, NY, USA

## Introduction

Mitigating the most damaging effects of climate change requires concerted and urgent action within this decade in both the energy and land sectors [1]. Recently, studies have highlighted the potential of tree cover restoration for mitigating climate change, in places where formerly forested landscapes have been lost or severely degraded [2–6]. Despite momentum to restore tree cover, improving the management of existing forests may represent a more cost-effective and rapidly deployable natural climate solution, and could sequester 0.1 to 2.3 PgCO<sub>2</sub>e per year [1, 5, 7•, 9••]. Given limited resources and the urgency of climate change, identifying near-term management actions that can maximize carbon stocks in managed forest stands is of paramount importance.

In natural forests, multiple studies have focused on how improved forest management practices, such as reduced-impact logging and liana control, can increase aboveground carbon stocks, relative to standard forestry practices [5, 8•, 10–12]. In planted forests, studies suggest that extending rotations to a biological rather than economical optimum can substantially increase time-averaged carbon stocks [5, 12]. Nonetheless, forestry practitioners commonly perform additional operations in planted stands such as fertilization and stand density management that directly influence plantation carbon stocks [13••, 15••]. Here, we consider the effects of two fertilization practice-intercropping of nitrogen (N)-fixing species and application of inorganic nitrogen, phosphorus, potassium (NPK) fertilizer-and one stand density management operation (thinning) on aboveground carbon stocks in forest plantations.

While the use of NPK fertilizers remains the most common fertilization practice [15••], interplanting of N-fixing plants as an alternative method has become increasingly prominent in mixed species plantations [16, 17, 18•, 19]. In addition to providing N-fixing benefits, planting of multiple species may confer additional ecological benefits, such as habitat for biodiversity and improved soil fertility [15••, 18•, 20, 21]. Furthermore, the use of conventional inorganic NPK fertilizers has potentially adverse environmental impacts from improper and over-use, including declines in soil fertility, elevated groundwater and surface water pollution, and increased GHG emissions from fertilizer production and use [22, 23].

The empirical effects of these fertilization techniques on aboveground carbon stocks in plantations has been well documented in multiple locations [24–29]. However, the magnitude of the effect has been rarely assessed at large scales [17, 30, 31] relevant for the design of forest-based climate solutions, with prior efforts constrained to regional

geographies [32] or specific tree species [16]. Furthermore, these prior large-scale meta-analyses are now more than 10 years old, and to the best of our knowledge, none has examined moderators of the fertilizing effect nor investigated how the effect size might vary with site and stand characteristics.

Similarly, the effects of thinning stand biodiversity [33, 34], soil carbon stocks [35, 36], soil microbial carbon [37], and even drought-related tree stress [38] have been studied at site-level scales. However, only a handful of studies have synthesized the changes in plantation aboveground carbon stocks after thinning operations in large-scale meta-analyses [39–41], and most of the previous investigations were plot-scale empirical studies [42–44]. Similar to fertilization treatments, few studies have investigated how thinning effects change with site characteristics, with most of the past efforts focusing on the impacts of stand age and time since treatment on the treatment's effect size [39, 40].

The global effects of these silvicultural treatments as well as the contexts under which they deliver the most carbon benefits still need to be documented. Our work builds on these previous empirical studies and meta-analyses. It aims to improve understanding of how prominent silvicultural practices impact aboveground carbon stocks in plantations globally. To do so, we systematically reviewed and compiled aboveground carbon measurements from interplanted, fertilized, and thinned tree plantations distributed across six continents and 18 countries. Using this dataset, we then quantified (i) how each treatment affected aboveground carbon stocks and (ii) how the effect size of each treatment varied with different environmental and management factors. In doing so, we provide insights on how fertilization and thinning treatments can be improved to promote carbon stocks in planted forests.

## Methods

### Literature Search and Data Collection

We conducted this meta-analysis using a recently published global database that compiles 4756 measurements of aboveground live tree carbon stocks in timber plantations, collected from 829 distinct sites across 278 studies [45••]. We subset this dataset to the 654 measurements of aboveground live tree biomass in timber plantations across 45 studies, 56 distinct sites, and 19 tree genera that included relevant management details. Full details on the monoculture plantation database compilation and data standardization processes are described in Bukoski et al. [45••]. In addition, we included 97 measurements of aboveground live trees biomass collected from one large compilation of aboveground carbon stocks in planted

forests [46••] that was found at a later date. We elected to include it as it substantially increased our sample size. In total, we collected 751 measurements of aboveground live tree biomass across 68 studies (Table S1.1, S1.2, and S1.3), 80 distinct sites, and 19 tree genera (Fig. 1).

For each of the three dominant silvicultural treatments—intercropping of N-fixing species, application of inorganic NPK fertilizer, and stand density management (i.e., thinning), we selected studies from our monoculture plantation database that had both (i) measurements of aboveground biomass from plots in which the treatment had been applied and (ii) measurements of aboveground biomass from control plots (without the treatment of interest). Only studies with imposed, replicated treatments at one or more sites were included in our dataset. Multiple comparisons within a single study (e.g., comparing different thinning intensities to a single unthinned control) were considered as distinct within-study observations, and in that case, treatment values were compared to the same control value. In total, there were 43 studies comprising 197 comparisons for intercropping of N-fixing plants, 17 studies comprising 164 comparisons for NPK fertilization, and 8 studies comprising 62 comparisons for thinning (the included studies are in Tables S1.1 & Table S1.2 & Table S1.3). Within the intercropped plots, we conservatively accounted for aboveground biomass of only the main tree crops to assess the fertilization effect of the interplanted N-fixing plants on the latter. Before running the analysis, we converted aboveground live tree biomass measurements to aboveground carbon using a 0.47 default conversion factor [47].


## Standardization of Treatment Effect Using Natural Log of the Response Ratio

For our meta-analysis, we used the natural log of the response ratio metric ( $\ln RR$ ) to standardize the effect of treatments on aboveground carbon across studies. Here,  $\ln RR$  reflects the change in aboveground carbon induced by each of our three treatments—interplanting of N-fixing species, application of NPK fertilizers, and thinning. We calculated  $\ln RR$  using Eq. (1):

$$\ln RR = \ln \frac{\bar{X}_t}{\bar{X}_c} = \ln \bar{X}_t - \ln \bar{X}_c \quad (1)$$

In Eq. (1),  $\bar{X}_t$  and  $\bar{X}_c$  are mean aboveground carbon in the paired treatment and control, respectively. A  $\ln RR$  value of 0 means the treatment did not induce any change in carbon compared to the control, while a positive value indicates the treatment had a positive effect on aboveground carbon, and a negative value indicates a decrease in aboveground carbon. To account for variation in sampling effort between studies, we weighted the effect sizes by the inverse of the sample variance for each response ratio. We calculated the sample variance using the standard error and number of replicates reported for each study [48]. When studies did not report the standard deviations associated with aboveground carbon measurements, we followed the methodology of Lajeunesse [49] and Koricheva [50]. This consisted of imputing the missing standard deviations by calculating the median coefficient of variation (ratio of standard deviation and mean) for each group (treatment or control) from

**Fig. 1** Map of locations of all sites included in the meta-analysis of intercropping N-fixing plants, NPK fertilization, and thinning effects on planted forests aboveground carbon stocks



studies that reported both means and standard deviations. We then multiplied the median coefficient of variation by the reported mean for either treatment or control groups for which standard deviations were missing. We performed a sensitivity analysis to test for any potential effects of these assumptions on our results (see Supplementary Information), which revealed that almost all results were robust to this approach, but we note where results were sensitive to the imputed standard deviations. Finally, we used funnel plots to confirm the absence of publication bias [51]. Response ratios were calculated using the {metaphor} package [52] in Program R v.1.4.1103 [53].

### Testing of Moderator Effects and Mixed-Effects Approach

We inserted several moderators (variables that influence the strength and/or form of a relationship between a predictor and a dependent variable [54]) into the mixed-effects model to assess how the treatment effect size varied with other factors hypothesized to influence aboveground carbon (e.g., genus or soil moisture regime). Our selection of moderator variables included both categorical and continuous moderators and sought to account for an array of environmental,

biological, and human factors, but was also limited by data availability (Table 1). Except soil moisture regime, all moderator variables were recorded using information reported in the studies themselves. To obtain soil moisture regimes, we intersected the geographic coordinates of each site with a map of global soil moisture regimes developed by USDA-NRCS [55]. This map of global soil moisture regimes was built using data taken from more than 22,000 climatic stations distributed around the world [55]. Soil moisture regime data were interpolated and rasterized on a 2-min grid cell [55].

Moderators were tested individually with separate models, such that the influence of a moderator on the effect size was determined using all available data for that particular moderator variable. Prior to analysis, we dropped categories of moderators for which we had only one observation. To account for the non-independence of multiple within-study observations, we inserted publication-level random effects into each of our models [50]. We fitted all models using restricted maximum likelihood estimation [50]. We determined the statistical significance of each moderator variable using an omnibus test of all model coefficients ( $p$ -value < 0.05) [52]. Prior to reporting our final results, we back-transformed ( $e^{\lnRR}$ ) the mean log response ratios and

**Table 1** Moderator variables inserted in the mixed-effects meta-analytic models for each treatment

Treatments	Moderator	Moderator type	Categories/units
Interplanting N-fixing plants	Previous land use	Categorical	Cropland, natural forest, plantation
	Tree genus	Categorical	<i>Alnus</i> , <i>Anacardium</i> , <i>Casuarina</i> , <i>Eucalyptus</i> , <i>Hymeronima</i> , <i>Pachira</i> , <i>Pinus</i> , <i>Populus</i> , <i>Pseudotsuga</i>
	Soil moisture regime	Categorical	Perudic, udic, ustic, xeric
	Intercropped plant genus	Categorical	<i>Acacia</i> , <i>Albizia</i> , <i>Alnus</i> , <i>Dalbergia</i> , <i>Enterolobium</i> , <i>Hippophae</i> , <i>Leucaena</i> , <i>Lupinus</i> , <i>Paraserianthes</i> , <i>Robinia</i> , <i>Salix</i>
	Wood type	Categorical	Hardwood, softwood
	Experimental design	Categorical	Additive, replacement
	Stand age	Continuous	Years
	Mean annual precipitation	Continuous	mm/year
Inorganic NPK fertilization	Previous land use	Categorical	Cropland, fire, natural forest, plantation
	Tree genus	Categorical	<i>Ailanthus</i> , <i>Eucalyptus</i> , <i>Macaranga</i> , <i>Picea</i> , <i>Pinus</i>
	Soil moisture regime	Categorical	Perudic, udic, ustic, xeric
	Wood Type	Categorical	Hardwood, softwood
	Stand age	Continuous	Years
	Time since fertilization	Continuous	Years
	Application method	Categorical	Continuous, pulse
	Mean annual precipitation	Continuous	mm/year
Thinning	Previous land use	Categorical	Cropland, natural forest, plantation
	Soil moisture regime	Categorical	Perudic, udic, ustic, xeric
	Wood Type	Categorical	Hardwood, softwood
	Tree genus	Categorical	<i>Acacia</i> , <i>Cunninghamia</i> , <i>Eucalyptus</i> , <i>Pinus</i>
	Time since thinning	Continuous	Years
	Mean annual precipitation	Continuous	mm/year
	Basal area removed	Continuous	Percent

their 95% confidence intervals and converted these values to percent change relative to the control.

Depending on data availability, we also tested how the effect size varied with either tree age or time since treatment using linear mixed-effects models. In all the interplanting treatment studies, N-fixing crops were planted at the same time as the main tree crop, and we therefore used stand age as our time variable. To limit the potential effects of sparse data at older ages, we dropped measurements above 20 years old before performing this regression. In total, 2 data points at 21 and 28 years old were dropped prior to this linear regression analysis. Of the inorganic fertilizer studies, 9 of 18 continuously applied inorganic fertilizer over the course of the study period and we used stand age as our time variable. However, 10 of 18 inorganic fertilizer treatment studies reported “time since treatment” data. For these observations, we also used mixed-effects regression models to test how the effect size varied with time since treatment in inorganically fertilized plots (Table 1). Limited data were available for older times since NPK fertilization. To limit the potential effects of sparse data at older times since treatment, we dropped measurements above a time since treatment of 5 years before running this regression. In total, two data points at 21 years old were dropped prior to this linear regression analysis. For our thinning observations, we assessed how the effect size varied with time since treatment in thinned plots as well (Table 1). We used all the time since thinning data to perform the regression analysis. Finally, we evaluated whether annual mean precipitation had an influence on the effect size of all three treatments using the same regression model type (Table 1).

### Greenhouse Gas Emissions from Inorganic Fertilizer Application

For each aboveground carbon measurement reported in NPK fertilization studies, we calculated the business as usual (BAU) associated amount of nitrous oxide (N<sub>2</sub>O) emitted (expressed in CO<sub>2</sub>e) for each ton of synthetic nitrogen (N) fertilizer applied, both in-field and upstream from fertilizer manufacturing itself [5]. We used an emissions factor of 2.54% for N fertilizer (11.9 MgCO<sub>2</sub>e per MgN applied) for in-field emissions and an upstream emissions factor of about 4 kgCO<sub>2</sub>e per kgN produced [5]. We extracted tonnage of N fertilizer directly from the studies included in this meta-analysis (Table S1.2). When the amount of N fertilizer used was reported as kilograms per tree, we converted to kilograms per hectare using the reported stand density value. We then compared the additional aboveground carbon gain induced by NPK fertilization (i.e., the difference in aboveground carbon between the treated and control plots) with emissions from N fertilizers use and manufacturing. We calculated the difference between additional carbon and N

fertilizer emissions and examined how this effect varied with the amount of N fertilizer applied using a linear regression model. Finally, we estimated the median net carbon balance of fertilized stands (expressed MgCO<sub>2</sub>e ha<sup>-1</sup>) across all studies included for the NPK fertilization treatment.


## Results

Our results showed effects on aboveground carbon that were (i) strongly positive and statistically significant for N-fixing species, (ii) strongly positive and statistically significant for the use of NPK fertilizer, and (iii) negative and statistically significant for thinning. Further, our incorporation of moderator variables provided additional nuance on how these silvicultural practices impact aboveground carbon in plantations. We provide additional details on each of the three treatments below.

### Interplanting of N-Fixing Species

Overall, intercropping of N-fixing plants in monoculture stands had a significantly positive effect (approximately +20%) on aboveground carbon of the primary species (Fig. 2) ( $p$ -value = 0.0031). All moderator variables were found to have significant effects on the relationship between interplanted N-fixing crops and aboveground carbon (Fig. 2). We found that “soil moisture regime” had a strong influence on the magnitude of the effect size ( $p < 0.001$ ) (Fig. 2). Interplanting of plantations growing on moister soils significantly increased aboveground carbon (+52% for perudic and +42% for ustic soils), but not in places with drier xeric soil moisture regimes (Table S5.1). Furthermore, N-fixing companion crops significantly increased the aboveground carbon stocks of the main tree crop when plantations occurred on former croplands (+67%,  $p < 0.0001$ ) (Fig. 2), rather than locations with tree cover. The genus of the main tree crop also influenced the magnitude of the effect size ( $p < 0.001$ ). Specifically, establishing *Eucalyptus* trees with N-fixing companion crops increased the latter aboveground carbon stocks by +25%, whereas other genera did not show a significant effect. Additionally, the genus of the intercropped species had a significant impact ( $p$ -value < 0.001) on the effect size as well (Fig. 2). Intercropping with *Leucaena*, *Albizia*, *Enterolobium*, and *Hippophae* induced a 52%, 80%, 87%, and 113% increase in the aboveground carbon of the main tree crop, respectively, whereas other intercropping genera did not have a significant effect (Fig. 2; Table S5.1). The type of wood of the main tree species also moderated the effect of the treatment ( $p$ -value = 0.0036). Hardwoods reacted positively to interplanted N-fixing crops (+25% in aboveground carbon stocks) (Table S5.1). Finally, the type

**Fig. 2** Meta-analysis results of the change in aboveground carbon of plantation trees in response to the interplanting of N-fixing plants. Error bars represent the 95% confidence intervals. Omnibus tests of significance for moderator variables are shown on the right side (NS means “not significant”). Results for the “Tree genus” moderator are only provided for genera with more than 5 observations (see Table S5.1). Results for the “Intercropped Genus” moderator are only provided for genera with significant effects or for genera with more than 5 observations (see Table S5.1). The number of observations in each category is shown in parenthesis



of experimental design influenced the magnitude of effect size ( $p$ -value < 0.001) (Fig. 2). The use of interplanted N-fixing crops appeared to be more beneficial when an additive design was adopted (+72%) than a replacement one (+16%) (Table S5.1).

However, our results also suggest that the effect of intercropping N-fixing species on aboveground carbon varied with stand age. By regressing  $\ln(\text{RR})$  on stand age, we identified a significant positive association with the treatment effect size ( $p$ -value < 0.0001) (Fig. 3). For every additional year that a stand was allowed to grow, the influence of intercropping N-fixing species on plantation aboveground carbon was increased by 3.6% (Fig. 3). Intercropping initially decreased aboveground carbon in the main tree crop, although not significantly, but its effect became positive when the stand was 3 years old and steadily increased over time from then on (+84% at 20 years old) (Fig. 3).


Finally, we did not identify a significant effect of mean annual precipitation, the other continuous variable in our model, on the magnitude of the effect size ( $p$ -value = 0.0872; Table S5.1).

## Inorganic NPK Fertilization

We found that inorganic NPK fertilizers significantly increased aboveground carbon by 44.5% overall ( $p$ -value < 0.001) (Fig. 4). Of the individual studies included in the meta-analysis, most reported a significant positive effect of NPK fertilization on aboveground carbon, whereas six studies found a non-significant effect. Similarly to interplanting of N-fixing species, all moderator variables were found to have significant effects on the relationship between NPK fertilizer and aboveground carbon.

We found that “tree genus” significantly influenced the change in aboveground carbon attributed to inorganic NPK fertilization (Fig. 4) ( $p$ -value < 0.001). The aboveground carbon of treated *Pinus* and *Eucalyptus* plantations were 51% ( $p$ -value < 0.001) and 37% ( $p$ -value < 0.001) higher in fertilized plots compared to the controls, respectively (Fig. 4; Table S5.2). In addition, both hardwoods and softwoods responded positively to the treatment; however, NPK fertilization had a slightly

**Fig. 3** Change in the effect size of interplanting N-fixing plants as a function of stand age. The significance of the regression is indicated by the  $p$ -value in the upper right as well as the intercept and slope values with their corresponding 95% confidence interval. The area shaded in blue around the regression lines indicate the 95% confidence interval



greater effect on aboveground carbon in hardwood (+46%) ( $p$ -value  $< 0.001$ ) relative to softwood plantations (+43%) ( $p$ -value  $< 0.001$ ) (Fig. 4; Table S5.2).

Soil moisture regime also had a strong influence on the magnitude of the effect size ( $p$ -value  $< 0.001$ ) (Fig. 4). NPK fertilization of plantations growing on moister soils significantly increased aboveground carbon (+82% for perudic and +42% for udic soils), but not in places with xeric and ustic soil moisture regimes (Table S5.2). It is worth noting that the influence on the effect size of the perudic soil moisture regime might be overestimated as only one study reported measurements for that soil moisture regime (Fig. 4). Furthermore, the methodology used to apply NPK fertilizers significantly influenced the magnitude of the fertilizing effect ( $p$ -value  $< 0.001$ ) (Fig. 4). The effect size appeared higher when fertilizers are applied episodically (+53%) rather than continuously (+36%) (Table S5.2). The type of previous land use also strongly influenced the treatment effect size and explained part of the heterogeneity in effect size across studies (Fig. 4) ( $p$ -value  $< 0.001$ ). Specifically, the use of inorganic fertilizers appeared to be more beneficial on lands that were previously plantations (+44%) or natural forests (+76%), than those cleared by fire or previously used for croplands (Table S5.2).

We found a negative, although not significant ( $p$ -value = 0.5), association between stand age and the magnitude of the effect size in fertilized stands (Table S5.2). However, when we used time since treatment rather than stand age, we found that the benefit of inorganic fertilizers on aboveground carbon significantly decline through time ( $p$ -value = 0.008) (Fig. 5). For every additional year since the

treatment, the effect size of inorganic fertilization on carbon stocks decreased by 6.66% (Fig. 5).


Mean annual precipitation also had a significant negative effect ( $p$ -value  $< 0.001$ ) on the magnitude of the change in aboveground carbon (Figure S3.1). For every additional millimeter of precipitation received per year, the treatment effect size was reduced by 0.03% (Figure S3.1).

Once we accounted for in-field and upstream fertilizer emissions, we found a negative median net carbon balance of fertilized stands across studies ( $-2 \text{ MgCO}_2\text{e}\cdot\text{ha}^{-1}$ ). In other words, fertilized stands were net emitters across studies (Figure S3.2), and the climate mitigation benefit of fertilizer declined with the amount of N fertilizer applied ( $p$ -value  $< 0.001$ ). The climate benefit became null at 0.55 MgN or greater applied per hectare (Fig. 6).

## Thinning

Across the different studies, thinning decreased standing aboveground carbon by about 34% ( $p$ -value  $< 0.001$ ) (Fig. 7). We found soil moisture regime to have a significant impact on the percent change in carbon from thinning ( $p$ -value  $< 0.001$ ) (Fig. 7). When thinning was conducted on xeric soils, carbon levels decreased significantly by 59% ( $p$ -value  $< 0.001$ ) (Table S5.3). We found type of previous land use to also have a significant influence on the magnitude of the effect size ( $p$ -value  $< 0.001$ ) (Fig. 7). Stands grown on former agricultural lands had aboveground carbon levels that were 55% lower ( $p$ -value  $< 0.001$ ) in thinned stands compared to controls (Table S5.3). When thinning was executed in stands that were established on former natural





**Fig. 4** Meta-analysis results of the change in aboveground carbon of plantation trees in response to NPK fertilization error bars represent the 95% confidence intervals. Omnibus tests of significance for


moderator variables are shown on the right side (NS means “not significant”). The number of observations in each category is shown in parenthesis

forestlands, the mean decrease in aboveground carbon was 44% ( $p$ -value = 0.0026) (Table S5.3). Finally, we also found that the type of wood influenced the magnitude of the effect size ( $p$ -value < 0.001) (Fig. 7). Both hardwood and softwood stands experienced large reductions in aboveground carbon due to thinning operations, which averaged a –33% change in aboveground carbon from thinning for both wood types.


Genus of tree did not significantly influence the magnitude of the effect size.

When we inserted the “time since treatment” continuous moderator variable in the model, we found that as time since thinning increased, the negative effect of thinning on carbon lessened ( $p$ -value < 0.001) (Fig. 8). For every additional year since the treatment, the effect size of thinning on

**Fig. 5** Change in the fertilizing effect size of NPK fertilizers as a function of time since treatment. The significance of the regression is indicated by the *p*-value in the upper right as well as the intercept and slope values with their corresponding 95% confidence interval. The area shaded in blue around the regression lines indicate the 95% confidence interval



**Fig. 6** Change in the net carbon balance of fertilized stands as a function of the amount of N applied. The significance of the regression is indicated by the *p*-value in the upper right as well as the intercept and slope values, and the coefficient of determination. The area shaded in blue around the regression lines indicate the 95% confidence interval




carbon stocks decreased by 4.6% (Fig. 8), suggesting that after 14 years the thinning effect disappears.


Mean annual precipitation also had a significant negative effect (*p*-value = 0.04) on the magnitude of the change in aboveground carbon (Figure S3.3). For every additional millimeter of precipitation received per year, the effect of thinning on carbon stocks was reduced by 0.03% (Figure S3.3). Finally, we found that as the percent of basal area removed increased, the negative effect size of thinning increased

(*p* < 0.001) (Table S5.3). For every additional percent of basal area removed, the effect of thinning on carbon stocks was increased by 1.3% (Table S5.3).

**Fig. 7** Meta-analysis results of the change in aboveground carbon of plantations from thinning. Error bars represent the 95% confidence intervals. Omnibus tests of significance for moderator variables are shown on the right side (NS means “not significant”). The number of observations in each category is shown in parenthesis



**Fig. 8** Change in the effect size of thinning operations as a function of time since thinning. The significance of the regression is indicated by the *p*-value in the lower right as well as the intercept and slope values with their corresponding 95% confidence interval. The area shaded in blue around the regression lines indicate the 95% confidence interval



## Discussion

Enhanced aboveground carbon sequestration in plantations via improved stand management is a prominent strategy for mitigating climate change. Our study quantified the effect size of three dominant silvicultural practices—intercropping N-fixing species, fertilization with inorganic NPK, and stand density management via thinning—on aboveground carbon stocks in plantations. We found that the magnitude of the effect size often depended on context and below we discuss the dominant drivers of variation and implications for landowners and land managers.

### Interplanting of N-Fixing Plants

We found that only some intercropping N-fixing species may be beneficial (in our case *Albizia*, *Enterolobium*, *Hippophae*, and *Leucaena* species) and/or primarily when the crop trees are mature. The species-specific effects may be mediated by compatibility of growth patterns [56, 57]. Growth compatibility can occur through at least three mechanisms: reduced competition for light via canopy stratification and crown complementarity, reduced competition for nutrients and water via root stratification, and direct or indirect growth facilitation [16, 58, 59]. All three mechanisms may explain our results. We found that intercropping with shade tolerant or moderately shade tolerant species (e.g., *Leucaena* spp., *Enterolobium* spp.) had greater effects on aboveground carbon relative to shade intolerant species (e.g., *Alnus* spp.). Root stratification couples trees with deep taproots (e.g., *Eucalyptus* spp.) that can acquire nutrients and water from lower soil horizons with species with shallow extensive horizontal root systems (e.g., *Acacia* spp.) [60]. Interplanting N-fixing plants may also confer benefits through chemical root stratification, which may facilitate mycorrhizal relationships between intercropped species and the main tree crops [16]. Furthermore, although we did not directly test these interactions, interplanting N-fixing legumes facilitate nutrient access for plantation trees by decreasing soil pH which promotes soil weathering and therefore general nutrient availability [61•]. Lastly, companion crops may inhibit the growth of competitors via allelopathic effects and/or eliminate local pathogens [58, 62].

Our results also showed that interplanting N-fixing plants become more beneficial through time. Early growth incompatibility and poor establishment of nitrogen-fixing companion crops could explain the negative or absent effect of intercropping early in the rotation [63, 64]. When crops are planted simultaneously, intense competition between them for resources such as light water or nutrients

may emerge soon after their establishment [56, 65••]. Conversely, when the N-fixing species is established a few years after the main crop, the former tends to compete with the latter to a much lesser degree for light and water [56, 65••, 66, 67]. As a result, complementarity between the crops increases and the main crop can allocate a greater proportion of aboveground growth using the additional N provided by the companion crop [65••].

Our analysis also showed that several environmental, biological, and human factors influence the magnitude of response of the major crops to interplanting of N-fixing species. The main tree crop benefited more from the treatment when it was performed on wetter soils. Soil moisture significantly affects N-fixation by controlling nodulation and nitrogenase activity [68]. Moist soils promote nodulation and nitrogenase activity while drier soils reduce the number of nodules produced and inhibit nitrogenase activity, resulting in very low rates of nitrogen fixation [68]. Wetter soils also promote the growth of companion crops themselves, which allows for more N-fixation and thus increases the amount of available nitrogen in the soil [68].

Furthermore, interplanting N-fixing crops was most beneficial when it was performed in plantations growing on lands that previously hosted agricultural crops, compared to those previously under tree cover. Agricultural lands often experience soil impoverishment, especially when intense cropping techniques are used [69]. Low nutrient concentration, especially for N, appears to favor complementarity between the main tree crop and its N-fixing companion crop [65••]. Nonetheless, since soil impoverishment also depends on the intensity of the land use, which was not captured here, this result should be interpreted with caution.

### Inorganic NPK Fertilization

The use of NPK fertilizers significantly increased the aboveground carbon stocks of plantations, which aligns with the findings of others [70, 71]. However, once we accounted for the CO<sub>2</sub>e emissions resulting from inorganic N fertilizer manufacturing and use, we found that higher levels of fertilizer application could negate and/or overwhelm the increase in aboveground carbon stocks induced by fertilization. This highlights that the climate mitigation potential of fertilized stands could be substantially overestimated if fertilizer emissions are not taken into consideration. This also emphasizes that inorganic fertilizers have a greenhouse gas cost that can exceed their carbon benefit, particularly when they are applied in large amounts [72]. To maximize the climate benefits of using synthetic fertilizers in planted forests, a holistic accounting of greenhouse gas emissions associated with producing and applying fertilizers is therefore needed [73].

We found that the benefit of NPK fertilization decreased with time. This phenomenon has been noticed in previous

studies and reviews [14, 74, 75] and several factors are believed to explain this trend. During the initial establishment phase, young seedlings generally do not have well developed root systems and providing easily accessible inorganic forms of nutrients can increase growth [18•, 74]. Over time, unfertilized seedlings develop their root systems and gain access to larger nutrient pools, which could explain the decrease in the magnitude of the effect size observed here [32, 74]. Furthermore, plantations trees increasingly rely on organic forms of N like glycine relative to inorganic forms (e.g.,  $\text{NH}_3$  &  $\text{NH}_4^+$ ) as they age [76], which could explain the lower effect size of fertilizers later in the rotation. Such a phenomenon might also apply in the context of the interplanting treatment discussed before. The effect of fertilizers on biomass carbon growth has also been suggested to decrease over time due to increased allocation of resources towards reproduction, which leads to lower nutrient uptake rates [77].

Our results showed that the benefit of NPK fertilization on growth declined as precipitation increased. High precipitation levels increase soil water content and, in some cases, push it beyond field capacity (i.e., soil becomes saturated) [78]. Saturated soils tend to be highly prone to leaching of very mobile nutrients such as N, which reduces their availability for plant uptake [15••, 78]. Our results also suggest that plantations growing on soils with udic moisture regimes experienced the largest increases in aboveground carbon when fertilized. Nutrients are generally more available when soil moisture levels are optimum and remain adequate for most of the year, as is the case for soils with an udic soil moisture regime [79].

Conversely to interplanting, the benefit of NPK fertilizers was higher when they were applied in plantations growing on lands that previously hosted tree plantations or natural forests. Harvested lands often experience soil impoverishment, especially when the whole tree harvesting method is employed [80, 81]. Indeed, harvest promotes nutrient mineralization and nitrification, making them more mobile and therefore more subject to leaching [82]. The removal of trees themselves also leads to a decrease in soil nutrient capital and can curb the growth of the next trees [83]. Fertilization is often used to compensate for the loss of nutrients [83], and poor soil post-harvest may explain the fertilizer benefit we observed. However, since this analysis did not account for the effect of land use intensity on soil impoverishment, this result should be interpreted with caution. Overall, these findings related to site conditions underline the fact that response of trees to fertilization is heavily site dependent and varies with site productivity and quality, and/or intensity of prior land use.

This study also revealed that plantation stands store more aboveground carbon when fertilizer applications are performed in pulses, rather than continuously. Aber et al. [84]

and Saarsalmi and Mälkönen [85] found similar results and attributed it to higher nutrient-use efficiency under pulse fertilization. Furthermore, several studies have stressed that continuous application of N favors faster N saturation, which reduces the positive effects of inorganic N-based fertilizers on biomass growth in the long-term [86]. Since N saturation is also believed to promote nitrate leaching from soils [87], increase emissions of  $\text{N}_2\text{O}$  [84], and induce large reduction in mycorrhizal symbionts [88], our results underline that careful handling of fertilizer operations is needed to maximize tree growth while minimizing negative environmental side effects. Nonetheless, this difference in growth responses to pulsed and continuous fertilization might also be caused by other chemical or physical soil variables (e.g., initial soil fertility, soil pH) which were not captured in this analysis.

## Thinning

Our results showed that thinning operations significantly lowered aboveground carbon in plantations, which aligns with the findings of others [40, 89]. Given that thinning is performed to reallocate resources and growing space to target trees of primary value and induces a trade-off in loss of stand-level carbon, this result was expected. Indeed, increased growth rates for individual trees post-thinning have been well documented [44, 90, 91]. We also identified an attenuation of the effect on overall stand aboveground carbon with increasing time since thinning (Fig. 8). Our results suggest that at the stand level, the negative effect of thinning on aboveground carbon lasts for approximately a decade before becoming attenuated.

Furthermore, we found that the negative effect of thinning on aboveground carbon might be amplified when performed in stands located in dry areas and growing on xeric soils. Non-thinned trees growing in stands on mesic and wetter soils tend to have access to more resources such as water and nutrients after treatment than the same trees growing on xeric soils, which could therefore induce a greater growth response to thinning operations and a weaker treatment effect in the former scenario than in the latter.

Thinning operations appear to have a more negative effect on stand carbon when they are performed on former agricultural lands. Nutrient and water availability are generally lower in those areas, and fertilization and irrigation are often needed to ensure afforestation/reforestation success [92]. As a result, thinned stands growing on former agricultural lands might have access to smaller nutrient and water pools that constrain post-thinning growth and therefore induce a larger thinning effect on stand-level aboveground carbon. Furthermore, former agricultural lands tend to be more compacted due to heavy machinery use [93], which could restrict the common increase in resource availability after thinning operations and further

accentuate the effect on aboveground carbon [43]. Overall, these results stress that response of stands to thinning is heavily site dependent. Stands on more productive sites will often respond quite differently to the same thinning treatment than a stand on lower quality sites [94].

### Caveats and Potential Limitations

Our study focuses on aboveground live tree carbon stocks only and did not examine changes in belowground carbon and soil organic carbon in response to fertilization and thinning. Belowground carbon and soils are substantial carbon pools in forest ecosystems. For example, soil carbon stocks are believed to represent 90% and 50% of the total carbon stock in boreal and tropical forests respectively [95]. Furthermore, in these forest types, roots are believed to store between 23 and 27% of total tree biomass carbon [96]. However, we omitted these pools due to little data availability and inconsistent field sampling methods across reviewed studies. This decision was further justified by the fact that changes in soil carbon stocks tend to be small relative to changes in the aboveground carbon pool and it takes longer to detect them [97]. Previous studies have found positive effects of NPK fertilization and interplanting N-fixing plants on soil carbon stocks in forest plantations [86, 98, 99]. Other studies have found neutral or negative effects of thinning on soil carbon stocks [35, 36, 40, 89, 100] and negative effects on belowground carbon as well [100]. If these results occur in other forest plantation management studies, focusing solely on carbon in aboveground biomass tissues may underestimate the impact of forest management actions such as fertilization on total ecosystem carbon stocks and overestimate it in the context of thinning. Additional empirical data on the impact of these management operations on plantations total carbon stocks as well as studies reporting responses of the different carbon pools are needed.

Data limitations also partly restricted our ability to assess the exact timing at which the positive effects of inorganic fertilizers on aboveground stand carbon stocks disappear. Although the data stopped at 5 years since treatment, if the decline through time continues following the same path, then we would predict the carbon benefits of inorganic fertilizers to disappear eight years after treatment. Further data on the impact of time on the effect size of fertilizers would therefore be valuable to refine our predictions. Moreover, data were heavily skewed towards younger ages across all of our focal silvicultural treatments. Long-term measurements of the effects of silvicultural treatments on plantation forest carbon stocks will greatly improve our understanding of the climate mitigation potential of these systems.

### Management Implications

This study reveals that the use of N-fixing companion crops as a fertilization technique is challenging and primarily depends heavily on growth complementarity between the crop and intercropped species. Knowledge of the silvics of individual species is a critical precursor to ensure successful growth benefits, and poorly implemented systems may induce null effects on growth. However, use of intercropping as the main fertilization technique can also reduce N<sub>2</sub>O emissions and nutrient volatilization, which are common issues associated with the application of inorganic NPK fertilizers [101]. Factors such as prior land use, soil moisture regime, and method of implementation are also likely to be key moderators of the effect of interplanted N-fixing plants on aboveground carbon. Knowledge of the site characteristics and history, as well as interactions between tree species seem therefore crucial to ensure successful associations.

Inorganic fertilizers, despite being growth catalyzers, can have detrimental effects on the environment which includes water eutrophication via nutrient leaching and runoffs, and air pollution via greenhouse gas (GHG) emissions from fertilizers manufacturing and nutrient volatilization. Thus, there are benefits to minimizing additions. Our results show that NPK fertilizers are not universally beneficial and their positive effects, when observed, decline through time. Furthermore, we found that GHG emissions resulting from N fertilizer manufacturing and in-field application could exceed the aboveground carbon gains induced by their use. Our results also stress that factors such as prior land use and soil moisture regime may be key moderators of the effect of fertilizers on aboveground carbon. Our findings emphasize therefore the need to apply the right rate of nutrients at the right times and in the right context. Landowners should carefully handle fertilization operations to maximize their carbon benefits and minimize their costs.

Next, our study suggests that the initial decrease in aboveground carbon caused by thinning operations could be compensated over time by the recovery of carbon post-thinning, particularly when coupled with extended rotation lengths. Indeed, by increasing rotation lengths, landowners can further attenuate the effects of thinning on stand carbon up to a point at which the initial loss of carbon is offset by the recovery of a productive and well-stocked stand [40, 42, 102]. From there, the individual trees in thinned stands could sequester additional carbon to levels as high or higher than unthinned stands of the same age [42, 102]. Critically, the likelihood of recovering unthinned carbon levels post-thinning will depend on the intensity of thinning and thinning technique employed, with more intensive thinning practices tending to result in longer or unachievable recovery of aboveground carbon [15••, 42, 91, 102]. For example, plots thinned from below may rapidly recover carbon

sequestration rates equivalent to control plots, while stands that experienced dominant and crown thinning operations may have longer periods to carbon recovery [42]. However, thinning can also confer important forest structural qualities important for management of forest carbon beyond maximization of standing aboveground carbon stocks [13••], such as conferring resistance to low or medium-intensity wild-fire in fire-prone landscapes and insect/pathogen outbreaks [13••, 103]. Further research on these co-benefits and the coupling of thinning with extended rotations is needed to validate the use of thinning as a forest carbon management tool.

Lastly, this meta-analysis underlines the need for forestry practitioners to be aware of site conditions, quality, and land use history before conducting thinning operations because their effects can vary from site to site, and they tend to strongly depend on local conditions. Additional studies on the influence of soil moisture regimes and previous land use on the magnitude of the effect size of thinning on stand aboveground carbon are needed to consolidate our results. Overall, our study underscores that selecting the appropriate species and treatments for each site is crucial to ensure an effective carbon management plan in forest plantations.

## Conclusion

Improved forest management has been highlighted as a key natural climate solution because of its ability to deliver climate benefits within short-time scales [1, 5, 7•, 8•]. Nonetheless, there is still substantial uncertainty regarding the forest practices that would help realize this mitigation potential and the context in which they would deliver the most climate benefits [1]. Our study provides additional considerations that help facilitate the use of improved management practices to increase plantations carbon stocks and hence mitigate climate change. By specifying the conditions in which fertilization and stand density management tend to be the most beneficial for carbon storage purposes, this study provides additional information to forest practitioners on how to use them as carbon management tools. Although not all management actions studied here provide substantial increases in carbon storage over the entire lifetime of plantation trees (e.g., interplanting N-fixing trees), they still may be desirable by enhancing monocultures biodiversity levels and resilience to disturbances such as pests or natural disasters [18•, 104, 105]. The latter tends to be crucial for the permanence and durability of carbon sequestered on the landscape, particularly under a changing climate [106•].

**Supplementary Information** The online version contains supplementary material available at <https://doi.org/10.1007/s40725-023-00182-5>.

**Acknowledgements** We thank Timothy Bowles and the Potts Group at U.C. Berkeley for valuable feedback on early drafts of this manuscript. Although their comments have greatly improved our study, the ideas and opinions expressed in our study do not necessarily reflect their own.

**Funding** We thank the Bezos Earth Fund for supporting C. H. M.'s time on this work. Grants from the following institutions to the Nature Conservancy partially funded S. C. C.-P.'s time on this work: the Children's Investment Fund Foundation, COmON Foundation, the Craig and Susan McCaw Foundation, the Doris Duke Charitable Foundation, Good Energies Foundation, and the Bezos Earth Fund.

**Data Availability** Data used for the study are published on Zenodo (<https://doi.org/10.5281/zenodo.7789868>)

## Declarations

**Conflict of Interest** M. D. P. is the Chief Science Officer for Carbon Direct Inc., a company combining science, technology, and capital to deliver quality CO<sub>2</sub> management at scale. M. D. P. is a shareholder in the company and thus stands to benefit financially from forest management targeted at climate change mitigation. The remaining authors declare no competing interests.

**Human and Animal Rights and Informed Consent** This article does not contain any studies with human or animal subjects performed by any of the authors.

**Open Access** This article is licensed under a Creative Commons Attribution 4.0 International License, which permits use, sharing, adaptation, distribution and reproduction in any medium or format, as long as you give appropriate credit to the original author(s) and the source, provide a link to the Creative Commons licence, and indicate if changes were made. The images or other third party material in this article are included in the article's Creative Commons licence, unless indicated otherwise in a credit line to the material. If material is not included in the article's Creative Commons licence and your intended use is not permitted by statutory regulation or exceeds the permitted use, you will need to obtain permission directly from the copyright holder. To view a copy of this licence, visit <http://creativecommons.org/licenses/by/4.0/>.

## References

Papers of particular interest, published recently, have been highlighted as:

- Of importance
  - Of major importance
1. IPCC. Climate Change 2022: Mitigation of Climate Change. Contribution of Working Group III to the Sixth Assessment Report of the Intergovernmental Panel on Climate Change. Cambridge University Press, Cambridge, UK and New York. 2022.
  2. Cook-Patton SC, Leavitt SM, Gibbs D, et al. Mapping carbon accumulation potential from global natural forest regrowth. *Nature*. 2020;585:545–50.
  3. Cook-Patton SC, Gopalakrishna T, Daigneault A, et al. Lower cost and more feasible options to restore forest cover in the contiguous United States for climate mitigation. *One Earth*. 2020;3:739–52.
  4. Di Sacco A, Hardwick KA, Blakesley D, et al. Ten golden rules for reforestation to optimize carbon sequestration,

- biodiversity recovery and livelihood benefits. *Glob Change Biol.* 2021;27:1328–48.
5. Griscom B, Adams J, Ellis P, et al. Natural climate solutions. *Proc Natl Acad Sci.* 2017. <https://doi.org/10.1073/pnas.1710465114>.
  6. Vincent J, Curran S, Ashton M. Forest restoration in low- and middle-income countries. *Annu Rev Environ Resour.* 2021. <https://doi.org/10.1146/annurev-environ-012220-020159>.
  - 7.● Cook-Patton S, Drever C, Griscom B, et al. Protect, manage and then restore lands for climate mitigation. *Nat Clim Chang.* 2021. <https://doi.org/10.1038/s41558-021-01198-0>. **(This paper presents a hierarchy of natural climate solutions for short- and long-term climate mitigation. It highlights the most cost-effective mitigation pathways, including improved forest management, that need to be implemented on the ground on a large scale to maximize carbon removal.)**
  - 8.● Roe S, Streck C, Beach R, et al. Land-based measures to mitigate climate change: potential and feasibility by country. *Glob Change Biol.* 2021;27:6025–58. **(This study provides one of the most recent estimates of the climate change mitigation potential of improved forest management practices. It highlights the key and cost-effective role that improved management practices could play in climate change mitigation.)**
  - 9.●● Walker WS, Gorelik SR, Cook-Patton SC, et al. The global potential for increased storage of carbon on land. *Proc Natl Acad Sci.* 2022;119:e2111312119. **(This paper modeled spatially explicit estimates of current carbon stocks, potential carbon storage (in reference and RCP8.5 climates), potential unrealized carbon storage in aboveground, belowground, and soil biomass. This paper also provides very recent estimates of the unrealized mitigation potential of improved forest management activities in boreal, temperate tropical, and global forests. It highlights the key mitigation role improved forest practices could play.)**
  10. Ellis PW, Gopalakrishna T, Goodman RC, et al. Reduced-impact logging for climate change mitigation (RIL-C) can halve selective logging emissions from tropical forests. *For Ecol Manage.* 2019;438:255–66.
  11. Finlayson C, Roopsind A, Griscom BW, Edwards DP, Freckleton RP. Removing climbers more than doubles tree growth and biomass in degraded tropical forests. *Ecol Evol.* 2022;12:e8758.
  12. Griscom BW, Busch J, Cook-Patton SC, et al. National mitigation potential from natural climate solutions in the tropics. *Phil Trans R Soc B: Biol Sci.* 2020;375:20190126.
  - 13.●● Kaarakka L, Cornett M, Domke G, Ontl T, Dee LE. Improved forest management as a natural climate solution: a review. *Ecol Solut Evid.* 2021;2:e12090. **(This study defines improved forest management and then provides a comprehensive review of improved forest management strategies, including thinning and extending rotation length, that could promote carbon sequestration rates in several carbon pools in managed forests.)**
  14. Mead DJ. Opportunities for improving plantation productivity. How much? How quickly? How realistic? *Biomass Bioenerg.* 2005;28:249–66.
  - 15.●● Ameray A, Bergeron Y, Valeria O, Montoro Girona M, Cavard X. Forest carbon management: a review of silvicultural practices and management strategies across boreal, temperate and tropical forests. *Curr Forestry Rep.* 2021;7:245–66. **(This paper reviews extensive and intensive forest management practices such as thinning, inorganic fertilization, and mixed plantations, used across tropical, temperate and boreal forests. It also explains their impacts on forest carbon fluxes and reservoirs.)**
  16. Forrester D, Bauhus J, Cowie A, Vanclay J. Mixed-species plantations of Eucalyptus with nitrogen-fixing trees: a review. *For Ecol Manage.* 2006;233:211–30.
  17. Piotto D. A meta-analysis comparing tree growth in monocultures and mixed plantations. *For Ecol Manage.* 2008;255:781–6.
  - 18.● Liu CLC, Kuchma O, Krutovsky KV. Mixed-species versus monocultures in plantation forestry: development, benefits, ecosystem services and perspectives for the future. *Glob Ecol Conserv.* 2018;15:e00419. **(This paper examines and compares the characteristics of pure and mixed plantations and the challenges associated with these two types of plantations. It also provides details on potential mechanisms explaining growth compatibility between different tree species.)**
  19. Nichols J, Rosemeyer M, Carpenter FL, Kettler J. Intercropping legume trees with native timber trees rapidly restores cover to eroded tropical pasture without fertilization. *For Ecol Manage.* 2001;152:195–209.
  20. Felton A, Lindbladh M, Brunet J, Fritz Ö. Replacing coniferous monocultures with mixed-species production stands: an assessment of the potential benefits for forest biodiversity in northern Europe. *For Ecol Manage.* 2010;260:939–47.
  21. Reverchon F, Bai SH, Liu X, Blumfield TJ. Tree Plantation Systems Influence Nitrogen Retention and the Abundance of Nitrogen Functional Genes in the Solomon Islands. *Front Microbiol.* 2015. <https://doi.org/10.3389/fmicb.2015.01439>
  22. Richards A, Forrester D, Bauhus J, Scherer-Lorenzen M. The influence of mixed tree plantations on the nutrition of individual species: a review. *Tree Physiol.* 2010;30:1192–208.
  23. Shrestha RK, Strahm BD, Sucre EB. Greenhouse gas emissions in response to nitrogen fertilization in managed forest ecosystems. *New Forest.* 2015;46:167–93.
  24. Ingerslev M, Hallbäck L. Above ground biomass and nutrient distribution in a limed and fertilized Norway spruce (*Picea abies*) plantation: Part II. Accumulation of biomass and nutrients. *For Ecol Manage.* 1999;119:21–38.
  25. Castaño-Díaz M, Barrio-Anta M, Afif-Khoury E, Cámara-Obregón A. Willow short rotation coppice trial in a former mining area in Northern Spain: effects of clone, fertilization and planting density on yield after five years. *Forests.* 2018;9:154.
  26. Hytönen J, Saarsalmi A. Biomass production of coppiced grey alder and the effect of fertilization. *Silva Fennica.* 2015;49:1–16.
  27. Batterman SA, Hedin LO, van Breugel M, Ransijn J, Craven DJ, Hall JS. Key role of symbiotic dinitrogen fixation in tropical forest secondary succession. *Nature.* 2013;502:224–7.
  28. Ewel JJ, Celis G, Schreeg L. Steeply increasing growth differential between mixture and monocultures of tropical trees. *Bio-tropica.* 2015;47:162–71.
  29. Huang Y, Chen Y, Castro-Izaguirre N, et al. Impacts of species richness on productivity in a large-scale subtropical forest experiment. *Sci.* 2018;362:80–3.
  30. Zhang Y, Chen HYH, Reich PB. Forest productivity increases with evenness, species richness and trait variation: a global meta-analysis. *J Ecol.* 2012;100:742–9.
  31. Elser JJ, Bracken MES, Cleland EE, Gruner DS, Harpole WS, Hillebrand H, Ngai JT, Seabloom EW, Shurin JB, Smith JE. Global analysis of nitrogen and phosphorus limitation of primary producers in freshwater, marine and terrestrial ecosystems. *Ecol Lett.* 2007;10:1135–42.
  32. Vadeboncoeur MA. Meta-analysis of fertilization experiments indicates multiple limiting nutrients in northeastern deciduous forests. *Can J For Res.* 2010;40:1766–80.
  33. Spake R, Yanou S, Yamaura Y, Kawamura K, Kitayama K, Doncaster C. Meta-analysis of management effects on biodiversity in plantation and secondary forests of Japan. *Conservation Science and Practice.* 2019;1:e14.
  34. Verschuyf J, Riffell S, Miller D, Wigley TB. Biodiversity response to intensive biomass production from forest thinning



- in North American forests – a meta-analysis. For *Ecol Manage.* 2011;261:221–32.
35. Gong C, Tan Q, Liu G, Xu M. Forest thinning increases soil carbon stocks in China. For *Ecol Manage.* 2021;482:118812.
  36. Boerner REJ, Huang J, Hart SC. Impacts of fire and fire surrogate treatments on forest soil properties: a meta-analytical approach. *J - Ecol Appl.* 2009;19:338–58.
  37. Holden SR, Treseder KK. A meta-analysis of soil microbial biomass responses to forest disturbances. *Front Microbiol.* 2013;4:163.
  38. Sohn JA, Saha S, Bauhus J. Potential of forest thinning to mitigate drought stress: a meta-analysis. For *Ecol Manage.* 2016;380:261–73.
  39. Lee J, Han S, Kim S, Lee S, Son Y, Son Y. A meta-analysis on the effect of forest thinning on diameter growth and carbon stocks in Korea. *J Korean For Soc.* 2015;104:527–35.
  40. Kalies EL, Haubensak KA, Finkral AJ. A meta-analysis of management effects on forest carbon storage. *J Sustain For.* 2016;35:311–23.
  41. Boerner REJ, Huang J, Hart SC. Fire, thinning, and the carbon economy: effects of fire and fire surrogate treatments on estimated carbon storage and sequestration rate. For *Ecol Manage.* 2008;255:3081–97.
  42. Hoover C, Stout S. The carbon consequences of thinning techniques: stand structure makes a difference. *J For.* 2007;105:266–70.
  43. Eriksson E. Thinning operations and their impact on biomass production in stands of Norway spruce and Scots pine. *Biomass Bioenerg.* 2006;30:848–54.
  44. Muñoz F, Rubilar R, Espinosa M, Cancino J, Toro J, Herrera M. The effect of pruning and thinning on above ground aerial biomass of *Eucalyptus nitens* (Deane & Maiden) Maiden. For *Ecol Manage.* 2008;255:365–73.
  45. ●● Bukoski JJ, Cook-Patton SC, Melikov C, Ban H, Chen JL, Goldman ED, Harris NL, Potts MD. Rates and drivers of aboveground carbon accumulation in global monoculture plantation forests. *Nat Commun.* 2022;13:4206. **(This paper provides a global monoculture carbon dataset that was used for this analysis and increases the scope and scale of current observations of carbon accumulation in global monoculture plantations by a factor of 60. It also presents estimates of the carbon accumulation potential of different species in pure tree plantations on a global scale. It also describes the main drivers of carbon accumulation in global monoculture forest plantations, such as past land use or tree species endemism. This global-scale analysis highlights key monoculture management considerations for carbon accumulation.)**
  46. ●● Feng Y, Schmid B, Loreau M, et al. Multispecies forest plantations outyield monocultures across a broad range of conditions. *Science.* 2022;376:865–8. **(This paper provides a global single-species and multispecies plantation carbon dataset that was used for this analysis. This study evaluates the impact of multi-species planting, specifically N-fixing species with non-N-fixing species, on stand height, DBH, and aboveground biomass. This study also highlights how the effect size of multispecies planting varies with stand characteristics such as age. This global scale analysis highlights key considerations for managing mixed plantations for carbon stocks.)**
  47. IPCC. Agriculture, forestry and other land use: forest Land. In: 2019 Refinement to the 2006 IPCC guidelines for national greenhouse gas inventories. 2019. vol. 4.
  48. Hedges LV, Gurevitch J, Curtis PS. The meta-analysis of response ratios in experimental ecology. *Ecology.* 1999;80:1150–6.
  49. Lajeunesse M. Recovering missing or partial data from studies: A survey of conversions and imputations for metaanalysis. In: *Handbook of meta-analysis in ecology and evolution.* Princeton University Press, Princeton, New Jersey, USA. 2013. pp 195–206.
  50. Koricheva J, Gurevitch J, Mengersen K. *Handbook of meta-analysis in ecology and evolution.* Princeton University Press; 2013.
  51. Sterne JAC, Sutton AJ, Ioannidis JPA, et al. Recommendations for examining and interpreting funnel plot asymmetry in meta-analyses of randomised controlled trials. *BMJ.* 2011;343:d4002.
  52. Viechtbauer W. Conducting meta-analyses in R with the metafor package. *J Stat Softw.* 2010;36:1–48.
  53. R Core Team. R: A language and environment for statistical computing. 2021.
  54. Baron RM, Kenny DA. The moderator-mediator variable distinction in social psychological research: conceptual, strategic, and statistical considerations. *J Pers Soc Psychol.* 1986;51:1173–82.
  55. USDA-NRCS, Soil Survey Division, World Soil Resources. Soil Moisture Regimes Map | NRCS Soils. 1997. <https://www.nature.com/scitable/content/global-map-of-soil-moisture-regimes-developed-59720131/>. Accessed 16 Feb 2022
  56. Haines SG, DeBell DS. Use of nitrogen-fixing plants to improve and maintain productivity of forest soils. In: *Impact of Intensive Harvesting on Forest Nutrient Cycling.* Syracuse, New York. 1979 pp. 279–302.
  57. Loreau M, Hector A. Partitioning selection and complementarity in biodiversity experiments. *Nature.* 2001;412:72–6.
  58. Oliveira N, del Río M, Forrester DI, Rodríguez-Soalleiro R, Pérez-Cruzado C, Cañellas I, Sixto H. Mixed short rotation plantations of *Populus alba* and *Robinia pseudoacacia* for biomass yield. For *Ecol Manage.* 2018;410:48–55.
  59. Williams LJ, Paquette A, Cavender-Bares J, Messier C, Reich PB. Spatial complementarity in tree crowns explains overyielding in species mixtures. *Nat Ecol Evol.* 2017;1:1–7.
  60. Khanna PK. Comparison of growth and nutrition of young monocultures and mixed stands of *Eucalyptus globulus* and *Acacia mearnsii*. For *Ecol Manage.* 1997;94:105–13.
  61. ●● Epihov DZ, Saltonstall K, Batterman SA, Hedin LO, Hall JS, van Breugel M, Leake JR, Beerling DJ. Legume–microbiome interactions unlock mineral nutrients in regrowing tropical forests. *Proc Natl Acad Sci.* 2021;118:e2022241118. **(This study explains how nitrogen-fixing trees promote mineral weathering of forest soils, which facilitates the release of available forms of nutrients such as nitrogen and phosphorus that can be taken up by non-nitrogen-fixing trees. This paper highlights the effects of nitrogen-fixing trees on the overall biogeochemical functioning of tropical forest ecosystems in a way that enhances their ability to assimilate and store atmospheric carbon.)**
  62. Binkley D, Sollins P, Bell R, Sachs D, Myrold D. Biogeochemistry of adjacent conifer and alder-conifer stands. *Ecology.* 1992;73:2022–33.
  63. Ghorbani M, Sohrabi H, Sadati E, Babaei F. Productivity and dynamics of pure and mixed-species plantations of *Populus deltoides* Bartr. ex Marsh and *Alnus subcordata* C A Mey. For *Ecol Manag.* 2018;409:890–8.
  64. Moore G, Bond B, Jones J. A comparison of annual transpiration and productivity in monoculture and mixed-species Douglas-fir and red alder stands. For *Ecol Manag.* 2011;262:2263–70.
  65. ●● Forrester DI. The spatial and temporal dynamics of species interactions in mixed-species forests: from pattern to process. For *Ecol Manag.* 2014;312:282–92. **(This paper reviews the spatial and temporal dynamics in mixed tree plantations, particularly between N-fixing and non-N-fixing species. It describes the different processes that drive these dynamics. It provides**

- key elements concerning the evolution of complementarity between N-fixing and non-fixing species as soil quality (fertility and water availability) decreases and as trees age.)**
66. le Maire G, Nouvellon Y, Christina M, Ponzoni F, Gonçalves J, Jean-Pierre B, Laclau J-P. Tree and stand light use efficiencies over a full rotation of single- and mixed-species *Eucalyptus grandis* and *Acacia mangium* plantations. *For Ecol Manage.* 2013;288:31–42.
  67. Balieiro FC, Fontes RLF, Dias LE, Franco AA, Campello EFC, de Faria SM. Accumulation and distribution of aboveground biomass and nutrients in pure and mixed stands of Guachapele and *Eucalyptus*. *J Plant Nutr.* 2002;25:2639–54.
  68. Danso SKA, Bowen GD, Sanginga N. Biological nitrogen fixation in trees in agro-ecosystems. *Plant Soil.* 1992;141:177–96.
  69. Tan ZX, Lal R, Wiebe KD. Global soil nutrient depletion and yield reduction. *J Sustain Agric.* 2005;26:123–46.
  70. Hytönen J, Saarsalmi A. Biomass production of coppiced grey alder and the effect of fertilization. *Silva Fennica.* 2015;49:1–16.
  71. Susanto D, Mulyati S, Purnomo H, Ruhayat D, Amirta R. Growth, biomass production and nutrient accumulation of *Macaranga gigantea* in response to NPK fertilizer application. *Nusantara Bioscience.* 2017;9:330–7.
  72. Ryan MG, Harmon ME, Birdsey RA, et al. A synthesis of the science on forests and carbon for U.S. Forests. *Ecological Society of America: Issues In Ecology.* 2010;13:1–16.
  73. McKinley DC, Ryan MG, Birdsey RA, et al. A synthesis of current knowledge on forests and carbon storage in the United States. *Ecol Appl.* 2011;21:1902–24.
  74. Trichet P, Bakker MR, Augusto L, Alazard P, Merzeau D, Saur E. Fifty years of fertilization experiments on *Pinus pinaster* in Southwest France: the importance of phosphorus as a fertilizer. *For Sci.* 2009;55:390–402.
  75. Carlson CA, Allan R, Soko S. Responses of *Eucalyptus* species to fertilizer applications made at planting on granitic parent materials in the Mpumalanga and Northern Provinces of South Africa. *South Afr For J.* 2001;191:29–37.
  76. Liu M, Xu F, Xu X, Wanek W, Yang X. Age alters uptake pattern of organic and inorganic nitrogen by rubber trees. *Tree Physiol.* 2018;38:1685–93.
  77. Genet H, Breda N, Dufrene E. Age-related variation in carbon allocation at tree and stand scales in beech and sessile oak Liebl using a chronosequence approach. *Tree Physiol.* 2010;30:2177–192.
  78. Lehmann J, Schroth G. Nutrient leaching. In: Schroth G, Sinclair FL, editors. *Trees, crops and soil fertility: concepts and research methods.* Wallingford: CABI; 2002. p. 151–66.
  79. Soil Survey Staff. *Illustrated guide to soil taxonomy, version 2.* US department of agriculture, national soil survey center. Lincoln. 2015.
  80. Hume AM, Chen HYH, Taylor AR. Intensive forest harvesting increases susceptibility of northern forest soils to carbon, nitrogen and phosphorus loss. *J Appl Ecol.* 2018;55:246–55.
  81. Bayle GK. Ecological and social impacts of eucalyptus tree plantation on the environment. *J Biodivers Conserv Bioresour Manag.* 2019;5:93–104.
  82. Bessaad A, Korboulewsky N. How much does leaf leaching matter during the pre-drying period in a whole-tree harvesting system? *For Ecol Manage.* 2020;477:118492.
  83. Walmsley J, Jones D, Reynolds B, Price M, Healey J. Whole tree harvesting can reduce second rotation forest productivity. *For Ecol Manage.* 2009;257:1104–11.
  84. Aber JD, Nadelhoffer KJ, Steudler P, Melillo JM. Nitrogen saturation in northern forest ecosystems. *Bioscience.* 1989;39:378–86.
  85. Saarsalmi A, Mälkönen E. Forest fertilization research in Finland: a literature review. *Scand J For Res.* 2001;16:514–35.
  86. Lucander K, Zanchi G, Akselsson C, Belyazid S. The effect of nitrogen fertilization on tree growth, soil organic carbon and nitrogen leaching—a modeling study in a steep nitrogen deposition gradient in Sweden. *Forests.* 2021;12:298.
  87. Lovett GM, Goodale CL. A new conceptual model of nitrogen saturation based on experimental nitrogen addition to an oak forest. *Ecosystems.* 2011;14:615–31.
  88. Högberg P, Johannisson C, Yarwood S, Callesen I, Näsholm T, Myrold DD, Högberg MN. Recovery of ectomycorrhiza after ‘nitrogen saturation’ of a conifer forest. *New Phytol.* 2011;189:515–25.
  89. Zhou D, Zhao SQ, Liu S, Oeding J. A meta-analysis on the impacts of partial cutting on forest structure and carbon storage. *Biogeosciences.* 2013;10:3691–703.
  90. Chase CW, Kimsey MJ, Shaw TM, Coleman MD. The response of light, water, and nutrient availability to pre-commercial thinning in dry inland Douglas-fir forests. *For Ecol Manage.* 2016;363:98–109.
  91. Lin J-C, Chiu C-M, Lin Y-J, Liu W. Thinning effects on biomass and carbon stock for Young Taiwan Plantations. *Sci Rep.* 2018. <https://doi.org/10.1038/s41598-018-21510-x>.
  92. Noulekoun F, Khamzina A, Naab J, Lamers J. Biomass allocation in five semi-arid afforestation species is driven mainly by ontogeny rather than resource availability. *Ann For Sci.* 2017. <https://doi.org/10.1007/s13595-017-0676-4>.
  93. Batey T. Soil compaction and soil management – a review. *Soil Use Manag.* 2009;25:335–45.
  94. Skovsgaard JP. Analysing effects of thinning on stand volume growth in relation to site conditions: a case study for even-aged Sitka spruce (*Picea sitchensis* (Bong.) Carr.). *For: Int J For Res.* 2009;82:87–104.
  95. Malhi Y, Meir P, Brown S. Forests, carbon and global climate. *Philos Transact Ser A Math Phys Eng Sci.* 2002;360:1567–91.
  96. Mokany K, Raison RJ, Prokushkin AS. Critical analysis of root: shoot ratios in terrestrial biomes. *Glob Change Biol.* 2006;12:84–96.
  97. Oldfield EE, Eagle AJ, Rubin RL, Rudek J, Sanderman J, Gordon DR. Crediting agricultural soil carbon sequestration. *Science.* 2022;375:1222–5.
  98. Forrester D, Pares A, O’Hara C, Khanna P, Bauhus J. Soil organic carbon is increased in mixed-species plantations of *Eucalyptus* and nitrogen-fixing acacia. *Ecosystems.* 2013;16:123–32.
  99. Voigtlaender M, Brandani CB, Caldeira DRM, et al. Nitrogen cycling in monospecific and mixed-species plantations of *Acacia mangium* and *Eucalyptus* at 4 sites in Brazil. *For Ecol Manage.* 2019;436:56.
  100. Zhang X, Guan D, Li W, Sun D, Jin C, Yuan F, Wang A, Wu J. The effects of forest thinning on soil carbon stocks and dynamics: a meta-analysis. *For Ecol Manage.* 2018;429:36–43.
  101. Ball BC, McTaggart IP, Scott A. Mitigation of greenhouse gas emissions from soil under silage production by use of organic manures or slow-release fertilizer. *Soil Use Manag.* 2004;20:287–95.
  102. Kaul M, Mohren GMJ, Dadhwal V. Carbon storage and sequestration potential of selected tree species in India. *Mitig Adapt Strat Glob Change.* 2010;15:489–510.
  103. D’Amato AW, Woodall CW, Weiskittel AR, Littlefield CE, Murray LT. Carbon conundrums: do United States’ current carbon market baselines represent an undesirable ecological threshold? *Glob Change Biol.* 2022. <https://doi.org/10.1111/gcb.16215>.
  104. Messier C, Bauhus J, Sousa-Silva R, et al. For the sake of resilience and multifunctionality, let’s diversify planted forests! *Conserv Lett.* 2022;15:e12829.
  105. Paquette A, Messier C. The effect of biodiversity on tree productivity: from temperate to boreal forests. *Glob Ecol Biogeogr.* 2011;20:170–80.

106. • Osuri AM, Gopal A, Raman TRS, DeFries R, Cook-Patton SC, Naeem S. Greater stability of carbon capture in species-rich natural forests compared to species-poor plantations. *Environ Res Lett.* 2020;15:034011. **(This study provides evidence of greater stability of carbon stocks in species-rich forests than in species-poor tree plantations. This study highlights**

**the need to diversify current monoculture tree plantations to make them more resilient and resistant to an uncertain future.)**

**Publisher's Note** Springer Nature remains neutral with regard to jurisdictional claims in published maps and institutional affiliations.