

UCLA

UCLA Previously Published Works

Title

Low-Intensity Walking Activity Is Associated With Better Health

Permalink

<https://escholarship.org/uc/item/8tp6g5xm>

Journal

Journal of Applied Gerontology, 33(7)

ISSN

0733-4648

Authors

Varma, Vijay R
Tan, Erwin J
Wang, Tao
[et al.](#)

Publication Date

2014-10-01

DOI

10.1177/0733464813512896

Peer reviewed



Published in final edited form as:

J Appl Gerontol. 2014 October ; 33(7): 870–887. doi:10.1177/0733464813512896.

Low-Intensity Walking Activity is Associated with Better Health

Vijay R. Varma, MPH^{1,2}, Erwin J. Tan, MD³, Tao Wang, PhD¹, Qian-Li Xue, PhD¹, Linda P. Fried, MD⁴, Christopher L. Seplaki, PhD⁵, Abby C. King, PhD⁶, Teresa E. Seeman, PhD⁷, George W. Rebok, PhD^{1,2}, and Michelle C. Carlson, PhD^{1,2}

¹Johns Hopkins Center on Aging and Health, Baltimore, MD

²Johns Hopkins, Bloomberg School of Public Health, Department of Mental Health, Baltimore, MD

³The Corporation for National and Community Service, Washington D.C.

⁴Mailman School of Public Health, Columbia University, New York, NY

⁵University of Rochester School of Medicine and Dentistry, Rochester, NY

⁶Stanford University School of Medicine, Palo Alto, CA

⁷School of Medicine, University of California, Los Angeles, CA

Abstract

Recommended levels of physical activity may represent challenging targets for many older adults at risk for disability, leading to the importance of evaluating whether low-intensity activity is associated with health benefits. We examined the cross-sectional association between low-intensity walking activity (<100 steps/minute) and health and physical function in a group of older adults. Participants (n=187; age: 66.8; 91.4% African-American; 76.5% female) wore a StepWatch Activity Monitor to measure components of low-intensity walking activity. Only 7% of participants met physical activity guidelines and moderate-intensity activity (>100 steps/min) contributed only 10% of total steps/day and two percent of total minutes/day. Greater amount, frequency, and duration of low-intensity activity were associated with better self-report and performance-based measures of physical function, better quality of life, and fewer depressive symptoms (ps<0.05). The cross-sectional relationship between low-intensity activity and health outcomes important to independent function suggest that we further explore the longitudinal benefits of low-intensity activity.

Keywords

mobility; physical activity; African Americans; low-intensity activity; walking

Regular physical activity is a Healthy People 2020 objective for all age groups (*Healthy People 2020*, 2010). More active compared to less active older adults, have higher levels of cardiorespiratory and muscular fitness, and higher levels of functional health (Physical

Activity Guidelines Advisory Committee Report, 2008). In order to achieve health benefits, current activity guidelines recommend at least 30 minutes of moderate-intensity aerobic activity five days a week or 20 minutes of vigorous-intensity aerobic activity three days a week (“How much physical activity do older adults need?,” 2011; Nelson et al., 2007; Physical Activity Guidelines Advisory Committee Report, 2008).

According to recent CDC surveillance data, less than 16 percent of older adults meet activity guidelines, and over 30 percent are inactive (report no leisure-time physical activity) (Macera et al., 2005). Evidence suggests that many older adults have difficulty initiating and adhering to exercise programs (Boyette, Sharon, & Brandon, 1997; Clark, 1999; B Resnick & Spellbring, 2000; Schutzer & Graves, 2004). This is of particular concern for older adults of low socioeconomic status (SES) who have lower baseline physical activity levels than higher SES older adults (“Facts about Physical Activity,” 2012; Marshall et al., 2007; Physical Activity Guidelines Advisory Committee Report, 2008), and fewer physical activity-related facilities due to restrictive environmental and neighborhood characteristics (Day, 2006; Parra-Medina et al.,; Powell, Slater, Chaloupka, & Harper, 2006).

While the current physical activity guidelines focus on moderate-intensity to vigorous physical activity, some studies suggest that low-intensity activity, which may include non-exercise, leisure-time activities (walking for pleasure), instrumental activities of daily living (e.g. walking related to housework or shopping), as well as low-intensity exercise may also yield health benefits (Blumenthal et al., 1988; Demakakos, Hamer, Stamatakis, & Steptoe, 2010; Hakim et al., 1998; Hu, Li, Colditz, Willett, & Manson, 2003; Pescatello, Murphey, & Costanzo, 2000; Physical Activity Guidelines Advisory Committee Report, 2008; Rippe, Ward, Porcari, & Freedson, 1988; Sattelmair et al., 2011). However, there is currently limited evidence for the relationship between low-intensity activity and health using objective physical activity measures (Duvivier et al., 2013; Lee et al., 2012).

The most recent U.S. Department of Health and Human Services Physical Activity Guidelines recommended additional observational and experimental studies to evaluate the nature of the benefits of low-intensity activity, particularly within understudied populations (Physical Activity Guidelines Advisory Committee Report, 2008). We therefore described walking activity in an older cohort of urban-dwelling adults at elevated risk for functional decline and disability due to age, low income, low education, and high number of chronic diseases (Freedman, Martin, & Schoeni, 2002; Seeman, Merkin, Crimmins, & Karlamangla, 2010), and examined components of low-intensity walking activity in daily life, including amount, duration, and frequency (Howley, 2001; Kesaniemi et al., 2001), and their cross-sectional associations with physical, mental, and cognitive health.

We report on baseline daily activity measured using an objective step activity device collected as an ancillary study within the Baltimore Experience Corps Trial (BECT). We hypothesized that: 1) the majority of participants would not meet activity guidelines, having minimal moderate-intensity walking activity; 2) greater low-intensity walking activity would be associated with better health.

Methods

Participants

The BECT is a randomized, controlled trial funded by the National Institute on Aging to evaluate the health benefits for older adults participating in Experience Corps Baltimore vs. a control group offered other low-service volunteer opportunities. Experience Corps is a high-intensity senior service program designed to place volunteers in elementary schools to improve the academic success of young children. Study design, sampling methodology, and recruitment have been described previously (Fried et al., 2004; Fried et al., 2013). Enrollment criteria included: aged ≥ 60 years, ≥ 24 on the Mini-Mental State Exam (MMSE) (Folstein, Folstein, & McHugh, 1975), and ability to read at a minimum 6th grade level (Wilkinson, 1993). Participants agreed to serve 15 or more hours/week for a full school year if randomized to EC. There were no exclusion criteria related to physical activity or physical function, including BMI, chronic diseases, or disability. A subset of participants consenting to the BECT ($n=195$) were simultaneously offered the SAM. All baseline measurements were taken prior to randomization to ensure that the intervention had no effects on the results. The study protocol was approved by the Johns Hopkins School of Medicine Institutional Review Board and each participant provided written, informed consent.

SAM Protocol

The SAM is a microprocessor-linked sensor that is worn on the dominant ankle and continuously measures step activity in daily life. The device measures the number of steps at one-minute intervals using acceleration, position, and timing information, and can therefore be used to differentiate between low-intensity and moderate to vigorous-intensity walking activity as well as characterize the amount, duration and frequency of daily walking activity. The SAM has been validated across a range of community dwelling older adult samples with varying functional limitations using self-report and objective measures (e.g. hand-tallied step counts and accelerometers) (Cavanaugh, Coleman, Gaines, Laing, & Morey, 2007; B. Resnick, Nahm, Orwig, Zimmerman, & Magaziner, 2001; Storti et al., 2008). The SAM is particularly sensitive in measuring activity at decreased gait speeds (Storti et al., 2008), and is well tolerated by older adults because it is small and placed on the ankle vs. the hip (Algase, Beattie, Leitsch, & Beel-Bates, 2003). The SAM, which is enclosed within a lightweight, pager-sized plastic case, was attached to the participant's right ankle and calibrated at the baseline evaluation by comparing hand-tallied step counts to one minute of walking activity at the participant's normal gait. We instructed participants to wear the SAM for three to seven days (excluding the first half day of data collection at the study visit when the SAM was placed on the participant's ankle) while keeping a wear time/activity diary at approximately one-hour intervals throughout these days. Participants were instructed to remove the SAM only when bathing, showering or swimming, and to replace the device immediately after. At the end of this period, participants were asked to mail back, in a pre-stamped, addressed envelope, the SAM, activity diary, and the Pittsburgh Sleep Quality Index (PSQI), which recorded wake and sleep times "during the past month" (Buysse, Reynolds, Monk, Berman, & Kupfer, 1989). The majority of participants wore the SAM during late summer and early fall, which reduced the influence of seasonal effects.

The SAM recorded stride counts within each 24-hour period producing 1440, one-minute intervals. We doubled stride counts to reflect step counts for both legs. If diary information was not available, we determined wake and sleep times using the PSQI, or imputed using the group's average wake and sleep times over available days.

We excluded days representing non-compliance according to the following four criteria: 1) less than 201 total steps/day; 2) days with less than a total of six hours of any activity between wake and sleep times; 3) days with six consecutive hours of inactivity (< 1 step) between wake and sleep times (Masse et al., 2005); and 4) days when participants recorded not following data collection protocol in their activity diaries. We conservatively elected not to identify and impute missing data (non-wear time) for days included in the analysis (Esliger, Copeland, Barnes, & Tremblay, 2005).

Of the 702 BECT participants, a subset of 195 were offered the SAM at baseline. Participants provided an average of 4.9 days of data. Based on the exclusion protocol detailed above, an average of 0.8 days (16.4% of data collected) were removed from analysis. Data for eight participants were excluded due to non-compliance, and a total of 187 participants comprised the final sample providing an average of 4.3 days (range: 1-9) each.

SAM Metrics

In order to characterize the proportion of participants meeting physical activity guidelines within our sample, we used the threshold of 10,000 steps/day developed in previous studies as a reasonable equivalent of U.S. physical activity guidelines (Tudor-Locke & Bassett, 2004; Tudor-Locke, Hatano, Pangrazi, & Kang, 2008). We classified participants who walked an average of < 10,000 steps/day as less active and ≥ 10,000 steps/day as active. Based on previous studies translating physical activity recommendations (30 minutes of moderate-intensity activity/day that can be split into three, 10-minute bouts/day) into a pedometer-based step goal (three bouts/day of 1000 steps in 10 minutes) (Marshall et al., 2009; Tudor-Locke, Sisson, Collova, Lee, & Swan, 2005), we additionally classified participants who completed < three bouts/day on average as less active, and ≥ three bouts/day as active.

Intensity ranges (effort associated with walking) included low-intensity (> 0 steps/min and < 100 steps/min) and moderate to vigorous-intensity (≥ 100 steps/min) based on studies translating laboratory measurements of oxygen consumption while walking into pedometer-based metrics (Marshall et al., 2009; Tudor-Locke et al., 2005). Metrics representing components of activity within intensity ranges are described below and summarized in Figure 1. All analyses (Table 2) were restricted to low-intensity activity (> 0 steps/min and < 100 steps/min), and excluded activity at moderate to vigorous-intensity (≥ 100 steps/min). All metrics were averaged across all valid days surveyed.

Activity amount was defined as the number of steps/day at low-intensity, and included total steps/day at > 0 and < 100 steps/min; steps at ≥ 100 steps/min were excluded. We included total number of steps/day across intensities and the number of steps/day at moderate to vigorous-intensity (≥ 100 steps/min) for comparison. Activity duration was defined as the number of minutes/day of low-intensity activity, and included total minutes/day at > 0 and

<100 steps/min; minutes at 100 steps/min were excluded. We included the number of minutes/day of moderate to vigorous-intensity activity (≥ 100 steps/min) for comparison. Activity frequency was defined as the number of bouts/day of 10-minute, low-intensity activity, and calculated by adding the number of times participants completed 10 consecutive minutes of low-intensity activity (> 0 steps/min and < 100 steps/min). This metric was based on U.S. Department of Health and Human Services physical activity recommendations stating that 30-minute bouts of activity can be obtained through three, 10-minute bouts that can be accumulated over the course of the day (Nelson et al., 2007; Physical Activity Guidelines Advisory Committee Report, 2008). We additionally calculated the number of bouts/day of 30-minute low-intensity activity, however this represented a low frequency event in our sample (mean = .56 bouts/ day, SD: .95).

Health Measures

We examined the association between low-intensity walking activity and baseline mobility, performance-based lower extremity function, depressive symptoms, quality of life, and global cognition.

Self-reported mobility measures included 1) difficulty walking one mile or several blocks; and 2) difficulty going up and down a flight of stairs (indoors) (Fried et al., 1991; Weiss, Fried, & Bandeen-Roche, 2007). Participants self-reported 'None', 'A little', 'Some', 'Quite a lot', or 'Cannot do' for each measure. Similar to previous studies, both mobility measures were coded as binary indicators so that odds ratios indicated any self-reported limitation ('A little', 'Some', 'Quite a lot', 'Cannot do') versus no limitation ('None'). Performance-based lower extremity function was measured using the Short Physical Performance Battery (SPPB). The SPPB assessed walking (timed four meter walk at the participant's usual pace), tests of balance (ability to stand in the side-by-side, semi-tandem, and tandem positions for 10 seconds), and strength (time to rise from a chair five times) (Guralnik et al., 2000). For each test, participants received a score of 0-4; zero indicated an inability to complete the test and 4 indicated the highest level of performance. Scores for all three performance-based measures were summed to create a total score ranging from zero to 12, with higher scores reflecting better lower-extremity function (Guralnik et al., 2000). Depressive symptoms were measured using the Geriatric Depression Scale (GDS) short form, a 15-item list developed specifically for use in older adult populations (higher scores represented more depressive symptoms) (Yesavage et al., 1982). Quality of life was measured by summing participants' self-report of any problems with mobility, self-care, usual activities, pain/discomfort, and anxiety/depression from the EuroQol questionnaire ("EuroQol--a new facility for the measurement of health-related quality of life. The EuroQol Group," 1990). Total scores ranged from zero to five with higher scores reflecting lower quality of life. Global cognition was measured using the MMSE (Folstein et al., 1975), a global test of cognitive function with scores ranging from 24-30 at baseline. All models were adjusted for confounders: age at baseline, gender and race.

Statistical Analysis

We first compared baseline socio-demographic and health characteristics between participants offered the SAMs and those in the larger BECT using chi-square tests for

dichotomous variables and t-tests for continuous variables (Table 1). Next, we confirmed that all three metrics of low-intensity activity were normally distributed. We additionally examined the Pearson correlations among low-intensity metrics.

We then examined the associations between the three metrics of low-intensity activity and health measures (Table 2). All analyses were restricted to low-intensity activity metrics, and excluded moderate or higher intensity activity. Logistic regression models were used to analyze the two binary health measures; model results were expressed as a unit increase in SAM metrics associated with an increased/ decreased odds of reporting “Difficulty with walking one mile or several blocks”, and “Difficulty with one flight of steps.” Linear regression models were used to analyze the four continuous health measures; model results were expressed as a unit increase in SAM metrics associated with an increase/ decrease in “Lower-extremity function (SPPB)”, “Depressive Symptoms (GDS)”, “Quality of Life”, and “Global Cognition (MMSE).” Estimated effect sizes of steps/day (amount) and minutes of activity (duration) on health measures were specifically expressed in units of 1000 steps/day and 10 minutes/day respectively based on previous studies as well as ease of interpretation (Sisson et al.). All statistical tests were two-sided and performed using Stata version 12 (StataCorp. 2011. Stata Statistical Software: Release 12. College Station, TX: StataCorp LP.). Regression diagnostics for outliers, normality of residuals, and checks for multicollinearity were performed by visual inspection of residual plots, and computation of variance inflation factors. Tobit regression and robust standard errors were used for all regression models where health measures had censoring or skew.

Results

Table 1 presents baseline characteristics of the study sample. Participants averaged 66.8 years, were predominately female (76.5%) and African-American (91.4%), with 36.9% reporting high school or less education, and 28.0% reporting household income less than \$15,000. Rates of chronic disease were high; 58.3% of participants were obese, 71.2% reported hypertension, 61.8% reported osteoarthritis, and 32.6% reported diabetes. Participants offered the SAMs did not vary significantly ($p < .05$) from those in the larger BECT on any of the above socio-demographic and health characteristics at baseline.

Considering physical activity guidelines (10,000 steps/day) 7.0% ($n=16$) were considered active and did meet guidelines. Based on Department of Health and Human Services guidelines of 30 minutes of moderate intensity activity/day, we found that 1.1% ($n=2$) met guidelines by completing three or more 10-minute moderate-intensity bouts/day. Participants completed a total of 7648.8 steps/day; the majority of walking activity was in the low-intensity range (6888.1 steps/day; 90.1% of total daily steps) with minimal activity in the moderate to vigorous-intensity range (760.7 steps/day; 9.9% of total daily steps). Participants additionally had, on average, only 6.9 minutes/day of moderate to vigorous-intensity activity (2.1% of total daily minutes of activity) with the majority of activity in the low-intensity range (321.9 minutes/day; 97.9% of total daily minutes of activity) (Figure 2). Participants averaged 10.3 bouts/day of 10-minute low-intensity activity. As expected, SAM metrics were highly correlated ($r = 0.87-0.88$; all $p < .001$).

Table 2 presents the associations between low-intensity activity metrics and health measures, after adjusting for age, gender and race. A 1000-step increase in steps/day at low-intensity was significantly associated with better physical function measures, including a 14% reduction in the odds of reporting difficulty walking one mile (OR=0.86, 95% CI: 0.77, 0.97); a 16% reduction in the odds of reporting difficulty with one flight of steps (OR=0.84, 95% CI: 0.73, 0.98); and a 0.18-point increase in SPPB (95% CI: 0.07, 0.28). Each additional bout of 10-minute activity was also associated with better physical function, including a 6% reduction in the odds of reporting difficulty walking one mile (OR=0.96, 95% CI: 0.93, 1.00); a 7% reduction in the odds of reporting difficulty with one flight of steps (OR=0.93, 95% CI: 0.87, 0.98); and a 0.08 point increase in SPPB (95% CI: 0.03, 0.13). A 10 minute increase in low-intensity minutes/day was associated with a 0.05 point increase in SPPB (95% CI: 0.02, 0.08). All low-intensity activity metrics were significantly associated with fewer depressive symptoms. Specifically, for each low-intensity measure, a 1000-step increase in steps/day, a 10 minute increase in low-intensity minutes/day, and an additional bout of 10-minute activity were each associated with, respectively, 0.09 (95% CI: -0.17, -0.01); .02 (95% CI: -0.05, -0.00); and .04 (95% CI: -0.07, -0.00) fewer depressive symptoms. These associations were significant despite the generally low levels of reported depressive symptoms (mean: 3.1; range: 2-10). For quality of life, a 1000-step increase in steps/day and an additional bout of 10-minute, low-intensity activity were also significantly associated with 0.07 (95% CI: -0.12, -0.02) and 0.03 (95% CI: -0.06, -0.01) points better, respectively. However, the MMSE was not associated with low-intensity activity although model estimates indicated a consistent trend between greater low-intensity activity and higher MMSE. MMSE scores were highly skewed (mean: 28.2; range: 24-30) due to trial enrollment criteria.

Discussion

In a community-based cohort of urban-dwelling, older adults, we observed that only seven percent met physical activity guidelines, with the majority of activity falling within the low intensity compared to the moderate to vigorous-intensity ranges. In cross-section, greater low-intensity walking activity was associated with better physical function and health, including better quality of life and fewer depressive symptoms. These findings provide evidence for the value of objectively measuring rates of low-intensity step activity in daily life, a range of activity that is much more common than moderate-intensity activity and may serve a target to maintain and promote health.

Moderate-intensity walking activity comprised only 10 percent of total steps/day and two percent of total minutes/day. By comparison, low-intensity activity was highly prevalent, possibly because it relates to routine daily functional and social activities. For example, activities related to low-intensity walking activity within our sample can include care-taking of grandchildren, walking to catch a bus, and church-related social activities. The step activity associated with these activities in the aggregate may represent health promoting behaviors that should be quantified particularly in minority older adults at elevated risk for functional decline and disability. The SAM was designed to precisely measure walking activity in older adult populations with varying levels of function (B. Resnick et al., 2001; Storti et al., 2008). Furthermore, it can be used to identify potentially meaningful metrics of

daily walking activity beyond total steps/day, including amount, duration and frequency within the low-intensity range, which may serve as targets in future designs of achievable and sustainable activity interventions in otherwise sedentary individuals.

One limitation of this study was its cross-sectional design. Given the BECT pre-post trial design, we will be able to further explore whether increasing levels of low-intensity activity may protect against subsequent functional, mobility, mental, and cognitive health declines. Another potential limitation was the placement of the SAM device on the ankle vs. hip. While this placement allows the SAM to detect steps in participants with altered gait, we did not capture non-ambulatory movement associated with trunk and upper body activities, a component of total daily physical activity. Also, physical activity guidelines recommend activity five days a week; averaging activity across all compliant days provided by participants to determine whether they met guidelines may have underestimated the proportion of participants who were physically active. Additionally, for those with very low levels of activity, exclusion criteria removing days with less than six hours of activity may have resulted in an underestimation of individuals who were very sedentary. However, the effect of any bias was likely minimal given the extremely low overall rates of moderate to vigorous-intensity activity. Finally, while the study sample represented an understudied and at-risk segment of the older adult population, a trial of high-intensity volunteer service may select for the more health conscious and physically active members of the community (Boughner, 2010; Lindsted, Fraser, Steinkohl, & Beeson, 1996). Therefore, study results may represent a conservative estimate of walking activity and health relative to the larger older adult population.

Conclusion

Measuring daily walking activity in a largely sedentary and at-risk older adult population demonstrated the importance of low-intensity levels of walking activity to numerous health measures, including physical function, mental health, and quality of life. These positive associations highlight the need to better characterize and understand how increasing daily levels of low-intensity walking in various ways may be associated with the prevention of disability and enhanced quality of life in the aging population.


References

- Algate DL, Beattie ER, Leitsch SA, Beel-Bates CA. Biomechanical activity devices to index wandering behavior in dementia. *Am J Alzheimers Dis Other Demen.* 2003; 18(2):85–92. [PubMed: 12708223]
- Blumenthal JA, Rejeski WJ, Walsh-Riddle M, Emery CF, Miller H, Roark S, et al. Comparison of high- and low-intensity exercise training early after acute myocardial infarction. *Am J Cardiol.* 1988; 61(1):26–30. [PubMed: 3337013]
- Boughner, RL., editor. *Encyclopedia of Research Design.* SAGE Publications; 2010.
- Boyette LW, Sharon BF, Brandon LJ. Exercise adherence for a strength training program in older adults. *J Nutr Health Aging.* 1997; 1(2):93–97. [PubMed: 16491532]
- Buysse DJ, Reynolds CF 3rd, Monk TH, Berman SR, Kupfer DJ. The Pittsburgh Sleep Quality Index: a new instrument for psychiatric practice and research. *Psychiatry Res.* 1989; 28(2):193–213. [PubMed: 2748771]

- Cavanaugh JT, Coleman KL, Gaines JM, Laing L, Morey MC. Using step activity monitoring to characterize ambulatory activity in community-dwelling older adults. *J Am Geriatr Soc.* 2007; 55(1):120–124. [PubMed: 17233695]
- Clark DO. Physical activity and its correlates among urban primary care patients aged 55 years or older. *J Gerontol B Psychol Sci Soc Sci.* 1999; 54(1):S41–48. [PubMed: 9934401]
- Day K. Active living and social justice: planning for physical activity in low-income, Black, and Latino communities. *Journal of the American Planning Association.* 2006; 72(1):88–99.
- Demakakos P, Hamer M, Stamatakis E, Steptoe A. Low-intensity physical activity is associated with reduced risk of incident type 2 diabetes in older adults: evidence from the English Longitudinal Study of Ageing. *Diabetologia.* 2010; 54(9):1877–1885. [PubMed: 20495973]
- Duvivier BM, Schaper NC, Bremers MA, van Crombrugge G, Menheere PP, Kars M, et al. Minimal intensity physical activity (standing and walking) of longer duration improves insulin action and plasma lipids more than shorter periods of moderate to vigorous exercise (cycling) in sedentary subjects when energy expenditure is comparable. *PLoS One.* 2013; 8(2):e55542. [PubMed: 23418444]
- Esliger DW, Copeland JL, Barnes JD, Tremblay MS. Standardizing and optimizing the use of accelerometer data for free-living physical activity monitoring. *Journal of Physical Activity and Health.* 2005; 2(3):366–383.
- The EuroQol Group. EuroQol--a new facility for the measurement of health-related quality of life. *Health Policy.* 1990; 16(3):199–208. [PubMed: 10109801]
- Facts about Physical Activity. Physical Activity. Aug 8. 2012 2013 Retrieved 2013, from <http://www.cdc.gov/physicalactivity/data/facts.html>
- Folstein MF, Folstein SE, McHugh PR. "Mini-mental state". A practical method for grading the cognitive state of patients for the clinician. *J Psychiatr Res.* 1975; 12(3):189–198. [PubMed: 1202204]
- Freedman VA, Martin LG, Schoeni RF. Recent Trends in Disability and Functioning Among Older Adults in the United States. *JAMA: The Journal of the American Medical Association.* 2002; 288(24):3137–3146. [PubMed: 12495394]
- Fried LP, Borhani NO, Enright P, Furberg CD, Gardin JM, Kronmal RA, et al. The Cardiovascular Health Study: design and rationale. *Ann Epidemiol.* 1991; 1(3):263–276. [PubMed: 1669507]
- Fried LP, Carlson MC, Freedman M, Frick KD, Glass TA, Hill J, et al. A social model for health promotion for an aging population: initial evidence on the Experience Corps model. *J Urban Health.* 2004; 81(1):64–78. [PubMed: 15047786]
- Fried LP, Carlson MC, McGill S, Seeman T, Xue QL, Frick K, et al. Experience Corps: A dual trial to promote the health of older adults and children's academic success. *Contemp Clin Trials.* 2013
- Guralnik JM, Ferrucci L, Pieper CF, Leveille SG, Markides KS, Ostir GV, et al. Lower extremity function and subsequent disability: consistency across studies, predictive models, and value of gait speed alone compared with the short physical performance battery. *J Gerontol A Biol Sci Med Sci.* 2000; 55(4):M221–231. [PubMed: 10811152]
- Hakim AA, Petrovitch H, Burchfiel CM, Ross GW, Rodriguez BL, White LR, et al. Effects of walking on mortality among nonsmoking retired men. *N Engl J Med.* 1998; 338(2):94–99. [PubMed: 9420340]
- Healthy People 2020. Office of Disease Prevention and Health Promotion. United States Department of Health and Human Services; Washington, DC: 2010.
- How much physical activity do older adults need?. Jan 12. 2011 Retrieved 12/1/2012, 2012, from <http://www.cdc.gov/physicalactivity/everyone/guidelines/olderadults.html>
- Howley ET. Type of activity: resistance, aerobic and leisure versus occupational physical activity. *Med Sci Sports Exerc.* 2001; 33(6 Suppl):S364–369. discussion S419–320. [PubMed: 11427761]
- Hu FB, Li TY, Colditz GA, Willett WC, Manson JE. Television watching and other sedentary behaviors in relation to risk of obesity and type 2 diabetes mellitus in women. *Jama.* 2003; 289(14):1785–1791. [PubMed: 12684356]
- Kesaniemi YK, Danforth E Jr, Jensen MD, Kopelman PG, Lefebvre P, Reeder BA. Dose-response issues concerning physical activity and health: an evidence-based symposium. *Med Sci Sports Exerc.* 2001; 33(6 Suppl):S351–358. [PubMed: 11427759]

- Lee PH, Nan H, Yu YY, McDowell I, Leung GM, Lam TH. For non-exercising people, the number of steps walked is more strongly associated with health than time spent walking. *J Sci Med Sport*. 2012
- Lindstedt KD, Fraser GE, Steinkohl M, Beeson WL. Healthy volunteer effect in a cohort study: temporal resolution in the Adventist Health Study. *J Clin Epidemiol*. 1996; 49(7):783–790. [PubMed: 8691229]
- Macera CA, Ham SA, Yore MM, Jones DA, Ainsworth BE, Kimsey CD, et al. Prevalence of physical activity in the United States: Behavioral Risk Factor Surveillance System. *Prev Chronic Dis*. 2005; 2(2):A17. [PubMed: 15888228]
- Marshall SJ, Jones DA, Ainsworth BE, Reis JP, Levy SS, Macera CA. Race/ethnicity, social class, and leisure-time physical inactivity. *Med Sci Sports Exerc*. 2007; 39(1):44–51. [PubMed: 17218883]
- Marshall SJ, Levy SS, Tudor-Locke CE, Kolkhorst FW, Wooten KM, Ji M, et al. Translating physical activity recommendations into a pedometer-based step goal: 3000 steps in 30 minutes. *Am J Prev Med*. 2009; 36(5):410–415. [PubMed: 19362695]
- Masse LC, Fuemmeler BF, Anderson CB, Matthews CE, Trost SG, Catellier DJ, et al. Accelerometer data reduction: a comparison of four reduction algorithms on select outcome variables. *Med Sci Sports Exerc*. 2005; 37(11 Suppl):S544–554. [PubMed: 16294117]
- Nelson ME, Rejeski WJ, Blair SN, Duncan PW, Judge JO, King AC, et al. Physical activity and public health in older adults: recommendation from the American College of Sports Medicine and the American Heart Association. *Circulation*. 2007; 116(9):1094–1105. [PubMed: 17671236]
- Parra-Medina D, Wilcox S, Wilson DK, Addy CL, Felton G, Poston MB. Heart Healthy and Ethnically Relevant (HHER) Lifestyle trial for improving diet and physical activity in underserved African American women. *Contemp Clin Trials*. 2010; 31(1):92–104. [PubMed: 19781665]
- Pescatello LS, Murphey D, Costanzo D. Low-intensity physical activity benefits blood lipids and lipoproteins in older adults living at home. *Age and Aging*. 2000; 29:433–439.
- Physical Activity Guidelines Advisory Committee Report. United States Department of Health and Human services; Washington D.C.: 2008.
- Powell LM, Slater S, Chaloupka FJ, Harper D. Availability of physical activity-related facilities and neighborhood demographic and socioeconomic characteristics: a national study. *Am J Public Health*. 2006; 96(9):1676–1680. [PubMed: 16873753]
- Resnick B, Nahm ES, Orwig D, Zimmerman SS, Magaziner J. Measurement of activity in older adults: reliability and validity of the Step Activity Monitor. *J Nurs Meas*. 2001; 9(3):275–290. [PubMed: 11881269]
- Resnick B, Spellbring A. The factors that influence exercise behavior in older adults. *Journal of Gerontological Nursing*. 2000; 26:34–42. [PubMed: 11111629]
- Rippe JM, Ward A, Porcari JP, Freedson PS. Walking for health and fitness. *Jama*. 1988; 259(18):2720–2724. [PubMed: 3282085]
- Sattelmair J, Pertman J, Ding EL, Kohl HW 3rd, Haskell W, Lee IM. Dose response between physical activity and risk of coronary heart disease: a meta-analysis. *Circulation*. 2011; 124(7):789–795. [PubMed: 21810663]
- Schutzer KA, Graves BS. Barriers and motivations to exercise in older adults. *Prev Med*. 2004; 39(5):1056–1061. [PubMed: 15475041]
- Seeman TE, Merkin SS, Crimmins EM, Karlamangla AS. Disability Trends Among Older Americans: National Health and Nutrition Examination Surveys, 1988-1994 and 1999-2004. *American Journal of Public Health*. 2010; 100(1):100–107. [PubMed: 19910350]
- Sisson SB, Camhi SM, Church TS, Tudor-Locke C, Johnson WD, Katzmarzyk PT. Accelerometer-determined steps/day and metabolic syndrome. *Am J Prev Med*. 38(6):575–582. [PubMed: 20494233]
- Storti KL, Pettee KK, Brach JS, Talkowski JB, Richardson CR, Kriska AM. Gait speed and step-count monitor accuracy in community-dwelling older adults. *Med Sci Sports Exerc*. 2008; 40(1):59–64. [PubMed: 18091020]
- Tudor-Locke C, Bassett DR Jr. How many steps/day are enough? Preliminary pedometer indices for public health. *Sports Med*. 2004; 34(1):1–8. [PubMed: 14715035]

- Tudor-Locke C, Hatano Y, Pangrazi RP, Kang M. Revisiting “how many steps are enough?”. *Med Sci Sports Exerc.* 2008; 40(7 Suppl):S537–543. [PubMed: 18562971]
- Tudor-Locke C, Sisson SB, Collova T, Lee SM, Swan PD. Pedometer-determined step count guidelines for classifying walking intensity in a young ostensibly healthy population. *Can J Appl Physiol.* 2005; 30(6):666–676. [PubMed: 16485518]
- Weiss CO, Fried LP, Bandeen-Roche K. Exploring the hierarchy of mobility performance in high-functioning older women. *J Gerontol A Biol Sci Med Sci.* 2007; 62(2):167–173. [PubMed: 17339641]
- Wilkinson, GS. WRAT3: Wide Range Achievement Test 3. Wide Range Inc; Wilmington, DE: 1993.
- Yesavage JA, Brink TL, Rose TL, Lum O, Huang V, Adey M, et al. Development and validation of a geriatric depression screening scale: a preliminary report. *J Psychiatr Res.* 1982; 17(1):37–49. [PubMed: 7183759]



Note. Walking activity over one day (24 hours)


	Metric	Definition
	Low-intensity activity	Steps under 100 steps/min
(I)	Amount	Total number of steps/day
(II)	Duration	Number of minutes of activity between 0-100 steps
(III)	Frequency	Number of bouts of continuous 10 minutes of activity

Figure 1.
Low-intensity activity metrics




Figure 2. Distribution of average number of minutes/day of activity by intensity (n=187) Moderate-intensity walking activity contributed to 9.9% of total steps/day and 2.1% of total minutes of activity/day *Note.* Low-intensity activity (<100 steps/min); **Moderate to vigorous-intensity** activity (≥ 100 steps/min)

Table 1

Baseline Characteristics Study Subjects (N=187)

Characteristic	N (%) or Mean \pm SD
Age (years)	66.8 \pm 5.6
Female	143 (76.5)
Race (African American)	171 (91.4)
Education (high school)	71 (36.9)
Income (\$15,000)	52 (28.0)
Chronic Disease	
Obesity (BMI \geq 30)	109 (58.3)
Hypertension	131 (71.2)
Osteoarthritis	113 (61.8)
Diabetes	60 (32.6)
Health Measures	
Difficulty walking 1 mile or several blocks	86 (46.0)
Difficulty with 1 flight of stairs	44 (23.5)
Lower extremity function (SPPB) ^a	8.6 \pm 2.1
Depressive symptoms (GDS) ^b	3.1 \pm 1.5
Quality of Life	1.0 \pm 1.1
Global Cognition (MMSE) ^c	28.3 \pm 1.52
% meeting physical activity guidelines	
<i>10,000 steps/day</i> ^d	
Active (\geq 10,000 steps/day)	13 (7.0)
<i>30 minutes of moderate intensity activity</i> ^e	
Active (\geq 30 minutes/day)	2(1.1)
Activity Metrics	
<i>Amount:</i> Steps/ day (total)	7648.8 \pm 3467.7
Steps/ day at low-intensity ^f	6888.1 \pm 2835.9
Steps/ day at moderate -intensity ^g	760.7 \pm 1121.6
<i>Duration:</i> Minutes of activity at low-intensity ^f	321.9 \pm 98.6
Minutes of activity at moderate to vigorous -intensity ^g	6.9 \pm 10.1
<i>Frequency:</i> Number of bouts of 10 min activity at low-intensity ^f	10.3 \pm 6.1

Note. SD = standard deviation

^aSPPB = Short Physical Performance Battery

^bGDS = Geriatric Depression Scale

^cMMSE = Mini-Mental State Exam

^d10,000 steps/day considered an estimate of daily recommended walking activity

^e30 minutes/day of moderate intensity activity (\geq 100 steps/min) considered an estimate of daily recommended walking activity

f low-intensity activity defined as walking activity at < 100 steps/min

g moderate to vigorous-intensity is defined as activity at 100 steps/min

Table 2

Associations between metrics of low-intensity activity and health measures

Health Measures	Low-intensity ^d activity metrics			
		Amount: Steps/day (per 1000 steps increase)	Duration: Minutes of activity (per 10 min increase)	Frequency: Number of bouts of 10 min activity
Difficulty walking one mile or several blocks	OR (95% CI)	0.86 (0.77, 0.97) ^{**}	0.98 (0.95, 1.01)	0.94 (0.89, 0.99) [*]
Difficulty going up or down a flight of stairs	OR (95% CI)	0.84 (0.73, 0.98) [*]	0.98 (0.95, 1.02)	0.93 (0.87, 0.98) [*]
Lower-extremity function (SPPB) ^b	β (95% CI)	0.18 (0.07, 0.28) ^{**}	0.05 (0.02, 0.08) ^{**}	0.08 (0.03, 0.13) ^{**}
Depressive Symptoms (GDS) ^c	β (95% CI)	-0.09 (-0.17, -0.01) [*]	-0.02 (-0.05, -0.00) [*]	-0.04 (-0.07, -0.00) [*]
Quality of Life	β (95% CI)	-0.07 (-0.12, -0.02) [*]	-0.02 (-0.03, -0.00)	-0.03 (-0.06, -0.01) [*]
Global Cognition (MMSE) ^d	β (95% CI)	0.06 (-0.05, 0.16)	0.02 (-0.01, 0.04)	0.00 (-0.04, 0.05)

Note. CI = confidence interval; OR = odds ratio; all models adjusted for covariates age, gender and race

* p < 0.05;

** p < 0.01

^a Low-intensity activity defined as walking activity at < 100 steps/min; all analyses were restricted to low-intensity activity metrics, and excluded moderate to vigorous-intensity activity (> 100 steps/min)

^b SPPB = Short Physical Performance Battery

^c GDS = Geriatric Depression Scale

^d MMSE = Mini-Mental State Exam