

Lawrence Berkeley National Laboratory

LBL Publications

Title

EFFECTS OF s-PROCESS BRANCHINGS ON STELLAR AND METEORITIC ABUNDANCES

Permalink

<https://escholarship.org/uc/item/8wp266hf>

Author

Norman, E.B.

Publication Date

1985-12-01

c.2



Lawrence Berkeley Laboratory

UNIVERSITY OF CALIFORNIA

RECEIVED
LAWRENCE
BERKELEY LABORATORY
FEB 18 1986
LIBRARY AND
DOCUMENTS SECTION


Presented at the 120 IAU Symposium,
Goa, India, December 3-7, 1985

EFFECTS OF s-PROCESS BRANCHINGS ON
STELLAR AND METEORITIC ABUNDANCES

E.B. Norman, K.T. Lesko, S.G. Crane,
R.M. Larimer, and A.E. Champagne

December 1985

TWO-WEEK LOAN COPY
*This is a Library Circulating Copy
which may be borrowed for two weeks.*



LBL-20298
c.2

DISCLAIMER

This document was prepared as an account of work sponsored by the United States Government. While this document is believed to contain correct information, neither the United States Government nor any agency thereof, nor the Regents of the University of California, nor any of their employees, makes any warranty, express or implied, or assumes any legal responsibility for the accuracy, completeness, or usefulness of any information, apparatus, product, or process disclosed, or represents that its use would not infringe privately owned rights. Reference herein to any specific commercial product, process, or service by its trade name, trademark, manufacturer, or otherwise, does not necessarily constitute or imply its endorsement, recommendation, or favoring by the United States Government or any agency thereof, or the Regents of the University of California. The views and opinions of authors expressed herein do not necessarily state or reflect those of the United States Government or any agency thereof or the Regents of the University of California.

97

EFFECTS OF s-PROCESS BRANCHINGS
ON STELLAR AND METEORITIC ABUNDANCES

E.B. Norman, K.T. Lesko, S.G. Crane, R.M. Larimer
Nuclear Science Division
Lawrence Berkeley Laboratory
University of California
Berkeley, CA 94710 U.S.A.

A.E. Champagne
Physics Department
Princeton University
Princeton, NJ 08544 U.S.A.

December 1985

2

EFFECTS OF s-PROCESS BRANCHINGS ON STELLAR AND METEORITIC ABUNDANCES*

E. B. Norman, K. T. Lesko, S. G. Crane, R. M. Larimer
 Nuclear Science Division, Lawrence Berkeley Laboratory
 Berkeley, CA 94720 U.S.A., and
 A. E. Champagne
 Physics Department, Princeton University
 Princeton, NJ 08544 U.S.A.

ABSTRACT. The level scheme and electromagnetic properties of ^{148}Pm have been studied using $^{149}\text{Sm}(d, ^3\text{He})$ and $^{148}\text{Nd}(p, n\gamma)$ reactions. Combining these measurements with estimates for E2/M1 decay branching ratios leads to the tentative conclusion that $^{148}\text{Pm}_{g,m}$ are in thermal equilibrium during the s-process. The branch at ^{148}Pm then leads to an inferred s-process neutron density of $3 \times 10^8 \text{ cm}^{-3}$.

At a number of points along the path of the slow neutron-capture process (s-process), nuclei are encountered which are sufficiently long-lived that neutron capture on these isotopes can compete with their normal beta decays. That such effects actually occur has been inferred from observations of both stellar and meteoritic elemental and isotopic abundances. The details of these s-process branchings depend sensitively on stellar environmental factors such as temperature, matter density, and neutron density. Thus, these branch points can provide specific information on the stellar conditions under which the s-process occurs.

Tomkin and Lambert¹ examined the effects of the s-process branch at ^{85}Kr on the elemental abundances they observed in the barium star HR 774. From their studies, it was concluded that in this star the s-process neutron density was in the range $10^6 \leq N(n) \leq 10^7 \text{ cm}^{-3}$.




Figure 1. The s-process path through the mass region $145 < A < 152$. Stable isotopes are indicated by shaded triangles. Beta decay half-lives of unstable isotopes are indicated.

Käppeler² has investigated the s-process branchings at ^{151}Sm , ^{185}W , and ^{204}Tl . From measurements and calculations of neutron-capture cross sections and measurements of meteoritic abundances, he has inferred an s-process neutron density on the order of $3 \times 10^8 \text{ cm}^{-3}$. Furthermore, a possible trend toward the requirement of higher neutron densities for the production of the heavier nuclei can be seen in Käppeler's results.

The branch at ^{148}Pm illustrated in Fig. 1 offers another opportunity to determine the s-process neutron density.³ The inferred value of this quantity depends sensitively on whether or not $^{148}\text{Pm}_{g,m}$ are in thermal equilibrium at the temperatures found in the helium-burning zones of red-giant stars. A possible mechanism by which such equilibration could occur is shown in Fig. 2. To decide this issue, the positions and decay modes of ^{148}Pm excited states must be known. Using the Princeton Q3D spectrograph we have observed the large number of low-lying ^{148}Pm excited states shown in Fig. 3. At LBL'S 88-Inch Cyclotron we have used γ -ray singles and γ - γ coincidences from the $^{148}\text{Nd}(p,n)^{148}\text{Pm}$ reaction to study the decay modes of these levels. A large number of γ -ray lines between 75 and 400 keV have been observed and coincidence relations for many transitions have been established. As yet, no direct link between $^{148}\text{Pm}_{g,m}$ (such as that shown in Fig. 2) has been observed. However, combining the observed high level density and γ -ray transitions with theoretical estimates of E2/M1 decay branching ratios leads to the tentative conclusion that $^{148}\text{Pm}_{g,m}$ are in thermal equilibrium during the s-process. The inferred neutron density using this branch point is then $3 \times 10^8 \text{ cm}^{-3}$.




Figure 2. Partial level scheme of ^{148}Pm illustrating the possibility of photoexcitation of $^{148}\text{Pm}_{g,m}$ to a higher lying state that decays to $^{148}\text{Pm}_g$.




Figure 3. Spectrum observed from the $^{149}\text{Sm}(d,^3\text{He})^{148}\text{Pm}$ reaction at 20 degrees. Energies of ^{148}Pm excited states are given in keV.

REFERENCES

- *Supported by U.S. Dept. of Energy contract No. DE-AC03-76SF00098, and U.S. National Science Foundation contract No. PHY83-04977.
- 1. J. Tomkin and D. L. Lambert. *Ap. J.* 273, 722 (1983).
- 2. F. Käppeler, *Proc. 2nd Wrkshp. Nucl. Astrophys.*, MPA 90, (1983) p.82.
- 3. F. Käppeler, *AIP Conf. Proc.* 125, 715 (1985).

This report was done with support from the Department of Energy. Any conclusions or opinions expressed in this report represent solely those of the author(s) and not necessarily those of The Regents of the University of California, the Lawrence Berkeley Laboratory or the Department of Energy.

Reference to a company or product name does not imply approval or recommendation of the product by the University of California or the U.S. Department of Energy to the exclusion of others that may be suitable.

*LAWRENCE BERKELEY LABORATORY
TECHNICAL INFORMATION DEPARTMENT
UNIVERSITY OF CALIFORNIA
BERKELEY, CALIFORNIA 94720*