

UC San Diego

UC San Diego Previously Published Works

Title

Redox regulation of mitochondrial fission, protein misfolding, synaptic damage, and neuronal cell death: potential implications for Alzheimer's and Parkinson's diseases

Permalink

<https://escholarship.org/uc/item/8xt645df>

Journal

Apoptosis: An International Journal on Programmed Cell Death, 15(11)

ISSN

1573-675X

Authors

Nakamura, Tomohiro
Lipton, Stuart A.

Publication Date

2010-11-01

DOI

10.1007/s10495-010-0476-x

Peer reviewed

Redox regulation of mitochondrial fission, protein misfolding, synaptic damage, and neuronal cell death: potential implications for Alzheimer's and Parkinson's diseases

Tomohiro Nakamura · Stuart A. Lipton

Published online: 23 February 2010

© The Author(s) 2010. This article is published with open access at Springerlink.com

Abstract Normal mitochondrial dynamics consist of fission and fusion events giving rise to new mitochondria, a process termed mitochondrial biogenesis. However, several neurodegenerative disorders manifest aberrant mitochondrial dynamics, resulting in morphological abnormalities often associated with deficits in mitochondrial mobility and cell bioenergetics. Rarely, dysfunctional mitochondrial occur in a familial pattern due to genetic mutations, but much more commonly patients manifest sporadic forms of mitochondrial disability presumably related to a complex set of interactions of multiple genes (or their products) with environmental factors (G × E). Recent studies have shown that generation of excessive nitric oxide (NO), in part due to generation of oligomers of amyloid- β (A β) protein or overactivity of the NMDA-subtype of glutamate receptor, can augment mitochondrial fission, leading to frank fragmentation of the mitochondria. *S*-Nitrosylation, a covalent redox reaction of NO with specific protein thiol groups, represents one mechanism contributing to NO-induced mitochondrial fragmentation, bioenergetic failure, synaptic damage, and eventually neuronal apoptosis. Here, we summarize our evidence in Alzheimer's disease (AD) patients and animal models showing that NO contributes to mitochondrial fragmentation via *S*-nitrosylation of dynamin-related protein 1 (Drp1), a protein involved in mitochondrial fission. These findings may provide a new target

for drug development in AD. Additionally, we review emerging evidence that redox reactions triggered by excessive levels of NO can contribute to protein misfolding, the hallmark of a number of neurodegenerative disorders, including AD and Parkinson's disease. For example, *S*-nitrosylation of parkin disrupts its E3 ubiquitin ligase activity, and thereby affects Lewy body formation and neuronal cell death.

Keywords *S*-Nitrosylation · Mitochondrial fragmentation · Dynamin-related protein 1 · β -Amyloid · Alzheimer's disease

Introduction

Brains with neurodegenerative diseases often manifest excessive generation of reactive nitrogen species (RNS) and reactive oxygen species (ROS), which can contribute to neuronal cell injury and death via a series of redox reactions [1–5]. While many intra- and extracellular molecules may participate in neuronal injury, accumulation of nitrosative stress due to excessive generation of nitric oxide (NO) appears to be a potential factor contributing to neuronal cell damage and death [6, 7]. A well-established model for NO production entails a central role of the *N*-methyl-D-aspartate (NMDA)-type glutamate receptors in nervous system. Excessive activation of NMDA receptors drives Ca²⁺ influx, which in turn activates neuronal NO synthase (nNOS) as well as the generation of ROS [8, 9]. Accumulating evidence suggests that NO can mediate both protective and neurotoxic effects by redox reactions with cysteine residues of target proteins to form *S*-nitrosothiols (SNOs), a process termed *S*-nitrosylation because of its effects on the chemical biology of protein function.

T. Nakamura · S. A. Lipton (✉)
Center for Neuroscience, Aging, and Stem Cell Research,
Sanford-Burnham Medical Research Institute, 10901 North
Torrey Pines Road, La Jolla, CA 92037, USA
e-mail: slipton@burnham.org

S. A. Lipton
Department of Neurosciences, University of California at San
Diego, 9500 Gilman Drive, La Jolla, CA 92037, USA

Importantly, normal mitochondrial respiration may also generate free radicals, principally ROS, and one such molecule, superoxide anion (O_2^-), reacts rapidly with free radical NO to form the very toxic product peroxynitrite ($ONOO^-$) [10, 11].

Production of NO from inducible NOS (iNOS) can also contribute to the pathogenesis of neurodegenerative diseases, including Alzheimer's disease (AD) [12, 13]. A classic feature of AD pathology is the generation of β -amyloid ($A\beta$) peptides. Recently, several lines of evidence have suggested that soluble oligomers of $A\beta$ represent the most toxic form of the peptide [14]. Consistent with this notion, $A\beta$ oligomers, but not fibrillar $A\beta$, induce high expression of iNOS in astrocytes and thus generation of NO [15]. Additionally, $A\beta$ is known to inhibit glutamate re-uptake, at least in part via generation of ROS; this can lead to pathological activation of NMDA receptors, thereby disturbing synaptic function in AD [16–18]. Excessive stimulation of NMDA receptors also leads to activation of nNOS, as discussed above, thus representing another source of NO emanating from $A\beta$ oligomers.

Dysfunction in mitochondria represents another hallmark of neurodegenerative diseases. For example, patients with early stage AD regularly exhibit declining mitochondrial energy metabolism and ATP production, which may subsequently cause synaptic loss and neuronal damage [19–21]. Interestingly, neurons in AD and other neurodegenerative brains often display abnormal mitochondrial morphology [22, 23].

Normally, mitochondria are known to continuously undergo fission and fusion (known as mitochondrial dynamics) to generate smaller organelles or elongated, tubular structures, respectively. This normal mitochondrial fission and fusion can facilitate formation of new mitochondria (biogenesis), repair of defective mitochondrial DNA through mixing, and redistribution of mitochondria to sites requiring high-energy production [24–26]. Conversely, an imbalance in fission or fusion initiates malfunctions in mitochondrial morphology and bioenergetics, and may thus contribute to neuronal injury during neurodegeneration [26–28]. Dysfunction in mitochondrial dynamics can result from either (i) rare genetic mutations in fission- or fusion-related genes, as occurs in Charcot-Marie-Tooth (CMT) Disease and Autosomal Dominant Optic Atrophy (ADOA) [29, 30], or (ii) posttranslational changes to the fission or fusion proteins. In particular, a posttranslational modification engendered by nitrosative/oxidative stress may well account for the more common sporadic cases of the disease [31, 32]. Hence, a key theme of this article is the hypothesis that excessive accumulation of nitrosative stress contributes to abnormal mitochondrial morphology in brains of neurodegenerative patients. In this review, we discuss a specific example, showing that *S*-nitrosylation of the mitochondrial

fission protein dynamin-related protein 1 (Drp1) contributes to excessive mitochondrial fission/fragmentation, synaptic injury, and neuronal apoptosis in neurodegenerative diseases such as AD. We also review evidence that *S*-nitrosylation of critical proteins can cause protein misfolding and contribute to the aggregation of aberrant proteins seen in many neurodegenerative disorders, including AD and Parkinson's disease (PD).

Mitochondrial fission/fusion machinery in nerve cells

Neurons are particularly vulnerable to mitochondrial defects because they require high levels of energy for their survival and specialized function. Mitochondrial biogenesis is required in regions that demand high concentrations of ATP, especially the synapse. The distribution of mitochondria at the nerve terminal can control synaptic transmission and structure [24, 33, 34].

In healthy neurons, the fission/fusion machinery proteins maintain mitochondrial integrity and insure their presence at critical locations. These proteins includes Drp1 and Fis1, acting as fission proteins, and Mitofusin (Mfn) and Opa1, operating as fusion proteins [35]. In both familial and sporadic neurodegenerative conditions, abnormal mitochondria regularly appear in the brain as a result of dysfunction in the fission/fusion machinery. Genetic mutations in Mfn2 can cause CMT disease, a hereditary peripheral neuropathy that affects both motor and sensory neurons [30, 36]. Mutations in Opa1 cause ADOA, characterized by the loss of retinal ganglion cells and the optic nerve, representing their axons [29]. Recently, Waterham et al. described a heterozygous, dominant-negative mutation of Drp1 in a patient whose symptoms were broadly similar to those of CMT neuropathy and ADOA [37]. Drp1 includes four distinct structural domains: an N-terminal GTPase domain, a dynamin-like middle domain, an insert B domain, and a C-terminal GED domain. The mutation (Ala 395 to Asp) was found in the middle domain of Drp1. This case study further suggested that a defect in mitochondrial fission may have more severe consequences than those of fusion defects, since the Drp1 mutation caused a much earlier onset (prenatal) and fatal outcome. Additionally, it is apparent that the balance between fission and fusion is critical for normal function of mitochondria and determination of phenotype in disease. These fission/fusion proteins are widely expressed in human tissues, clearly supporting the notion that neurons are particularly sensitive to mitochondrial dysfunction.

Dysregulation of mitochondrial dynamics in AD

An estimated 26 million people globally have AD, which is thought to be the most common form of dementia. In AD

brains, neuronal loss in the hippocampus and cerebral cortex mainly accounts for the cognitive decline. Degen-erating AD brains contain aberrant accumulations of mis-folded, aggregated proteins— $A\beta$ and tau—which can adversely affect neuronal connectivity and plasticity, and trigger cell death signaling pathways. These aggregates are recognized as either intracellular neurofibrillary tangles, which contain hyperphosphorylated tau, or extracellular plaques, which contain $A\beta$. β -Secretase and γ -secretase proteolytically cleave amyloid precursor protein (APP) in its transmembrane region to generate $A\beta$. It is currently thought that soluble oligomers of misfolded protein are pathogenic and that the large aggregates may actually be an attempt by the cell to wall off the aberrant proteins (although such aggregates could potentially be toxic by location or if not contained by the proper chaperones). Interestingly, a recent study showed that an N-terminal fragment of APP may also contribute to neurodegeneration in AD models [38].

Emerging evidence suggests that mitochondrial dys-function plays a prominent role in the pathogenesis of AD [39]. Analyses of autopsy and biopsy samples revealed that mitochondria isolated from AD brains exhibit diminished respiratory capacity [20], and that AD neurons contain a number of mitochondria with fractured cristae [40]. Additionally, electron-microscopic studies have described an increase in mitochondrial fragmentation in human AD brains [22, 23]. In cell-based experiments, $A\beta$ production resulted in the appearance of fragmented and abnormally distributed mitochondria [27, 41], suggesting that $A\beta$ (possibly in the form of soluble oligomers) may trigger excessive mitochondrial fission in AD patients. Pathologi-cal forms of tau may also contribute to mitochondrial fragmentation in AD brains since expression of caspase-cleaved tau induced mitochondrial fission in a calcineurin-dependent manner [42].

Similarly, dysfunction in mitochondrial integrity is associated with PD [43]. For instance, the Parkinsonian neurotoxins, rotenone and 1-methyl-4-phenyl-pyridinium ion (MPP⁺), which inhibit complex I of the mitochondrial electron transport chain, can induce excessive mitochon-drial fragmentation and cell death [27]. Additionally, multiple groups recently observed that a deficiency in familial PD-related proteins, such as parkin and PINK1, led to the appearance of mitochondrial pathology [44–47]. Exogenous expression of mitochondrial fusion proteins, Mfn2 and OPA1, or dominant negative Drp1 rescued the altered mitochondrial morphology, suggesting that parkin or PINK1 deficiency promoted mitochondrial fragmenta-tion [46]. Interestingly, Drp1 seems to activate autophagy/mitophagy pathways for morphologic remodeling of mitochondria in PINK1-deficient neuroblastoma cells [48]. Taken together, dysregulation of mitochondrial dynamics

may contribute to a common pathway leading to the patho-genesis of several neurodegenerative diseases, including AD and PD.

NMDA receptor signaling pathways induce generation of RNS/ROS

The amino acid glutamate is the major excitatory neuro-transmitter in the brain. Excitatory neurotransmission plays an important role in the normal development and plasticity of synapses, and for some forms of learning or memory; however, excessive activation of glutamate receptors is implicated in neuronal damage in many neurological dis-orders ranging from acute hypoxic-ischemic brain injury to chronic neurodegenerative diseases. Among several fami-lies of glutamate receptors, the NMDA receptor has long attracted attention because it has several properties that set it apart from the other receptors. One such characteristic is that NMDA receptor-coupled channels are highly perme-able to Ca^{2+} , triggering Ca^{2+} entry after ligand binding if the cell is first depolarized in order to relieve channel block by Mg^{2+} [49, 50]. Subsequent binding of Ca^{2+} to various intracellular molecules can lead to many significant con-sequences. In particular, excessive activation of NMDA receptors leads to the production of damaging free radicals (e.g., NO and ROS) and other enzymatic processes, con-tributing to cell injury and death [6, 11, 51–54]. It is cur-rently thought that overstimulation of extrasynaptic NMDA receptors mediates this neuronal damage, while, in con-trast, synaptic activity predominantly activates survival pathways [55–57]. Intense hyperstimulation of excitatory receptors leads to necrotic cell death, but milder or chronic overstimulation can result in apoptotic or other forms of cell death [53, 54, 58]. Furthermore, $A\beta$ oligomers induce neuronal synaptic damage via an NMDA receptor-depend-ent mechanism [59].

Excessive activation of glutamate receptors has been implicated in neuronal damage in many neurological dis-orders. John Olney coined the term “excitotoxicity” to describe this phenomenon [60, 61]. This form of toxicity is mediated at least in part by excessive activation of NMDA-type receptors [6, 7, 62], resulting in excessive Ca^{2+} influx through a receptor’s associated ion channel. Increased levels of neuronal Ca^{2+} , in conjunction with the Ca^{2+} -binding protein CaM, trigger the activation of nNOS and subsequent generation of NO from the amino acid L-argi-nine [8, 63]. NO is a gaseous free radical (thus highly diffusible) and a key molecule that plays a vital role in normal signal transduction but in excess can lead to neu-ronal cell damage and death. The diversity of NO effects on neuronal survival can also be caused by the formation of different NO species or intermediates: NO radical (NO[•]),

nitrosonium cation (NO^+), nitroxyl anion (NO^- , with high energy singlet and lower energy triplet forms) [11].

Nitrosative stress regulates protein misfolding and neuronal cell death

Early investigations indicated that NO participates in cellular signaling pathways, which regulate broad aspects of brain function, including synaptic plasticity, normal development, and neuronal cell death [51, 64–66]. In general, NO exerts physiological and some pathophysiological effects via stimulation of guanylate cyclase to form cyclic guanosine-3',5'-monophosphate (cGMP) or through *S*-nitrosylation of regulatory protein thiol groups [9, 11, 67–70]. *S*-Nitrosylation is the covalent redox reaction of an NO group with a critical cysteine thiol/sulfhydryl (RSH or, more properly, thiolate anion, RS^-) to form an *S*-nitrosothiol derivative (R-SNO). Such modification modulates the function of a broad spectrum of mammalian, plant, and microbial proteins. In general, a consensus motif of amino acids comprised of nucleophilic residues (generally an acid and a base) surround a critical cysteine, which increases the cysteine sulfhydryl's susceptibility to *S*-nitrosylation [71, 72]. Our group first identified the physiological relevance of *S*-nitrosylation by showing that NO and related RNS exert paradoxical effects via redox-based mechanisms—NO is neuroprotective via *S*-nitrosylation of NMDA receptors (as well as other subsequently discovered targets, including caspases), and yet can also be neurodestructive by formation of peroxynitrite (or, as later discovered, reaction with additional molecules such as MMP-9, parkin, peroxiredoxin II, Drp1, protein-disulfide isomerase (PDI), and GAPDH) [11, 73–80]. Over the past decade, accumulating evidence has suggested that *S*-nitrosylation can regulate the biological activity of a great variety of proteins, in some ways akin to phosphorylation [11, 72, 79–92]. Chemically, NO is often a good “leaving group,” facilitating further oxidation of critical thiols to disulfide bonds among neighboring (vicinal) cysteine residues or, via reaction with ROS, to sulfenic (–SOH), sulfinic (–SO₂H), or sulfonic (–SO₃H) acid derivatization of the protein [79, 82, 83, 93]. Alternatively, *S*-nitrosylation may possibly produce a nitroxyl disulfide, in which the NO group is shared by close or vicinal cysteine thiols [94].

In prior work, analyses of mice deficient in either nNOS or iNOS confirmed that NO is an important mediator of cell injury and death after excitotoxic stimulation; excessive NO generated from nNOS or iNOS is detrimental to neuronal survival [95, 96]. In addition, inhibition of NOS activity ameliorates the progression of disease pathology in animal models of AD, PD, and Amyotrophic Lateral Sclerosis, suggesting that excess generation of NO plays a

pivotal role in the pathogenesis of several neurodegenerative diseases [97–100].

A common theme in many neurodegenerative disorders is the finding of abnormal aggregates of misfolded proteins. Recent findings have implied that NO-related species may significantly participate in the process of protein misfolding through protein *S*-nitrosylation under degenerative conditions. Specifically, we and others recently mounted physiological and chemical evidence that *S*-nitrosylation modulates the (i) ubiquitin E3 ligase activity of parkin [81, 82, 101], and (ii) chaperone and isomerase activities of PDI [83], contributing to protein misfolding and neurotoxicity in models of neurodegenerative disorders (Fig. 1). Mutations in the parkin gene can cause autosomal recessive juvenile Parkinsonism (ARJP), accounting for some cases of hereditary PD manifest in young patients with onset beginning anywhere from the teenage years through the 40s [102–104]. As an E3 ubiquitin ligase, parkin catalyzes formation of polyubiquitin chains on substrate proteins (e.g., synphilin-1 and Pael-R); these ubiquitin chains represent the signal for proteasomal degradation of the proteins. Clearance of misfolded or aberrant proteins requires, at least in part, the activity of the ubiquitin-proteasome system (UPS). We and others found evidence for *S*-nitrosylation of parkin (forming SNO-parkin) not only in rodent models of PD but also in the brains of human patients with PD and the related α -synucleinopathy, diffuse Lewy body disease (DLBD). SNO-parkin initially stimulates ubiquitin E3 ligase activity, resulting in enhanced ubiquitination as observed in Lewy bodies, followed by a decrease in enzyme activity, producing a futile cycle of dysfunctional UPS [82, 101, 105]. Moreover, *S*-nitrosylation appears to compromise the neuroprotective effect of parkin [81]. These mechanisms involve *S*-nitrosylation of critical cysteine residues in the first RING domain of parkin [82]. Nitrosative and oxidative stress can also alter the solubility of parkin via posttranslational modification of cysteine residues, which may concomitantly compromise its protective function [106–108].

During protein folding in the ER, PDI can introduce disulfide bonds into proteins (oxidation), break disulfide bonds (reduction), and catalyze thiol/disulfide exchange (isomerization), thus facilitating disulfide bond formation, rearrangement reactions, and structural stability [109]. In many neurodegenerative disorders as well as in cerebral ischemia, the accumulation of immature and denatured proteins results in ER dysfunction [110–113], and upregulation of PDI represents an adaptive response known as the unfolded protein response (UPR), which promotes protein refolding and may contribute to neuroprotection [111, 114–116]. In a recent study, we reported that the *S*-nitrosylation of PDI (to form SNO-PDI) disrupts this neuroprotective role [83]. We found that PDI is




Fig. 1 Schema of S-nitrosylated species contributing to the accumulation of aberrant proteins and neuronal damage. NMDAR hyperactivation triggers generation of NO/ROS. S-Nitrosylation of parkin (forming SNO-PARK) and PDI (forming SNO-PDI) can contribute to neuronal cell injury in part by triggering accumulation of misfolded proteins

S-nitrosylated in the brains of virtually all cases examined of sporadic AD and PD. SNO-PDI formation led to the accumulation of polyubiquitinated/misfolded proteins and activation of the UPR. Moreover, S-nitrosylation abrogated the inhibitory effect of PDI on aggregation of proteins observed in Lewy body inclusions [83, 117]. S-Nitrosylation of PDI also prevented its attenuation of neuronal cell death triggered by ER stress, misfolded proteins, or proteasome inhibition.

Nitrosative stress can also result in defects in mitochondrial function. For example, NO affects mitochondrial respiration by reversibly inhibiting complexes I and IV [118, 119]. Mitochondria thus compromised will release ROS, and this in turn could contribute to brain aging and/or pathological conditions associated with neurodegenerative diseases. Additionally, increased nitrosative and oxidative stress can elicit dysfunction of mitochondrial dynamics (fission and fusion events) [27, 120, 121]. However, until recently little was known regarding the molecular and pathogenic mechanisms by which NO contributes to the formation of fragmented mitochondria. Our recent findings


have shed light on the molecular events underlying this relationship, particularly in AD. Specifically, we recently discovered physiological and chemical evidence that S-nitrosylation modulates the GTPase activity of Drp1, thus contributing to mitochondrial fragmentation, bioenergetic impairment, synaptic damage, and eventually frank neuronal loss in models of AD.

S-Nitrosylation of Drp1 mediates mitochondrial fission and neurotoxicity in cell models of AD

In addition to the rare genetic mutations seen in the genes encoding mitochondrial fission and fusion proteins, recent studies have demonstrated that posttranslational modification of these molecules can contribute to altered mitochondria dynamics. For example, phosphorylation, ubiquitination, sumoylation, and proteolytic cleavage of Drp1 regulate mitochondrial fission by affecting Drp1 activity, at least in cell culture systems [122–129]. Excessive activation of mitochondrial fission or fusion proteins by posttranslational modification was posited to contribute to neurodegeneration by compromising mitochondrial function. Interestingly, along these lines, we recently reported that excessive NO can also lead to S-nitrosylation of Drp1 at Cys644 [31]. Cys644 resides within the GTPase effector domain (GED) of Drp1, which influences both GTPase activity and oligomer formation of Drp1 [130–133]. S-Nitrosylation of Drp1 (forming SNO-Drp1) induces formation of Drp1 dimers, which function as building blocks for tetramers and higher order structures of Drp1, and activates Drp1 GTPase activity; however, substitution of Cys644 to Ala (C644A) abrogated these effects of NO.

We further demonstrated that exposure to oligomeric A β peptide results in formation of SNO-Drp1 in cell culture models (Fig. 2). Moreover, we and others have observed that Drp1 is S-nitrosylated in the brains of virtually all cases of sporadic AD [23, 31]. In order to determine the consequences of S-nitrosylated Drp1 in neurons, we exposed cultured cerebrocortical neurons to the physiological NO donor, SNO-C, or to A β oligomers and found that both induced SNO-Drp1 formation and led to the accumulation of fragmented mitochondria. Moreover, mutation of a specific cysteine residue in Drp1 (C644A) prevented these effects of SNO-C or A β on mitochondrial fragmentation, consistent with the notion that SNO-Drp1 triggered excessive mitochondria fission or fragmentation. Finally, in response to A β , SNO-Drp1—induced mitochondrial fragmentation caused synaptic damage, an early characteristic feature of AD, and eventually apoptotic neuronal cell death. Importantly, blockade of Drp1 nitrosylation (using the Drp1(C644A) mutant) prevented A β -mediated synaptic loss and neuronal cell death,

Fig. 2 Possible mechanism whereby *S*-nitrosylated species contribute to excessive mitochondrial fragmentation and neuronal damage. NMDAR hyperactivation triggers generation of NO and subsequent *S*-nitrosylation of neuronal proteins, contributing to synaptic damage and eventually neuronal death. Soluble oligomers of A β oligomers, known to be a key mediator of AD pathogenesis, can facilitate neuronal NO production in both NMDAR-dependent and -independent manners. *S*-Nitrosylation of Drp1 (forming SNO-Drp1) can contribute to synaptic damage and neuronal cell death by triggering excessive mitochondrial fission and bioenergetic impairment



suggesting that SNO-Drp1 may represent a potential therapeutic target to protect neurons and their synapses in AD.

In addition to AD, SNO-Drp1 may affect the pathogenesis of other neurodegenerative disorders, such as Huntington's disease (HD), as we have recently observed nitrosylation of Drp1 in HD brains as well as in AD brains. An expanded CAG repeat in the huntingtin (htt) gene is the cause of HD. A potential mechanism for neurodegeneration in HD that is triggered by mutant Htt (mtHtt) involves mitochondrial dysfunction. Evidence for this includes reduced activity of respiratory complexes, decreased mitochondrial membrane potential, and changes in mitochondrial ultrastructure [134, 135]. Interestingly, expression of mtHtt induces mitochondrial fragmentation in Hela cells [136], and exposure to the mitochondrial complex II inhibitor 3-NP, which reproduces many of the pathological features of HD, leads to mitochondria fission [137]. Furthermore, accumulating evidence suggests that excitotoxic

pathways, which increase NO, may contribute to the pathophysiology of HD [138]. Hence, it is tempting to postulate that SNO-Drp1 may be involved in the mitochondrial fragmentation and neuronal injury observed in HD.

Conclusions

Excessive nitrosative and oxidative stress triggered by overstimulation of NMDA receptors and/or A β production may result in mitochondrial dysfunction, thus contributing to synaptic damage and neuronal loss in sporadic forms of neurodegenerative diseases. Additionally, nitrosylation reactions can lead to aberrant and misfolded proteins in neurodegenerative conditions. For example, *S*-nitrosylation of parkin and PDI can contribute to the accumulation of such aggregated proteins. Our recent identification of

SNO-Drp1, triggered by oligomeric $A\beta$, provides a mechanistic link between free radical production, abnormal mitochondrial morphology, and neuronal cell injury in neurodegenerative disorders such as AD. Elucidation of this new pathway may lead to the development of additional therapeutic approaches to prevent excessive mitochondrial fission by targeted disruption or prevention of SNO-Drp1.

Acknowledgments This work was supported in part by NIH Grants P01 HD29587, R01 EY05477, R01 EY09024, the American Parkinson's Disease Association, San Diego Chapter, and an Ellison Senior Scholars Award in Aging (to S.A.L.).

Open Access This article is distributed under the terms of the Creative Commons Attribution Noncommercial License which permits any noncommercial use, distribution, and reproduction in any medium, provided the original author(s) and source are credited.

References

- Lin MT, Beal MF (2006) Mitochondrial dysfunction and oxidative stress in neurodegenerative diseases. *Nature* 443:787–795
- Barnham KJ, Masters CL, Bush AI (2004) Neurodegenerative diseases and oxidative stress. *Nat Rev Drug Discov* 3:205–214
- Muchowski PJ (2002) Protein misfolding, amyloid formation, and neurodegeneration: a critical role for molecular chaperones? *Neuron* 35:9–12
- Emerit J, Edeas M, Bricaire F (2004) Neurodegenerative diseases and oxidative stress. *Biomed Pharmacother* 58:39–46
- Beal MF (2001) Experimental models of Parkinson's disease. *Nat Rev Neurosci* 2:325–334
- Lipton SA, Rosenberg PA (1994) Excitatory amino acids as a final common pathway for neurologic disorders. *N Engl J Med* 330:613–622
- Lipton SA (2006) Paradigm shift in neuroprotection by NMDA receptor blockade: memantine and beyond. *Nat Rev Drug Discov* 5:160–170
- Bredt DS, Hwang PM, Glatt CE, Lowenstein C, Reed RR, Snyder SH (1991) Cloned and expressed nitric oxide synthase structurally resembles cytochrome P-450 reductase. *Nature* 351:714–718
- Garthwaite J, Charles SL, Chess-Williams R (1988) Endothelium-derived relaxing factor release on activation of NMDA receptors suggests role as intercellular messenger in the brain. *Nature* 336:385–388
- Beckman JS, Beckman TW, Chen J, Marshall PA, Freeman BA (1990) Apparent hydroxyl radical production by peroxynitrite: implications for endothelial injury from nitric oxide and superoxide. *Proc Natl Acad Sci USA* 87:1620–1624
- Lipton SA, Choi YB, Pan ZH et al (1993) A redox-based mechanism for the neuroprotective and neurodestructive effects of nitric oxide and related nitroso-compounds. *Nature* 364:626–632
- Medeiros R, Prediger RD, Passos GF et al (2007) Connecting TNF-alpha signaling pathways to iNOS expression in a mouse model of Alzheimer's disease: relevance for the behavioral and synaptic deficits induced by amyloid beta protein. *J Neurosci* 27:5394–5404
- Wang Q, Rowan MJ, Anwyl R (2004) Beta-amyloid-mediated inhibition of NMDA receptor-dependent long-term potentiation induction involves activation of microglia and stimulation of inducible nitric oxide synthase and superoxide. *J Neurosci* 24:6049–6056
- Haass C, Selkoe DJ (2007) Soluble protein oligomers in neurodegeneration: lessons from the Alzheimer's amyloid beta-peptide. *Nat Rev Mol Cell Biol* 8:101–112
- White JA, Manelli AM, Holmberg KH, Van Eldik LJ, Ladu MJ (2005) Differential effects of oligomeric and fibrillar amyloid-beta 1–42 on astrocyte-mediated inflammation. *Neurobiol Dis* 18:459–465
- Li S, Hong S, Shepardson NE, Walsh DM, Shankar GM, Selkoe D (2009) Soluble oligomers of amyloid beta protein facilitate hippocampal long-term depression by disrupting neuronal glutamate uptake. *Neuron* 62:788–801
- Matos M, Augusto E, Oliveira CR, Agostinho P (2008) Amyloid-beta peptide decreases glutamate uptake in cultured astrocytes: involvement of oxidative stress and mitogen-activated protein kinase cascades. *Neuroscience* 156:898–910
- Trotti D, Danbolt NC, Volterra A (1998) Glutamate transporters are oxidant-vulnerable: a molecular link between oxidative and excitotoxic neurodegeneration? *Trends Pharmacol Sci* 19:328–334
- Liang WS, Reiman EM, Valla J et al (2008) Alzheimer's disease is associated with reduced expression of energy metabolism genes in posterior cingulate neurons. *Proc Natl Acad Sci USA* 105:4441–4446
- Parker WD Jr, Parks J, Filley CM, Kleinschmidt-DeMasters BK (1994) Electron transport chain defects in Alzheimer's disease brain. *Neurology* 44:1090–1096
- Reddy PH (2007) Mitochondrial dysfunction in aging and Alzheimer's disease: strategies to protect neurons. *Antioxid Redox Signal* 9:1647–1658
- Baloyannis SJ (2006) Mitochondrial alterations in Alzheimer's disease. *J Alzheimers Dis* 9:119–126
- Wang X, Su B, Lee HG et al (2009) Impaired balance of mitochondrial fission and fusion in Alzheimer's disease. *J Neurosci* 29:9090–9103
- Chen H, Chan DC (2006) Critical dependence of neurons on mitochondrial dynamics. *Curr Opin Cell Biol* 18:453–459
- Frederick RL, Shaw JM (2007) Moving mitochondria: establishing distribution of an essential organelle. *Traffic* 8:1668–1675
- Knott AB, Perkins G, Schwarzenbacher R, Bossy-Wetzel E (2008) Mitochondrial fragmentation in neurodegeneration. *Nat Rev Neurosci* 9:505–518
- Barsoum MJ, Yuan H, Gerencser AA et al (2006) Nitric oxide-induced mitochondrial fission is regulated by dynamin-related GTPases in neurons. *EMBO J* 25:3900–3911
- Bossy-Wetzel E, Barsoum MJ, Godzik A, Schwarzenbacher R, Lipton SA (2003) Mitochondrial fission in apoptosis, neurodegeneration and aging. *Curr Opin Cell Biol* 15:706–716
- Delettre C, Lenaers G, Griffoin JM et al (2000) Nuclear gene OPA1, encoding a mitochondrial dynamin-related protein, is mutated in dominant optic atrophy. *Nat Genet* 26:207–210
- Zuchner S, Mersiyanova IV, Muglia M et al (2004) Mutations in the mitochondrial GTPase mitofusin 2 cause Charcot-Marie-Tooth neuropathy type 2A. *Nat Genet* 36:449–451
- Cho DH, Nakamura T, Fang J et al (2009) S-Nitrosylation of Drp1 mediates beta-amyloid-related mitochondrial fission and neuronal injury. *Science* 324:102–105
- Westermann B (2009) Nitric oxide links mitochondrial fission to Alzheimer's disease. *Sci Signal* 2:pe29
- Li H, Chen Y, Jones AF et al (2008) Bcl-xL induces Drp1-dependent synapse formation in cultured hippocampal neurons. *Proc Natl Acad Sci USA* 105:2169–2174
- Li Z, Okamoto K, Hayashi Y, Sheng M (2004) The importance of dendritic mitochondria in the morphogenesis and plasticity of spines and synapses. *Cell* 119:873–887

35. Youle RJ, Karbowski M (2005) Mitochondrial fission in apoptosis. *Nat Rev Mol Cell Biol* 6:657–663
36. Kijima K, Numakura C, Izumino H et al (2005) Mitochondrial GTPase mitofusin 2 mutation in Charcot-Marie-Tooth neuropathy type 2A. *Hum Genet* 116:23–27
37. Waterham HR, Koster J, van Roermund CW, Mooyer PA, Wanders RJ, Leonard JV (2007) A lethal defect of mitochondrial and peroxisomal fission. *N Engl J Med* 356:1736–1741
38. Nikolaev A, McLaughlin T, O’Leary DD, Tessier-Lavigne M (2009) APP binds DR6 to trigger axon pruning and neuron death via distinct caspases. *Nature* 457:981–989
39. Wang X, Su B, Zheng L, Perry G, Smith MA, Zhu X (2009) The role of abnormal mitochondrial dynamics in the pathogenesis of Alzheimer’s disease. *J Neurochem* 109(Suppl 1):153–159
40. Hirai K, Aliev G, Nunomura A et al (2001) Mitochondrial abnormalities in Alzheimer’s disease. *J Neurosci* 21:3017–3023
41. Wang X, Su B, Siedlak SL et al (2008) Amyloid-beta overproduction causes abnormal mitochondrial dynamics via differential modulation of mitochondrial fission/fusion proteins. *Proc Natl Acad Sci USA* 105:19318–19323
42. Quintanilla RA, Matthews-Roberson TA, Dolan PJ, Johnson GV (2009) Caspase-cleaved tau expression results in mitochondrial dysfunction in cortical neurons. Implications for the pathogenesis of Alzheimer disease. *J Biol Chem* 284:18754–18766
43. Van Laar VS, Berman SB (2009) Mitochondrial dynamics in Parkinson’s disease. *Exp Neurol* 218:247–256
44. Exner N, Treske B, Paquet D et al (2007) Loss-of-function of human PINK1 results in mitochondrial pathology and can be rescued by parkin. *J Neurosci* 27:12413–12418
45. Greene JC, Whitworth AJ, Kuo I, Andrews LA, Feany MB, Pallanck LJ (2003) Mitochondrial pathology and apoptotic muscle degeneration in *Drosophila* parkin mutants. *Proc Natl Acad Sci USA* 100:4078–4083
46. Lutz AK, Exner N, Fett ME et al (2009) Loss of parkin or PINK1 function increases DRP1-dependent mitochondrial fragmentation. *J Biol Chem* 284:22938–22951
47. Poole AC, Thomas RE, Andrews LA, McBride HM, Whitworth AJ, Pallanck LJ (2008) The PINK1/Parkin pathway regulates mitochondrial morphology. *Proc Natl Acad Sci USA* 105:1638–1643
48. Dagda RK, Cherra SJ III, Kulich SM, Tandon A, Park D, Chu CT (2009) Loss of PINK1 function promotes mitophagy through effects on oxidative stress and mitochondrial fission. *J Biol Chem* 284:13843–13855
49. Nowak L, Bregestovski P, Ascher P, Herbet A, Prochiantz A (1984) Magnesium gates glutamate-activated channels in mouse central neurones. *Nature* 307:462–465
50. Mayer ML, Westbrook GL, Guthrie PB (1984) Voltage-dependent block by Mg^{2+} of NMDA responses in spinal cord neurones. *Nature* 309:261–263
51. Dawson VL, Dawson TM, London ED, Bredt DS, Snyder SH (1991) Nitric oxide mediates glutamate neurotoxicity in primary cortical cultures. *Proc Natl Acad Sci USA* 88:6368–6371
52. Lafon-Cazal M, Pietri S, Culcasi M, Bockaert J (1993) NMDA-dependent superoxide production and neurotoxicity. *Nature* 364:535–537
53. Bonfoco E, Krainc D, Ankarcrona M, Nicotera P, Lipton SA (1995) Apoptosis and necrosis: two distinct events induced, respectively, by mild and intense insults with N-methyl-D-aspartate or nitric oxide/superoxide in cortical cell cultures. *Proc Natl Acad Sci USA* 92:7162–7166
54. Budd SL, Tanneti L, Lishnak T, Lipton SA (2000) Mitochondrial and extramitochondrial apoptotic signaling pathways in cerebrocortical neurons. *Proc Natl Acad Sci USA* 97:6161–6166
55. Hardingham GE, Fukunaga Y, Bading H (2002) Extrasynaptic NMDARs oppose synaptic NMDARs by triggering CREB shut-off and cell death pathways. *Nat Neurosci* 5:405–414
56. Papadia S, Stevenson P, Hardingham NR, Bading H, Hardingham GE (2005) Nuclear Ca^{2+} and the cAMP response element-binding protein family mediate a late phase of activity-dependent neuroprotection. *J Neurosci* 25:4279–4287
57. Papadia S, Soriano FX, Leveille F et al (2008) Synaptic NMDA receptor activity boosts intrinsic antioxidant defenses. *Nat Neurosci* 11:476–487
58. Ankarcrona M, Dypbukt JM, Bonfoco E et al (1995) Glutamate-induced neuronal death: a succession of necrosis or apoptosis depending on mitochondrial function. *Neuron* 15:961–973
59. Shankar GM, Bloodgood BL, Townsend M, Walsh DM, Selkoe DJ, Sabatini BL (2007) Natural oligomers of the Alzheimer amyloid-beta protein induce reversible synapse loss by modulating an NMDA-type glutamate receptor-dependent signaling pathway. *J Neurosci* 27:2866–2875
60. Olney JW (1969) Brain lesions, obesity, and other disturbances in mice treated with monosodium glutamate. *Science* 164:719–721
61. Olney JW, Wozniak DF, Farber NB (1997) Excitotoxic neurodegeneration in Alzheimer disease. New hypothesis and new therapeutic strategies. *Arch Neurol* 54:1234–1240
62. Chen HS, Lipton SA (2006) The chemical biology of clinically tolerated NMDA receptor antagonists. *J Neurochem* 97:1611–1626
63. Abu-Soud HM, Stuehr DJ (1993) Nitric oxide synthases reveal a role for calmodulin in controlling electron transfer. *Proc Natl Acad Sci USA* 90:10769–10772
64. O’Dell TJ, Hawkins RD, Kandel ER, Arancio O (1991) Tests of the roles of two diffusible substances in long-term potentiation: evidence for nitric oxide as a possible early retrograde messenger. *Proc Natl Acad Sci USA* 88:11285–11289
65. Bredt DS, Snyder SH (1994) Nitric oxide: a physiologic messenger molecule. *Annu Rev Biochem* 63:175–195
66. Schuman EM, Madison DV (1994) Locally distributed synaptic potentiation in the hippocampus. *Science* 263:532–536
67. Isaacs AM, Senn DB, Yuan M, Shine JP, Yankner BA (2006) Acceleration of amyloid beta-peptide aggregation by physiological concentrations of calcium. *J Biol Chem* 281:27916–27923
68. Stamler JS, Simon DI, Osborne JA et al (1992) S-Nitrosylation of proteins with nitric oxide: synthesis and characterization of biologically active compounds. *Proc Natl Acad Sci USA* 89:444–448
69. Lei SZ, Pan ZH, Aggarwal SK et al (1992) Effect of nitric oxide production on the redox modulatory site of the NMDA receptor-channel complex. *Neuron* 8:1087–1099
70. Kandel ER, O’Dell TJ (1992) Are adult learning mechanisms also used for development? *Science* 258:243–245
71. Stamler JS, Toone EJ, Lipton SA, Sucher NJ (1997) (S)NO signals: translocation, regulation, and a consensus motif. *Neuron* 18:691–696
72. Hess DT, Matsumoto A, Kim SO, Marshall HE, Stamler JS (2005) Protein S-nitrosylation: purview and parameters. *Nat Rev Mol Cell Biol* 6:150–166
73. Melino G, Bernassola F, Knight RA, Corasaniti MT, Nistico G, Finazzi-Agro A (1997) S-Nitrosylation regulates apoptosis. *Nature* 388:432–433
74. Tanneti L, D’Emilia DM, Lipton SA (1997) Suppression of neuronal apoptosis by S-nitrosylation of caspases. *Neurosci Lett* 236:139–142
75. Dimmeler S, Haendeler J, Nehls M, Zeiher AM (1997) Suppression of apoptosis by nitric oxide via inhibition of interleukin-1beta-converting enzyme (ICE)-like and cysteine

- protease protein (CPP)-32-like proteases. *J Exp Med* 185:601–607
76. Mannick JB, Hausladen A, Liu L et al (1999) Fas-induced caspase denitrosylation. *Science* 284:651–654
 77. Kim WK, Choi YB, Rayudu PV et al (1999) Attenuation of NMDA receptor activity and neurotoxicity by nitroxyl anion, NO. *Neuron* 24:461–469
 78. Choi YB, Tenneti L, Le DA et al (2000) Molecular basis of NMDA receptor-coupled ion channel modulation by S-nitrosylation. *Nat Neurosci* 3:15–21
 79. Gu Z, Kaul M, Yan B et al (2002) S-Nitrosylation of matrix metalloproteinases: signaling pathway to neuronal cell death. *Science* 297:1186–1190
 80. Hara MR, Agrawal N, Kim SF et al (2005) S-nitrosylated GAPDH initiates apoptotic cell death by nuclear translocation following Siah1 binding. *Nat Cell Biol* 7:665–674
 81. Chung KK, Thomas B, Li X et al (2004) S-Nitrosylation of parkin regulates ubiquitination and compromises parkin's protective function. *Science* 304:1328–1331
 82. Yao D, Gu Z, Nakamura T et al (2004) Nitrosative stress linked to sporadic Parkinson's disease: S-nitrosylation of parkin regulates its E3 ubiquitin ligase activity. *Proc Natl Acad Sci USA* 101:10810–10814
 83. Uehara T, Nakamura T, Yao D et al (2006) S-nitrosylated protein-disulphide isomerase links protein misfolding to neurodegeneration. *Nature* 441:513–517
 84. Jaffrey SR, Erdjument-Bromage H, Ferris CD, Tempst P, Snyder SH (2001) Protein S-nitrosylation: a physiological signal for neuronal nitric oxide. *Nat Cell Biol* 3:193–197
 85. Haendeler J, Hoffmann J, Tischler V, Berk BC, Zeiher AM, Dimmeler S (2002) Redox regulatory and anti-apoptotic functions of thioredoxin depend on S-nitrosylation at cysteine 69. *Nat Cell Biol* 4:743–749
 86. Sliskovic I, Raturi A, Mutus B (2005) Characterization of the S-denitrosation activity of protein disulfide isomerase. *J Biol Chem* 280:8733–8741
 87. Stamler JS, Singel DJ, Loscalzo J (1992) Biochemistry of nitric oxide and its redox-activated forms. *Science* 258:1898–1902
 88. Stamler JS (1994) Redox signaling: nitrosylation and related target interactions of nitric oxide. *Cell* 78:931–936
 89. Stamler JS, Lamas S, Fang FC (2001) Nitrosylation: the prototypic redox-based signaling mechanism. *Cell* 106:675–683
 90. Lipton SA, Choi YB, Takahashi H et al (2002) Cysteine regulation of protein function—as exemplified by NMDA-receptor modulation. *Trends Neurosci* 25:474–480
 91. Wang G, Moniri NH, Ozawa K, Stamler JS, Daaka Y (2006) Nitric oxide regulates endocytosis by S-nitrosylation of dynamin. *Proc Natl Acad Sci USA* 103:1295–1300
 92. Kang-Decker N, Cao S, Chatterjee S et al (2007) Nitric oxide promotes endothelial cell survival signaling through S-nitrosylation and activation of dynamin-2. *J Cell Sci* 120:492–501
 93. Stamler JS, Hausladen A (1998) Oxidative modifications in nitrosative stress. *Nat Struct Biol* 5:247–249
 94. Houk KN, Hietbrink BN, Bartberger MD et al (2003) Nitroxyl disulfides, novel intermediates in transnitrosation reactions. *J Am Chem Soc* 125:6972–6976
 95. Huang Z, Huang PL, Panahian N, Dalkara T, Fishman MC, Moskowitz MA (1994) Effects of cerebral ischemia in mice deficient in neuronal nitric oxide synthase. *Science* 265:1883–1885
 96. Iadecola C, Zhang F, Casey R, Nagayama M, Ross ME (1997) Delayed reduction of ischemic brain injury and neurological deficits in mice lacking the inducible nitric oxide synthase gene. *J Neurosci* 17:9157–9164
 97. Hantraye P, Brouillet E, Ferrante R et al (1996) Inhibition of neuronal nitric oxide synthase prevents MPTP-induced parkinsonism in baboons. *Nat Med* 2:1017–1021
 98. Przedborski S, Jackson-Lewis V, Yokoyama R, Shibata T, Dawson VL, Dawson TM (1996) Role of neuronal nitric oxide in 1-methyl-4-phenyl-1,2,3,6-tetrahydropyridine (MPTP)-induced dopaminergic neurotoxicity. *Proc Natl Acad Sci USA* 93:4565–4571
 99. Liberatore GT, Jackson-Lewis V, Vukosavic S et al (1999) Inducible nitric oxide synthase stimulates dopaminergic neurodegeneration in the MPTP model of Parkinson disease. *Nat Med* 5:1403–1409
 100. Chabrier PE, Demerle-Pallardy C, Auguet M (1999) Nitric oxide synthases: targets for therapeutic strategies in neurological diseases. *Cell Mol Life Sci* 55:1029–1035
 101. Lipton SA, Nakamura T, Yao D, Shi ZQ, Uehara T, Gu Z (2005) Comment on “S-nitrosylation of parkin regulates ubiquitination and compromises parkin's protective function”. *Science* 308:1870; author reply 1870
 102. Kitada T, Asakawa S, Hattori N et al (1998) Mutations in the parkin gene cause autosomal recessive juvenile parkinsonism. *Nature* 392:605–608
 103. Shimura H, Hattori N, Kubo S et al (2000) Familial Parkinson disease gene product, parkin, is a ubiquitin-protein ligase. *Nat Genet* 25:302–305
 104. Cookson MR (2005) The biochemistry of Parkinson's disease. *Annu Rev Biochem* 74:29–52
 105. Lim KL, Chew KC, Tan JM et al (2005) Parkin mediates non-classical, proteasomal-independent ubiquitination of synphilin-1: implications for Lewy body formation. *J Neurosci* 25:2002–2009
 106. Wang C, Ko HS, Thomas B et al (2005) Stress-induced alterations in parkin solubility promote parkin aggregation and compromise parkin's protective function. *Hum Mol Genet* 14:3885–3897
 107. Wong ES, Tan JM, Wang C et al (2007) Relative sensitivity of parkin and other cysteine-containing enzymes to stress-induced solubility alterations. *J Biol Chem* 282:12310–12318
 108. LaVoie MJ, Cortese GP, Ostaszewski BL, Schlossmacher MG (2007) The effects of oxidative stress on parkin and other E3 ligases. *J Neurochem* 103:2354–2368
 109. Lyles MM, Gilbert HF (1991) Catalysis of the oxidative folding of ribonuclease A by protein disulfide isomerase: dependence of the rate on the composition of the redox buffer. *Biochemistry* 30:613–619
 110. Hu BR, Martone ME, Jones YZ, Liu CL (2000) Protein aggregation after transient cerebral ischemia. *J Neurosci* 20:3191–3199
 111. Conn KJ, Gao W, McKee A et al (2004) Identification of the protein disulfide isomerase family member PDIP in experimental Parkinson's disease and Lewy body pathology. *Brain Res* 1022:164–172
 112. Rao RV, Bredesen DE (2004) Misfolded proteins, endoplasmic reticulum stress and neurodegeneration. *Curr Opin Cell Biol* 16:653–662
 113. Atkin JD, Farg MA, Turner BJ et al (2006) Induction of the unfolded protein response in familial amyotrophic lateral sclerosis and association of protein-disulfide isomerase with superoxide dismutase 1. *J Biol Chem* 281:30152–30165
 114. Tanaka S, Uehara T, Nomura Y (2000) Up-regulation of protein-disulfide isomerase in response to hypoxia/brain ischemia and its protective effect against apoptotic cell death. *J Biol Chem* 275:10388–10393
 115. Ko HS, Uehara T, Nomura Y (2002) Role of ubiquilin associated with protein-disulfide isomerase in the endoplasmic reticulum in stress-induced apoptotic cell death. *J Biol Chem* 277:35386–35392
 116. Hetz C, Russelakis-Carneiro M, Walchli S et al (2005) The disulfide isomerase Grp58 is a protective factor against prion neurotoxicity. *J Neurosci* 25:2793–2802

117. Chung KK, Zhang Y, Lim KL et al (2001) Parkin ubiquitinates the alpha-synuclein-interacting protein, synphilin-1: implications for Lewy-body formation in Parkinson disease. *Nat Med* 7:1144–1150
118. Cleeter MW, Cooper JM, Darley-Usmar VM, Moncada S, Schapira AH (1994) Reversible inhibition of cytochrome c oxidase, the terminal enzyme of the mitochondrial respiratory chain, by nitric oxide. Implications for neurodegenerative diseases. *FEBS Lett* 345:50–54
119. Clementi E, Brown GC, Feelisch M, Moncada S (1998) Persistent inhibition of cell respiration by nitric oxide: crucial role of S-nitrosylation of mitochondrial complex I and protective action of glutathione. *Proc Natl Acad Sci USA* 95:7631–7636
120. Bossy-Wetzel E, Lipton SA (2003) Nitric oxide signaling regulates mitochondrial number and function. *Cell Death Differ* 10:757–760
121. Yuan H, Gerencser AA, Liot G et al (2007) Mitochondrial fission is an upstream and required event for bax foci formation in response to nitric oxide in cortical neurons. *Cell Death Differ* 14:462–471
122. Wasiak S, Zunino R, McBride HM (2007) Bax/Bak promote sumoylation of DRP1 and its stable association with mitochondria during apoptotic cell death. *J Cell Biol* 177:439–450
123. Breckenridge DG, Kang BH, Kokel D, Mitani S, Staehelin LA, Xue D (2008) *Caenorhabditis elegans* drp-1 and fis-2 regulate distinct cell-death execution pathways downstream of ced-3 and independent of ced-9. *Mol Cell* 31:586–597
124. Karbowski M, Neutzner A, Youle RJ (2007) The mitochondrial E3 ubiquitin ligase MARCH5 is required for Drp1 dependent mitochondrial division. *J Cell Biol* 178:71–84
125. Nakamura N, Kimura Y, Tokuda M, Honda S, Hirose S (2006) MARCH-V is a novel mitofusin 2- and Drp1-binding protein able to change mitochondrial morphology. *EMBO Rep* 7:1019–1022
126. Yonashiro R, Ishido S, Kyo S et al (2006) A novel mitochondrial ubiquitin ligase plays a critical role in mitochondrial dynamics. *EMBO J* 25:3618–3626
127. Chang CR, Blackstone C (2007) Cyclic AMP-dependent protein kinase phosphorylation of Drp1 regulates its GTPase activity and mitochondrial morphology. *J Biol Chem* 282:21583–21587
128. Cribbs JT, Strack S (2007) Reversible phosphorylation of Drp1 by cyclic AMP-dependent protein kinase and calcineurin regulates mitochondrial fission and cell death. *EMBO Rep* 8:939–944
129. Taguchi N, Ishihara N, Jofuku A, Oka T, Mihara K (2007) Mitotic phosphorylation of dynamin-related GTPase Drp1 participates in mitochondrial fission. *J Biol Chem* 282:11521–11529
130. Low HH, Lowe J (2006) A bacterial dynamin-like protein. *Nature* 444:766–769
131. Zhu PP, Patterson A, Stadler J, Seeburg DP, Sheng M, Blackstone C (2004) Intra- and intermolecular domain interactions of the C-terminal GTPase effector domain of the multimeric dynamin-like GTPase Drp1. *J Biol Chem* 279:35967–35974
132. Ramachandran R, Surka M, Chappie JS et al (2007) The dynamin middle domain is critical for tetramerization and higher-order self-assembly. *EMBO J* 26:559–566
133. Pitts KR, McNiven MA, Yoon Y (2004) Mitochondria-specific function of the dynamin family protein DLP1 is mediated by its C-terminal domains. *J Biol Chem* 279:50286–50294
134. Gu M, Gash MT, Mann VM, Javoy-Agid F, Cooper JM, Schapira AH (1996) Mitochondrial defect in Huntington's disease caudate nucleus. *Ann Neurol* 39:385–389
135. Panov AV, Gutekunst CA, Leavitt BR et al (2002) Early mitochondrial calcium defects in Huntington's disease are a direct effect of polyglutamines. *Nat Neurosci* 5:731–736
136. Wang H, Lim PJ, Karbowski M, Monteiro MJ (2009) Effects of overexpression of huntingtin proteins on mitochondrial integrity. *Hum Mol Genet* 18:737–752
137. Liot G, Bossy B, Lubitz S, Kushnareva Y, Sejbuk N, Bossy-Wetzel E (2009) Complex II inhibition by 3-NP causes mitochondrial fragmentation and neuronal cell death via an NMDA- and ROS-dependent pathway. *Cell Death Differ* 16:899–909
138. Bossy-Wetzel E, Petrilli A, Knott AB (2008) Mutant huntingtin and mitochondrial dysfunction. *Trends Neurosci* 31:609–616