

# Lawrence Berkeley National Laboratory

## Recent Work

### Title

POSITRON SPECTRUM FROM THE DECAY OF THE U-MESON

### Permalink

<https://escholarship.org/uc/item/9195q24w>

### Authors

Sagane, Ryokichi  
Dudziak, Walter F.  
Vedder, James.

### Publication Date

1954-05-25

UNCLASSIFIED

UNIVERSITY OF  
CALIFORNIA

*Radiation  
Laboratory*

TWO-WEEK LOAN COPY

This is a Library Circulating Copy  
which may be borrowed for two weeks.  
For a personal retention copy, call  
Tech. Info. Division, Ext. 5545

BERKELEY, CALIFORNIA

## **DISCLAIMER**

This document was prepared as an account of work sponsored by the United States Government. While this document is believed to contain correct information, neither the United States Government nor any agency thereof, nor the Regents of the University of California, nor any of their employees, makes any warranty, express or implied, or assumes any legal responsibility for the accuracy, completeness, or usefulness of any information, apparatus, product, or process disclosed, or represents that its use would not infringe privately owned rights. Reference herein to any specific commercial product, process, or service by its trade name, trademark, manufacturer, or otherwise, does not necessarily constitute or imply its endorsement, recommendation, or favoring by the United States Government or any agency thereof, or the Regents of the University of California. The views and opinions of authors expressed herein do not necessarily state or reflect those of the United States Government or any agency thereof or the Regents of the University of California.

UNIVERSITY OF CALIFORNIA

Radiation Laboratory

Contract No. W-7405-eng-48

POSITRON SPECTRUM FROM THE DECAY OF THE  $\mu$ -MESON  
Ryokichi Sagane, Walter F. Dudziak, and James Vedder

May 25, 1954

Berkeley, California

POSITRON SPECTRUM FROM THE DECAY OF THE  $\mu$ -MESON

Ryokichi Sagane, Walter F. Dudziak, and James Vedder

Radiation Laboratory, Department of Physics  
University of California, Berkeley, California

May 25, 1954

The positron spectrum (Fig. 1) produced by the decay of the  $\mu$ -meson has been studied in detail, and also with much improved accuracy, with a 40 in. (pole base diameter) spiral-orbit spectrometer.

The experimental method that was adopted is similar to the one reported by one of us in 1951.<sup>1</sup> The 340-Mev deflected proton beam was used to produce  $\pi^+$ -mesons in the target. Some of the created  $\pi^+$ -mesons decayed inside the target into  $\mu$ -mesons that in turn disintegrated into positrons. Thus the target mounted coaxially with the symmetric magnetic field was the source of positrons. The energy spectrum of these created positrons was analyzed by means of the spiral-orbit spectrometer.<sup>2</sup>

Positrons were measured with momentum resolutions of  $\pm 0.6 \sim \pm 1.8\%$  at half intensity. They were detected as quadruple coincidences of signals from four plastic crystals (two of these crystals were  $1/4$  by 1.5 by 2.5 in., and the other two,  $1/4$  by 3 by 3 in.). The counts were taken following a 2- $\mu$  sec delay relative to the proton pulses and with four consecutive gates each having a 2- $\mu$  sec width. This enabled us not only to check the half life but also to reduce background, which usually came in as accidentals.




Fig. 1. Positron Spectrum from "Thick"-Target Data.

-4-

Measurements were made on four thin Be targets of different diameters and thicknesses, as the resolution is directly related to the effective diameter of the cylindrical or tubular targets, and also as the energy absorption of the positrons through the target itself is the other major correction to be applied. In addition to these, two thick targets (C and Al, each 1 1/2-in. outside diameter by 4-in. long) were studied. Each experiment was repeated from 4 to 10 times, and reproducibility of the results was checked.

The accuracy of each magnetic field setting was kept better than 0.1% by means of a proton nuclear resonance method during each measurement. As a result, the accuracy of the absolute  $H\rho$  value of each measured point is considered to be better than 0.2%.

The results are summarized as follows:

- (A) All the data (most of which have a reasonably small statistical error) can be fitted best to the theoretical curve with

$$\rho = 0.23 \begin{array}{l} + 0.03 \\ - 0.05 \end{array}$$

(the constant introduced by Dr. Michel<sup>3</sup>), if proper corrections for absorption and resolution are applied, as is shown in Fig. 1.

- (B) As illustrated in Fig. 2, all thin-target experiments show a definite sharp cutoff at the energy maximum.
- (C) The absolute value of this maximum energy has been calculated as  $52.8 \pm 0.2$  Mev.
- (D) This corresponds to a  $\mu$ -meson mass value of  $m_{\mu} = 207 \pm 0.8 m_e$ .




Fig. 2. Behavior of Positron Spectrum Near  $E_{max}$  from Four "Thin" Be Targets.



-6-

- (E) With the aid of the fundamental mass ratio<sup>7</sup>  $\frac{\pi^+}{\mu^+} = 1.321 \pm 0.002$ , the mass of the  $\pi^+$ -meson can be established. This value is

$$m_{\pi^+} = 273.4 \pm 1.1 m_e.$$

- (F) The data obtained in 1951<sup>1</sup> were found in good agreement with the present data. (Unfortunately, the mass value of the  $\mu$ -meson at that time was too high, and the results, therefore, were interpreted to show  $\rho = 0$ .)
- (G) Since we have measured directly the cutoff at  $E_{\max}$  from our experiment, our present results are not dependent on some other mass measurements.
- (H) The very good agreement of the mass value of the  $\mu$ -meson obtained with the assumption of the range energy relation<sup>7</sup> gives strong indication that the assumed relation is good for mesons to this order of accuracy.
- (I) From our present data, we are not able to exclude the possibility of the spin values of  $3/2$  for the  $\mu$ -meson.

We have been informed of the values  $\rho = 0.6, 0.5$  and  $\sim 0$ , obtained in the recent measurements (each with a different method) from Columbia,<sup>4</sup> Massachusetts Institute of Technology,<sup>5</sup> and Los Alamos.<sup>6</sup> As the discrepancies of the  $\rho$  values are quite serious, during the last five months, we have made a close examination of the possible systematic errors that we might have overlooked. Our present analysis of our data cannot account for this serious difference in  $\rho$  value.

We would like to acknowledge the interest, and particularly the effort in obtaining the funds for the construction of the 40-inch spiral-orbit spectrometer, of Professor E. O. Lawrence and Dr. Walter Barkas.

## REFERENCES

1. R. Sagane, W. Gardner and H. Hubbard, Phys. Rev. 82, 557, (1951).
2. Miyamoto, Proc. Phys. Math. Soc. Japan 24, 676 (1942).
3. L. Michel, Proc. Phys. Soc. A. 63, 514 (1950).
4. Lederman and Sargent, private communication.
5. J. H. Vilain and R. W. Williams, Phys. Rev. 92, 1586 (1953).
6. F. Harrison, C. Cowan, Jr. and F. Reines, Nucleonics 12, 3, 44 (1954).
7. F. M. Smith, W. Birnbaum, and W. Barkas, Phys. Rev. 91, 765 (1953).