

UC San Diego

UC San Diego Previously Published Works

Title

Regulation of epidermal growth factor receptor degradation by heterotrimeric G alpha s protein

Permalink

<https://escholarship.org/uc/item/91p9443d>

Journal

Molecular Biology of the Cell, 15(12)

ISSN

1059-1524

Authors

Bin, Z
Lavoie, C
Tang, T D
[et al.](#)

Publication Date

2004-12-01

Peer reviewed

Regulation of Epidermal Growth Factor Receptor Degradation by Heterotrimeric G α s Protein

Bin Zheng,* Christine Lavoie,* Ting-Dong Tang, Phuong Ma, Timo Meerloo, Anthony Beas, and Marilyn G. Farquhar[†]

Department of Cellular and Molecular Medicine, University of California, San Diego, La Jolla, CA 92093-0651

Submitted June 3, 2004; Revised September 15, 2004; Accepted September 17, 2004
Monitoring Editor: Juan S. Bonifacino

Heterotrimeric G proteins have been implicated in the regulation of membrane trafficking, but the mechanisms involved are not well understood. Here, we report that overexpression of the stimulatory G protein subunit (G α s) promotes ligand-dependent degradation of epidermal growth factor (EGF) receptors and Texas Red EGF, and knock-down of G α s expression by RNA interference (RNAi) delays receptor degradation. We also show that G α s and its GTPase activating protein (GAP), RGS-PX1, interact with hepatocyte growth factor-regulated tyrosine kinase substrate (Hrs), a critical component of the endosomal sorting machinery. G α s coimmunoprecipitates with Hrs and binds Hrs in pull-down assays. By immunofluorescence, exogenously expressed G α s colocalizes with myc-Hrs and GFP-RGS-PX1 on early endosomes, and expression of either Hrs or RGS-PX1 increases the localization of G α s on endosomes. Furthermore, knock-down of both Hrs and G α s by double RNAi causes greater inhibition of EGF receptor degradation than knock-down of either protein alone, suggesting that G α s and Hrs have cooperative effects on regulating EGF receptor degradation. These observations define a novel regulatory role for G α s in EGF receptor degradation and provide mechanistic insights into the function of G α s in endocytic sorting.

INTRODUCTION

Heterotrimeric G proteins serve as important molecular switches that relay extracellular signals from G protein-coupled receptors (GPCRs) on the cell membrane to downstream effectors (Gilman, 1987; Neves *et al.*, 2002). Besides their plasma membrane location, heterotrimeric G proteins also are found on membranes of intracellular compartments along both the endocytic and secretory pathways where indirect evidence suggests they play several roles in membrane trafficking (Bomsel and Mostov, 1992; Helms, 1995; Nurnberg and Ahnert-Hilger, 1996; Stow and Heimann, 1998). One of the prototypical heterotrimeric G proteins, G α s, the stimulatory subunit of heterotrimeric G proteins, has been suggested to regulate endocytic trafficking. Reagents that activate G α s, e.g., cholera toxin and a peptide mimicking the interacting region of G α s with the β 2-adrenergic receptor, block endosome-endosome and phagosome-endosome fusion *in vitro* (Colombo *et al.*, 1992, 1994; Beron *et al.*, 1995). Cholera toxin and recombinant G α s proteins also have been found to promote transcytosis of the poly-

meric IgA receptor through endosomes in polarized epithelial cells (Bomsel and Mostov, 1993).

Although the molecular basis for the function of G α s in signal transduction at the plasma membrane has been well characterized, little is known about the mechanisms whereby G α s influences endocytic trafficking. Our recent discovery of RGS-PX1 has provided a putative link between G α s and endocytic trafficking (Zheng *et al.*, 2001). RGS-PX1, a member of the regulator of G protein signaling (RGS) protein family (De Vries *et al.*, 2000; Hollinger and Hepler, 2002), functions as a GTPase activating protein (GAP) for G α s through its conserved RGS domain that interacts specifically with G α s, but no other G α protein (Zheng *et al.*, 2001). RGS-PX1 is also known as sorting nexin 13 (SNX13) and serves as an SNX protein, through its phosphatidylinositol-binding phoX (PX) domain. This domain is shared by SNX proteins that are involved in protein sorting in endosomes (Haft *et al.*, 1998; Worry and Dixon, 2002). We showed previously that RGS-PX1 is a functional SNX protein that is localized on endosomes and delays epidermal growth factor (EGF) receptor degradation, probably at the steps of endosome sorting and lysosome targeting (Zheng *et al.*, 2001). The fact that RGS-PX1 can bind G α s and affect EGF receptor trafficking suggested that G α s also might be involved in regulating of EGF receptor endocytosis and down-regulation.

The EGF receptor represents the classical model system to study mechanisms of ligand-induced receptor endocytosis and down-regulation in mammalian cells (Carpenter, 2000; Katzmann *et al.*, 2002; Sorkin and Von Zastrow, 2002). On ligand binding, EGF receptors are rapidly internalized via clathrin-coated pits and delivered to early endosomes where the majority of the receptors are sorted into the luminal vesicles of late endosomes or multivesicular bodies (MVBs) and targeted for degradation in lysosomes (Wishart *et al.*,

Article published online ahead of print. Mol. Biol. Cell 10.1091/mbc.E04-06-0446. Article and publication date are available at www.molbiolcell.org/cgi/doi/10.1091/mbc.E04-06-0446.

* These authors contributed equally to this work.

[†] Corresponding author. E-mail address: mfarquhar@ucsd.edu.

Abbreviations used: EGF, epidermal growth factor; EGFR, epidermal growth factor receptor; ESCRT, endosomal sorting complexes required for transport; GAP, GTPase activating protein; GPCR, G protein-coupled receptor; Hrs, hepatocyte growth factor-regulated tyrosine kinase substrate; MVB, multivesicular body; PFA, paraformaldehyde; PX, phoX domain; RGS, regulator of G protein signaling; RNAi, RNA interference; siRNA, small-interfering RNA; SNX, sorting nexin.

2001; Katzmann *et al.*, 2002; Sorkin and Von Zastrow, 2002; Stahl and Barbieri, 2002; Raiborg *et al.*, 2003). Hepatocyte growth factor-regulated tyrosine kinase substrate (Hrs) has been demonstrated to be a central player in this sorting pathway (Raiborg and Stenmark, 2002; Clague and Urbe, 2003; Raiborg *et al.*, 2003). Hrs is found in specialized clathrin-coated microdomains of early endosomes that are enriched in mono-ubiquitinated receptors targeted for lysosomal degradation (Raiborg *et al.*, 2001b; Sachse *et al.*, 2002). Hrs also recruits the endosomal sorting complexes required for transport (ESCRT) complexes to endosomes through its interaction with Tsg101 in ESCRT complex I and regulates the formation of MVBs (Bache *et al.*, 2003a; Katzmann *et al.*, 2003; Lu *et al.*, 2003). In this study, we have investigated the effects of $G_{\alpha s}$ on EGF receptor degradation. Our findings establish a novel role for $G_{\alpha s}$ in regulating endocytic trafficking and sorting.

MATERIALS AND METHODS

Materials

Mammalian expression vector pcDNA3.1 containing $G_{\alpha s}$ long (L) and short (S) splicing variants were obtained from Guthrie cDNA Resource Center (Sayre, PA). Mammalian expression vector pXER-EGFR encoding the EGF receptor was obtained from Dr. Gordon Gill (University of California, San Diego, CA). pGas-green fluorescent protein (GFP) construct, expressing a $G_{\alpha s}$ -GFP fusion protein with GFP inserted between the helical and GTPase domains, was obtained from Dr. Mark Rasenick (University of Illinois, Chicago, IL) (Yu and Rasenick, 2002). The pcDNA3-myc-Hrs construct was obtained from Dr. A Beans (University of Texas Medical School, Houston, TX). The GFP-tagged RGS-PX1 construct containing residues 257–957 of human RGS-PX1 was described previously (Zheng *et al.*, 2001). The FLAG-tagged RGS-PX1 construct was prepared by inserting the cDNA encoding human RGS-PX1 (residues 51–957) into p3XFLAG-CMV-10 (Sigma-Aldrich, St. Louis, MO).

Antibodies

Affinity-purified rabbit IgG against $G_{\alpha s}$ used for immunoblotting was obtained from Calbiochem (San Diego, CA). Rabbit antibodies against Rab5 were provided by Dr. Angela Wandinger-Ness (University of New Mexico, Albuquerque, NM). Other antibodies were obtained from the following sources: monoclonal antibodies (mAbs) against actin and FLAG (M2) (Sigma-Aldrich), myc (Cell Signaling Technology, Beverly, MA), and GFP (BD Biosciences Clontech, Palo Alto, CA), and polyclonal antibodies against EGF receptor (Santa Cruz Biotechnology, Santa Cruz, CA) and GFP and Hrs (Alexis Biochemicals, San Diego, CA).

Cell Culture and Transfection

Human embryonic kidney (HEK)293T cells (obtained from Dr. Alexandra Newton, University of California, San Diego, CA), and Cos7 cells were maintained in Dulbecco's modified Eagle's high glucose medium (DMEM) (Invitrogen, Carlsbad, CA) containing 10% fetal calf serum (FCS) (Hyclone Laboratories, Logan, UT), penicillin, and streptomycin. HEK293 cells were transfected using calcium phosphate as described previously (Zheng *et al.*, 2000). Cos7 cells were transfected using FuGENE 6 (Roche Diagnostics, Indianapolis, IN) according to the manufacturer's instructions.

Epidermal Growth Factor Receptor (EGFR) Degradation Assays

HEK293 cells in six-well plates were transfected with pXER-EGFR together with pcDNA3.1- $G_{\alpha s}$ -L and pcDNA3.1- $G_{\alpha s}$ -S (1:1), or pcDNA3.1 empty vector. Twenty-four hours after transfection, cells were serum starved overnight in DMEM with 0.5% fetal bovine serum (FBS) and then incubated in the presence or absence of 100 nM EGF (Molecular Probes, Eugene, OR) at 37°C. Cells were lysed in Laemmli sample buffer or in 50 mM Tris, pH 7.6, 150 mM NaCl, 1% Triton, and Complete protease inhibitor (Roche Diagnostics) followed by immunoblotting with antibodies to EGFR, $G_{\alpha s}$, and actin. EGF receptor degradation was quantified by densitometry (three independent experiments) by using Quantity One software (Bio-Rad, Hercules, CA).

Immunofluorescence

Cos7 cells were fixed in 3% paraformaldehyde (PFA) in 100 mM phosphate buffer, pH 7.4, for 30 min, permeabilized with 0.1% Triton X-100 for 10 min, blocked with 10% fetal calf serum for 30 min, and incubated with primary antibodies for 1 h at 25°C, followed by Alexa Fluor 594-conjugated goat




Figure 1. Overexpression of $G_{\alpha s}$ promotes degradation of EGFR in HEK293 cells. (A) HEK293 cells were transfected with pXER-EGFR together with pcDNA3.1- $G_{\alpha s}$ ($G_{\alpha s}$) or control vector (control) for 24 h, serum starved overnight, and then treated with 100 nM EGF for 0 or 60 min, followed by immunoblotting with antibodies against EGF receptor, actin, or $G_{\alpha s}$. $G_{\alpha s}$ is seen as two bands representing the long and short forms of $G_{\alpha s}$. Data shown are representative of at least three independent experiments. (B) Quantification of EGF receptor degradation. Results from three independent experiments were analyzed by Quantity One software (Bio-Rad). When cells transfected with control vectors are stimulated with EGF, ~50% of the receptors seen at 0 min are degraded by 60 min after adding EGF. In cells transfected with $G_{\alpha s}$, degradation is enhanced as ~80% of the receptors are degraded by 60 min. Data presented as percentage of total EGF receptor at 0 min in control cells.

anti-mouse F(ab')₂ and/or Alexa 488 goat anti-rabbit F(ab')₂ (Molecular Probes) for 1 h. Some specimens were permeabilized with 0.05% saponin for 1 min at 4°C before fixation. Specimens were analyzed using a Zeiss Axiophot equipped with a Hamamatsu Orca ER charge-coupled device (CCD) or by deconvolution microscopy by using an Applied Precision (Issaquah, WA) Delta Vision imaging system coupled to an S100 fluorescence microscope (Carl Zeiss, Thornwood, NY). For cross-sectional images of cells, stacks were obtained with 200-nm step width. Deconvolution was done on an SGI workstation (Mountain View, CA) by using Delta Vision reconstitution software, and images were processed as Tiff files by using Photoshop 7.0 (Adobe Systems, San Jose, CA).

Uptake of Texas Red EGF

Cos7 cells were transfected with pcDNA3, $G_{\alpha s}$ -GFP, or myc-Hrs for 12 h. After serum starvation for 3 h, cells were incubated in DMEM containing 0.4 μ g/ml Texas Red EGF (Molecular Probes) in 0.5% FBS for 10 min at 37°C and washed and incubated in DMEM containing 0.5% FBS for up to 1 h at 37°C.

For semiquantitative analysis of bound and internalized Texas Red EGF, all images were captured with the exact same settings. Control cells and cells




Figure 2. Overexpression of Gas-GFP promotes degradation of Texas Red. (A) Cos7 cells transfected with pGas-GFP or control vector were incubated with Texas Red EGF for 10 min and chased for 30 or 60 min. Cells expressing Gas-GFP (traced in white) and those expressing control vector showed similar levels of Texas Red EGF at 0-min chase. However, after 30- or 60-min chase there is considerably less Texas Red EGF remaining in cells expressing Gas-GFP. (B) Semiquantitative representation of the data shown in A. In cells transfected with control vector ~30% of the Texas Red EGF is degraded at 30 min and 80% by 60 min, whereas in cells expressing Gas-GFP ~70% are degraded at 30 min and ~95% at 60 min. Average integrated intensity of Texas Red EGF pixels per cell were measured as described in *Materials and Methods*. Data are expressed as the mean \pm SE of three experiments.

expressing the GFP constructs were traced using Photoshop. For each cell, the number and intensity of positive pixels (pixels with grayscale values between 75 and 255) was determined using Image J software (National Institutes of Health, Bethesda, MD; <http://rsb.info.nih.gov/ij>). Results were calculated as

the total number of positive pixels per condition multiplied by the cumulative pixel intensity divided by the number of cells. Ten to 30 cell profiles were measured for each condition, and the results are displayed as the mean of three separate experiments.

Immunogold Labeling

HEK293 cells were fixed in 4% PFA alone or 4% PFA containing 0.2% glutaraldehyde in 0.1 M phosphate buffer, pH 7.4, overnight, pelleted in 10% gelatin in phosphate buffer, cryoprotected, and snap frozen in liquid nitrogen. Ultrathin cryosections (70–90 nm) were cut at -100°C on a Leica Ultracut UCT with an EM FCS cryoattachment (Leica, Bannockburn, IL) by using a Diatome diamond knife (Diatome US, Fort Washington, PA), picked up with a 1:1 mixture of 2.3 M sucrose and 2% methyl cellulose (15 cp), and transferred onto Formvar- and carbon-coated copper grids. Sections were blocked and incubated with primary antibodies for 2 h at room temperature, followed by gold conjugated goat anti-rabbit IgG and gold conjugated goat anti-mouse IgG (Amersham Biosciences, Piscataway, NJ) for 1 h. Sections were contrasted for 10 min with 2% neutral uranyl acetate and stained for 10 min with 0.2% uranyl acetate in 1.8% methyl cellulose on ice. Grids were viewed and photographed using a Philips CM-10 transmission electron microscope (FEI, Hillsboro, OR) equipped with a 794 Multiscan CCD camera (Gatan, Pleasanton, CA).

Coimmunoprecipitation

HEK293 cells were plated in 60-mm plates and transfected with various constructs. After 48 h, cells were lysed in 1% Triton X-100 in phosphate-buffered saline (PBS) buffer containing protease inhibitors (0.12 mg/ml phenylmethylsulfonyl fluoride, 2 mg/ml leupeptin, and 1 mg/ml aprotinin) at 4°C for 30 min and centrifuged at $15,000 \times g$ for 10 min. Cell lysates were incubated with primary antibodies overnight at 4°C , followed by incubation with protein A- or G-Sepharose (Oncogene, San Diego, CA) for an additional 1 h at 4°C . Beads were washed (three times) with lysis buffer and boiled in Laemmli sample buffer, and bound immune complexes were analyzed by SDS-PAGE and immunoblotting.

For coimmunoprecipitation of Hrs and *Gas* from cytosolic and membrane fractions, HEK293 cells were scraped into cold PBS containing protease inhibitors and homogenized by 10 passages through a 28 1/2-gauge needle. Nuclei and unbroken cells were removed by centrifugation, and postnuclear supernatants were centrifuged at $100,000 \times g$ for 1 h at 4°C to prepare cytosolic (supernatant, S100) and membrane (pellet, P100) fractions (Zheng *et al.*, 2000). Membrane pellets were lysed in 1% Triton X-100 in PBS containing protease inhibitors for 1 h, centrifuged ($15,000 \times g$ for 10 min), and the membrane lysates and cytosolic fractions were used for immunoprecipitation.

Immunoblotting

Protein samples were separated on 10% SDS-PAGE and transferred to polyvinylidene difluoride membranes (Millipore, Bedford, MA). Membranes were blocked in Tris-buffered saline (20 mM Tris-HCl, pH 7.5, 500 mM NaCl) containing 0.1% Tween 20 and 5% nonfat milk and incubated with primary antibodies for 2 h at room temperature or overnight at 4°C , followed by incubation with horseradish peroxidase-conjugated goat anti-rabbit or anti-mouse IgG (Bio-Rad) and enhanced chemiluminescence detection (Pierce Chemical, Rockford, IL).

In Vitro Glutathione S-Transferase (GST) Pull-Down Assays

Full-length rat Hrs cDNA and a human RGS-PX1 fragment (PXC) containing the PX domain and the C-terminus (residues 526–957) were amplified by polymerase chain reaction (PCR) and subcloned into pGEX-KG (Amersham Biosciences). GST fusion proteins were expressed in *Escherichia coli* BL21 and purified on glutathione-Sepharose 4B (Pharmacia, Piscataway, NJ) beads according to the manufacturer's instructions. ^{35}S -labeled, in vitro translation products of *Gas* or Hrs were prepared by using the TNT T7 rabbit reticulocyte Quick Coupled Transcription/Translation system (Promega, San Luis Obispo, CA) in the presence of [^{35}S]methionine (1000 Ci/mmol, in vivo cell labeling grade; Amersham Biosciences), pcDNA3.1-*Gas*-L and pcDNA3.1-*Gas*-S (1:1 ratio) or pcDNA3-myc-Hrs. For pull-down assays, GST fusion proteins ($\sim 75 \mu\text{g}$) immobilized on beads were incubated with in vitro-translated products in 20 mM Tris-HCl, pH 8.0, 2 mM MgSO_4 , 6 mM β -mercaptoethanol, 5% glycerol, and 0.01% C12E10, in the presence of protease inhibitors for 2 h at 4°C , and washed four times with the same buffer. GST pull-down assays on brain lysates (5 mg) were performed using a lysis buffer containing 20 mM HEPES, pH 7.4, 150 mM NaCl, 300 mM sucrose, 1% Triton X-100, and 0.01% C12E10 as described previously (Zheng *et al.*, 2001). Bound proteins were eluted with Laemmli sample buffer, resolved by SDS-PAGE, and visualized by autoradiography.

RNA Interference

The following small-interfering RNA (siRNA) oligos synthesized by Dharmacon Research (Lafayette, CO) were used for RNAi knock-down of *Gas* and Hrs (Bache *et al.*, 2003b): *Gas*-sense, 5'-GGC GCA GCG UGA GGC CAA CdTdT; *Gas*-antisense, 5'-GUU GGC CUC ACG CUG CGC CdTdT; and Hrs-sense, 5'-CGA CAA GAA CCC ACA CGU CdTdT; Hrs-antisense, 5'-GAC GUG UGG GUU CUU GUC GdTdT. All oligos were designed based on human sequences. Scrambled RNA oligos (scramble II duplex; Dharmacon

Research) were used as controls. Cos7 cells in six-well plates (30% confluent; 1.5 ml of normal culture medium without antibiotics per well) were transfected with 1 μl of 75 μM siRNA duplex and 8 μl of Oligofectamine reagent (Invitrogen) according to the manufacturer's instructions. The cells were analyzed 72 h after transfection. For double RNAi experiments, the total RNAi oligos were kept the same among different wells by addition of scrambled RNAi oligos.

RESULTS

Gas Overexpression Promotes Degradation of EGF Receptors and Texas Red EGF

RGS-PX1 acts both as a GAP that regulates the activity of *Gas* and as a SNX involved in the down-regulation of the EGF receptor (Zheng *et al.*, 2001). These dual activities suggested that RGS-PX1 could link *Gas* to EGF receptor sorting at endosomes and that *Gas* also might be involved in EGF




Figure 3. Depletion of *Gas* expression by RNAi delays degradation of the EGF receptor. Cos7 cells were transfected with 37.5 nM control or *Gas*-specific siRNA oligos by using Oligofectamine. After 3 d, cells were treated with 100 nM EGF for 60 min, lysed, and analyzed by immunoblotting with antibodies against EGF receptor, *Gas*, and actin. The level of *Gas* is reduced to $<5\%$ of control levels in cells transfected with oligos specific for *Gas*. In cells transfected with control siRNA, 70% of the EGF receptors are degraded after 60-min stimulation with EGF. In cells transfected with *Gas* siRNA, degradation is delayed as only 34% of the receptors are degraded after 60 min. Data presented as percentages of the amount of EGFR at 0 min in each group of cells. Data shown are representative of at least three independent experiments. (B) Quantification of EGF receptor degradation. Results from three independent experiments were analyzed by Quantity One software. Data presented as percent of total EGFR present at 0 min in control cells.




Figure 4. RGS-PX1 interacts with Hrs. (A) In vitro translated, ^{35}S -labeled Hrs binds to GST-RGS-PX1(PXC) but not to GST alone. GST-RGS-PX1(PXC) and GST alone ($\sim 75 \mu\text{g}$ each) immobilized on glutathione beads were incubated with in vitro-translated Hrs. Bound proteins were separated by SDS-PAGE and detected by autoradiography. Input equals 3% of total in vitro translation product. (B) Myc-Hrs (top, lane 4) coimmunoprecipitates with FLAG-RGS-PX1 in cells transfected with both proteins but not in those transfected with myc-Hrs alone (top, lane 3). HEK293 cells were transfected with FLAG-RGS-PX1 and myc-Hrs or myc-Hrs alone, and immunoprecipitation (IP) was carried out on lysates (lanes 1 and 2) with anti-FLAG mouse IgG, followed by immunoblotting (IB) of immunoprecipitates with anti-myc. (C) FLAG-RGS-PX1 coimmunoprecipitates with myc-Hrs. Immunoprecipitation was carried out with anti-myc on lysates from HEK293 cells transfected with myc-Hrs alone (lane 1), FLAG-RGS-PX1 alone (lane 3), or both FLAG-RGS-PX1 and myc-Hrs (lane 2), followed by immunoblotting with anti-FLAG IgG.

receptor down-regulation. To find out whether this is the case, we transiently transfected HEK293 cells with EGF receptor and either *Gas* or control vector and determined the kinetics of EGF receptor degradation. As shown in Figure 1, A and B, cells transfected with *Gas* contained less EGF receptors ($\sim 20\%$) at steady state than control cells, suggesting *Gas* expression enhances basal turnover of EGFR. Similarly, ligand-induced degradation of the receptor was enhanced in cells transfected with *Gas*, because 80% of the receptors were degraded by 60 min after adding EGF (Figure 1, A and B), whereas in cells transfected with control vector, only 50% of the receptors had been degraded.

Next, we used immunofluorescence to evaluate the effects of overexpressing *Gas* on the uptake and degradation of Texas Red EGF. Cos7 cells transfected with *Gas*-GFP (Yu and Rasenick, 2002) or empty vector were incubated with Texas Red EGF for 10 min followed by incubation in the absence of ligand for 30 or 60 min. As shown in Figure 2A, the levels of Texas Red EGF were similar in cells transfected with *Gas*-GFP and control vector after 10-min incubation with Texas Red EGF, suggesting that *Gas* overexpression does not impair internalization of EGF. However, at 30 and 60 min "chase," cells expressing *Gas*-GFP contained significantly less Texas Red EGF than those transfected with empty vector (Figure 2A) or GFP alone (our unpublished data). Semiquantitative analysis of the amount of Texas Red EGF remaining (Figure 2B) revealed that in cells transfected with control vector, 66% remained at 30 min and 20% at 60 min, whereas only 30 and 7% remained at the same times in cell expressing *Gas*-GFP. As a control, we also transfected

Hrs into Cos7 cells and found, consistent with previous reports (Raiborg *et al.*, 2001b; Bishop *et al.*, 2002; Urbe *et al.*, 2003), that overexpression of Hrs strongly inhibited Texas Red EGF degradation (our unpublished data). We also examined the effect of *Gas* overexpression on the uptake of transferrin-Alexa594 in Cos7 cells and found no difference in transferrin uptake between cells transfected with *Gas*-GFP and GFP alone (our unpublished data). Together, these results indicate that overexpression of *Gas* promotes specific degradation of EGF receptors and their ligands.

Depletion of *Gas* Delays Degradation of EGF Receptors

We further evaluated the effects of knocking down endogenous *Gas* protein levels in Cos7 cells on ligand-induced degradation of EGF receptors. We found that siRNA oligos designed specifically for *Gas* blocked EGF-dependent receptor degradation (Figure 3, A and B). In cells transfected with scrambled siRNA, 70% of the receptors were degraded after 60-min stimulation with EGF, whereas in cells transfected with *Gas* siRNA degradation was delayed as only 34% of the EGFR had been degraded (Figure 3, A and B). In addition, cells transfected with *Gas* siRNA had higher levels (140%) of EGF receptor at steady state than cells transfected with scrambled siRNA (Figure 3, A and B), suggesting the basal level of EGF receptor turnover is delayed by knockdown of *Gas*. Thus, we have shown by three different approaches that *Gas* regulates EGF receptor degradation.




Figure 5. Interaction between Hrs and Gas. (A) In vitro-translated, ^{35}S -labeled Gas binds to GST-Hrs but not to GST alone. GST-Hrs and GST proteins ($\sim 75 \mu\text{g}$ each) immobilized on glutathione beads were incubated with in vitro-translated, [^{35}S]Gas as in Figure 4. Input equals 3% of total in vitro translation product. (B) Endogenous Gas from rat brain lysates binds to GST-Hrs but not to GST. GST-Hrs and GST immobilized on glutathione beads were incubated with rat brain lysates ($\sim 5 \text{ mg}$). Bound proteins were immunoblotted with anti-Gαs IgG. Input equals 3% of total brain lysate. (C) Myc-Hrs coimmunoprecipitates with Gas-GFP (lane 4). Lysates (lanes 1 and 2) from HEK293 cells transfected with Gas-GFP or GFP together with myc-Hrs were immunoprecipitated with anti-GFP, followed by immunoblotting with anti-myc and anti-GFP antibodies. (D) Gas and Hrs are found in approximately equal amounts in both membrane (P100, lane 2) and cytosolic (S100, lane 1) fractions. Gas coimmunoprecipitates with myc-Hrs predominantly ($>95\%$) from membrane fractions (lane 4, bottom). Very little Gas is coprecipitated with myc-Hrs from the cytosolic fraction (lane 3, bottom). Cytosolic (S100, lane 1) and membrane (P100, lane 2) fractions prepared from HEK293 cells transfected with Gas and myc-Hrs were immunoprecipitated with anti-myc (myc, lanes 3 and 4) or control (ctrl, lanes 5 and 6) mouse IgGs, followed by immunoblotting with anti-Gαs and anti-myc antibodies.

RGS-PX1 Interacts with Hrs In Vivo and In Vitro

Next, we investigated whether RGS-PX1 or Gas delays EGF receptor degradation by interacting with components of the endosomal sorting machinery. We reasoned that RGS-PX1 might bind Hrs, an endosomal protein required for efficient degradation of EGF receptors, because Hrs has been shown to interact with SNX1 (Chin *et al.*, 2001), the founding member of the SNX protein family that shares strong sequence homology with the C-terminal PX domain and coiled-coil region of RGS-PX1 (Kurten *et al.*, 1996; Zheng *et al.*, 2001).

When we incubated ^{35}S -labeled, in vitro-translated Hrs with GST-RGS-PX1(PXC), a GST fusion protein containing the PX domain and C-terminal coiled-coil region of RGS-PX1 that is homologous to SNX1, Hrs bound to GST-RGS-PX1(PXC), but not to GST alone (Figure 4A). We further tested whether RGS-PX1 coimmunoprecipitates with Hrs in HEK293 cells transfected with myc-tagged Hrs and FLAG-tagged RGS-PX1. We found that when immunoprecipitation was carried out with anti-FLAG IgG, myc-Hrs coprecipitated with FLAG-RGS-PX1 (Figure 4B). Similarly, when anti-myc IgG was used, myc-Hrs coprecipitated with FLAG-




Figure 6. Colocalization of Gas-GFP with myc-Hrs on early endosomes. (A–C) In Cos7 cells transfected with Gas-GFP alone, Gas is distributed on the plasma membrane (arrow, A) and on small vesicular structures (arrowheads, A). Hrs is distributed on early endosomes throughout the cell (B). Merged image (yellow) shows occasional overlap in the vesicular distribution of Gas-GFP and Hrs (arrowheads and inset, C). (D–I) In cells transfected with Myc-Hrs, which promotes formation of large, clustered endosomes Gas-GFP is distributed on the plasma membrane (arrow, D) and on the enlarged endosomes (arrowheads and inset, D and G). Myc-Hrs (arrowheads and inset, E) and Rab5 (arrowheads and inset, H) are also present on these endosomes. Gas-GFP colocalizes (yellow) with Myc-Hrs (arrowheads and inset, F) and Rab5 (arrowhead and inset, I). Cos7 cells were transfected with Gas-GFP alone (A–C) or together with Myc-Hrs (D–I) and permeabilized with saponin before fixation to release the cytosolic proteins and facilitate the detection of membrane-associated pools of Gas and Hrs. Cells were then fixed with 3% PFA, permeabilized, and double labeled with mouse anti-GFP (A, D, and G), anti-Hrs (B), or anti-myc (E) IgG or rabbit anti-rab5 (H) IgG and analyzed by deconvolution immunofluorescence microscopy. Bar, 2 μ m.

RGS-PX1 in cells cotransfected with both proteins (Figure 4C). These findings support the conclusion that RGS-PX1 interacts with Hrs both *in vitro* and *in vivo*.

Gas Interacts with Hrs In Vivo and In Vitro

Given that we have previously shown that RGS-PX1 binds to and serves as a GAP for Gas (Zheng *et al.*, 2001), we next asked whether Gas also interacts with Hrs in pull-down and immunoprecipitation assays. We found that 35 S-labeled, *in*

vitro-translated Gas bound to GST-Hrs, but not to GST alone (Figure 5A) and that GST-Hrs, but not GST alone, was able to pull-down endogenous Gas from brain lysates (Figure 5B). Similarly, when we transfected myc-tagged Hrs together with Gas-GFP into HEK293 cells and carried out immunoprecipitation with anti-GFP IgG, myc-Hrs coprecipitated with Gas-GFP (Figure 5C). Because Hrs and Gas have been found in both membrane and cytosolic fractions, we investigated where they interact. We found that Gas and




Figure 7. Immunogold localization of Hrs and *Gas* in HEK293 cells. (A). *Gas*-GFP (10-nm gold) is localized to numerous coated tubules (arrowheads) and endosomes (asterisks). (B and C). Myc-Hrs (5-nm gold, arrows) and *Gas*-GFP (10-nm gold) colocalize in coated domains of early endosomes (asterisks). HEK293 cells transfected with myc-Hrs and *Gas*-GFP were fixed either in 4% PFA (A) or a mixture of 4% PFA and 0.2% glutaraldehyde (B and C) and prepared for ultrathin cryosectioning as described in *Materials and Methods*. Ultrathin cryosections were labeled with anti-myc mAb (5-nm gold) and polyclonal anti-GFP (10-nm gold). Bar, 100 nm.

Hrs were equally distributed between membrane (P100) and cytosolic fractions (S100) in HEK293 cells expressing myc-tagged Hrs together with pcDNA3.1-*Gas* (Figure 5D, lanes 1 and 2). However, the majority of the *Gas* (>95%) coimmunoprecipitated with myc-Hrs from membrane fractions (Figure 5D, lane 4). These results indicate that *Gas* interacts with Hrs and that the interaction takes place largely on membranes, presumably on endosomal membranes as both Hrs (Komada *et al.*, 1997; Raiborg *et al.*, 2001a) and RGS-PX1 (Zheng *et al.*, 2001) are localized on early endosomes.

Gas, RGS-PX1, and Hrs Colocalize on Early Endosomes

To determine the localization of *Gas* and whether it colocalizes with Hrs and RGS-PX1 on endosomes, we carried out indirect immunofluorescence and deconvolution analysis on Cos7 cells expressing *Gas*-GFP alone or *Gas* together with RGS-PX1 and Hrs. Because roughly 50% of both *Gas* and Hrs are found in cytosolic fractions (Figure 5D), we permeabilized the cells before fixation to release cytosolic proteins and facilitate the detection of membrane-associated pools of




Figure 8. Colocalization of *Gas* with RGS-PX1 on early endosomes. GFP-RGS-PX1 is found on endosomes (arrowheads and insets, A and D) and partially colocalizes with *Gas*-WT (arrowheads and inset, B and C) on endosomes loaded with Texas Red EGF (arrowheads and inset, E and F). Cos7 cells were transfected with GFP-RGS-PX1 and *Gas*-WT. In D–F, cells also were incubated with Texas Red EGF for 15 min at 37°C. Cells were permeabilized with saponin, fixed with 3% PFA, double labeled with mouse anti-GFP mAb (A and D), and rabbit anti-*Gas* (B) IgG, and analyzed as described in Figure 6. Bar, 2 μ m.

Gas and *Hrs*. In cells transfected with *Gas*-GFP alone, *Gas*-GFP showed fine, punctate staining throughout the cytoplasm that partially overlapped with the early endosome markers *Hrs* (Figure 6, A–C) and Rab5 (our unpublished data).

In cells expressing both *Gas*-GFP and myc-*Hrs*, these two proteins strongly colocalized in endosomes (Figure 6, D–F). In agreement with previous reports (Raiborg *et al.*, 2001b; Bishop *et al.*, 2002; Urbe *et al.*, 2003), overexpression of *Hrs* resulted in enlarged, clustered endosomes. Furthermore, *Gas*-GFP and Rab5 colocalized on these enlarged endosomes (Figure 6, G–I). More *Gas* colocalized with *Hrs* on these endosomes (Figure 6, D–F) compared with cells transfected with *Gas*-GFP alone (Figure 6, A–C), suggesting expression of *Hrs* causes more *Gas* to translocate to early endosomes. By immunogold labeling, *Gas*-GFP and myc-*Hrs* colocalized in coated microdomains of these enlarged endosomes (Figure 7). Thus, the immunofluorescence results and the coimmunoprecipitation assays together indicate the *Gas* binds *Hrs* on early endosomes.

In cells cotransfected with untagged *Gas* and GFP-RGS-PX1, GFP-RGS-PX1 and *Gas* colocalized on endosomes (Figure 8, A–C) that also were labeled with Texas Red EGF after 15 min uptake (Figure 8, D–F). This is consistent with its early endosome localization reported previously. Again, compared with cells expressing *Gas*-GFP alone (Figure 6, A–C), more *Gas* seemed to be localized on early endosomes

(Figure 8, A–C), suggesting expression of RGS-PX1, as well as *Hrs*, causes more *Gas* to translocate to early endosomes.

Gas, RGS-PX1, and *Hrs* Form a Coprecipitable Complex

The ability of both RGS-PX1 and *Gas* to interact with *Hrs* and the colocalization of these three proteins on early endosomes suggested that they might be present in the same protein complex. To test this possibility, we performed coimmunoprecipitation experiments by using HEK293 cells transfected with myc-*Hrs*, FLAG-RGS-PX1, and *Gas*. When immunoprecipitation was carried out with an anti-myc IgG, both FLAG-RGS-PX1 and *Gas* coprecipitated with myc-*Hrs* (Figure 9, lane 3). Similarly, anti-FLAG IgG was able to bring down both myc-*Hrs* and *Gas* (Figure 9, lane 2). These results suggest that *Gas*, RGS-PX1, and *Hrs* form a coprecipitable protein complex.

Knockdown of Both *Gas* and *Hrs* Further Delays EGF Receptor Degradation

The interaction between *Gas* and *Hrs* suggests *Gas* may function together with *Hrs* in the endosomal sorting and down-regulation of the EGF receptor. To test this hypothesis, we performed double RNAi experiment to knock-down the expression of both *Gas* and *Hrs* (Figure 10, A and B). In cells transfected with both *Gas* and *Hrs* RNAi oligos, ~25% of the EGFR was degraded at 30 min after adding EGF, whereas in cells transfected with either *Gas* or *Hrs* RNAi




Figure 9. $G_{\alpha s}$, FLAG-RGS-PX1, and myc-Hrs form a coprecipitable complex. HEK293 cells were transfected with pcDNA3- $G_{\alpha s}$, FLAG-RGS-PX1, and myc-Hrs. Lysates were immunoprecipitated with anti-FLAG (lane 2), anti-myc (lane 3), or control (ctrl) (lane 4) mouse IgGs, followed by immunoblotting with anti- $G_{\alpha s}$ (top), anti-FLAG (middle), or anti-myc (bottom) IgG. $G_{\alpha s}$ (top) coprecipitates with both FLAG-RGS-PX1 (lane 2) and myc-Hrs (lane 3).

oligos alone, ~50% of the EGF receptors had been degraded at 30 min. These results together with the interaction between Hrs and $G_{\alpha s}$ strongly suggest that $G_{\alpha s}$ cooperates with Hrs in regulating EGF receptor degradation.

DISCUSSION

Our work presented here demonstrates a regulatory role for the heterotrimeric $G_{\alpha s}$ protein in EGF receptor trafficking and down-regulation. We find that expression of $G_{\alpha s}$ accelerates degradation of both EGF receptors and Texas Red EGF, whereas depletion of $G_{\alpha s}$ by RNAi delays their degradation. We also show that $G_{\alpha s}$ forms a complex with RGS-PX1 and Hrs that seems to cause more $G_{\alpha s}$ to translocate to early endosomes. Based on these findings, we propose the following model (Figure 11) for the function of $G_{\alpha s}$ on endosomes: 1) in the presence of RGS-PX1, $G_{\alpha s}$ translocates from the plasma membrane or cytoplasm to early endosomes after EGF binding, where it forms a complex with RGS-PX1 and Hrs; and 2) through interaction with Hrs, $G_{\alpha s}$ regulates endosomal sorting and hence modulates down-regulation of the EGF receptor.

Traditionally, heterotrimeric G proteins have been considered to be largely associated with the cell membrane. Our study indicates early endosomes represent a novel intracellular location for $G_{\alpha s}$. This is in keeping with previous implications that $G_{\alpha s}$ plays a role in endosomal functions, such as early endosome fusion, phagosome-endosome fusion, and transcytosis of pIgR (Colombo *et al.*, 1992, 1994; Bomsel and Mostov, 1993; Beron *et al.*, 1995). Consistent with our localization data, more recently it was shown that endogenous $G_{\alpha s}$ also can be found in rat liver endosomes based on cell fractionation and immunofluorescence studies (Van Dyke, 2004).

How translocation of $G_{\alpha s}$ to endosomes is triggered is still an open question. There are two possible scenarios. First, activation of a GPCR linked to $G_{\alpha s}$ could stimulate translocation. It has been reported that activation of β -adrenergic receptors or cholera toxin treatment, $G_{\alpha s}$ dissociates from the cell membrane into the cytoplasm (Ransnas *et al.*, 1989; Levis and Bourne, 1992; Wedegaertner and Bourne, 1994; Wedegaertner *et al.*, 1996; Yu and Rasenick, 2002). Alterna-

tively, activation of the EGF receptor by EGF could trigger the translocation of $G_{\alpha s}$. It has been shown that $G_{\alpha s}$ is tyrosine phosphorylated by the EGF receptor in vitro and in response to EGF stimulation in vivo (Liebmann *et al.*, 1996; Poppleton *et al.*, 1996). Conceivably, this phosphorylation event might be related to the change in the subcellular localization of $G_{\alpha s}$. Furthermore, $G_{\alpha s}$ has been shown to interact directly with the juxtamembrane region (50 aa) of the EGF receptor in both yeast two-hybrid and coimmunoprecipitation assays; this interaction was suggested to be responsible for the activation of adenylyl cyclase by EGF stimulation in cardiomyocytes (Nair *et al.*, 1990; Sun *et al.*, 1997). Intriguingly, the juxtamembrane region of the EGF receptor contains a dileucine motif that is required for efficient sorting of receptors to lysosomes (Lin *et al.*, 1986; Kil *et al.*, 1999; Bao *et al.*, 2000). The juxtamembrane region also includes a protein kinase C phosphorylation site, and phosphorylation of the EGF receptor by protein kinase C has been shown to switch receptors from the degradation to




Figure 10. Simultaneous knockdown of both $G_{\alpha s}$ and Hrs causes a delay in EGF receptor degradation greater than knock-down of either $G_{\alpha s}$ or Hrs alone. Cos7 cells were transfected with 75 nM, $G_{\alpha s}$ siRNA alone, Hrs siRNA alone, both $G_{\alpha s}$ and Hrs siRNA, or with control siRNA oligos by using Oligofectamine. After 3 d, cells were treated with 100 nM EGF for 30 min, lysed, and analyzed by immunoblotting with antibodies against EGF receptor, $G_{\alpha s}$, Hrs, and actin. In cells transfected with Hrs or $G_{\alpha s}$ RNAi alone, 50–55% of the EGF receptors have been degraded after 30 min, whereas in those transfected with both $G_{\alpha s}$ and Hrs siRNA, degradation is delayed as only 25% of the receptors have been degraded. Data shown are representative of at least three independent experiments. (B) Quantification of EGF receptor degradation. Results from three independent experiments were analyzed by Quantity One software. Data presented as percent of total EGFR at 0 min in each group of cells.




Figure 11. Model of the proposed function of Gas in EGF receptor degradation. After EGF stimulation Gas is recruited to early endosomes where it forms a complex with RGS-PX1 and Hrs on the endosomal membrane. Together with Hrs, Gas promotes the sorting of ubiquitinated (Ub) EGFRs into the luminal vesicles of MVBs and hence facilitates their degradation.

recycling pathways (Lin *et al.*, 1986; Kil *et al.*, 1999; Bao *et al.*, 2000). It would be interesting to know whether these sorting motifs are involved in the binding of Gas to the EGF receptor.

In this work, we have used both overexpression and RNAi knock-down approaches to demonstrate the role of Gas in EGF receptor degradation. Previously, it has been shown that overexpression of G α i1 inhibited internalization of low-density lipoprotein and transferring, possibly by binding free G $\beta\gamma$ subunits and forming inactive heterotrimers (Lin *et al.*, 1998). Although overexpression of Gas may cause similar sequestration of G $\beta\gamma$ subunits, our RNAi knock-down results strongly suggest that Gas plays a direct role in regulating EGF receptor degradation. Whether free G $\beta\gamma$ also is involved in the degradation of EGF receptor directly remains to be investigated.

As a core component of the endosome sorting machinery, Hrs is evolutionarily conserved in eukaryotes. In budding yeast *Saccharomyces cerevisiae*, the Hrs homolog Vps27 is one of the “class E” vacuolar protein sorting (Vps) proteins required for formation of MVBs, sorting of membrane proteins into MVBs, or budding into MVBs (Vida and Emr, 1995; Katzmann *et al.*, 2003). It is noteworthy that *S. cerevisiae* does not seem to have a Gas homolog. The two heterotrimeric G proteins encoded in *S. cerevisiae*, GPA1 and GPA2, are closer to the G α i rather than the Gas subfamily of mammalian G proteins in amino acid sequence. As for RGS-PX1, its putative homolog in *S. cerevisiae*, Mdm1, contains a PX-associated (PXA) domain of unknown function and a PX domain that binds to phosphatidylinositol-3-phosphate (McConnell and Yaffe, 1992; Yu and Lemmon, 2001). However, Mdm1 does not have a homologous RGS domain, and no functional link between Mdm1 and MVB sorting has been reported to date. The absence of a Gas homolog in *S. cerevisiae* and the lack of an RGS domain in Mdm1 lead us to propose that Gas serves as a regulatory module in the endosome sorting machinery in higher organisms, rather than a evolutionarily conserved core component like Hrs.

Depletion of Hrs by RNAi in mammalian cells was shown to decrease the membrane association of the ESCRT complex, reduce the number of MVBs, and disrupt lysosomal targeting of EGF receptors, leading to impaired EGF receptor down-regulation (Bache *et al.*, 2003a,b). We report here that depletion of Gas by RNAi, similar to Hrs, delays deg-

radation of EGF receptors. Moreover, simultaneous depletion of Gas and Hrs by double RNAi further inhibited EGF receptor degradation compared with depletion of Gas or Hrs alone. These results, together with our observation that Gas interacts with Hrs, suggest that Gas and Hrs act together to promote ligand-dependent degradation of EGF receptors. We have previously found that overexpression of RGS-PX1 slowed EGF receptor degradation (Zheng *et al.*, 2001), an effect of RGS-PX1 that could be explained by its GAP activity on Gas. Alternatively, overexpression of RGS-PX1 might have a dominant-negative effect through its interaction with Hrs. Although we have shown that Hrs can form a coimmunoprecipitable complex with RGS-PX1 and Gas, there is also the possibility that some complexes may contain Hrs and RGS-PX1 only, or Hrs and Gas only, and that Gas may promote EGF receptor degradation by competing RGS-PX1 from Hrs.

Hrs has more recently been shown to regulate degradation of other receptors, including the G protein-coupled receptors CXCR4 (Marchese *et al.*, 2003) and DOR (Hislop *et al.*, 2004) and *Drosophila* Notch and Patched receptors (Jekely and Rorth, 2003), supporting a general role of Hrs in regulating endosomal sorting and degradation of cell surface receptors. It would be of interest to investigate whether the regulatory function of Gas in sorting EGF receptors can be extended to other receptors, especially those coupled to heterotrimeric G proteins.

Unlike its positive role in endosomal sorting, Hrs has recently been suggested to prevent endosome fusion. Recombinant Hrs proteins were found to inhibit homotypic fusion of early endosomes, probably by binding to SNAP-25, thereby inhibiting the formation of a SNARE protein complex containing syntaxin 13, SNAP-25, and VAMP2 (Sun *et al.*, 2003; Yan *et al.*, 2004). Gas has similarly been suggested to negatively regulate endosomal fusion based on the observation that activation of Gas by either cholera toxin or a Gas stimulatory peptide blocked endosomal fusion *in vitro* (Colombo *et al.*, 1994).

In summary, our findings support a previously unappreciated role of Gas in endocytic trafficking and down-regulation of the EGF receptor. Further studies are required to define the precise role of Gas in endosomal sorting in general, to understand the mechanisms involved in the translocation of Gas to early endosomes, and to unravel the differ-

ences in the regulation of G α s functions at the plasma membrane and early endosomes.

ACKNOWLEDGMENTS

We thank Drs. Gordon Gill, Mark Rasenick, and Angela Wandinger-Ness for kindly providing reagents. This work is supported by National Institutes of Health grants CA-100768 and DK-17780 to M.G.F. B.Z. and C.L. were supported in part by postdoctoral fellowships from the American Heart Association and the Canadian Institute of Health Research, respectively. T.T. and A.B. are members of the Molecular Pathology and Biomedical Sciences Graduate Programs, respectively, University of California, San Diego.

REFERENCES

- Bache, K.G., Brech, A., Mehlum, A., and Stenmark, H. (2003a). Hrs regulates multivesicular body formation via ESCRT recruitment to endosomes. *J. Cell Biol.* **162**, 435–442.
- Bache, K.G., Raiborg, C., Mehlum, A., and Stenmark, H. (2003b). STAM and Hrs are subunits of a multivalent ubiquitin-binding complex on early endosomes. *J. Biol. Chem.* **278**, 12513–12521.
- Bao, J., Alroy, I., Waterman, H., Schejter, E.D., Brodie, C., Gruenberg, J., and Yarden, Y. (2000). Threonine phosphorylation diverts internalized epidermal growth factor receptors from a degradative pathway to the recycling endosome. *J. Biol. Chem.* **275**, 26178–26186.
- Beron, W., Colombo, M.I., Mayorga, L.S., and Stahl, P.D. (1995). In vitro reconstitution of phagosome-endosome fusion: evidence for regulation by heterotrimeric GTPases. *Arch. Biochem. Biophys.* **317**, 337–342.
- Bishop, N., Horman, A., and Woodman, P. (2002). Mammalian class E vps proteins recognize ubiquitin and act in the removal of endosomal protein-ubiquitin conjugates. *J. Cell Biol.* **157**, 91–101.
- Bomsel, M. and Mostov, K.E. (1992). Role of heterotrimeric G proteins in membrane traffic. *Mol. Biol. Cell* **3**, 1317–1328.
- Bomsel, M. and Mostov, K.E. (1993). Possible role of both the alpha and beta gamma subunits of the heterotrimeric G protein, Gs, in transcytosis of the polymeric immunoglobulin receptor. *J. Biol. Chem.* **268**, 25824–25835.
- Carpenter, G. (2000). The EGF receptor: a nexus for trafficking and signaling. *Bioessays* **22**, 697–707.
- Chin, L.S., Raynor, M.C., Wei, X., Chen, H.Q., and Li, L. (2001). Hrs interacts with sorting nexin 1 and regulates degradation of epidermal growth factor receptor. *J. Biol. Chem.* **276**, 7069–7078.
- Clague, M.J. and Urbe, S. (2003). Hrs function: viruses provide the clue. *Trends Cell Biol.* **13**, 603–606.
- Colombo, M.I., Mayorga, L.S., Casey, P.J., and Stahl, P.D. (1992). Evidence of a role for heterotrimeric GTP-binding proteins in endosome fusion. *Science* **255**, 1695–1697.
- Colombo, M.I., Mayorga, L.S., Nishimoto, I., Ross, E.M., and Stahl, P.D. (1994). Gs regulation of endosome fusion suggests a role for signal transduction pathways in endocytosis. *J. Biol. Chem.* **269**, 14919–14923.
- De Vries, L., Zheng, B., Fischer, T., Elenko, E., and Farquhar, M.G. (2000). The regulator of G protein signaling family. *Annu. Rev. Pharmacol. Toxicol.* **40**, 235–271.
- Gilman, A.G. (1987). G proteins: transducers of receptor-generated signals. *Annu. Rev. Biochem.* **56**, 615–649.
- Haft, C.R., de la Luz Sierra, M., Barr, V.A., Haft, D.H., and Taylor, S.I. (1998). Identification of a family of sorting nexin molecules and characterization of their association with receptors. *Mol. Cell Biol.* **18**, 7278–7287.
- Helms, J.B. (1995). Role of heterotrimeric GTP binding proteins in vesicular protein transport: indications for both classical and alternative G protein cycles. *FEBS Lett.* **369**, 84–88.
- Hislop, J.N., Marley, A., and Von Zastrow, M. (2004). Role of mammalian vacuolar protein-sorting proteins in endocytic trafficking of a non-ubiquitinated G protein-coupled receptor to lysosomes. *J. Biol. Chem.* **279**, 22522–22531.
- Hollinger, S. and Hepler, J.R. (2002). Cellular regulation of RGS proteins: modulators and integrators of G protein signaling. *Pharmacol. Rev.* **54**, 527–559.
- Jekely, G. and Rorth, P. (2003). Hrs mediates downregulation of multiple signalling receptors in *Drosophila*. *EMBO Rep.* **4**, 1163–1168.
- Katzmann, D.J., Odorizzi, G., and Emr, S.D. (2002). Receptor downregulation and multivesicular-body sorting. *Nat. Rev. Mol. Cell Biol.* **3**, 893–905.
- Katzmann, D.J., Stefan, C.J., Babst, M., and Emr, S.D. (2003). Vps27 recruits ESCRT machinery to endosomes during MVB sorting. *J. Cell Biol.* **162**, 413–423.
- Kil, S.J., Hobert, M., and Carlin, C. (1999). A leucine-based determinant in the epidermal growth factor receptor juxtamembrane domain is required for the efficient transport of ligand-receptor complexes to lysosomes. *J. Biol. Chem.* **274**, 3141–3150.
- Komada, M., Masaki, R., Yamamoto, A., and Kitamura, N. (1997). Hrs, a tyrosine kinase substrate with a conserved double zinc finger domain, is localized to the cytoplasmic surface of early endosomes. *J. Biol. Chem.* **272**, 20538–20544.
- Kurten, R.C., Cadena, D.L., and Gill, G.N. (1996). Enhanced degradation of EGF receptors by a sorting nexin, SNX1. *Science* **272**, 1008–1010.
- Levis, M.J. and Bourne, H.R. (1992). Activation of the alpha subunit of Gs in intact cells alters its abundance, rate of degradation, and membrane avidity. *J. Cell Biol.* **119**, 1297–1307.
- Liebmann, C., Graness, A., Boehmer, A., Kovalenko, M., Adomeit, A., Steinmetzer, T., Nurnberg, B., Wetzker, R., and Boehmer, F.D. (1996). Tyrosine phosphorylation of G α and inhibition of bradykinin-induced activation of the cyclic AMP pathway in A431 cells by epidermal growth factor receptor. *J. Biol. Chem.* **271**, 31098–31105.
- Lin, C.R., Chen, W.S., Lazar, C.S., Carpenter, C.D., Gill, G.N., Evans, R.M., and Rosenfeld, M.G. (1986). Protein kinase C phosphorylation at Thr 654 of the unoccupied EGF receptor and EGF binding regulate functional receptor loss by independent mechanisms. *Cell* **44**, 839–848.
- Lin, H.C., Duncan, J.A., Kozasa, T., and Gilman, A.G. (1998). Sequestration of the G protein beta gamma subunit complex inhibits receptor-mediated endocytosis. *Proc. Natl. Acad. Sci. USA* **95**, 5057–5060.
- Lu, Q., Hope, L.W., Brasch, M., Reinhard, C., and Cohen, S.N. (2003). TSG101 interaction with HRS mediates endosomal trafficking and receptor downregulation. *Proc. Natl. Acad. Sci. USA* **100**, 7626–7631.
- Marchese, A., Raiborg, C., Santini, F., Keen, J.H., Stenmark, H., and Benovic, J.L. (2003). The E3 ubiquitin ligase AIP4 mediates ubiquitination and sorting of the G protein-coupled receptor CXCR4. *Dev. Cell* **5**, 709–722.
- McConnell, S.J., and Yaffe, M.P. (1992). Nuclear and mitochondrial inheritance in yeast depends on novel cytoplasmic structures defined by the MDM1 protein. *J. Cell Biol.* **118**, 385–395.
- Nair, B.G., Parikh, B., Milligan, G., and Patel, T.B. (1990). Gs alpha mediates epidermal growth factor-elicited stimulation of rat cardiac adenylate cyclase. *J. Biol. Chem.* **265**, 21317–21322.
- Neves, S.R., Ram, P.T., and Iyengar, R. (2002). G protein pathways. *Science* **296**, 1636–1639.
- Nurnberg, B., and Ahnert-Hilger, G. (1996). Potential roles of heterotrimeric G proteins of the endomembrane system. *FEBS Lett.* **389**, 61–65.
- Poppleton, H., Sun, H., Fulgham, D., Bertics, P., and Patel, T.B. (1996). Activation of G α by the epidermal growth factor receptor involves phosphorylation. *J. Biol. Chem.* **271**, 6947–6951.
- Raiborg, C., Bache, K.G., Mehlum, A., Stang, E., and Stenmark, H. (2001a). Hrs recruits clathrin to early endosomes. *EMBO J.* **20**, 5008–5021.
- Raiborg, C., Bremnes, B., Mehlum, A., Gillooly, D.J., D'Arrigo, A., Stang, E., and Stenmark, H. (2001b). FYVE and coiled-coil domains determine the specific localisation of Hrs to early endosomes. *J. Cell Sci.* **114**, 2255–2263.
- Raiborg, C., Rusten, T.E., and Stenmark, H. (2003). Protein sorting into multivesicular endosomes. *Curr. Opin. Cell Biol.* **15**, 446–455.
- Raiborg, C. and Stenmark, H. (2002). Hrs and endocytic sorting of ubiquitinated membrane proteins. *Cell Struct. Funct.* **27**, 403–408.
- Ransnas, L.A., Svoboda, P., Jasper, J.R., and Insel, P.A. (1989). Stimulation of beta-adrenergic receptors of S49 lymphoma cells redistributes the alpha subunit of the stimulatory G protein between cytosol and membranes. *Proc. Natl. Acad. Sci. USA* **86**, 7900–7903.
- Sachse, M., Urbe, S., Oorschot, V., Strous, G.J., and Klumperman, J. (2002). Bilayered clathrin coats on endosomal vacuoles are involved in protein sorting toward lysosomes. *Mol. Biol. Cell* **13**, 1313–1328.
- Sorkin, A. and Von Zastrow, M. (2002). Signal transduction and endocytosis: close encounters of many kinds. *Nat. Rev. Mol. Cell Biol.* **3**, 600–614.
- Stahl, P.D. and Barbieri, M.A. (2002). Multivesicular bodies and multivesicular endosomes: the “ins and outs” of endosomal traffic. *Sci STKE* **2002**, PE32.
- Stow, J.L. and Heimann, K. (1998). Vesicle budding on Golgi membranes: regulation by G proteins and myosin motors. *Biochim. Biophys. Acta* **1404**, 161–171.
- Sun, H., Chen, Z., Poppleton, H., Scholich, K., Mullenix, J., Weipz, G.J., Fulgham, D.L., Bertics, P.J., and Patel, T.B. (1997). The juxtamembrane, cyto-

- solic region of the epidermal growth factor receptor is involved in association with alpha-subunit of Gs. *J. Biol. Chem.* 272, 5413–5420.
- Sun, W., Yan, Q., Vida, T.A., and Bean, A.J. (2003). Hrs regulates early endosome fusion by inhibiting formation of an endosomal SNARE complex. *J. Cell Biol.* 162, 125–137.
- Urbe, S., Sachse, M., Row, P.E., Preisinger, C., Barr, F.A., Strous, G., Klumperman, J., and Clague, M.J. (2003). The UIM domain of Hrs couples receptor sorting to vesicle formation. *J. Cell Sci.* 116, 4169–4179.
- Van Dyke, R.W. (2004). Heterotrimeric G protein subunits are located on rat liver endosomes. *BMC Physiol.* 4, 1.
- Vida, T.A. and Emr, S.D. (1995). A new vital stain for visualizing vacuolar membrane dynamics and endocytosis in yeast. *J. Cell Biol.* 128, 779–792.
- Wedegaertner, P.B. and Bourne, H.R. (1994). Activation and depalmitoylation of Gs alpha. *Cell* 77, 1063–1070.
- Wedegaertner, P.B., Bourne, H.R., and von Zastrow, M. (1996). Activation-induced subcellular redistribution of Gs alpha. *Mol. Biol. Cell* 7, 1225–1233.
- Wishart, M.J., Taylor, G.S., and Dixon, J.E. (2001). Phoxy lipids: revealing PX domains as phosphoinositide binding modules. *Cell* 105, 817–820.
- Worby, C.A. and Dixon, J.E. (2002). Sorting out the cellular functions of sorting nexins. *Nat. Rev. Mol. Cell. Biol.* 3, 919–931.
- Yan, Q., Sun, W., McNew, J.A., Vida, T.A., and Bean, A.J. (2004). Ca²⁺ and NSF differentially regulate disassembly of SNARE complexes on early endosomes. *J. Biol. Chem.*
- Yu, J.W. and Lemmon, M.A. (2001). All phox homology (PX) domains from *Saccharomyces cerevisiae* specifically recognize phosphatidylinositol 3-phosphate. *J. Biol. Chem.* 276, 44179–44184.
- Yu, J.Z. and Rasenick, M.M. (2002). Real-time visualization of a fluorescent G(alpha)(s): dissociation of the activated G protein from plasma membrane. *Mol. Pharmacol.* 61, 352–359.
- Zheng, B., Chen, D., and Farquhar, M.G. (2000). MIR16, a putative membrane glycerophosphodiester phosphodiesterase, interacts with RGS16. *Proc. Natl. Acad. Sci. USA* 97, 3999–4004.
- Zheng, B., Ma, Y.C., Ostrom, R.S., Lavoie, C., Gill, G.N., Insel, P.A., Huang, X.Y., and Farquhar, M.G. (2001). RGS-PX1, a GAP for GalphaS and sorting nexin in vesicular trafficking. *Science* 294, 1939–1942.