UC Davis UC Davis Previously Published Works

Title

In Vivo Multimodal Imaging of Drusenoid Lesions in Rhesus Macaques

Permalink

https://escholarship.org/uc/item/9440q298

Journal

Scientific Reports, 7(1)

ISSN

2045-2322

Authors

Yiu, Glenn
Tieu, Eric
Munevar, Christian
<u>et al.</u>

Publication Date

2017

DOI

10.1038/s41598-017-14715-z

Copyright Information

This work is made available under the terms of a Creative Commons Attribution License, available at <u>https://creativecommons.org/licenses/by/4.0/</u>

Peer reviewed

SCIENTIFIC REPORTS

Received: 6 June 2017 Accepted: 16 October 2017 Published online: 03 November 2017

OPEN In Vivo Multimodal Imaging of **Drusenoid Lesions in Rhesus** Macaques

Glenn Yiu¹, Eric Tieu¹, Christian Munevar², Brittany Wong³, David Cunefare³, Sina Farsiu³, Laura Garzel⁴, Jeffrey Roberts^{2,4} & Sara M. Thomasy²

Nonhuman primates are the only mammals to possess a true macula similar to humans, and spontaneously develop drusenoid lesions which are hallmarks of age-related macular degeneration (AMD). Prior studies demonstrated similarities between human and nonhuman primate drusen based on clinical appearance and histopathology. Here, we employed fundus photography, spectral domain optical coherence tomography (SD-OCT), fundus autofluorescence (FAF), and infrared reflectance (IR) to characterize drusenoid lesions in aged rhesus macaques. Of 65 animals evaluated, we identified lesions in 20 animals (30.7%). Using the Age-Related Eye Disease Study 2 (AREDS2) grading system and multimodal imaging, we identified two distinct drusen phenotypes - 1) soft drusen that are larger and appear as hyperreflective deposits between the retinal pigment epithelium (RPE) and Bruch's membrane on SD-OCT, and 2) hard, punctate lesions that are smaller and undetectable on SD-OCT. Both exhibit variable FAF intensities and are poorly visualized on IR. Eyes with drusen exhibited a slightly thicker RPE compared with control eyes ($+3.4 \mu m$, P=0.012). Genetic polymorphisms associated with drusenoid lesions in rhesus monkeys in ARMS2 and HTRA1 were similar in frequency between the two phenotypes. These results refine our understanding of drusen development, and provide insight into the absence of advanced AMD in nonhuman primates.

In humans, drusen are focal deposits of extracellular debris located between the retinal pigment epithelium (RPE) and Bruch's membrane, and may occur with normal aging or as a sign of age-related macular degeneration (AMD) - the leading cause of vision loss in the elderly. Drusen are primarily composed of lipids such as phosphatidylcholine, esterified cholesterol, and unesterified cholesterols¹⁻³, as well as various proteins including vitronectin, crystallins, amyloid proteins, apolipoproteins, immunoglobulins, and complement components⁴⁻⁶. The advent of ocular imaging technologies has led to more refined classification of different drusen phenotypes, including soft drusen, cuticular drusen, and reticular pseudodrusen. Soft drusen are larger, yellow-white mounds that are histologically external to the RPE basement membrane. Cuticular drusen are smaller, densely-packed, and more uniform in size. They were previously thought to be nodular thickenings of the RPE basement membrane⁷, but were subsequently found to be external to the basal lamina of the RPE with a similar composition as soft drusen⁸. Reticular pseudodrusen are located in the subretinal space above the RPE^{9,10}, and may be distinguished by their location on histology and optical coherence tomography (OCT)¹¹. These different drusen phenotypes are associated with certain genotypic variants such as those in the complement factor H (CFH) and age-related maculopathy susceptibility (ARMS2) loci, and may have differential propensity for progression to RPE atrophy and late-stage AMD^{12,13}.

Most laboratory animals such as rodents are poor models of AMD and drusen due to their relative short lifespan and lack of a macula. In contrast, nonhuman primates are the only mammals that possess a true macula comparable to humans, and spontaneously develop drusenoid lesions similar to those seen in AMD. Early studies reported a high prevalence of drusenoid deposits in free-ranging colonies of rhesus macaques (Macaca mulatta) from Cayo Santiago in Puerto Rico, where drusenoid pathology was noted in 30-74% of animals¹⁴⁻¹⁷. Subsequent studies of other colonies within the continental U.S. demonstrated a lower prevalence of 5-47%¹⁸⁻²⁰, likely due

¹Department of Ophthalmology & Vision Science, University of California, Davis, Sacramento, California, USA. ²Department of Surgical and Radiological Sciences, School of Veterinary Medicine, University of California, Davis, Davis, California, USA. ³Department of Biomedical Engineering, Duke University, Durham, North Carolina, USA. ⁴California National Primate Research Center, Davis, California, USA. Correspondence and requests for materials should be addressed to G.Y. (email: gyiu@ucdavis.edu)

to factors such as reduced sunlight exposure from indoor housing, more regular diet, age and genetic differences between colonies, as well as different methods of detecting and classifying the lesions.

Prior studies of drusenoid deposits in nonhuman primates were mostly qualitative, relying on clinical examination, fundus photography, and histological analysis to characterize the lesions. The deposits were classified as "hard" or "soft" based on funduscopic appearance, followed by post-mortem assessment of their histopathology. These studies revealed increasing frequency of drusenoid deposits with age^{14,21}, and similar molecular composition as human drusen²². Rhesus monkeys and humans were also found to share genetic polymorphisms in genes such as ARMS2 and HTRA1 (high temperature-requirement factor A1) that are associated with drusen formation^{23–25}. These studies suggest that the drusenoid deposits in nonhuman primates may offer significant insight into drusen formation in humans, but more refined, quantitative methods are necessary to characterize these lesions in monkeys for translational research.

In this study, we surveyed aged rhesus macaques (age >19 years) at the California National Primate Research Center (CNPRC) to identify animals with drusenoid lesions. We used the validated age-related eye disease study 2 (AREDS2) system and digital image segmentation for grading drusen from fundus photographs in human studies^{26,27}, then employed *in vivo* ocular imaging modalities, including blue-peak fundus autofluorescence (FAF), infrared reflectance (IR), and spectral domain-OCT (SD-OCT), to characterize the drusenoid lesions in rhe-sus macaques *in vivo*. We identified two morphologic phenotypes of drusenoid deposits, which can be distinguished by different imaging features, and investigated potential associations between drusen phenotype with single-nucleotide polymorphisms in ARMS2 and HTRA1 which are known to be associated with macular drusen in rhesus monkeys. Comparing our results with prior studies of drusen prevalence in other primate colonies, we speculate on the pathogenesis of drusen formation in rhesus macaques, and the absence of advanced AMD in nonhuman primates.

Methods

Subjects. The CNPRC is accredited by the Association for Assessment and Accreditation of Laboratory Animal Care (AAALAC) International. All studies using rhesus macaques (Macaca mulatta) followed the guidelines of the Association for Research in Vision and Ophthalmology Statement for the Use of Animals in Ophthalmic and Vision Research, complied with the National Institutes of Health (NIH) Guide for the Care and Use of Laboratory Animals, and were approved by the University of California, Davis Institutional Animal Care and Use Committee. Complete ophthalmic examinations were performed in aged rhesus macaques (\geq 19 years) undergoing routine semi-annual physical examinations. The animals were sedated with intramuscular injection of ketamine hydrochloride, midazolam, and dexmedetomidine, followed by pupillary dilation with tropicamide (Bausch & Lomb, Tampa, FL) and cycloplegia with cyclopentolate (Akorn, Lake Forest, IL). All subjects underwent slit lamp biomicroscopy, dilated fundus biomicroscopy, streak retinoscopy, A-scan biometry, and rebound tonometry. After the initial survey, multimodal imaging of the macula was performed on all animals with drusenoid lesions identified on fundus examination, as well as a selection of age-matched control animals, including fundus photography, blue-peak FAF, IR imaging, and SD-OCT. Animal eyes that showed any other retinal or choroidal lesions on exam were excluded from this study. For all animals, age, sex, cycloplegic refraction (diopters), intraocular pressure (mmHg) measured by rebound tonometry, and axial length (mm) on A-scan biometry were recorded.

Multimodal Ocular Imaging. Fundus photography was performed using the CF-1 Retinal Camera (Canon, Tokyo, Japan) with a 50-degree wide-angle lens. Blue-peak FAF and IR imaging were obtained simultaneously with SD-OCT using the Spectralis HRA+OCT platform (Heidelberg Engineering, Heidelberg, Germany). Confocal scanning laser ophthalmoscopy was used to capture $30^{\circ} \times 30^{\circ}$ autofluorescence images using an excitation light of 488 nm and a long-pass barrier filter starting at 500 nm, along with a $20^{\circ} \times 20^{\circ}$ SD-OCT volume scan with 1024 A-scans per B-scan and raster line spacing of $51 \,\mu$ m, centered on the fovea, in high-resolution mode. Infrared reflectance images were captured using an 820 nm diode laser, along with a $30^{\circ} \times 5^{\circ}$ SD-OCT raster scan with 1536 A-scans per B-scan and 234 μ m spacing between B-scans, in high-resolution enhanced-depth imaging (EDI) mode. 25 scans were averaged for each B-scan, using the Heidelberg eye tracking Automatic Real-Time (ART) software. All imaging was performed by the first author (GY) at CNPRC. Animals were monitored by a trained technician and a CNPRC veterinarian (LG) at all times.

Image Analysis. Digital color fundus photographs were evaluated by two experienced image graders (GY, ET) using ImageJ (version 1.49v; National Institutes of Health, Bethesda, MD). Drusenoid lesions were categorized using the Age-Related Eye Disease Study 2 (AREDS2) system for classification of AMD, which has been previously validated for grading of digital fundus photographs in human clinical trials with high reproducibility26. Grading involves a modified grid template adapted from the Early Treatment Diabetic Retinopathy Study (ETDRS)²⁸, with a set of graduated circles used to estimate maximum drusen size and total area involved by pigment abnormalities and drusen (Fig. 1A). The grid template was calibrated in these animals based on the ocular dimensions of the optic disc in rhesus macaques, which measures 1400 µm vertically and 1000 µm horizontally²⁹, compared to $1800 \,\mu\text{m}$ in both dimensions in humans²⁶. The radii of the modified grid template circles remain as 1/3 DD, 1 DD, and 2 DD based on the vertical disc diameter, but are equivalent to $467 \mu m$, $1400 \mu m$, and $2800 \mu m$ respectively (Fig. 1B). Rhesus and human studies can be compared by applying a scaling constant of 0.778 for linear measurements, and 0.605 for area measurements. Lesions graded include area of hyperpigmentation or hypopigmentation; maximum size and area of drusen within the grid; presence of calcified drusen, drusenoid pigment epithelial detachment, or reticular drusen; presence, area, configuration, and center-involvement of geographic RPE atrophy; and presence of neovascular AMD (Supplemental Table 1). The details of the grading criteria have been extensively described^{26,30}. Any discrepancies in grading were resolved by open adjudication



Figure 1. Grading and quantification of drusenoid lesions in rhesus macaques from fundus photographs. A representative fundus photograph of a left rhesus macaque eye with large soft drusen is shown overlaid with the modified Early Treatment Diabetic Retinopathy Study (ETDRS) grid template (**A**), along with a diagram of the graduated measurement circles used for estimating the drusen size and area based on a modified Age-Related Eye Disease Study 2 (AREDS2) grading system (**B**). Both the grid template and measurement circles are calibrated to the vertical dimension of the rhesus optic nerve, which is 1400 μ m in adult animals (compared to 1800 μ m in human adults). The same representative fundus photograph is shown with manual segmentation of drusenoid lesions (yellow outlines), as well as the inner and outer rings (yellow circles) (**C**). Scatterplots show the inverse relationship between number of drusenoid lesions and average lesion size (**D**), as well as the association between AREDS2 drusen area grade with total segmented lesion area (**E**) and between AREDS2 drusen size grade with largest segmented lesion size (**F**).

between the graders. Manual drusen segmentation of the fundus photographs was then performed by the two graders by outlining individual lesions within the inner and outer ETDRS circle using the freehand selection tool in ImageJ as previously described²⁷ and calibrated to the vertical diameter of the optic disc. All measurements were determined from the mean of the two graders' measurements, and include the number of drusenoid lesions, total area of lesions, average lesion size, and size of largest lesion (Fig. 1C).

The SD-OCT images were exported from Heidelberg Explorer software (version 1.8.6.0, Heidelberg Engineering, Heidelberg, Germany) to Duke Optical Coherence Tomography Retinal Analysis Program (DOCTRAP, version 62.0)^{31,32}, a custom image segmentation software designed using MATLAB (Mathworks, Natick, MA). For thickness measurements of retinal and choroidal layers, segmentation boundaries were automatically determined by DOCTRAP, followed by manual refinement by two experienced graders (CM, BW) along a 3 mm horizontal segment centered on the fovea. Measured layers include nerve fiber layer (NFL), ganglion cell/inner plexiform layers (GCL/IPL), inner nuclear layer (INL), outer plexiform layer (OPL), outer nuclear layer (ONL), photoreceptor inner and outer segments (IS/OS), retinal pigment epithelium (RPE) which includes any drusenoid complex between the RPE and Bruch's membrane, as well as the choriocapillaris (CC), and outer choroid (OC). The GCL/IPL were measured as a single complex due to difficulty in distinguishing between these layers even in normal retinal scans. The total retinal thickness was measured from the internal limiting membrane to Bruch's membrane, and total choroidal thickness was measured from the mean of the two graders' measurements, and then averaged across the central 3mm for thickness comparisons.

Correlation in drusen size and area between left and right eyes was determined using two-tailed Pearson correlation. Differences in chorioretinal layer thickness measurements and differences in lesion number, size, and area were compared using linear regression analysis with generalized estimating equations to account for two eyes evaluated from each animal. The association of AREDS2 grading with segmented measurements was also determined using linear regression with generalized estimating equations.

Genotype Analysis. Genomic DNA from blood or buffy coat was extracted with a commercial kit (DNeasy Blood and Tissue Kit; Qiagen, Hilden, Germany) following the manufacturer's protocol, and quantified using a spectrophotometer (NanoDrop 2000c; Thermo Fisher Scientific, Waltham, MA, USA). The region of interest was amplified using the following primers HTRA1-Forward 5'-TATCACTTCACTGTGGGTCCGG-3', HTRA1-Reverse 5'-GCGATTCGCGTCCTTCAAACTA-3', ARMS2-Exon1-Forward 5'-GATGGCAGCTGGCTTGGCAAGG, ARMS2-Exon1-Reverse 5'-GGGGTAAGGCCTGATCATCTGCA-3' with a high-fidelity DNA polymerase (Phusion or Q5; New England Biolabs, Ipswich, MA, USA) and purified using a PCR purification kit (QIAquick; Qiagen). Sanger sequencing of PCR products was performed by the University of California, Davis DNA Sequencing facility (Davis, CA, USA). Association of single nucleotide polymorphisms with drusen phenotype was determined using a Fisher's Exact test. All statistical analyses were performed using SPSS Statistics (version 22, IBM).

Results

Demographics and Ocular Characteristics. Of 65 aged macaques (>19 years) evaluated at CNPRC, we identified drusenoid lesions by dilated fundus biomicroscopy in both eyes of 20 animals (30.7%). Of these animals, 10 were male, and mean age was 23.3 ± 2.7 years (range 19.3 to 29.4 years). Among the 40 eyes with drusenoid lesions, mean intraocular pressure (IOP) was 19.1 ± 4.4 mmHg, and mean axial length was 19.8 ± 0.7 mm. We also selected 8 age-matched animals with no ocular findings as normal controls for ocular imaging, with a mean age of 23.9 ± 2.9 years (range 21.3 to 28.2 years), of which 3 were male. Among the 16 control eyes, mean IOP and axial length were 17.1 ± 4.5 mmHg and 19.7 ± 1.0 mm, respectively. No statistical differences in age (P=0.609), sex (P=0.401), IOP (P=0.298), and axial length (P=0.945) were observed between the two groups of animals.

AREDS2 Classification of Drusenoid Lesions on Fundus Photography. Among the 40 eyes noted to have drusenoid lesions on fundus examination, 34 eyes had fundus photographs with adequate quality for grading of drusenoid lesions (Table 1). Using the AREDS2 classification for drusen size, the majority of lesions were classified as "small" (grade 2, \leq 59 µm diameter) or "large" (grade 4, >116 µm diameter), with similar proportions of each. A majority of eyes with drusenoid lesions had a total lesion area between grading circles I-2 (0.089 mm²) and O-2 (0.297 mm²). Left and right eyes of individual animals showed a strong correlation in drusen size (r = 0.989; P < 0.001) and area (r = 0.970; P < 0.001). No cases of calcified drusen, drusenoid pigment epithelial detachment, or reticular drusen were identified in any of the fundus photographs. There was also no evidence of hyperpigmentation, hypopigmentation, geographic atrophy, or neovascular AMD noted in any eye in this cohort of macaques.

We noted that drusenoid lesions in rhesus macaques adapted either a "soft" drusen or "hard" punctate appearance, based on the original AREDS terminology for drusen type. When classified in this manner, eyes with predominantly soft drusen showed significantly larger drusen size (P < 0.001) and a trend toward greater total lesion area (P = 0.102; Table 1). Some eyes with soft drusen also showed hard punctate lesions, while eyes with predominantly hard lesions rarely showed any soft drusen. Fellow eyes showed perfect correlation (r = 1.000; P < 0.001) in predominant lesion phenotype classified in this manner. Lesions were occasionally noted outside of the central macula, although only one animal exhibited any significant drusen (area ≥ 0.297 mm²) beyond the 5.6mm outer circle of the ETDRS grid (Table 1).

Quantification of Drusenoid Lesions on Fundus Photography. Of eyes with digital fundus photographs, 21 were found to be adequate for image segmentation of individual drusenoid lesions. Quantitative drusen segmentation showed an inverse association between lesion size and number of lesions (Fig. 1D). Eyes

	Any Drusen (n=40)	Soft Drusen (n = 16)	Hard Lesions (n = 18)	P-value
Drusen size (largest)				<0.001*
0: None	0	0	0	
1: Questionable	0	0	0	
2: Definite, diameter $< C-0^{\&}$	17	0	17	
3: Diameter > C-0, but < C-1	1	0	1	
4: Diameter > C-1, but < C-2	15	15	0	
5: Diameter > C-2	1	1	0	
8: Cannot grade	6	0	0	
Drusen area [#]				0.102
0: Questionable, or total area < C-0	0	0	0	
1: Area \geq C-0, but $<$ C-1	0	0	0	
2: Area \geq C-1, but < C-2	2	0	2	
3: Area \geq C-2, but < I-2	4	1	3	
4: Area \geq I-2, but < O-2	16	7	9	
5: Area \geq O-2, but < ½ DA	7	3	4	
6: Area \geq ½ DA, but < 1DA	2	1	1	
7: Area \geq 1DA	3	3	0	
8: Cannot grade	6	0	0	
Drusen outside grid				cannot compute
0: None, questionable, or area < O-2	33	15	18	
5: Area ≥ O-2	1	1	0	
8: Cannot grade	6	0	0	

Table 1. Fundus photograph grading of drusenoid lesions in Rhesus macaques using the AREDS2 classification system. *Statistically significant. *Area within modified Early Treatment of Diabetic Retinopathy Study (ETDRS) grid. & The circular areas for grading measured in square millimeters are: C-0=0.003 mm², C-1=0.011 mm², C-2=0.043 mm², I-2=0.089 mm², O-2=0.297 mm², 1/2 disc area (DA)=0.77 mm², 1 DA=1.54 mm².

with primarily soft drusen showed fewer number of lesions (P = 0.006) and larger average lesion size (P < 0.001), with a trend toward greater total lesion area (P = 0.297; Table 2). AREDS2 drusen area and size grading correlated well with the total lesion area and size of largest lesion as measured from image segmentation (P < 0.001 for both) (Fig. 1E,F). Hence, both AREDS2 grading and quantitative measurements from drusen segmentation demonstrate the distinct size and distribution of these two lesion phenotypes.

Comparison of Drusenoid Lesions on OCT and FAF. Image analysis of soft and hard lesions showed distinct characteristics on multimodal ocular imaging. Soft drusen (Fig. 2A) have a variable appearance on FAF imaging, with some lesions appearing more hyper-autofluorescent, and others having a more iso-autofluorescent appearance (Fig. 2B). Near IR imaging of the ocular fundus in rhesus macaques showed a mottled appearance corresponding to the underlying choroidal vascular architecture, and the drusenoid deposits are difficult to distinguish (Fig. 2C). On SD-OCT, the soft deposits appear as dome-shaped hyperreflective deposits located between the RPE and Bruch's membrane (Fig. 2D–G). Deposits show variable autofluorescence intensities, with some larger lesions appearing more hyper-autofluorescent (Fig. 2D), while some less elevated ones show a less intense autofluorescence signature and may appear as focal thickening of the RPE itself (Fig. 2E–G). There was no clear disruption of the outer retinal layers, such as the external limiting membrane (ELM) or inner segment-outer segment (IS-OS) junction; or changes to the underlying choriocapillaris, which appear in rhesus macaques as a distinct hyporeflective band as our group has described recently³⁵. The cone outer-segment tip (COST) lines, also known as the interdigitation zone, were variably visible in different animals, and showed no consistent difference between normal subjects and those with either drusen phenotype.

Hard punctate lesions (Fig. 3A) also have a variable appearance on FAF imaging, with some lesions appearing more hyper-autofluorescent while others are not as readily discernable (Fig. 3B). Rarely, these lesions may appear on near IR imaging (Fig. 3C), but most of the lesions could not be distinguished by this modality. Interestingly, the small punctate lesions were uniformly undetectable on even high density SD-OCT scans, with no visible disruptions of the RPE, outer retinal layers, or choroidal vasculature (Fig. 3D–G). Hence, these hard lesions do not appear to represent true sub-RPE deposits as in AMD drusen.

Retinal and Choroidal Layer Thicknesses. To compare retinal and choroidal thicknesses, we employed a custom, semi-automated segmentation software to measure the average thicknesses of individual layers across the central 3mm along a horizontal SD-OCT line scan through the fovea (Fig. 4A–B). Among the animals, 23 eyes with drusen and 14 eyes from age-matched controls had SD-OCT images of adequate quality for thickness measurements. Among these eyes, total retinal and choroidal thickness, as well as the thicknesses of individual layers, were similar between the two groups (Fig. 4C–D), with the exception of the RPE which was statistically

	Any Drusen (n=21)	Soft Drusen (n=9)	Hard Lesions (n = 12)	P-value
Number of Lesions	$181.5 \pm 174.7^{\text{\#}}$	67.7 ± 44.9	266.9 ± 188.4	0.006*
Total Area of Lesions (mm ²)	0.418 ± 0.414	0.568 ± 0.558	0.305 ± 0.233	0.297
Average Lesion Size (mm ²)	0.004 ± 0.004	0.007 ± 0.004	0.001 ± 0.001	< 0.001*
Size of Largest Lesion (mm ²)	0.015 ± 0.015	0.027 ± 0.016	0.006 ± 0.005	0.001*

Table 2. Fundus photograph quantification of drusenoid lesions in Rhesus macaques using digital imagesegmentation. *Statistically significant. #All values expressed as mean \pm standard deviation.





thicker in eyes with drusenoid lesions by approximately $3.4 \,\mu m$ (P = 0.012; Table 3). When comparing eyes with predominantly soft drusen versus hard punctate lesions (Fig. 4E,F), there were no significant differences in any of the retinal or choroidal layers (Table 3).

Genotype of Animals with Drusenoid Lesions. Genetic polymorphisms in the ARMS2 and HTRA1 loci associated with drusenoid lesions in rhesus macaques have been independently confirmed in colony animals from the Oregon National Primate Research Center, the Caribbean Primate Research Center (CPRC), and the German Primate Center (Deutsche Primatenzentrum or DPZ) in Gottingen, the majority of which originated from the CPRC colony from Cayo Santiago^{23–25}. In our cohort of animals, the high-risk allele in the ARMS2 locus (SNP position 10682 T > C) occurred at a higher frequency (70.7%) than previously reported at CPRC (20.3%) or DPZ (49.2%), with no significant difference between eyes with and without drusenoid lesions (P = 1.000; Table 4). The risk allele in HTRA1 (SNP position 17076 C > A) occurred at much lower frequency (4.6%) than the other colonies (21.0% to 60.0%), and also showed no association with presence of drusenoid lesions (P = 0.176). When comparing animals with the soft drusen and hard lesions, we found no difference in risk allele distribution in either the ARMS2 (P = 0.793) or HTRA1 (P = 0.471) loci. Our results suggest that the genetic factors previously



Figure 3. Multimodal imaging of hard punctate lesions in rhesus macaques. Fundus photographs (**A**), as well as fundus autofluorescence (**B**), infrared reflectance (**C**), and spectral-domain optical coherence tomography (**D**–**G**) images of a right eye of a rhesus macaque with hard, punctate lesions are shown, with corresponding lesions labeled with yellow, red, green, and blue circles (**A**–**C**) and lines (**D**–**G**). The colored dashed lines in (**B**) correspond to the line-scan segment shown in (**D**–**G**). Scale bars: $500 \,\mu$ m.

.....

found to confer increased risk of developing drusen in other rhesus colonies, may not contribute as strongly in our cohort of animals at CNPRC.

Discussion

Nonhuman primates are the only mammals that spontaneously develop drusenoid lesions similar to those in humans, and thus have significant translational potential as an animal model for developing therapies for AMD. Ocular imaging technologies are often employed in macaques for preclinical testing of novel treatments or drugs, but rarely for understanding the pathophysiology of age-related retinal changes that mimic human disease. In this study, we employed validated measures of drusenoid lesions from fundus photographs, as well as a battery of ocular imaging modalities employed in human clinical studies, to provide the first robust, quantitative characterization of drusenoid lesions in rhesus macaques. We describe two distinct clinical phenotypes of "soft" drusen and "hard" punctate lesions that can be distinguished by their size on fundus photographs and appearance on SD-OCT. We also provide important normative macular thickness values of retinal and choroidal layers on SD-OCT, and found a slightly thicker RPE in macaque eyes with drusenoid lesions. Taken together, these data provide an important framework for research in drusen formation and AMD using rhesus macaques.

The AREDS2 system for grading AMD lesions is derived from major clinical trials sponsored by the National Eye Institute to study the effects of nutritional antioxidant supplements including lutein and zeaxanthin in humans with AMD. This validated system is highly reproducible and predictive of 5-year risk for progression to advanced AMD in humans, which is defined by the presence of center-involving RPE atrophy or neovascular AMD³⁶. Interestingly, neither form of advanced AMD has ever been clearly documented in nonhuman primates, and we did not identify any cases in our survey of the rhesus colony at CNPRC. One explanation is that the occurrence is so rare that it has not yet been observed in animals under study. The drusen grading in most of our cohort was equivalent to an AREDS severity scale of 4, which in humans correspond to a rate of only 4.8% for progression to advanced AMD over 5 years³⁰. Many of the animals in the study are near the end of their lifespan, and thus may not survive long enough for this progression to occur. Rhesus macaques also do not have the same genetic risk factors for AMD as humans. Among a selection of 42 human DNA markers associated with drusen, 24 have been identified in rhesus monkeys, of which only 7 were polymorphic³⁷, and only the ARMS2/HTRA1





loci have been found to be associated with drusen in these animals²³⁻²⁵. In addition, our study animals are partially housed indoors (mean outdoor time $53.6 \pm 34.0\%$), and may have a healthier diet and lower sun exposure compared to their free-ranging counterparts. Studies have shown that rhesus monkeys fed a diet lacking macular pigment carotenoids lutein and zeaxanthin are more susceptible to blue-light damage and may develop drusen at an earlier age^{38,39}, suggesting that standard laboratory diets low in fat and rich in xanthophylls may limit drusen progression also. In fact, we found only a 30.7% prevalence of drusenoid lesions in our rhesus colony at CNPRC compared to the 47% noted from the same colony surveyed 22 years ago²⁰, and up to rates as high as 74% at other sites¹⁷, despite surveying an older cohort of animals (Table 5). From an anatomic standpoint, macaques have very dark uveal pigmentation⁴⁰, which in humans is protective against AMD⁴¹. None of the animals we examined demonstrated pigmentary changes in the macula, which are also strongly-associated with AMD progression in humans³⁰. In contrast, macular pigmentary changes have been more frequently reported in free-ranging animal colonies with incidences of up to $32\%^{15,18}$. Thus, developing nonhuman primate models with more advanced

	Normal Eyes (n = 14)	Any Drusen (n = 23)	P-value
Total retinal thickness	292.6 ± 21.4	314.0 ± 69.7	0.198
Nerve fiber layer (NFL)	15.7 ± 1.6	17.8 ± 4.6	0.051
Ganglion cell/inner plexiform layer (GCL/IPL)	51.5 ± 10.2	57.5 ± 15.7	0.244
Inner nuclear layer (INL)	29.8 ± 7.9	31.8 ± 7.7	0.515
Outer plexiform layer (OPL)	21.0 ± 3.0	23.1 ± 6.0	0.182
Outer nuclear layer (ONL)	83.3 ± 6.7	88.5 ± 17.4	0.253
Photoreceptor inner segments (IS)	33.0 ± 1.9	35.1±7.6	0.208
Photoreceptor outer segments (OS)	41.7 ± 4.5	40.3 ± 10.4	0.599
Retinal pigment epithelium (RPE)	16.5 ± 2.7	19.9 ± 4.6	0.012*
Choriocapillaris (CC)	20.2 ± 4.9	19.6 ± 3.3	0.728
Outer choroid (C)	158.8 ± 23.2	173.2 ± 40.5	0.202
Total choroidal thickness	179.1 ± 25.0	192.8 ± 41.7	0.246
	Soft Drusen (n = 12)	Hard Lesions (n = 11)	P-value
Total retinal thickness	Soft Drusen (n = 12) 306.8±15.4	Hard Lesions (n = 11) 321.8 ± 101.4	P-value 0.630
Total retinal thickness Nerve fiber layer (NFL)	Soft Drusen (n = 12) 306.8±15.4 16.7±1.5	Hard Lesions (n = 11) 321.8±101.4 19.2±6.3	P-value 0.630 0.202
Total retinal thickness Nerve fiber layer (NFL) Ganglion cell/inner plexiform layer (GCL/IPL)	Soft Drusen (n = 12) 306.8 ± 15.4 16.7 ± 1.5 55.1 ± 6.8	Hard Lesions (n=11) 321.8±101.4 19.2±6.3 60.0±21.9	P-value 0.630 0.202 0.487
Total retinal thickness Nerve fiber layer (NFL) Ganglion cell/inner plexiform layer (GCL/IPL) Inner nuclear layer (INL)	Soft Drusen (n = 12) 306.8 ± 15.4 16.7 ± 1.5 55.1 ± 6.8 31.1 ± 3.9	Hard Lesions (n = 11) 321.8 ± 101.4 19.2 ± 6.3 60.0 ± 21.9 32.6 ± 10.6	P-value 0.630 0.202 0.487 0.667
Total retinal thickness Nerve fiber layer (NFL) Ganglion cell/inner plexiform layer (GCL/IPL) Inner nuclear layer (INL) Outer plexiform layer (OPL)	Soft Drusen (n = 12) 306.8±15.4 16.7±1.5 55.1±6.8 31.1±3.9 21.3±2.8	Hard Lesions (n = 11) 321.8±101.4 19.2±6.3 60.0±21.9 32.6±10.6 25.0±8.0	P-value 0.630 0.202 0.487 0.667 0.143
Total retinal thickness Nerve fiber layer (NFL) Ganglion cell/inner plexiform layer (GCL/IPL) Inner nuclear layer (INL) Outer plexiform layer (OPL) Outer nuclear layer (ONL)	Soft Drusen (n = 12) 306.8±15.4 16.7±1.5 55.1±6.8 31.1±3.9 21.3±2.8 90.8±5.1	Hard Lesions (n = 11) 321.8 ± 101.4 19.2 ± 6.3 60.0 ± 21.9 32.6 ± 10.6 25.0 ± 8.0 85.9 ± 24.9	P-value 0.630 0.202 0.487 0.667 0.143 0.528
Total retinal thickness Nerve fiber layer (NFL) Ganglion cell/inner plexiform layer (GCL/IPL) Inner nuclear layer (INL) Outer plexiform layer (OPL) Outer nuclear layer (ONL) Photoreceptor inner segments (IS)	Soft Drusen (n = 12) 306.8±15.4 16.7±1.5 55.1±6.8 31.1±3.9 21.3±2.8 90.8±5.1 33.6±0.8	Hard Lesions (n = 11) 321.8 ± 101.4 19.2 ± 6.3 60.0 ± 21.9 32.6 ± 10.6 25.0 ± 8.0 85.9 ± 24.9 36.7 ± 11.0	P-value 0.630 0.202 0.487 0.667 0.143 0.528 0.354
Total retinal thickness Nerve fiber layer (NFL) Ganglion cell/inner plexiform layer (GCL/IPL) Inner nuclear layer (INL) Outer plexiform layer (OPL) Outer nuclear layer (ONL) Photoreceptor inner segments (IS) Photoreceptor outer segments (OS)	Soft Drusen (n = 12) 306.8±15.4 16.7±1.5 55.1±6.8 31.1±3.9 21.3±2.8 90.8±5.1 33.6±0.8 38.5±4.4	Hard Lesions (n = 11) 321.8 ± 101.4 19.2 ± 6.3 60.0 ± 21.9 32.6 ± 10.6 25.0 ± 8.0 85.9 ± 24.9 36.7 ± 11.0 42.2 ± 14.4	P-value 0.630 0.202 0.487 0.667 0.143 0.528 0.354 0.415
Total retinal thickness Nerve fiber layer (NFL) Ganglion cell/inner plexiform layer (GCL/IPL) Inner nuclear layer (INL) Outer plexiform layer (OPL) Outer nuclear layer (ONL) Photoreceptor inner segments (IS) Photoreceptor outer segments (OS) Retinal pigment epithelium (RPE)	Soft Drusen (n = 12) 306.8 ± 15.4 16.7 ± 1.5 55.1 ± 6.8 31.1 ± 3.9 21.3 ± 2.8 90.8 ± 5.1 33.6 ± 0.8 38.5 ± 4.4 19.7 ± 2.9	Hard Lesions (n = 11) 321.8 ± 101.4 19.2 ± 6.3 60.0 ± 21.9 32.6 ± 10.6 25.0 ± 8.0 85.9 ± 24.9 36.7 ± 11.0 42.2 ± 14.4 20.1 ± 6.1	P-value 0.630 0.202 0.487 0.667 0.143 0.528 0.354 0.415 0.844
Total retinal thickness Nerve fiber layer (NFL) Ganglion cell/inner plexiform layer (GCL/IPL) Inner nuclear layer (INL) Outer plexiform layer (OPL) Outer nuclear layer (ONL) Photoreceptor inner segments (IS) Photoreceptor outer segments (OS) Retinal pigment epithelium (RPE) Choriocapillaris (CC)	Soft Drusen (n = 12) 306.8 ± 15.4 16.7 ± 1.5 55.1 ± 6.8 31.1 ± 3.9 21.3 ± 2.8 90.8 ± 5.1 33.6 ± 0.8 38.5 ± 4.4 19.7 ± 2.9 19.9 ± 3.3	Hard Lesions (n = 11) 321.8 ± 101.4 19.2 ± 6.3 60.0 ± 21.9 32.6 ± 10.6 25.0 ± 8.0 85.9 ± 24.9 36.7 ± 11.0 42.2 ± 14.4 20.1 ± 6.1 19.3 ± 3.5	P-value 0.630 0.202 0.487 0.667 0.143 0.528 0.354 0.415 0.844 0.663
Total retinal thicknessNerve fiber layer (NFL)Ganglion cell/inner plexiform layer (GCL/IPL)Inner nuclear layer (INL)Outer plexiform layer (OPL)Outer nuclear layer (ONL)Photoreceptor inner segments (IS)Photoreceptor outer segments (OS)Retinal pigment epithelium (RPE)Choriocapillaris (CC)Outer choroid (C)	Soft Drusen (n = 12) 306.8 ± 15.4 16.7 ± 1.5 55.1 ± 6.8 31.1 ± 3.9 21.3 ± 2.8 90.8 ± 5.1 33.6 ± 0.8 38.5 ± 4.4 19.7 ± 2.9 19.9 ± 3.3 16.7 ± 38.0	Hard Lesions (n = 11) 321.8 ± 101.4 19.2 ± 6.3 60.0 ± 21.9 32.6 ± 10.6 25.0 ± 8.0 85.9 ± 24.9 36.7 ± 11.0 42.2 ± 14.4 20.1 ± 6.1 19.3 ± 3.5 179.7 ± 44.0	P-value 0.630 0.202 0.487 0.667 0.143 0.528 0.354 0.415 0.844 0.663

 Table 3. Retinal and choroidal layer thicknesses of age-matched normal eyes and eyes with drusenoid lesions in rhesus macaques. *Statistically significant.

		Frequ		
	Genotype	Normal Eyes	Any Drusen	P-value
ARMS2	TT	4	1	1.000
SNP 10682	CT	19	9	
	CC	22	10	
HTRA1	CC	43	18	0.176
SNP 17076	AC	0	2	
	AA	2	0	
		Soft Drusen	Hard Lesions	
ARMS2	TT	1	0	0.793
SNP 10682	CT	3	3	
	CC	4	6	
HTRA1	CC	7	9	0.471
SNP 17076	AC	1	0	
	AA	0	0	

 Table 4.
 Frequencies of risk alleles in ARMS2 and HTRA1 associated with drusen in rhesus macaques.

 Abbreviations: SNP, Single-nucleotide polymorphisms.

•

stages of AMD may require long-term dietary and environmental modifications that better represent human behavior. Interestingly, Japanese macaques (Macaca fuscata) have an early-onset form of drusen with a dominant mode of inheritance⁴². Cynomologus macaques (Macaca fascicularis) also develop drusen in an age-dependent manner with a potential link to hematologic markers⁴³. Thus, the quantitative and imaging techniques used in our study may provide the necessary platform to study other nonhuman primate models of AMD in a more rigorous manner.

Modern ocular imaging technologies such as SD-OCT and FAF provide new ways to characterize drusen phenotypes in humans and macaques. SD-OCT of soft drusen in rhesus monkeys appear as dome-shaped hyper-reflective deposits under the RPE, which is very similar to those seen in AMD⁴⁴⁻⁴⁷. However, the soft drusen in macaques appear more uniform in morphology, and lack any reflective drusen substructures or hyperreflective foci in the overlying outer retina which are predictive of progression to geographic atrophy in humans^{47,48}. We also did not find any reticular pseudodrusen, which are more strongly associated with geographic atrophy⁴⁹, in any of the aged animals in our colony. The absence of these drusen abnormalities may further explain the absence

	Location	Total Animals	Age Range	Drusen Frequency	Pigment Change
El Mofty et al. 1978 ¹⁵	CPRC (free-ranging)	105	1-21	30%	32%
Stafford et al. 1984 ¹⁹	Various (most indoor)	574	10-31	5.9%	NR
Engel et al. 1988 ¹⁷	CPRC (free-ranging)	29	13-27	74%	NR
Monaco <i>et al.</i> 1990 ¹⁸	Naval Aerospace Research Lab (indoor & free-ranging)	100	1-13	31%	10%
Hope <i>et al.</i> 1992 ¹⁴	CPRC (free-ranging)	246	1-30	57.7%	NR
Olin <i>et al.</i> 1995 ²⁰	CRPRC & WRPRC (indoor)	62	20-33	47%	NR
Gouras et al. 2008 ²¹	NIH & ONPRC (indoor)	121	10-39	61%	NR
Current study 2017	CNPRC (indoor & outdoor)	61	19-30	30.7%	0%

Table 5. Summary of published studies of drusen frequency in various primate research facilities. Abbreviations: CPRC, Caribbean Primate Research Center; CRPRC, California Regional Primate Research Center; WRPRC, Wisconsin Regional Primate Research Center; NIH, National Institute of Health; ONPRC, Oregon National Primate Research Center; NR, not recorded.

of any advanced AMD phenotype in nonhuman primates. Although we noted extramacular drusen in a selection of animals, their association with AMD progression in humans remain unclear. Peripheral drusen are more frequently observed in AMD patients, but may not be associated with major genetic risk alleles for AMD such as CFH and ARMS2⁵⁰. Further studies using ultrawide-field imaging may better characterize the role of extramacular drusen in rhesus macaques⁵¹.

In this study, we also described hard, punctate lesions in rhesus macaques that are undetectable on SD-OCT. We cannot exclude the possibility that the lesions are so small that they are below the detection threshold of SD-OCT, although this is unlikely since the imaging system has an axial resolution of 3.87 µm per pixel, a transverse resolution of 4.82 µm per pixel, and a line scan density of 51 µm between B-scans, compared to the large number of punctate lesions sampled which on average were approximately 10 µm in diameter on fundus photographs. Interestingly, similar small hypopigmented lesions have been described on histology in macaques as a form of lipoidal degeneration where RPE cells become filled with vacuoles and lipid droplets, rather than true sub-RPE deposits as seen with AMD drusen⁵²⁻⁵⁴. It is unclear, however, whether these punctate lesions represent a separate entity or are precursors to larger soft drusen. We noted that eyes with mostly soft drusen also have punctate lesions, but those with predominantly hard lesions rarely exhibit any soft drusen. Animals with soft drusen are also older than those with mainly hard drusen (25.2 ± 2.5 years vs. 22.3 ± 2.2 years; P = 0.024). Finally, while eyes with drusenoid lesions demonstrated RPE thickening, there was no significant difference in RPE thickness between animals with soft drusen or hard lesions. Together, these evidences suggest that the two drusen phenotypes may be different manifestations of the same pathophysiological process, and that hard, punctate lesions may occur at an earlier stage. This is further supported by the similar distribution of risk alleles in ARMS2 and HTRA1 between animals with the two drusen phenotypes. Since SD-OCT imaging affords the ability to monitor morphologic changes of these lesions over time, future longitudinal studies using SD-OCT and fundus photography may enhance our understanding of drusen progression in nonhuman primates.

FAF imaging provides another measure of disease activity in retinal conditions, although the precise source of the autofluorescence remains controversial^{55–57}. Many hypothesize that blue-peak FAF is a measure of lipofuscin, autofluorescent pigment granules in RPE cells that result from lysosomal digestion of photoreceptor outer segments⁵⁸. Rhesus monkeys deficient in lutein, zeaxanthin, and omega-3 fatty acids, for example, may exhibit elevated FAF intensities and are also associated with higher incidences of drusen lesions⁵⁹. However, the FAF signature of macaque drusen have never been clearly explored. In humans, drusen may exhibit variable FAF patterns, likely due to differences in drusen size, composition, and effects on overlying RPE and photoreceptors^{60,61}. In our study, different drusenoid lesions exhibit distinct levels of FAF, and may represent variable amounts and types of fluorophores within the lesions themselves or different stages of lipofuscin accumulation or RPE degeneration. Thus, including FAF imaging with longitudinal studies may further our insight into the pathophysiology of drusen development and RPE atrophy in these animals^{62,63}.

Our study employs a systematic approach using multimodal ocular imaging to characterize and monitor drusenoid lesions in the aged rhesus colony at CNPRC. In contrast to prior reports from other primate centers where the prevalence of drusenoid lesions is higher, the current study is limited by the smaller cohort of affected animals at CNPRC. Compared to human subjects, ocular imaging of macaques is more challenging, with occasional ocular pathologies such as corneal scars or senile cataracts limiting our ability to attain high-resolution images in every animal. Also, while the AREDS2 grading system has been validated in human studies, the modified form calibrated from the anatomic dimensions of the macaque fundus still require further assessment. Nevertheless, using the tools developed in this research, future studies to characterize the progression of druse-noid lesions in rhesus macaques will be imperative toward establishing a powerful nonhuman primate model of AMD.

References

- 1. Curcio, C. A., Millican, C. L., Bailey, T. & Kruth, H. S. Accumulation of cholesterol with age in human Bruch's membrane. Investigative ophthalmology & visual science 42, 265–274 (2001).
- Curcio, C. A. *et al.* Esterified and unesterified cholesterol in drusen and basal deposits of eyes with age-related maculopathy. *Exp Eye Res* 81, 731–741, https://doi.org/10.1016/j.exer.2005.04.012 (2005).

- Curcio, C. A., Johnson, M., Huang, J. D. & Rudolf, M. Aging, age-related macular degeneration, and the response-to-retention of apolipoprotein B-containing lipoproteins. *Progress in retinal and eye research* 28, 393–422, https://doi.org/10.1016/j. preteyeres.2009.08.001 (2009).
- Crabb, J. W. et al. Drusen proteome analysis: an approach to the etiology of age-related macular degeneration. Proceedings of the National Academy of Sciences of the United States of America 99, 14682–14687, https://doi.org/10.1073/pnas.222551899 (2002).
- Malek, G., Li, C. M., Guidry, C., Medeiros, N. E. & Curcio, C. A. Apolipoprotein B in cholesterol-containing drusen and basal deposits of human eyes with age-related maculopathy. *Am J Pathol* 162, 413–425, https://doi.org/10.1016/S0002-9440(10)63836-9 (2003).
- Mullins, R. F., Russell, S. R., Anderson, D. H. & Hageman, G. S. Drusen associated with aging and age-related macular degeneration contain proteins common to extracellular deposits associated with atherosclerosis, elastosis, amyloidosis, and dense deposit disease. *FASEB J* 14, 835–846 (2000).
- Gass, J. D., Jallow, S. & Davis, B. Adult vitelliform macular detachment occurring in patients with basal laminar drusen. Am J Ophthalmol 99, 445–459 (1985).
- Russell, S. R., Mullins, R. F., Schneider, B. L. & Hageman, G. S. Location, substructure, and composition of basal laminar drusen compared with drusen associated with aging and age-related macular degeneration. Am J Ophthalmol 129, 205–214 (2000).
- Zweifel, S. A., Spaide, R. F., Curcio, C. A., Malek, G. & Imamura, Y. Reticular pseudodrusen are subretinal drusenoid deposits. Ophthalmology 117, 303–312 e301, https://doi.org/10.1016/j.ophtha.2009.07.014 (2010).
- Rudolf, M. *et al.* Sub-retinal drusenoid deposits in human retina: organization and composition. *Exp Eye Res* 87, 402–408, https:// doi.org/10.1016/j.exer.2008.07.010 (2008).
- Heiferman, M. J. *et al.* Reticular Pseudodrusen on Infrared Imaging Are Topographically Distinct from Subretinal Drusenoid Deposits on En Face Optical Coherence Tomography. *Retina* 35, 2593–2603, https://doi.org/10.1097/IAE.00000000000666 (2015).
- Zhou, Q. et al. Pseudodrusen and Incidence of Late Age-Related Macular Degeneration in Fellow Eyes in the Comparison of Age-Related Macular Degeneration Treatments Trials. Ophthalmology 123, 1530–1540, https://doi.org/10.1016/j.ophtha.2016.02.043 (2016).
- Smith, R. T. et al. Complement factor H 402H variant and reticular macular disease. Arch Ophthalmol 129, 1061–1066, https://doi. org/10.1001/archophthalmol.2011.212 (2011).
- 14. Hope, G. M. et al. A primate model for age related macular drusen. Br J Ophthalmol 76, 11-16 (1992).
- 15. El-Mofty, A., Gouras, P., Eisner, G. & Balazs, E. A. Macular degeneration in rhesus monkey (Macaca mulatta). *Exp Eye Res* 27, 499–502 (1978).
- Dawson, W. W., Ulshafer, R. J., Engel, H. M., Hope, G. M. & Kessler, M. J. Macular disease in related rhesus monkeys. Doc Ophthalmol 71, 253–263 (1989).
- Engel, H. M., Dawson, W. W., Ulshafer, R. J., Hines, M. W. & Kessler, M. J. Degenerative changes in maculas of rhesus monkeys. Ophthalmologica 196, 143–150 (1988).
- Monaco, W. A. & Wormington, C. M. The rhesus monkey as an animal model for age-related maculopathy. Optometry and vision science: official publication of the American Academy of Optometry 67, 532–537 (1990).
- 19. Stafford, T. J., Anness, S. H. & Fine, B. S. Spontaneous degenerative maculopathy in the monkey. Ophthalmology 91, 513–521 (1984).
- Olin, K. L. et al. Trace element status and free radical defense in elderly rhesus macaques (Macaca mulatta) with macular drusen. Proc Soc Exp Biol Med 208, 370–377 (1995).
- 21. Gouras, P. *et al.* Drusenoid maculopathy in rhesus monkeys (Macaca mulatta): effects of age and gender. *Graefes Arch Clin Exp* Ophthalmol 246, 1395–1402, https://doi.org/10.1007/s00417-008-0910-8 (2008).
- Umeda, S. *et al.* Molecular composition of drusen and possible involvement of anti-retinal autoimmunity in two different forms of macular degeneration in cynomolgus monkey (Macaca fascicularis). *FASEB J* 19, 1683–1685, https://doi.org/10.1096/fj.04-3525fje (2005).
- Francis, P. J. et al. Rhesus monkeys and humans share common susceptibility genes for age-related macular disease. Hum Mol Genet 17, 2673–2680, https://doi.org/10.1093/hmg/ddn167 (2008).
- Singh, K. K., Krawczak, M., Dawson, W. W. & Schmidtke, J. Association of HTRA1 and ARMS2 gene variation with drusen formation in rhesus macaques. *Exp Eye Res* 88, 479–482, https://doi.org/10.1016/j.exer.2008.10.019 (2009).
- Pahl, L. et al. Characterization of the 10q26-orthologue in rhesus monkeys corroborates a functional connection between ARMS2 and HTRA1. Exp Eye Res 98, 75–78, https://doi.org/10.1016/j.exer.2012.03.007 (2012).
- Danis, R. P. et al. Methods and reproducibility of grading optimized digital color fundus photographs in the Age-Related Eye Disease Study 2 (AREDS2 Report Number 2). Investigative ophthalmology & visual science 54, 4548–4554, https://doi.org/10.1167/iovs.13-11804 (2013).
- Yehoshua, Z. et al. Comparison of drusen area detected by spectral domain optical coherence tomography and color fundus imaging. Investigative ophthalmology & visual science 54, 2429–2434, https://doi.org/10.1167/iovs.12-11569 (2013).
- Classification of diabetic retinopathy from fluorescein angiograms. ETDRS report number 11. Early Treatment Diabetic Retinopathy Study Research Group. Ophthalmology 98, 807–822 (1991).
- Jonas, J. B. & Hayreh, S. S. Ophthalmoscopic appearance of the normal optic nerve head in rhesus monkeys. Investigative ophthalmology & visual science 41, 2978-2983 (2000).
- Davis, M. D. et al. The Age-Related Eye Disease Study severity scale for age-related macular degeneration: AREDS Report No. 17. Arch Ophthalmol 123, 1484–1498, https://doi.org/10.1001/archopht.123.11.1484 (2005).
- Chiu, S. J., Toth, C. A., Rickman, B. C., Izatt, J. A. & Farsiu, S. Automatic segmentation of closed-contour features in ophthalmic images using graph theory and dynamic programming. *Biomed. Opt. Express* 3, 1127–1140 (2012).
- Farsiu, S. et al. Quantitative Classification of Eyes with and without Intermediate Age-related Macular Degeneration Using Optical Coherence Tomography. Ophthalmology 121, 162–172, https://doi.org/10.1016/j.ophtha.2013.07.013 (2014).
- Vuong, V. S. et al. Repeatability of Choroidal Thickness Measurements on Enhanced Depth Imaging Optical Coherence Tomography Using Different Posterior Boundaries. Am J Ophthalmol 169, 104–112, https://doi.org/10.1016/j.ajo.2016.06.023 (2016).
- 34. Yiu, G. *et al.* Characterization of the choroid-scleral junction and suprachoroidal layer in healthy individuals on enhanced-depth imaging optical coherence tomography. *JAMA ophthalmology* 132, 174–181, https://doi.org/10.1001/jamaophthalmol.2013.7288 (2014).
- Yiu, G. et al. Effect of Uveal Melanocytes on Choroidal Morphology in Rhesus Macaques and Humans on Enhanced-Depth Imaging Optical Coherence Tomography. Investigative ophthalmology & visual science (2016).
- 36. Vitale, S. et al. Evaluating the Validity of the Age-Related Eye Disease Study Grading Scale for Age-Related Macular Degeneration: AREDS2 Report 10. JAMA ophthalmology 134, 1041–1047, https://doi.org/10.1001/jamaophthalmol.2016.2383 (2016).
- Singh, K. K., Ristau, S., Dawson, W. W., Krawczak, M. & Schmidtke, J. Mapping of a macular drusen susceptibility locus in rhesus macaques to the homologue of human chromosome 6q14-15. *Exp Eye Res* 81, 401–406, https://doi.org/10.1016/j.exer.2005.02.011 (2005).
- 38. Neuringer, M. Nonhuman Primate Models of Macular Degeneration (conference paper) in Association for Research in Vision & Ophthalmology Annual Meeting. Seattle, WA (2016).

- Barker, F. M. 2nd *et al.* Nutritional manipulation of primate retinas, V: effects of lutein, zeaxanthin, and n-3 fatty acids on retinal sensitivity to blue-light-induced damage. *Investigative ophthalmology & visual science* 52, 3934–3942, https://doi.org/10.1167/ iovs.10-5898 (2011).
- Yiu, G. *et al.* Effect of Uveal Melanocytes on Choroidal Morphology in Rhesus Macaques and Humans on Enhanced-Depth Imaging Optical Coherence Tomography. *Investigative ophthalmology & visual science* 57, 5764–5771, https://doi.org/10.1167/iovs.16-20070 (2016).
- Frank, R. N., Puklin, J. E., Stock, C. & Canter, L. A. Race, iris color, and age-related macular degeneration. *Transactions of the American Ophthalmological Society* 98, 109–115; discussion 115–107 (2000).
- Pennesi, M. E. et al. Measuring cone density in a Japanese macaque (Macaca fuscata) model of age-related macular degeneration with commercially available adaptive optics. Adv Exp Med Biol 801, 309–316, https://doi.org/10.1007/978-1-4614-3209-8_39 (2014).
- Nishiguchi, K. M. et al. Analysis of Macular Drusen and Blood Test Results in 945 Macaca fascicularis. PloS one 11, e0164899, https://doi.org/10.1371/journal.pone.0164899 (2016).
- Gregori, G. et al. Spectral domain optical coherence tomography imaging of drusen in nonexudative age-related macular degeneration. Ophthalmology 118, 1373–1379, https://doi.org/10.1016/j.ophtha.2010.11.013 (2011).
- Yehoshua, Z. *et al.* Natural history of drusen morphology in age-related macular degeneration using spectral domain optical coherence tomography. *Ophthalmology* 118, 2434–2441, https://doi.org/10.1016/j.ophtha.2011.05.008 (2011).
- Spaide, R. F. & Curcio, C. A. Drusen characterization with multimodal imaging. *Retina* 30, 1441–1454, https://doi.org/10.1097/ IAE.0b013e3181ee5ce8 (2010).
- Ouyang, Y., Heussen, F. M., Hariri, A., Keane, P. A. & Sadda, S. R. Optical coherence tomography-based observation of the natural history of drusenoid lesion in eyes with dry age-related macular degeneration. *Ophthalmology* 120, 2656–2665, https://doi. org/10.1016/j.ophtha.2013.05.029 (2013).
- Veerappan, M. *et al.* Optical Coherence Tomography Reflective Drusen Substructures Predict Progression to Geographic Atrophy in Age-related Macular Degeneration. *Ophthalmology* 123, 2554–2570, https://doi.org/10.1016/j.ophtha.2016.08.047 (2016).
- Gliem, M. et al. Quantitative Fundus Autofluorescence in Early and Intermediate Age-Related Macular Degeneration. JAMA ophthalmology 134, 817–824, https://doi.org/10.1001/jamaophthalmol.2016.1475 (2016).
- Ersoy, L. et al. Extramacular drusen are highly associated with age-related macular degeneration, but not with CFH and ARMS2 genotypes. Br J Ophthalmol 100, 1047–1051, https://doi.org/10.1136/bjophthalmol-2015-306806 (2016).
- 51. Writing Committee for the, O. P. R. S. *et al.* Peripheral Retinal Changes Associated with Age-Related Macular Degeneration in the Age-Related Eye Disease Study 2: Age-Related Eye Disease Study 2 Report Number 12 by the Age-Related Eye Disease Study 2 Optos PEripheral RetinA (OPERA) Study Research Group. *Ophthalmology* 124, 479–487, https://doi.org/10.1016/j.ophtha.2016.12.004 (2017).
- Fine, B. S. & Kwapien, R. P. Pigment epithelial windows and drusen: an animal model. *Investigative ophthalmology & visual science* 17, 1059–1068 (1978).
- 53. Feeney-Burns, L., Malinow, R., Klein, M. L. & Neuringer, M. Maculopathy in cynomolgus monkeys. A correlated fluorescein angiographic and ultrastructural study. Arch Ophthalmol 99, 664–672 (1981).
- Anderson, M., Dawson, W. W., Gonzalez-Martinez, J. & Curcio, C. A. Drusen and lipid-filled retinal pigment epithelium cells in a rhesus macula. Vet Ophthalmol 9, 201–207, https://doi.org/10.1111/j.1463-5224.2006.00463.x (2006).
- Smith, R. T., Farsiu, S. & Allingham, M. Understanding RPE LipofuscinLetters. Investigative Ophthalmology & Visual Science 57, 6766–6766, https://doi.org/10.1167/iovs.16-21081 (2016).
- Sparrow, J. R. & Boulton, M. RPE lipofuscin and its role in retinal pathobiology. *Experimental Eye Research* 80, 595–606, https://doi. org/10.1016/j.exer.2005.01.007 (2005).
- Rudolf, M. *et al.* Histologic Basis of Variations in Retinal Pigment Epithelium Autofluorescence in Eyes with Geographic Atrophy. Ophthalmology 120, 821–828, https://doi.org/10.1016/j.ophtha.2012.10.007 (2013).
- Delori, F. C. et al. In vivo fluorescence of the ocular fundus exhibits retinal pigment epithelium lipofuscin characteristics. Investigative Ophthalmology & Visual Science 36, 718–729 (1995).
- McGill, T. J., Renner, L. M. & Neuringer, M. Elevated Fundus Autofluorescence in Monkeys Deficient in Lutein, Zeaxanthin, and Omega-3 Fatty Acids. Investigative ophthalmology & visual science 57, 1361–1369, https://doi.org/10.1167/iovs.15-18596 (2016).
- 60. Bindewald, A. et al. Classification of fundus autofluorescence patterns in early age-related macular disease. Investigative ophthalmology & visual science 46, 3309-3314, https://doi.org/10.1167/iovs.04-0430 (2005).
- Landa, G., Rösen, R. B., Pilavas, J. & Garcia, P. M. Drusen characteristics revealed by spectral-domain optical coherence tomography and their corresponding fundus autofluorescence appearance in dry age-related macular degeneration. *Ophthalmic Res* 47, 81–86, https://doi.org/10.1159/000324988 (2012).
- Allingham, M. J. et al. Semiautomatic segmentation of rim area focal hyperautofluorescence predicts progression of geographic atrophy due to dry age related macular degeneration. Investigative Ophthalmology & Visual Science 57, 2283–2289 (2016).
- 63. Holz, F. G., Bellman, C., Staudt, S., Schütt, F. & Völcker, H. E. Fundus Autofluorescence and Development of Geographic Atrophy in Age-Related Macular Degeneration. *Investigative Ophthalmology & Visual Science* 42, 1051–1056 (2001).

Acknowledgements

We thank Dr. Philip Kass at UC Davis School of Veterinary Medicine for input with statistical analyses. This study was supported by the California National Primate Research Center base grant NIH P510D011107. GY is supported by NIH K08 EY026101, the E. Matilda Ziegler Foundation for the Blind, Alcon Research Institute, ARVO Foundation, Barr Foundation for Retinal Research, and the California National Primate Research Center. SF is supported by NIH P30 EY005722. ST is supported by NIH K08 EY021142. No funding organizations had any role in the design or conduct of this research. The content is solely the responsibility of the authors and does not necessarily represent the official views of the funding agencies.

Author Contributions

G.Y. designed, gathered, and analyzed all the experimental data, along with support from E.T. (genotyping); C.M., B.W., D.C., and S.F. (image processing); and L.G., J.R., and S.T. (animal support). G.Y. wrote the main manuscript. G.Y., E.T., D.C., S.F., L.G., J.R., and S.T. critically reviewed the manuscript.

Additional Information

Supplementary information accompanies this paper at https://doi.org/10.1038/s41598-017-14715-z.

Competing Interests: The authors declare that they have no competing interests.

Publisher's note: Springer Nature remains neutral with regard to jurisdictional claims in published maps and institutional affiliations.

Open Access This article is licensed under a Creative Commons Attribution 4.0 International License, which permits use, sharing, adaptation, distribution and reproduction in any medium or format, as long as you give appropriate credit to the original author(s) and the source, provide a link to the Creative Commons license, and indicate if changes were made. The images or other third party material in this article are included in the article's Creative Commons license, unless indicated otherwise in a credit line to the material. If material is not included in the article's Creative Commons license and your intended use is not permitted by statutory regulation or exceeds the permitted use, you will need to obtain permission directly from the copyright holder. To view a copy of this license, visit http://creativecommons.org/licenses/by/4.0/.

© The Author(s) 2017