

UC San Diego

UC San Diego Previously Published Works

Title

Association of Subjective Social Status With Life's Simple 7s Cardiovascular Health Index Among Hispanic/Latino People: Results From the HCHS/SOL

Permalink

<https://escholarship.org/uc/item/94c1999r>

Journal

Journal of the American Heart Association, 10(16)

ISSN

2047-9980

Authors

Piedra, Lissette M
Andrade, Flavia CD
Hernandez, Rosalba
[et al.](#)

Publication Date

2021-08-17

DOI

10.1161/jaha.119.012704






Copyright Information

This work is made available under the terms of a Creative Commons Attribution-NonCommercial-NoDerivatives License, available at <https://creativecommons.org/licenses/by-nc-nd/4.0/>

Peer reviewed

ORIGINAL RESEARCH

Association of Subjective Social Status With Life's Simple 7s Cardiovascular Health Index Among Hispanic/Latino People: Results From the HCHS/SOL

Lisette M. Piedra , PhD, MSW; Flavia C. D. Andrade, PhD, MA, MS; Rosalba Hernandez, PhD, MS; Krista M. Perreira, PhD; Linda C. Gallo , PhD; Hector M. González, PhD; Sara Gonzalez, PhD, MPH; Jianwen Cai , PhD; Jinsong Chen, PhD; Sheila F. Castañeda, PhD; K. M. Tabb, PhD; Gregory A. Talavera, MD, MPH; Ramon A. Durazo-Arvizu , PhD; Martha L. Daviglus , MD, PhD

BACKGROUND: Evidence suggests that subjective (perceived) social status (SSS) may predict health outcomes more strongly than objective social status, but little is known about the relationship between SSS and cardiovascular health (CVH). This study focuses on this relationship among diverse Hispanic/Latino adults because while poor CVH profiles are prevalent in this population, immigration complicates attempts to measure their social status.

METHODS AND RESULTS: We analyzed baseline HCHS/SOL (Hispanic Community Health Study/Study of Latinos) data on 15 374 Hispanic/Latino adults aged 18 to 74 years in 2008 to 2011. SSS was assessed using the McArthur Scale, a 10-rung “social ladder.” CVH was based on levels of 7 metrics defined by the American Heart Association. Linear and logistic regressions were used to examine cross-sectional associations of SSS with CVH (overall and single metrics) after adjusting for objective social status, demographic, and health factors. Less than half of the population (46%) had Ideal scores in ≥ 4 metrics of CVH. In multivariable-adjusted models, an increase in SSS was associated with a higher overall CVH score ($\beta=0.04$; 95% CI, 0.01–0.06) and greater likelihood of Ideal levels of body mass index, physical activity, and fasting blood glucose levels. Nativity and time in the United States modified the association between SSS and Ideal smoking.

CONCLUSIONS: Subjective measures of social status can enhance an understanding of CVH among Hispanic/Latino people. Future studies should explore the stability of SSS over time in comparison with objective social status and the mechanisms through which SSS may influence CVH.

Key Words: cardiovascular health ■ Hispanic/Latino ■ subjective social status

Subjective measures of social status (SSS) quantify how people perceive their position in the social hierarchy.¹ Self-perceived social inequalities can exacerbate psychological distress and negatively affect health beyond absolute socioeconomic standing.^{2–5} A meta-analysis that pooled 9 studies of adults aged ≥ 18 years found that low SSS increases the odds of coronary artery disease, hypertension, diabetes mellitus, dyslipidemia, and obesity, with a modest

attenuation after adjusting for objective measures of social status (OSS).⁶ Although OSS explained some of the association between SSS and these outcomes, the association between SSS and dyslipidemia remained. However, while these studies disaggregated their findings by race, none of these included a Hispanic/Latino group.

Evidence suggests traditional OSS measures of socioeconomic disparities have a weaker connection

Correspondence to: Lisette M. Piedra, PhD, MSW, School of Social Work, University of Illinois at Urbana-Champaign, 1010 West Nevada St., Urbana, IL 61801. E-mail: lpiedra@illinois.edu

For Sources of Funding and Disclosures, see page 14.

© 2021 The Authors. Published on behalf of the American Heart Association, Inc., by Wiley. This is an open access article under the terms of the Creative Commons Attribution-NonCommercial-NoDerivs License, which permits use and distribution in any medium, provided the original work is properly cited, the use is non-commercial and no modifications or adaptations are made.

JAHA is available at: www.ahajournals.org/journal/jaha

CLINICAL PERSPECTIVE

What Is New?

- This study examines the relationship between subjective social status (SSS) and cardiovascular health among a diverse sample of Latino people.
- Evidence suggests that higher levels of SSS are associated with cardiovascular health.

What Are the Clinical Implications?

- SSS could prove to be a protective factor, especially at the population level, where small differences that compound over time matter.
- SSS could enhance an understanding of cardiovascular health among Hispanic/Latino people.
- When evaluating cardiovascular health, clinicians should consider patients' SSS along with other factors.

Nonstandard Abbreviations and Acronyms

CVH	cardiovascular health
HCHS/SOL	Hispanic Community Health Study/ Study of Latinos
OSS	objective social status
PCS	Physical Component Summary

to health outcomes for the Hispanic/Latino population, especially for the foreign-born, than other ethnic/racial groups in the United States.^{7,8} Evidence is particularly mixed with respect to the impact of education.^{9,10} In one study, investigators found that education correlated with a number of health-related variables among non-Hispanic White people but not for people of Mexican origin in the United States.¹⁰ They offered a number of migratory-related explanations, including the fact that higher socioeconomic status (SES) in Mexico is associated with more unhealthy behaviors, such as smoking. However, in a recent study that used data from Mexico and the United States, Beltrán-Sánchez and colleagues⁷ found steeper educational gradients among people living in urban areas in Mexico compared with those living outside metropolitan regions in 5 of 6 biomarkers of metabolic syndrome. They argue that the flatter SES gradients observed in the United States may partially be explained by the presence of a sizeable immigrant population of rural Mexicans with low levels of education.

Markides and Eschbach⁸ argue that the *meaning* of traditional OSS factors—low education, income, occupational status, and overall SES—for Hispanic/Latino

subgroups merits further scrutiny. Factors such as discrimination related to citizenship status, country of origin, and level of acculturation may influence perceptions of economic standing such that measures used traditionally for Americans of European ancestry with multigenerational lineage in the United States do not apply to Hispanic/Latino people. Furthermore, traditional OSS measures may have little ability to assess the socioeconomic position of Hispanic/Latino adults who work in informal labor markets¹¹ or who have departed the workforce.¹² In any case, the existing divergences calls for a comprehensive approach to understanding cardiovascular health (CVH) in Hispanic/Latino people that surpasses conventional accounting by OSS.¹³

Emerging evidence supports an approach that focuses on how SSS influences CVH, given the connection between chronic stress and negative emotion and low SSS. A study based on small laboratory experiments with non-Hispanic White participants suggests that chronic stressors associated with low SSS may overactivate the hypothalamic-pituitary-adrenal axis and stimulate proinflammatory immune mediators, which seems to weaken cortisol responses.¹⁴ Indeed, people with low SSS produce more proinflammatory cytokines such as interleukin-6 than those with high SSS.¹⁵ Other laboratory-based studies have shown that increased stress-related allostatic load can, over time, suppress immune system functioning and lead to adverse cardiometabolic consequences.^{16–19}

Research on the association between SSS and CVH on Hispanic/Latino people is particularly thin. Not only are poor CVH profiles prevalent in this population,^{20,21} but immigration complicates attempts to measure their social status. For example, both Latino immigrants and migrants from Puerto Rico may lose income and social status when they begin residence in one of the 50 states, but they may also gain a kind of social mobility unattainable in their place of origin.¹² Migration creates multiple socioeconomic frames of reference that may influence behaviors and health outcomes that remain undetectable by OSS.²² One study that employed several measures of SSS and subjective well-being found variation in the coefficients, indicating that immigrants maintain simultaneous points of reference in both the United States and their country of origin for SSS.²² In another study, investigators observed that while prolonged stays in the United States have the undesirable effects (ie, perceived discrimination and diminished social networks), SSS increased with resident time, and this improvement was strongly associated with lower odds of depression and anxiety.⁵

The current study used data from the HCHS/SOL (Hispanic Community Health Study/Study of Latinos) to examine the relationship between SSS and CVH²³ in

adults from varying Hispanic/Latino backgrounds, independent of OSS. We expected SSS to be positively associated with CVH, such that adults with higher SSS will display more favorable CVH profiles than those with lower SSS. In addition, we tested whether nativity and, for immigrants, length of stay in the United States modifies associations between SSS and CVH. Given that immigrants might rely on multiple socioeconomic frames of reference, we expected to find a significant effect modification by nativity status such that foreign-born individuals would show a stronger association between SSS and CVH than US-born respondents, with the strongest association among those who migrated <10 years earlier.

METHODS

Study Population and Data Source

The current investigation uses data from the HCHS/SOL, a community-based prospective cohort study of 16 415 self-identified Hispanic/Latino adults aged 18 to 74 years, which has examined the prevalence of risk and protective factors of cardiovascular disease across Hispanic/Latino subgroups.^{24,25} Details of the recruitment and study protocol for HCHS/SOL have been published previously.^{24,25} Because of the sensitive nature of the data collected for this study, requests to access the data set from qualified researchers trained in human subject confidentiality protocols may be sent to HCHS/SOL at <https://sites.csc.unc.edu/hchs/>. Participants were enrolled from randomly selected households across 4 US urban areas (Chicago, IL; Miami, FL; Bronx, NY; and San Diego, CA) with the baseline examination held from 2008 to 2011.²⁴ Briefly, the cohort was selected through a stratified 2-stage area probability sample design that provided diversity with regard to SES and national origin or background.²⁴ In the first stage, households were sampled within geographic clusters (ie, census blocks), with oversampling of clusters most likely to be Hispanic/Latino according to the proportion of the population found to be Hispanic/Latino in the 2000 decennial census. In the second stage, households with a Hispanic/Latino surname were selected at a higher rate than other addresses. The HCHS/SOL cohort consists of first- through third-generation immigrant (or, in the case of Puerto Rico, migrant) Hispanic/Latino adults from diverse backgrounds (Mexican, Puerto Rican, Dominican, Cuban, and Central and South American). Potential participants were excluded if they planned to move within 6 months, were on active-duty military service, or were physically unable to attend the baseline examination. The baseline clinical examination included comprehensive biological, behavioral, and sociodemographic assessments.²² The institutional

review boards at the data coordinating center and at each field center approved this study, and all subjects gave written consent.

A total of 16 415 individuals participated in the baseline assessment. The analytic sample was limited to participants with complete data for the main variables of interest: (1) subjective social status, (2) metrics required to compute a cardiovascular health score (Life's Simple 7), and (3) preselected covariates. Although none were missing data on Life's Simple 7, data from 1041 (6.8%) individuals were excluded because of missing data on selected variables (154 on SSS, 5 on OSS, and 882 on additional variables). This resulted in a final analytic sample of 15 374 Hispanic/Latino adults. Participants excluded from the analysis did not differ from the included sample on SSS levels, language primarily spoken, sex, nativity/US length of stay, or discrimination experiences. However, those excluded tended to have lower OSS (less educated, lower income, less likely to be employed) and were older. They also had worse CVH composite scores but showed no significant differences in individual CVH indicators. The final sample with complete data is restricted to 15 374 individuals.

Outcome Variables

Cardiovascular Health: Life's Simple 7

AHA definitions²³ were used to operationalize CVH metrics: diet, physical activity, smoking status, body mass index (BMI), cholesterol, blood pressure, and fasting glucose.²⁰ Detailed information on the scoring of CVH has been reported previously.²⁰ Each indicator was coded categorically as Poor (0 points), Intermediate (1 point), or Ideal (2 points). A composite CVH score was calculated by summing across the 7 indicators (scores range from 0 to 14; higher scores indicate better CVH).²² Finally, a dichotomous Life's Simple 7 CVH cut point (≥ 4 Ideal indicators) was generated, which has been associated with cardioprotection and reduced 20-year incidence of coronary heart disease.^{26–29} These variables have been used in previous studies of HCHS/SOL.²⁰

Protocols to measure CVH indicators have been described elsewhere.²⁵ Briefly, 4 healthy lifestyle factors and 3 biomarkers were used. Former and current smoking status was self-reported. Two 24-hour dietary recalls were used to evaluate dietary intake across 5 food categories (ie, fruits/vegetables, fish, grains, sweetened beverages, and sodium). Physical activity was obtained through a modified version of the Global Physical Activity Questionnaire, which asks for self-reported activities during work, transport, and leisure.^{30,31} BMI, measured as kilograms per meter squared, was calculated from staff-ascertained measures of weight (nearest 0.1 kg) and height (nearest

centimeter). Biomarkers collected were total cholesterol, fasting blood glucose, and blood pressure. After a 12-hour fast, blood was drawn to obtain lipid profiles and fasting glucose values. Total cholesterol was measured using a cholesterol oxidase enzymatic method; fasting blood glucose was obtained by means of the hexokinase enzymatic method (Roche Diagnostics, Indianapolis, IN). Systolic and diastolic blood pressure were attained by averaging 3 readings taken with participants in a seated position using an automatic sphygmomanometer. In addition to the measurements of fasting glucose, cholesterol, and blood pressure, we considered self-reported medication use as reported in the medical history questionnaire to identify those with preexisting conditions.

Independent Variable

Subjective Social Status

The MacArthur Scale of Subjective Social Status³² captures self-perceived socioeconomic standing and pictorially depicts a 10-rung “social ladder” on which respondents are asked to subjectively rank their social standing using the general US population as a reference group (scores range from 1 to 10; higher scores indicate higher SSS). The participant is asked: “At the top of the ladder are the people who are the best off—those who have the most money, the most education, and the most respected jobs. At the bottom are the people who are the worst off—who have the least money, least education, and the least respected jobs or no jobs.” Participants verbally indicate the rung (from 1 to 10) that best depicts their current social standing (see Figure). In addition to treating SSS as a continuous measure, tertiles were created based on the distribution of scores to test for possible threshold effects (as clinically based cutoffs are unavailable). The MacArthur Scale has undergone previous psychometric testing with documented adequacy for validity and reliability in English^{32–34} and has been translated into Spanish.

Potential Confounders

Sociodemographic Covariates

Covariates included baseline age (in years), sex (male or female), Hispanic/Latino background, insurance status (yes/no), nativity (born in one of the 50 US states or born in a US territory or foreign country), length of stay in the United States, marital status (single, married/living with a partner, or separated/divorced/widowed), and study center. Nativity was combined with length of stay, and 3 groups were generated: born in the United States (including US territories), foreign-born immigrated <10 years ago, and foreign-born immigrated ≥10 years before. The 10-year cutoff aligns

with the definition of acculturation, in which lower acculturation was demarcated as residing in the United States for <10 years and higher acculturation as living in the United States for ≥10 years.^{35,36} Puerto Rico was the only US territory represented in the sample. It was treated as part of the United States because Puerto Ricans have automatic citizenship status when they enter the US mainland. We also included a question of perceived discrimination: “How often do people treat you unfairly because you are Hispanic or Latino?” (never or sometimes/always).

Objective Social Status

Three measures were used: (1) educational attainment (less than high school, high school graduate/general education degree, greater than high school); (2) annual income (<\$20 000, \$20 000 to \$50 000, >\$50 000, not reported); and (3) employment status (employed [full or part time] or unemployed).

Physical Health

We used the Physical Component Summary (PCS) of the Short-Form 12-Item Health Survey.³⁷ Because the PCS omits information specific to coronary disease, we included a question that asked participants to disclose the presence of prevalent coronary heart disease (yes or no).

Mental Health

The Center for Epidemiologic Studies Depression Scale was used to capture mental health. The 10-item Center for Epidemiologic Studies Depression Scale measures depressive symptoms with possible scores in the range of 0 to 30, and scores ≥10 are indicative of “significant” symptomatology.³⁸

Analytic Procedures

All analyses were weighted to account for selection probability and the complex sampling design, except for the correlation analyses. Analyses were performed using Statistical Analysis Software version 9.2 (SAS Institute, Cary, NC) and STATA SE 14.0 (StataCorp LP, College Station, TX). Descriptive statistics such as means, SDs, and percentages were used to summarize and compare characteristics for the overall sample by tertiles of SSS levels. Chi-squared tests were used to compare categorical variables, and *t* tests were used for continuous variables.

To assess the relationship between SSS and CVH, we first calculated prevalence estimates for counts of Ideal CVH indicators by tertiles of SSS and tested whether Ideal CVH scores were similarly distributed across SSS tertiles. Finally, multivariable regression models were used to assess the relationship of SSS



Figure. The MacArthur Scale of subjective social status.

with CVH, independent of OSS. Linear regression models were conducted for the overall CVH score and logistic regression models for the overall CVH score based on the ≥ 4 Ideal criterion and for the 2-level category for each of the CVH metric items.

The modeling procedures to test the relationship between SSS and CVH included 4 models. In model 1, SSS was adjusted for basic demographic variables—age, sex, Hispanic/Latino background, and study site. Model 2 adjusted for all covariates in model 1 and

added 3 OSS measures (ie, educational attainment, annual income, and employment status). Model 3 included all covariates in model 2 plus nativity/US length of stay, insurance status, and perceived discrimination. These variables address social factors that have a bearing on a person's sense of stability and receptiveness in the home context. Model 4 adjusted for all covariates included in model 3 and added variables that influence long-term health—prevalent coronary heart disease, mental health (Center for Epidemiologic Studies Depression Scale—continuous), and self-perceived health status (Short-Form 12-Item Health Survey PCS).

An interaction term for nativity/US length of stay and SSS was added to the models to examine whether the association between SSS and cardiovascular health differs by either of these factors. A significant regression coefficient for this interaction term at the 0.05 significance level would suggest that nativity/US length of stay modifies the relationship between SSS and CVH.

Although our sample allows for the inclusion of a number of Hispanic/Latino subgroups, the size of the Mexican population in this country and the unique political relationship of Puerto Rico with the United States, which confers automatic citizenship status to its migrants, creates the need for sensitivity tests to ensure that these groups are not driving the results. Sensitivity analyses were performed to assess whether the results were robust when considering subgroups of Hispanics/Latinos. In the first set of analyses, the sample was restricted to Mexicans only. Next, the analysis used the entire sample but excluded Puerto Ricans to test whether their inclusion as a US-born group affected the results.

RESULTS

Participant Characteristics

Baseline demographic, socioeconomic, and health characteristics are summarized in Table 1. Average SSS was 4.4 (SE, 0.0), with about 60% reporting having a high school education or lower. About 42% lived in households with annual income below \$20 000, and almost half (49%) were not employed. There were statistical differences in educational level, income, and employment by SSS levels ($P<0.001$). Across individual metrics, Ideal status was achieved at a rate of 1.6% for healthy eating, 23.3% for BMI, 49.8% for blood pressure, 53.5% for total cholesterol, 66.7% for fasting glucose, 66.8% for physical activity, and 76.3% for smoking. Less than half (46%) of the sample had ≥ 4 Ideal scores across all 7 metrics.

Prevalence data for Ideal CVH health and metrics by SSS are shown in Table 2. Prevalence of all CVH indicators differed across SSS levels ($P<0.05$),

save for prevalence of healthy diet. Prevalence data for Ideal CVH health and metrics by OSS indicators have been published elsewhere.²⁰ Briefly, that study found significant variations in CVH criterion prevalence by household income and education. Those who reported higher incomes were 15% more likely to meet favorable CVH criterion than those with lower income levels. Education followed a similar trend; those who reported college or more were 19% more likely to meet CVH criterion than high school noncompleters.

Association of SSS and CVH

Table 3 displays the parameter estimates and odds ratios for the association between SSS and CVH scores and metrics across 4 models. In model 1, each unit increase in SSS was positively associated with overall CVH score and Ideal levels of smoking, BMI, physical activity, and fasting blood glucose levels ($P<0.01$). Once adjusted for OSS (model 2), higher SSS remained positively associated with overall CVH score, BMI, and fasting glucose ($P<0.05$). These associations remained in model 3 after further adjustment for nativity/length of stay, marital status, insurance status, and perceived discrimination. SSS was associated with increases in overall CVH score ($\beta=0.04$; 95% CI, 0.01–0.06; $P<0.01$) and with higher odds of having an Ideal BMI (odds ratio [OR], 1.04; 95% CI, 1.01–1.07; $P<0.05$), Ideal physical activity (OR, 1.03; 95% CI, 1.0–1.07; $P<0.05$), and Ideal fasting glucose (OR, 1.04; 95% CI, 1.01–1.07; $P<0.05$). However, when analyses were further adjusted for mental and physical health indicators (model 4), the association between SSS and overall CVH became nonsignificant. In model 4, higher SSS remained positively associated with BMI (OR, 1.04; 95% CI, 1.00–1.07; $P<0.05$) and fasting glucose (OR, 1.03; 95% CI, 1.00–1.07; $P<0.05$).

Sensitivity analyses focused on those of Mexican descent and excluding Puerto Ricans yielded similar results to that of pooled analyses (results not shown). For Mexicans, each unit increase of SSS was associated with better overall CVH ($\beta=0.05$; 95% CI, 0.01–0.10; $P<0.05$). In addition, increases in SSS were associated with higher odds of having Ideal BMI (OR, 1.08; 95% CI, 1.02–1.13; $P<0.01$), Ideal physical activity (OR, 1.06; 95% CI, 1.01–1.12; $P<0.05$), and Ideal fasting glucose (OR, 1.09, 95% CI, 1.04–1.16, $P<0.01$). When we excluded Puerto Ricans from the analysis, we found an association between SSS and overall CVH ($\beta=0.04$; 95% CI, 0.01–0.06; $P<0.01$), Ideal BMI (OR, 1.05; 95% CI, 1.01–1.09; $P<0.05$) and Ideal fasting glucose (OR, 1.04; 95% CI, 1.01–1.08; $P<0.05$), much like those of the pooled analyses.

Table 1. Characteristics of the HCHS/SOL Target Population by SSS Category

Variables	All Mean or % (SE)	SSS			P Value
		Low (1–3) Mean or % (SE)	Moderate (4–7) Mean or % (SE)	High (8–10) Mean or % (SE)	
N	15 374	5104	9509	761	
SSS	4.38 (0.03)	2.31 (0.02)	5.02 (0.02)	8.58 (0.04)	<0.001
Education					
Less than HS	32.2 (0.7)	40.8 (1.1)	28.8 (0.8)	22.3 (2.1)	<0.001
HS	28.4 (0.6)	27.8 (1.0)	29.2 (0.7)	21.1 (2.1)	
More than HS	39.4 (0.8)	31.4 (1.0)	41.9 (1.0)	56.7 (2.8)	
Income, USD					
<20 000	41.8 (0.9)	57.9 (1.1)	35.4 (1.0)	24.9 (2.1)	<0.001
20 000–50 000	37.3 (0.7)	27.3 (1.0)	42.3 (0.8)	34.6 (2.7)	
>50 000	12.0 (0.8)	3.8 (0.5)	14.3 (0.9)	32.1 (3.2)	
Not reported	8.9 (0.4)	11.0 (0.7)	7.95 (0.4)	8.40 (1.5)	
Employment status					
Employed	51.1 (0.7)	42.8 (1.0)	54.5 (0.9)	59.7 (2.4)	
Age, y	41.0 (0.3)	43.7 (0.4)	39.7 (0.3)	41.1 (0.8)	
<50	70.2 (0.7)	63.7 (1.1)	73.3 (0.7)	70.0 (2.5)	<0.001
50–60	17.5 (0.5)	20.7 (0.8)	16.0 (0.6)	16.6 (1.7)	
>60	12.4 (0.5)	15.6 (0.8)	10.7 (0.5)	13.5 (1.9)	
Sex					
Male	48.0 (0.6)	47.6 (0.9)	48.3 (0.7)	47.2 (2.7)	
Female	52.0 (0.6)	52.5 (0.9)	51.7 (0.7)	52.8 (2.7)	
Hispanic background					
Mexican	37.5 (1.7)	32.7 (1.9)	40.3 (1.7)	40.8 (3.8)	<0.001
Cuban	20.0 (1.7)	23.5 (2.2)	18.7 (1.6)	15.4 (2.2)	
Puerto Rican	15.9 (0.8)	15.9 (1.0)	15.7 (0.8)	17.8 (2.0)	
Dominican	9.7 (0.7)	13.0 (1.1)	8.3 (0.7)	7.7 (1.5)	
Central American	7.4 (0.6)	8.1 (0.7)	7.3 (0.6)	5.5 (1.0)	
South American	4.9 (0.3)	3.7 (0.4)	5.3 (0.4)	7.0 (1.4)	
Other	4.1 (0.3)	3.2 (0.4)	4.4 (0.4)	5.9 (1.1)	
Nativity/length of stay					
US born	23.0 (0.8)	18.1 (1.1)	25.1 (0.9)	25.9 (2.1)	<0.001
Foreign born <10 y	27.6 (1.0)	30.7 (1.4)	26.6 (1.02)	21.0 (2.9)	
Foreign born ≥10 y	49.4 (0.8)	51.1 (1.2)	48.25 (0.9)	53.0 (2.9)	
Site					
Bronx	27.8 (1.5)	34.2 (2.0)	25.0 (1.5)	25.0 (2.7)	<0.001
Chicago	16.2 (1.0)	13.1 (1.0)	17.4 (1.1)	20.0 (2.4)	
Miami	29.3 (2.2)	31.6 (2.6)	28.5 (2.1)	25.9 (3.1)	<0.001
San Diego	26.6 (1.8)	21.1 (1.7)	29.1 (2.0)	29.2 (4.2)	
Marital status					
Single	34.4 (0.7)	33.7 (1.1)	34.62 (0.8)	35.9 (2.5)	<0.001
Married/living with a partner	49.2 (0.8)	46.2 (1.1)	50.58 (0.9)	49.7 (2.6)	
Separated/divorced/widowed	16.4 (0.5)	20.1 (0.8)	14.80 (0.6)	14.4 (1.7)	
Health insurance					
Yes	50.2 (0.9)	48.4 (1.3)	50.25 (1.1)	60.47 (2.7)	<0.001

(Continued)

Table 1. Continued

Variables	All Mean or % (SE)	SSS			P Value
		Low (1–3) Mean or % (SE)	Moderate (4–7) Mean or % (SE)	High (8–10) Mean or % (SE)	
Perceived discrimination					
Never	50.9 (0.8)	49.2 (1.2)	51.4 (0.9)	54.8 (2.4)	0.057
Some/often/always	49.1 (0.8)	50.8 (1.2)	48.6 (0.9)	45.2 (2.4)	
SF-12 Health Survey					
Physical (PCS)	50.0 (0.1)	48.4 (0.3)	50.7 (0.2)	51.2 (0.5)	<0.001
Prevalent CHD					
Yes	4.6 (0.3)	5.5 (0.4)	4.4 (0.4)	4.1 (0.9)	0.162
CES-D (continuous)	7.0 (0.1)	8.1 (0.2)	6.6 (0.09)	5.5 (0.2)	<0.001

CES-D indicates Center for Epidemiologic Studies Depression Scale; CHD, coronary heart disease; HCHS/SOL, Hispanic Community Health Study/Study of Latinos; HS, high school; PCS, Physical Component Summary; SF-12, Short-Form 12-Item Health Survey; SSS, subjective social status; and USD, US dollars.

Effect Modification by Nativity/Length of Stay

Results indicate that the positive association of SSS with CVH was only statistically significant for US-born and foreign-born people living in the United States for ≥ 10 years ($P < 0.05$; results not shown), but it was not significant for foreign-born people living in the United States for < 10 years. This pattern remained in models 2 to 4, but estimates were no longer statistically significant in model 4 for foreign-born living in the United States for ≥ 10 years. Next, models examined whether nativity/length of stay modified the association between SSS and CVH measures. Results indicate that nativity/length of stay modified the association between SSS and Ideal smoking ($P < 0.05$; results not shown). The interaction terms for SSS and foreign-born people with < 10 years were negative, indicating that shorter length of stay reduces the odds of having Ideal smoking. The interaction terms for SSS and foreign-born people with ≥ 10 years' residence were not statistically significant.

DISCUSSION

This study contributes to the growing evidence that subjective measures of social status can enhance an understanding of CVH by examining data from Hispanic/Latino people. Our results indicate that higher SSS is associated with higher odds of having Ideal CVH indicators, which is consistent with a meta-analysis that found lower SSS to be associated with increased odds of cardiovascular disease among other ethnic groups (not including Hispanic/Latino people).⁶ Our findings also resonate with results of Perreira et al⁵ that showed SSS improved with length of stay in the United States and that higher SSS was associated with lower odds of mental health problems.

Prior studies that examined SSS with Hispanic/Latino immigrants and their US-born counterparts have relied on self-reported measures of physical health. In addition, most studies that examined SSS with Hispanic/Latino immigrants and their US-born counterparts have focused on a singular Hispanic/Latino heritage group (usually Mexican) in one geographic location^{39,40} (studies that use data from the National Latino and Asian American Study^{12,41} are the exception). By using HCHS/SOL data, we were able to examine a diverse and large sample of Hispanic/Latino people and to discern the distribution of SSS and its associations with empirical measures of cardiovascular health. The data offered diversity of heritage backgrounds, geographic residency, and SES that reflects the overall demographics one would expect in a national sample⁴² (with the exception of age).⁴³

In addition, our use of the Life's Simple 7, which includes objective and subjective data, enabled us to examine a more complete profile of overall CVH and its constituent parts. For example, we examined the associations for Mexicans only and for the entire sample excluding Puerto Ricans (analyses not shown). We found that SSS was consistently associated with Ideal BMI and Ideal fasting glucose. When we excluded Puerto Ricans from the analysis, to test whether including this group as US born would alter the results, we found an association between SSS and overall CVH, similar to when focusing exclusively on Mexicans. Analyses restricted to Mexicans only also found a positive association between SSS with Ideal physical activity and overall CVH. For Mexicans, who are at a higher risk of metabolic conditions, the fact that higher SSS has such an effect on CVH and physical activity highlights the need for targeted policy interventions. We speculate that given the lower levels of education and income as well the higher

Table 2. Prevalence Estimates of Ideal CVH by SSS and OSS Indicators (N=15 374)

	≥4 Ideals % (SE)	Smoking % (SE)	BMI % (SE)	Physical Activity % (SE)	Healthy Diet % (SE)	Total Cholesterol % (SE)	Blood Pressure % (SE)	Fasting Glucose % (SE)
Total	45.97 (0.67)	76.3 (0.6)	23.3 (0.5)	66.8 (0.7)	1.6 (0.2)	53.5 (0.6)	49.8 (0.7)	66.7 (0.7)
SSS tertiles								
Low (1–3)	39.73 (1.09)	73.5 (0.9)	20.1 (0.8)	62.6 (1.1)	1.9 (0.3)	51.5 (1.0)	44.7 (1.2)	61.2 (1.1)
Moderate (4–7)	48.92 (0.77)	77.5 (0.7)	24.6 (0.7)	68.5 (0.7)	1.4 (0.2)	54.8 (0.8)	52.4 (0.9)	69.2 (0.7)
High (8–10)	46.92 (2.33)	79.6 (2.0)	25.1 (2.3)	70.8 (2.4)	2.4 (0.9)	49.8 (2.4)	47.4 (2.5)	69.2 (2.4)
P value*	<0.001	<0.001	<0.001	<0.001	0.16	0.01	<0.001	<0.001
Education								
Less than HS	37.76 (1.05)	73.58 (0.99)	19.69 (0.86)	62.17 (1.04)	1.61 (0.22)	49.78 (1.04)	43.30 (1.19)	58.44 (1.06)
HS	50.44 (1.19)	74.78 (1.07)	24.09 (1.00)	69.61 (1.11)	1.45 (0.28)	59.60 (1.15)	53.68 (1.16)	70.16 (1.13)
More than HS	49.48 (1.05)	79.71 (0.81)	25.57 (0.89)	68.59 (0.99)	1.71 (0.26)	52.23 (1.07)	52.22 (1.01)	71.00 (0.98)
P value*	<0.001	<0.001	<0.001	<0.001	0.775	<0.001	<0.001	<0.001
Income, USD								
<20 000	42.23 (0.98)	73.37 (0.89)	21.36 (0.74)	63.32 (0.89)	1.58 (0.23)	51.94 (0.94)	46.48 (1.01)	64.73 (0.87)
20 000–50 000	48.26 (0.99)	77.59 (0.86)	23.49 (0.86)	70.66 (0.94)	1.69 (0.20)	55.43 (1.02)	52.62 (1.06)	66.91 (0.97)
>50 000	50.23 (1.92)	83.36 (1.35)	23.25 (1.51)	72.42 (1.67)	2.19 (0.62)	49.58 (2.07)	51.12 (1.94)	70.62 (1.98)
P value*	<0.001	<0.001	0.129	<0.001	0.512	0.01	<0.001	0.009
Employment status								
Employed full- or part-time	50.84 (0.85)	78.60 (0.78)	23.55 (0.72)	74.18 (0.75)	1.37 (0.16)	54.94 (0.86)	52.99 (0.92)	70.07 (0.79)
Not employed	40.89 (0.95)	73.97 (0.84)	22.95 (0.78)	59.13 (0.90)	1.85 (0.25)	52.07 (0.86)	46.38 (1.07)	63.20 (1.01)
P value*	<0.001	<0.001	0.544	<0.001	0.084	0.014	<0.001	<0.001

BMI indicates body mass index; CVH, cardiovascular health; HS, high school; OSS, objective markers of social status; SSS, subjective social status; and USD, US dollars.

*Chi-square test was used.

Table 3. Parameter Estimates and 95% CIs for the Association Between 1-Unit Increase in SSS and 1-Unit Increase in OSS (Education, Income, and Employment Status) and CVH Scores and Metrics (N=15 374)

	CVH	≥4 Ideals	Smoking	BMI	Physical Activity	Healthy Diet	Total Cholesterol	Blood Pressure	Fasting Glucose
	Beta (95% CI)	OR (95% CI)	OR (95% CI)	OR (95% CI)	OR (95% CI)	OR (95% CI)	OR (95% CI)	OR (95% CI)	OR (95% CI)
Model 1									
SSS	0.07 (0.04 to 0.10)	1.03 (1.00 to 1.06)	1.07 (1.04 to 1.11)	1.04 (1.01 to 1.08)	1.05 (1.02 to 1.08)	0.97 (0.87 to 1.09)	0.98 (0.95 to 1.00)	1.00 (0.97 to 1.04)	1.06 (1.02 to 1.09)
P value	<0.001	0.050	<0.001	0.009	0.002	0.657	0.089	0.828	<0.001
Model 2									
SSS	0.04 (0.01 to 0.07)	1.01 (0.98 to 1.04)	1.03 (0.99 to 1.06)	1.04 (1.01 to 1.08)	1.03 (1.00 to 1.06)	0.95 (0.85 to 1.06)	0.99 (0.96 to 1.02)	0.99 (0.95 to 1.02)	1.04 (1.01 to 1.07)
P value	0.003	0.588	0.130	0.009	0.074	0.367	0.513	0.464	0.023
Education (ref: less than HS)									
HS	0.22 (0.11 to 0.33)	1.17 (1.02 to 1.34)	1.20 (1.03 to 1.40)	1.00 (0.86 to 1.17)	1.18 (1.04 to 1.35)	1.24 (0.75 to 2.04)	1.03 (0.90 to 1.18)	1.11 (0.97 to 1.26)	1.17 (1.02 to 1.34)
P value	<0.001	0.027	0.019	0.977	0.010	0.404	0.647	0.125	0.023
More than HS	0.39 (0.27 to 0.50)	1.34 (1.17 to 1.53)	1.41 (1.21 to 1.65)	1.18 (1.01 to 1.38)	1.22 (1.07 to 1.39)	1.28 (0.90 to 1.83)	0.92 (0.80 to 1.07)	1.26 (1.11 to 1.43)	1.38 (1.20 to 1.60)
P value	<0.001	<0.001	<0.001	0.034	0.004	0.173	0.278	<0.001	<0.001
Income, USD (Ref: <20 000)									
20 000–50 000	0.13 (0.03 to 0.23)	1.04 (0.92 to 1.17)	1.22 (1.06 to 1.39)	1.05 (0.93 to 1.19)	1.11 (0.99 to 1.24)	1.05 (0.71 to 1.55)	1.05 (0.93 to 1.19)	1.12 (0.98 to 1.28)	0.95 (0.84 to 1.08)
P value	0.014	0.582	0.004	0.444	0.083	0.814	0.392	0.097	0.461
>50 000	0.26 (0.09 to 0.44)	1.12 (0.91 to 1.38)	1.77 (1.41 to 2.22)	1.00 (0.82 to 1.22)	1.02 (0.83 to 1.25)	1.34 (0.70 to 2.55)	0.81 (0.67 to 0.99)	1.06 (0.85 to 1.34)	1.15 (0.92 to 1.44)
P value	0.003	0.269	<0.001	0.997	0.865	0.371	0.042	0.597	0.232
Not reported	0.22 (0.08 to 0.37)	1.23 (1.01 to 1.51)	1.20 (0.98 to 1.45)	1.54 (1.29 to 1.84)	0.95 (0.80 to 1.13)	0.62 (0.22 to 1.73)	1.13 (0.94 to 1.37)	1.09 (0.90 to 1.33)	1.21 (1.01 to 1.46)
P value	0.003	0.044	0.072	<0.001	0.591	0.363	0.189	0.389	0.044
Employment status (Ref: employed)									
Not employed	−0.30 (−0.39 to −0.20)	0.77 (0.68 to 0.87)	0.81 (0.71 to 0.93)	1.03 (0.92 to 1.16)	0.64 (0.58 to 0.71)	1.39 (0.93 to 2.09)	1.07 (0.96 to 1.19)	0.90 (0.79 to 1.03)	0.84 (0.74 to 0.95)
P value	<0.001	<0.001	0.002	0.601	<0.001	0.111	0.213	0.134	0.006
Model 3									
SSS	0.04 (0.01 to 0.06)	1.01 (0.98 to 1.04)	1.02 (0.99 to 1.06)	1.04 (1.01 to 1.07)	1.03 (1.00 to 1.07)	0.95 (0.85 to 1.06)	0.99 (0.96 to 1.02)	0.99 (0.95 to 1.02)	1.04 (1.01 to 1.07)
P value	0.006	0.658	0.267	0.016	0.041	0.318	0.455	0.485	0.021

(Continued)

Table 3. Continued

	CVH	≥4 Ideals	Smoking	BMI	Physical Activity	Healthy Diet	Total Cholesterol	Blood Pressure	Fasting Glucose
	Beta (95% CI)	OR (95% CI)	OR (95% CI)	OR (95% CI)	OR (95% CI)	OR (95% CI)	OR (95% CI)	OR (95% CI)	OR (95% CI)
Education (Ref: <high school)									
HS	0.23 (0.12 to 0.33)	1.01 (0.98 to 1.04)	1.23 (1.05 to 1.43)	1.00 (0.86 to 1.17)	1.18 (1.03 to 1.34)	1.28 (0.77 to 2.12)	1.03 (0.90 to 1.18)	1.11 (0.97 to 1.26)	1.17 (1.02 to 1.34)
<i>P</i> value	<0.001	0.616	0.009	0.964	0.015	0.336	0.689	0.117	0.023
More than HS	0.40 (0.29 to 0.52)	1.17 (1.02 to 1.35)	1.47 (1.25 to 1.72)	1.18 (1.01 to 1.38)	1.20 (1.05 to 1.37)	1.34 (0.94 to 1.92)	0.91 (0.79 to 1.05)	1.27 (1.12 to 1.45)	1.39 (1.20 to 1.60)
<i>P</i> value	<0.001	0.024	<0.001	0.042	0.009	0.108	0.186	<0.001	<0.001
Income, USD (Ref: <20 000)									
20 000–50 000	0.13 (0.03 to 0.23)	1.04 (0.92 to 1.17)	1.21 (1.05 to 1.38)	1.10 (0.97 to 1.25)	1.14 (1.01 to 1.28)	1.07 (0.73 to 1.57)	1.06 (0.93 to 1.19)	1.09 (0.95 to 1.25)	0.96 (0.85 to 1.10)
<i>P</i> value	0.009	0.558	0.007	0.142	0.034	0.737	0.377	0.212	0.594
>50 000	0.27 (0.11 to 0.44)	1.12 (0.91 to 1.37)	1.73 (1.36 to 2.21)	1.04 (0.85 to 1.27)	1.07 (0.87 to 1.31)	1.41 (0.72 to 2.76)	0.78 (0.64 to 0.96)	1.04 (0.82 to 1.31)	1.18 (0.94 to 1.47)
<i>P</i> value	0.001	0.297	<0.001	0.713	0.544	0.309	0.017	0.751	0.153
Not reported	0.23 (0.08 to 0.38)	1.22 (1.00 to 1.50)	1.19 (0.98 to 1.45)	1.49 (1.24 to 1.79)	0.97 (0.82 to 1.16)	0.82 (0.22 to 1.73)	1.12 (0.93 to 1.36)	1.11 (0.91 to 1.36)	1.21 (1.01 to 1.46)
<i>P</i> value	0.003	0.053	0.073	<0.001	0.743	0.360	0.240	0.294	0.040
Employment status (Ref: employed)									
Not employed	−0.28 (−0.38 to −0.19)	0.77 (0.69 to 0.87)	0.83 (0.73 to 0.95)	1.02 (0.91 to 1.15)	0.64 (0.58 to 0.71)	1.44 (0.95 to 2.19)	1.05 (0.94 to 1.17)	0.92 (0.80 to 1.05)	0.84 (0.74 to 0.96)
<i>P</i> value	<0.001	<0.001	0.005	0.715	<0.001	0.084	0.368	0.191	0.008
Model 4									
SSS	0.02 (−0.00 to 0.05)	1.00 (0.97 to 1.03)	1.01 (0.97 to 1.04)	1.04 (1.00 to 1.07)	1.03 (0.99 to 1.06)	0.94 (0.84 to 1.05)	0.99 (0.96 to 1.02)	0.98 (0.95 to 1.02)	1.03 (1.00 to 1.07)
<i>P</i> value	0.078	0.963	0.619	0.032	0.100	0.250	0.400	0.379	0.037
Education (Ref: <high school)									
HS	0.18 (0.07 to 0.29)	1.00 (0.96 to 1.03)	1.19 (1.02 to 1.40)	0.98 (0.85 to 1.15)	1.15 (1.01 to 1.32)	1.25 (0.76 to 2.07)	1.02 (0.89 to 1.17)	1.09 (0.96 to 1.24)	1.15 (1.00 to 1.32)
<i>P</i> value	<0.001	0.782	0.025	0.845	0.033	0.375	0.752	0.181	0.045
More than HS	0.33 (0.21 to 0.44)	1.14 (0.99 to 1.31)	1.40 (1.18 to 1.65)	1.14 (0.97 to 1.33)	1.16 (1.01 to 1.33)	1.29 (0.90 to 1.84)	0.90 (0.78 to 1.04)	1.24 (1.09 to 1.41)	1.35 (1.17 to 1.56)
<i>P</i> value	<0.001	0.069	<0.001	0.112	0.031	0.163	0.144	<0.001	<0.001
Income, USD (Ref: <20 000)									
20 000–50 000	0.05 (−0.05 to 0.15)	0.98 (0.86 to 1.11)	1.14 (0.99 to 1.30)	1.06 (0.94 to 1.21)	1.09 (0.97 to 1.23)	1.02 (0.69 to 1.49)	1.04 (0.92 to 1.18)	1.07 (0.93 to 1.23)	0.93 (0.81 to 1.06)

(Continued)

Table 3. Continued

	CVH	≥4 Ideals	Smoking	BMI	Physical Activity	Healthy Diet	Total Cholesterol	Blood Pressure	Fasting Glucose
	Beta (95% CI)	OR (95% CI)	OR (95% CI)	OR (95% CI)	OR (95% CI)	OR (95% CI)	OR (95% CI)	OR (95% CI)	OR (95% CI)
P value	0.310	0.722	0.072	0.352	0.144	0.937	0.507	0.354	0.278
>50 000	0.13 (−0.03 to 0.29)	1.00 (0.82 to 1.24)	1.61 (1.26 to 2.06)	0.97 (0.79 to 1.19)	0.99 (0.80 to 1.22)	1.28 (0.65 to 2.53)	0.77 (0.62 to 0.94)	0.99 (0.78 to 1.26)	1.10 (0.89 to 1.37)
P value	0.119	0.963	<0.001	0.799	0.918	0.472	0.011	0.957	0.387
Not reported	0.16 (0.01 to 0.31)	1.16 (0.94 to 1.43)	1.14 (0.93 to 1.38)	1.44 (1.20 to 1.74)	0.94 (0.79 to 1.12)	0.60 (0.21 to 1.68)	1.11 (0.91 to 1.34)	1.10 (0.90 to 1.34)	1.18 (0.98 to 1.42)
P value	0.032	0.156	0.204	<0.001	0.472	0.328	0.299	0.351	0.088
Employment status (Ref: employed)									
Not employed	−0.18 (−0.28 to −0.09)	0.83 (0.73 to 0.93)	0.88 (0.77 to 1.01)	1.07 (0.95 to 1.20)	0.67 (0.61 to 0.75)	1.52 (1.01 to 2.31)	1.07 (0.95 to 1.19)	0.94 (0.82 to 1.07)	0.89 (0.78 to 1.01)
P value	<0.001	0.002	0.067	0.297	<0.001	0.046	0.253	0.355	0.068

Model 1: adjusted for age, sex, Hispanic/Latino background, and study site; model 2: model 1+ OSS (education, income, and employment); model 3: model 2+nativity, marital status, insurance status, and perceived discrimination; model 4: model 3+prevalent coronary heart disease, Center for Epidemiologic Studies Depression Scale (continuous), physical health. BMI indicates body mass index; CVH, cardiovascular health; HS, high school; OR, odds ratio; OSS, objective markers of social status; SSS, subjective social status; and USD, US dollars.

rates of being undocumented among Mexicans compared with those from other Latino/Hispanic backgrounds, SSS may be capturing features of social context associated with CVH better than traditional measures of SES.

Even so, we found only partial evidence for our second hypothesis that nativity/US length of stay modifies the association between SSS and CVH. For the foreign born, longer US stays and higher SSS showed indirect benefits to CVH. Nativity/US length of stay moderated the relationship between SSS and CVH, particularly for Ideal smoking and Ideal blood pressure. A shorter length of stay reduces the magnitude of the positive association between SSS and CVH.

Smoking is a highly changeable behavior; people can readily improve their CVH profiles by quitting. However, whether someone smokes merits some discussion, as it is arguably best understood by considering *both* OSS and SSS measures. Kaplan et al⁴⁴ found that Hispanic/Latino people with low SES were more likely to smoke, less likely to have quit, and less disposed to have used nonprescription quit products compared with those with higher income and education levels. However, they also found that smoking was more common for those born in the United States and who held a higher level of acculturation to the dominant US culture, particularly among women. Merzel et al⁴⁵ examined smoking cessation among Hispanic/Latino people and found few differences in socioeconomic characteristics by sex. Instead, they found that younger and more acculturated women had lower odds of sustaining cessation. These studies suggest that the influence of SSS on health behaviors may be complex.

This study has several limitations worth considering. Given the cross-sectional design, we cannot infer causal relationships between SSS and CVH. The association was evident in an older sample of Hispanics/Latinos (median age, 41) and reflects a snapshot in time, not a longitudinal assessment. This study also did not evaluate differences in associations between SSS and CVH by Hispanic/Latino background. Such evaluations may prove telling, as our analysis of those of Mexican background illustrated a strong association between SSS and CVH. We did exclude participants who were missing data (n=1041). However, our analysis of the exclusions (<10% the sample) showed few differences. We also suspect that in model 4, we may be inducing the estimation of the total effect toward the null by overadjusting for previous health conditions.⁴⁶ A test of this (not shown) found that when adjusting for mental health (Center for Epidemiologic Studies Depression Scale) or physical health (PCS) separately, the results remained very similar. PCS showed a slightly larger attenuation in the coefficients, but given that the coefficients are

somewhat small, further studies are needed to examine this issue.

Another issue that arose relates to the evaluation of clinical relevance when only small changes in CVH are observed as SSS increases. Model 3 (Table 3) shows a 1-unit increase in SSS is associated with a 0.04-unit improvement in CVH. Given that our CVH metric ranges from 0 to 14, a substantial 5-rung jump in SSS would increase CVH by a mere fifth of a point. Such a modest change in CVH falls well below the detection threshold for CVD mortality risk²⁷ and raises the question of clinical relevance. However, from the perspective of public health, which reckons with cumulative effects over the life span, a different construal can be made. Although the effect of SSS may be considered “very small,” such relatively small effects can have meaningful implications if the effects compound or accumulate over time to magnify the effect.^{47–49} It is reasonable to think that SSS could have such compounding effects because one’s perceived social status could influence changes in the social and physical environment in ways that affect health.⁴⁸ Moreover, just like disease outcomes, CVH is likely determined by multiple factors, so any one variable may show limited associations. Thus, at the population level, small differences can have a compounding effect over time.^{49,50} The accumulated effects of not smoking, regular exercise, and a heart-healthy diet on CVH occur incrementally over time, and in this regard, SSS might play a protective role. For example, as we discussed earlier, higher SSS scores coincided with less smoking among foreign-born people living in the United States, a health promoting behavior that reaps dividends over time.

In contrast to measures used to evaluate CVH, procedures to appraise SSS were limited in scope. While there is overwhelming evidence for associations between SES indicators and various maladies and hard clinical end points, the same is not true for SSS. In addition, the MacArthur Scale assessed only the participants’ self-perceived social status relative to the general United States, not their community. Some studies have examined associations of proximal (one’s community) and distal (one’s country) SSS with CVH^{51,52} and found that the distinction makes a difference. People’s distal status may not coincide with their proximal status, and it remains unknown whether status in close social networks (ie, a leadership role in church) might buffer the effects of a relatively lower status when considering a national reference group. Small clinical studies point in this direction. One study found that low SSS relative to the community was associated with impaired vasodilation of the brachial artery in young to middle-aged adults.⁵¹ In another study, people who ranked themselves with low SSS with regard to their country and community showed reduced activity in beta-adrenergic receptors,⁵² which mediate vasodilation, heart rate

increase, and immune functions.^{52,53} Over time, such impairment has been implicated in the pathophysiology of cardiovascular disease.⁵⁴

Future studies should explore the stability of SSS over time in comparison with OSS. We know that trajectories of exposure, in which OSS features prominently, are useful predictors of health-related outcomes of interest.^{48,55} For instance, one study examined the effects of both lifetime socioeconomic trajectory and cumulative disadvantage from childhood on the cognitive performance of Mexican American older adults found that compared with those with continuous low SES throughout the life course, those with more advantaged lifetime SES trajectories experienced fewer cognitive declines.⁴⁸ Chronic exposure to low SSS might result in larger effect sizes when it comes to CVH, but we will not know until these types of studies are performed; trajectory studies simply are not available for measures of SSS. In addition, more studies are needed to explore the mechanisms through which SSS may influence CVH, and it would be useful to establish clinical end points.

CONCLUSIONS

CVH arises in a milieu that is partially biological and behavioral. Among possible factors that can influence CVH, social status remains constant. The role that OSS plays is well established and increasing evidence suggests SSS provides meaningful additional information, especially at the population level, in which small differences can have a cumulative effect over time. Yet investigators continue to rely solely on OSS measures. Despite being a relatively simple measure, SSS remains underused in health studies and when it is used, its application is limited to one point in time. The inclusion of it and other complementary measures could provide a more robust picture, including its objective and subjective features, for how social status influences health *over time*. Such an approach would also help reveal how inequitable social arrangements affect long-term health and facilitate actions to reduce disparities in CVH. Moreover, attention to SSS could yield innovative social interventions that help elevate one’s sense of social status. At the population level, a small positive difference in SSS could have large returns on health outcomes even if OSS remains modest over the life course. Such an approach would also help reveal how inequitable social arrangements affect long-term health and facilitate actions to reduce disparities in CVH.

ARTICLE INFORMATION

Received March 20, 2019; accepted May 12, 2021.

Affiliations

School of Social Work, University of Illinois at Urbana-Champaign, Urbana, IL (L.M.P., F.C.A., R.H., K.M.T.); School of Medicine, University of North Carolina at Chapel Hill, NC (K.M.P.); Department of Psychology, San Diego State University, Chula Vista, CA (L.C.G., S.F.C., G.A.T.); Department of Neurosciences, University of California, San Diego, La Jolla, CA (H.M.G.); Department of Epidemiology and Population Health, Albert Einstein College of Medicine, Bronx, NY (S.G.); Department of Biostatistics, University of North Carolina at Chapel Hill, NC (J.C.); Department of Medicine, University of Illinois at Chicago, IL (J.C., R.A.D., M.L.D.); and Feinberg School of Medicine, Northwestern University, Chicago, IL (M.L.D.).

Acknowledgments

The authors thank the staff and participants of HCHS/SOL for their important contributions. A complete list of staff and investigators is available on the study Web site: <http://www.csc.unc.edu/hchs/>.

Sources of Funding

The Hispanic Community Health Study/Study of Latinos was carried out as a collaborative study supported by contracts from the National Heart, Lung, and Blood Institute (NHLBI) to the University of North Carolina (N01-HC65233), University of Miami (N01-HC65234), Albert Einstein College of Medicine (N01-HC65235), Northwestern University (N01-HC65236), and San Diego State University (N01-HC65237). The following institutes/centers/offices contribute to the HCHS/SOL through a transfer of funds to the NHLBI: National Center on Minority Health and Health Disparities, the National Institute of Deafness and Other Communications Disorders, the National Institute of Dental and Craniofacial Research, the National Institute of Diabetes and Digestive and Kidney Diseases, the National Institute of Neurological Disorders and Stroke, and the Office of Dietary Supplements. Its contents are solely the responsibility of the authors and do not necessarily represent the official views of the NIH.

Disclosures

None.

REFERENCES

- Davis JA. Status symbols and the measurement of status perception. *Sociometry*. 1956;19:154–165. DOI: 10.2307/2785629.
- Marmot M. Social determinants of health inequalities. *Lancet*. 2005;365:1099–1104. DOI: 10.1016/S0140-6736(05)71146-6.
- Wilkinson RG. Health, hierarchy, and social anxiety. *Ann N Y Acad Sci*. 1999;896:48–63. DOI: 10.1111/j.1749-6632.1999.tb08104.x.
- Demakakos P, Nazroo J, Breeze E, Marmot M. Socioeconomic status and health: the role of subjective social status. *Soc Sci Med*. 2008;67:330–340. DOI: 10.1016/j.socscimed.2008.03.038.
- Perreira KM, Gotman N, Isasi CR, Arguelles W, Castañeda SF, Daviglus ML, Giachello AL, Gonzalez P, Penedo FJ, Salgado H, et al. Mental health and exposure to the United States: key correlates from the Hispanic Community Health Study of Latinos. *J Nerv Ment Dis*. 2015;203:670. DOI: 10.1097/NMD.0000000000000350.
- Tang KL, Rashid R, Godley J, Ghali WA. Association between subjective social status and cardiovascular disease and cardiovascular risk factors: a systematic review and meta-analysis. *BMJ Open*. 2016;6:e010137. DOI: 10.1136/bmjopen-2015-010137.
- Beltrán-Sánchez H, Palloni A, Riosmena F, Wong R. SES gradients among Mexicans in the United States and in Mexico: a new twist to the Hispanic paradox? *Demography*. 2016;53:1555–1581. DOI: 10.1007/s13524-016-0508-4.
- Markides KS, Eschbach K. Hispanic paradox in adult mortality in the United States. In: Rogers R, Crimmins E, eds. *International Handbook of Adult Mortality*. Dordrecht: Springer; 2011:227–240.
- Turra CM, Goldman N. Socioeconomic differences in mortality among US adults: insights into the Hispanic paradox. *J Gerontol B Psychol Sci Soc Sci*. 2007;62:S184–S192. DOI: 10.1093/geronb/62.3.S184.
- Goldman N, Kimbro RT, Turra CM, Pebley AR. Socioeconomic gradients in health for White and Mexican-origin populations. *Am J Public Health*. 2006;96:2186–2193. DOI: 10.2105/AJPH.2005.062752.
- Venkatess SA. *Off the Books: The Underground Economy of the Urban Poor*. Cambridge, MA: Harvard University Press; 2006.
- Alcántara C, Chen C-N, Alegría M. Do post-migration perceptions of social mobility matter for Latino immigrant health? *Soc Sci Med*. 2014;101:94–106. DOI: 10.1016/j.socscimed.2013.11.024.
- Beltrán-Sánchez H, Palloni A, Riosmena F, Wong R. SES gradients among Mexicans in the United States and in Mexico: a new twist to the Hispanic paradox? *Demography*. 2016;53:1555–1581. DOI: 10.1007/s13524-016-0508-4.
- Gruenewald TL, Kemeny ME, Aziz N. Subjective social status moderates cortisol responses to social threat. *Brain Behav Immun*. 2006;20:410–419. DOI: 10.1016/j.bbi.2005.11.005.
- Derry HM, Fagundes CP, Andridge R, Glaser R, Malarkey WB, Kiecolt-Glaser JK. Lower subjective social status exaggerates interleukin-6 responses to a laboratory stressor. *Psychoneuroendocrinology*. 2013;38:2676–2685. DOI: 10.1016/j.psyneuen.2013.06.026.
- Seeman T, Epel E, Gruenewald T, Karlamangla A, McEwen BS. Socio-economic differentials in peripheral biology: cumulative allostatic load. *Ann N Y Acad Sci*. 2010;1186:223–239. DOI: 10.1111/j.1749-6632.2009.05341.x.
- Adler NE, Tan JX. Commentary: tackling the health gap: the role of psychosocial processes. *Int J Epidemiol*. 2017;46:1329–1331. DOI: 10.1093/ije/dyx167.
- Dickerson SS, Gable SL, Irwin MR, Aziz N, Kemeny ME. Social-evaluative threat and proinflammatory cytokine regulation: an experimental laboratory investigation. *Psychol Sci*. 2009;20:1237–1244. DOI: 10.1111/j.1467-9280.2009.02437.x.
- Dickerson SS, Kemeny ME. Acute stressors and cortisol responses: a theoretical integration and synthesis of laboratory research. *Psychol Bull*. 2004;130:355–391. DOI: 10.1037/0033-2909.130.3.355.
- González HM, Tarraf W, Rodríguez CJ, Gallo LC, Sacco RL, Talavera GA, Heiss G, Kizer JR, Hernandez R, Davis S, et al. Cardiovascular health among diverse Hispanics/Latinos: Hispanic Community Health Study/Study of Latinos (HCHS/SOL) results. *Am Heart J*. 2016;176:134–144. DOI: 10.1016/j.ahj.2016.02.008.
- Daviglus ML, Pirzada A, Talavera GA. Cardiovascular disease risk factors in the Hispanic/Latino population: lessons from the Hispanic Community Health Study/Study of Latinos (HCHS/SOL). *Prog Cardiovasc Dis*. 2014;57:230–236. DOI: 10.1016/j.pcad.2014.07.006.
- Gelatt J. Looking down or looking up: status and subjective well-being among Asian and Latino immigrants in the United States. *Int Migr Rev*. 2013;47:39–75. DOI: 10.1111/imre.12013.
- Lloyd-Jones DM, Hong Y, Labarthe D, Mozaffarian D, Appel LJ, Van Horn L, Greenland K, Daniels S, Nichol G, Tomaselli GF, et al. Defining and setting national goals for cardiovascular health promotion and disease reduction the American Heart Association's Strategic Impact Goal through 2020 and beyond. *Circulation*. 2010;121:586–613. DOI: 10.1161/CIRCULATIONAHA.109.192703.
- LaVange LM, Kalsbeek WD, Sorlie PD, Avilés-Santa LM, Kaplan RC, Barnhart J, Liu K, Giachello A, Lee DJ, Ryan J, et al. Sample design and cohort selection in the Hispanic Community Health Study/Study of Latinos. *Ann Epidemiol*. 2010;20:642–649. DOI: 10.1016/j.annepidem.2010.05.006.
- Sorlie PD, Avilés-Santa LM, Wassertheil-Smolser S, Kaplan RC, Daviglus ML, Giachello AL, Schneiderman N, Raj L, Talavera G, Allison M, et al. Design and implementation of the Hispanic Community Health Study/Study of Latinos. *Ann Epidemiol*. 2010;20:629–641. DOI: 10.1016/j.annepidem.2010.03.015.
- Folsom AR, Yatsuya H, Nettleton JA, Lutsey PL, Cushman M, Rosamund WD, ARIC Study Investigators. Community prevalence of ideal cardiovascular health, by the American Heart Association definition, and relationship with cardiovascular disease incidence. *J Am Coll Cardiol*. 2011;57:1690–1696. DOI: 10.1016/j.jacc.2010.11.041.
- Ford ES, Greenland KJ, Hong Y. Ideal cardiovascular health and mortality from all causes and diseases of the circulatory system among adults in the United States. *Circulation*. 2012;125:987–995. DOI: 10.1161/CIRCULATIONAHA.111.049122.
- Laitinen TT, Pakkala K, Magnussen CG, Viikari JSA, Oikonen M, Taittonen L, Mikkilä V, Jokinen E, Hutri-Kähönen N, Laitinen T, et al. Ideal cardiovascular health in childhood and cardiometabolic outcomes in adulthood: the Cardiovascular Risk in Young Finns Study. *Circulation*. 2012;125:1971–1978. DOI: 10.1161/CIRCULATIONAHA.111.073585.
- Rasmussen-Torvik LJ, Shay CM, Abramson JG, Friedrich CA, Nettleton JA, Prizment AE, Folsom AR. Ideal cardiovascular health is inversely associated with incident cancer: the Atherosclerosis Risk In Communities

- study. *Circulation*. 2013;127:1270–1275. DOI: 10.1161/CIRCULATIONAHA.112.001183.
30. LaMonte MJ, Durstine JL, Addy CL, Irwin ML, Ainsworth BE. Physical activity, physical fitness, and Framingham 10-year risk score: the cross-cultural activity participation study. *J Cardiopulm Rehabil*. 2001;21:63–70. DOI: 10.1097/00008483-200103000-00001.
 31. Arredondo EM, Sotres-Alvarez D, Stoutenberg M, Davis SM, Crespo NC, Carnethon MR, Castañeda SF, Isasi CR, Espinoza RA, Daviglus ML, et al. Physical activity levels in US Latino/Hispanic adults: results from the Hispanic Community Health Study/Study of Latinos. *Am J Prev Med*. 2016;50:500–508. DOI: 10.1016/j.amepre.2015.08.029.
 32. Adler N, Stewart J; Psychosocial_Working_Group. The MacArthur Scale of Subjective Social Status [printable version]. Published 2007. Available at: <https://www.macses.ucsf.edu/research/psychosocial/usladder.php>. Accessed October 25, 2016.
 33. Adler NE, Epel ES, Castellazzo G, Ickovics JR. Relationship of subjective and objective social status with psychological and physiological functioning: preliminary data in healthy white women. *Health Psychol*. 2000;19:586–592. DOI: 10.1037/0278-6133.19.6.586.
 34. Ostrove JM, Adler NE, Kuppermann M, Washington AE. Objective and subjective assessments of socioeconomic status and their relationship to self-rated health in an ethnically diverse sample of pregnant women. *Health Psychol*. 2000;19:613–618. DOI: 10.1037/0278-6133.19.6.613.
 35. Thomson MD, Hoffman-Goetz L. Defining and measuring acculturation: a systematic review of public health studies with Hispanic populations in the United States. *Soc Sci Med*. 2009;69:983–991. DOI: 10.1016/j.socscimed.2009.05.011.
 36. Commodore-Mensah Y, Ukonu N, Obisesan O, Kumi Aboagye J, Agyemang C, Reilly CM, Dunbar SB, Okosun IS. Length of residence in the United States is associated with a higher prevalence of cardiometabolic risk factors in immigrants: a contemporary analysis of the National Health Interview Survey. *J Am Heart Assoc*. 2016;5:e004059. DOI: 10.1161/JAHA.116.004059.
 37. Ware J Jr, Kosinski M, Keller SD. A 12-Item Short-Form Health Survey: construction of scales and preliminary tests of reliability and validity. *Med Care*. 1996;34:220–233. DOI: 10.1097/00005650-199603000-00003.
 38. Radloff LS. The CES-D scale: a self-report depression scale for research in the general population. *Appl Psychol Meas*. 1977;1:385–401. DOI: 10.1177/014662167700100306.
 39. Fleuriet KJ, Sunil TS. Perceived social stress, pregnancy-related anxiety, depression and subjective social status among pregnant Mexican and Mexican American women in south Texas. *J Health Care Poor Underserved*. 2014;25:546–561. DOI: 10.1353/hpu.2014.0092.
 40. Franzini L, Fernandez-Esquer ME. The association of subjective social status and health in low-income Mexican-origin individuals in Texas. *Soc Sci Med*. 2006;63:788–804. DOI: 10.1016/j.socscimed.2006.01.009.
 41. Garza JR, Glenn BA, Mistry RS, Ponce NA, Zimmerman FJ. Subjective social status and self-reported health among US-born and immigrant Latinos. *J Immigr Minor Health*. 2017;19:108–119. DOI: 10.1007/s10903-016-0346-x.
 42. Andrade FCD, Viruell-Fuentes E. Latinos and the changing demographic landscape: key dimensions for infrastructure building. In: Buki LP, Piedra LM, eds. *Creating Infrastructures for Latino Mental Health*. New York: Springer; 2011:3–30.
 43. Lopez MH, Krogstad JM, Flores A. Key facts about young Latinos, one of the nation's fastest-growing populations. Pew Research Center; Published 2018. Available at: <http://www.pewresearch.org/fact-tank/2018/09/13/key-facts-about-young-latinos/>. Accessed May 12, 2020.
 44. Kaplan RC, Bangdiwala SI, Barnhart JM, Castañeda SF, Gellman MD, Lee DJ, Pérez-Stable EJ, Talavera GA, Youngblood ME, Giachello AL. Smoking among U.S. Hispanic/Latino adults: the Hispanic Community Health Study/Study of Latinos. *Am J Prev Med*. 2014;46:496–506. DOI: 10.1016/j.amepre.2014.01.014.
 45. Merzel CR, Isasi CR, Strizich G, Castañeda SF, Gellman M, Maisonet Giachello AL, Lee DJ, Penedo FJ, Perreira KM, Kaplan RC. Smoking cessation among US Hispanic/Latino adults: findings from the Hispanic Community Health Study/Study of Latinos (HCHS/SOL). *Prev Med*. 2015;81:412–419. DOI: 10.1016/j.ypmed.2015.10.006.
 46. Schisterman EF, Cole SR, Platt RW. Overadjustment bias and unnecessary adjustment in epidemiologic studies. *Epidemiology*. 2009;20:488–495. DOI: 10.1097/EDE.0b013e3181a819a1.
 47. Roberts BW, Kuncel NR, Shiner R, Caspi A, Goldberg LR. The power of personality: the comparative validity of personality traits, socioeconomic status, and cognitive ability for predicting important life outcomes. *Perspect Psychol Sci*. 2007;2:313–345. DOI: 10.1111/j.1745-6916.2007.00047.x.
 48. Haan MN, Zeki Al-Hazzouri A, Aiello AE. Life-span socioeconomic trajectory, nativity, and cognitive aging in Mexican Americans: the Sacramento Area Latino Study on Aging. *J Gerontol B Psychol Sci Soc Sci*. 2011;66B:i102–i110. DOI: 10.1093/geronb/gbq071.
 49. Abelson RP. A variance explanation paradox: when a little is a lot. *Psychol Bull*. 1985;97:129. DOI: 10.1037/0033-2909.97.1.129.
 50. Rosenthal R. How are we doing in soft psychology? *Am Psychol*. 1990;45:775. DOI: 10.1037/0003-066X.45.6.775.
 51. Cooper DC, Milic MS, Mills PJ, Bardwell WA, Ziegler MG, Dimsdale JE. Endothelial function: the impact of objective and subjective socioeconomic status on flow-mediated dilation. *Ann Behav Med*. 2010;39:222–231. DOI: 10.1007/s12160-010-9181-9.
 52. Euteneuer F, Mills PJ, Rief W, Ziegler MG, Dimsdale JE. Subjective social status predicts in vivo responsiveness of β -adrenergic receptors. *Health Psychol*. 2012;31:525. DOI: 10.1037/a0025990.
 53. Mills PJ, Dimsdale JE. The promise of adrenergic receptor studies in psychophysiology research II: applications, limitations, and progress. *Psychosom Med*. 1993;55:448–457. DOI: 10.1097/00006842-199309000-00008.
 54. Triposkiadis F, Karayannis G, Giamouzis G, Skoularigis J, Louridas G, Butler J. The sympathetic nervous system in heart failure physiology, pathophysiology, and clinical implications. *J Am Coll Cardiol*. 2009;54:1747–1762. DOI: 10.1016/j.jacc.2009.05.015.
 55. Wu S, An S, Li W, Lichtenstein AH, Gao J, Kris-Etherton PM, Wu Y, Jin C, Huang S, Hu FB, et al. Association of trajectory of cardiovascular health score and incident cardiovascular disease. *JAMA Netw Open*. 2019;2:e194758. DOI: 10.1001/jamanetworkopen.2019.4758.