

Lawrence Berkeley National Laboratory

Recent Work

Title

THE REACTION OF HEPTASULFUR BODE WITH DIBORANE

Permalink

<https://escholarship.org/uc/item/94h7m22g>

Author

Mendelsohn, Marshall H.

Publication Date

1971-12-01

c.1

RECEIVED
LAWRENCE
RADIATION LABORATORY

FEB 4 1972

LIBRARY AND
DOCUMENTS SECTION

THE REACTION OF HEPTASULFUR IMIDE
WITH DIBORANE

Marshall H. Mendelsohn and William L. Jolly

December 1971

AEC Contract No. W-7405-eng-48

For Reference

Not to be taken from this room



DISCLAIMER

This document was prepared as an account of work sponsored by the United States Government. While this document is believed to contain correct information, neither the United States Government nor any agency thereof, nor the Regents of the University of California, nor any of their employees, makes any warranty, express or implied, or assumes any legal responsibility for the accuracy, completeness, or usefulness of any information, apparatus, product, or process disclosed, or represents that its use would not infringe privately owned rights. Reference herein to any specific commercial product, process, or service by its trade name, trademark, manufacturer, or otherwise, does not necessarily constitute or imply its endorsement, recommendation, or favoring by the United States Government or any agency thereof, or the Regents of the University of California. The views and opinions of authors expressed herein do not necessarily state or reflect those of the United States Government or any agency thereof or the Regents of the University of California.

Contribution from the Chemistry Department, University of California, and
the Inorganic Materials Research Division, Lawrence Berkeley Laboratory,
Berkeley, California 94720

The Reaction of Heptasulfur Imide with Diborane

By Marshall H. Mendelsohn and William L. Jolly*

It is known that heptasulfur imide reacts with boron trichloride¹ and boron tribromide to give S_7BNCl_2 ¹ and S_7NBr_2 ², respectively. In this note we report the results of our study of the reaction of heptasulfur imide with diborane. This study was undertaken to determine whether the reaction gives an analogous product (S_7NBH_2), a borane adduct (S_7NBH_3), or a diborane derivative such as $S_7NB_2H_5$.

Results and Discussion

The reaction of heptasulfur imide with diborane was carried out in various ether solvents, at either room temperature or 0°. The experimental results, summarized in Table I, show that, with diborane in excess, one-half mole of diborane reacts per mole of S_7NH and that one mole of hydrogen is formed per mole of S_7NH . This stoichiometry corresponds to the following reaction:




Table I.

Stoichiometry of the Reaction of S_7NH with B_2H_6

Solvent	Temperature, °C	S_7NH , mmol	B_2H_6 , mmol	measured H_2/S_7NH	time of meas., hrs	B_2H_6 consumed / S_7NH
monoglyme ^a	23	0.288	0.41	1.11 ^b	17	0.49
monoglyme	23	0.992	0.72	1.35 ^b	64.5	-
monoglyme	0	0.574	0.516	1.0 ^c	66	0.495
	23			1.15 ^b	24	
THF ^a	23	0.386	0.356	0.99	67	0.505
THF	23	0.616	0.50	1.13 ^b	65.5	-
THF	23	0.567	0.264	1.09 ^{b,d}	145	-
THF	0	0.611	0.593	1.02	497	0.52
THF	0	0.302	0.0716	1.31 ^{b,d}	978	-
dioxane	23	0.505	0.633	2.20 ^b	319	-
diethyl ether ^a	0	0.575	0.292	0.98	260	-


^a reaction mixture not stirred

^b traces of H_2 still being evolved, due to slow decomposition

^c stopped purposely to pump out excess B_2H_6

^d listed as H_2/B_2H_6

When the reaction is carried out in diethyl ether at 0°, some of the product precipitates as a white, crystalline ether adduct, $S_7NBH_2 \cdot Et_2O$. Similar adducts can be obtained from the other ethers by vacuum evaporation of the solvent and removal of excess diborane. The compounds undergo rapid hydrolysis with excess water at room temperature, as follows:




The ^{11}B nmr spectrum of S_7NBH_2 in diethyl ether solution consists of a 1:2:1 triplet due to coupling with the two protons of the BH_2 group. The Raman spectrum of the diethyl ether adduct has lines at 2419 and 2409 cm^{-1} , characteristic of terminal BH_2 groups, and lines at 766 and 750 cm^{-1} , typical S-N stretching frequencies.

As can be noted from Table I, the only products that were stable (as determined by hydrogen evolution) were those obtained from diethyl ether and tetrahydrofuran. The diethyl ether adduct is stable at room temperature in the absence of air and moisture, and the product from tetrahydrofuran is stable at 0°. Hydrogen, hydrogen sulfide, and ether were identified as decomposition products of the tetrahydrofuran and monoglyme adducts at room temperature.

A white solid pyridine adduct, $S_7NBH_2 \cdot C_5H_5N$ was isolated from the reaction of $S_7NBH_2 \cdot Et_2O$ with pyridine. Pure samples of this compound were stable indefinitely in vacuo at 0° and could be kept in an ordinary closed bottle at room temperature for as long as 5 days without decomposition. However, impure samples decomposed within a day or two.

When the reaction of heptasulfur imide and diborane is carried out with an excess of S_7NH in tetrahydrofuran at 0° , the reaction does not stop with the formation of S_7NBH_2 , as evidenced by the continued slow evolution of hydrogen. Presumably the following subsequent reaction occurs:



However, the reaction was incomplete after 1 month and was not further studied.

Experimental Section

Reagents and procedure. - Heptasulfur imide was prepared by the usual method³ and was purified by one recrystallization from methanol followed by repeated recrystallizations from carbon tetrachloride until the solutions were colorless and the solid melted at $113-114^\circ$ (lit. 113.5°).⁴ The infrared spectrum agreed with that reported in the literature.⁴ Diborane was made by the reaction of potassium borohydride with phosphoric acid⁵ and purified by vacuum distillation through a -112° trap. Tetrahydrofuran, diethyl ether, and dioxane were distilled from lithium aluminum hydride immediately before use. Monoglyme (1,2-dimethoxyethane) was distilled first from calcium hydride and then from lithium aluminum hydride. Pyridine was distilled from potassium hydroxide, and carbon tetrachloride was distilled from phosphorus pentoxide.

Approximately 10 ml of solvent was vacuum-distilled into a 50-ml flask containing a known weight of heptasulfur imide. A known amount of diborane was condensed into the reaction vessel, and the vessel was allowed to warm to either room temperature or 0°. At various times the evolved hydrogen was collected using a Toepler pump, measured in a gas buret, and identified by mass spectroscopy. To isolate the crystalline product $S_7NBH_2 \cdot Et_2O$, an all-glass system was employed. The apparatus was sealed off after addition of the reactants and was provided with a break-seal for removal of hydrogen and a side arm for washing the product by decantation. The product was then removed in a dry box. Anal. Calcd for $S_7NBH_2 \cdot C_4H_{10}O$: C, 14.76; H, 3.69; N, 4.30; S, 69.01. Found: C, 14.86; H, 3.69; N, 4.45; S, 68.68.

A sample of S_7NBH_2 prepared in tetrahydrofuran was treated with a 500-fold excess of distilled water. The evolved hydrogen was collected and measured as described above. The precipitated S_7NH was collected by filtration, air-dried, and then weighed. It was identified by its infrared spectrum.⁴ The boric acid formed was determined by a standard double titration method.⁶ Anal. Calcd for S_7NBH_2 prepared from 0.386 mmol of S_7NH : H_2 , 0.772 mmol; $B(OH)_3$, 0.386 mmol; S_7NH , 0.386 mmol. Found: H_2 , 0.773 mmol; $B(OH)_3$, 0.393 mmol; S_7NH , 0.31 mmol.

Raman spectrum. - The following lines were observed for the diethyl ether adduct ($cm^{-1} \pm 3 cm^{-1}$): 2969(w), 2930(w-m), 2419(w), 2409(sh), 1471(w), 1452(w-m), 1442(sh), 1324(w), 1181(w-m), 1091(w-m), 1005(w), 870(w-m), 851(sh), 832(w), 766(w-m), 750(w-m), 602(m), 476(vs), 458(m), 437(m), 292(s), 251(m), 213(vs), 201(s), 178(m), and 153(s). The Raman

spectrometer consisted of a Spectra-Physics 125 helium-neon laser with line-filter (35-mW) and a Spex 1401 monochromator equipped with an ITT FW-130 photomultiplier tube.

Nmr spectra.- The samples for the ^{11}B nmr spectra were prepared and recorded in 12 mm od tubes containing diethyl ether, in which the compound is only slightly soluble. For this reason the spectra were computer-averaged over 10-150 scans. The spectrum consisted of a 1:2:1 triplet centered 33.5 ppm upfield from an external reference peak of boron trichloride, with $J_{\text{B-H}} = 136$ Hz. A Varian model V4311 high resolution nmr spectrometer operated at 8.134 MHz was used for the boron spectra. Proton spectra of samples prepared from monoglyme and diethyl ether were recorded in CS_2 using a Varian A-60 spectrometer. Although the samples decomposed in CS_2 , the spectra did indicate the presence of coordinated the/ether. For the sample from diethyl ether, a triplet at 1.98 ppm and a quarter at 4.75 ppm, relative to TMS, were observed, while from monoglyme two singlets at 4.63 ppm and 4.67 ppm and a complex pattern centered at 4.87 ppm were noted. The complexity of the latter spectrum indicates that only one of the oxygen atoms of monoglyme was coordinated to S_7NBH_2 .

X-ray diffraction.- Some preliminary X-ray diffraction data on a single crystal of $\text{S}_7\text{NBH}_2 \cdot \text{C}_4\text{H}_{10}\text{O}$ correspond to the space group P_{nma} and the unit cell dimensions (\AA) $\underline{a} = 15.80$, $\underline{b} = 13.16$, and $\underline{c} = 6.75$.

Preparation and characterization of $\text{S}_7\text{NBH}_2 \cdot \text{C}_5\text{H}_5\text{N}$.- The pyridine adduct is prepared by addition of excess, dry pyridine to the solid $\text{S}_7\text{NBH}_2 \cdot \text{Et}_2\text{O}$. The solution is evaporated to a volume of 0.2-0.3 ml, and dry CCl_4 is

then added. A white solid is precipitated and is washed with a small amount of chilled CCl_4 . The infrared spectrum of a Nujol mull recorded with a Perkin-Elmer Model 137B Infracord Spectrophotometer consisted of the following peaks ($\text{cm}^{-1} \pm 10 \text{ cm}^{-1}$): 2415(m), 2353(sh), 1618(m), 1186(m), 1157(w), 1139(sh), 1131(s), 1100(m-s), 1025(w), 1013(w), 977(w), 961(sh), 953(m), 943(s), 929(w), 917(m), 775(m), 768(m), 751(s), 687(s). A sample melting at $87.5\text{-}88^\circ$ was analyzed. Anal. Calcd for $\text{S}_7\text{NBH}_2 \cdot \text{C}_5\text{H}_5\text{N}$: C, 18.17; H, 2.14; N, 8.48; S, 67.92; B, 3.3. Found: C, 18.97; H, 2.08; N, 8.62; S, 67.98; B, 3.2 (as a B_2O_3 residue). A proton nmr spectrum showed the protons on the pyridine ring, but we were unable to see the protons attached to boron.

Acknowledgment.- We wish to thank Dr. James Shear at the U. S. Department of Agriculture Western Regional Research Laboratory for recording the Raman spectrum, Dr. Allan Zalkin of the Lawrence Berkeley Laboratory for the X-ray diffraction data, and Dr. James Neely and John Illige for recording the ^{11}B nmr spectra. This research was supported by the U. S. Atomic Energy Commission.

References

- (1) H. G. Heal, J. Inorg. Nuclear Chem., 20, 165 (1961).
- (2) H. G. Heal, J. Chem. Soc., 4442 (1962).
- (3) M. Becke-Goehring and E. Fluck, Inorg. Syn., 8, 103 (1966).
- (4) H. G. Heal and J. Kane, Inorg. Syn., 11, 193 (1969).
- (5) A. D. Norman and W. L. Jolly, Inorg. Syn., 11, 15 (1969).
- (6) A. I. Vogel, "Quantitative Inorganic Analysis," Wiley, New York, 1961, pp. 252-4.

LEGAL NOTICE

This report was prepared as an account of work sponsored by the United States Government. Neither the United States nor the United States Atomic Energy Commission, nor any of their employees, nor any of their contractors, subcontractors, or their employees, makes any warranty, express or implied, or assumes any legal liability or responsibility for the accuracy, completeness or usefulness of any information, apparatus, product or process disclosed, or represents that its use would not infringe privately owned rights.

TECHNICAL INFORMATION DIVISION
LAWRENCE BERKELEY LABORATORY
UNIVERSITY OF CALIFORNIA
BERKELEY, CALIFORNIA 94720