

UC Berkeley

UC Berkeley Previously Published Works

Title

All-sky Search for Time-integrated Neutrino Emission from Astrophysical Sources with 7 yr of IceCube Data

Permalink

<https://escholarship.org/uc/item/95z9x104>

Journal

The Astrophysical Journal, 835(2)

ISSN

0004-637X

Authors

Aartsen, MG
Abraham, K
Ackermann, M
[et al.](#)

Publication Date

2017-02-01

DOI

10.3847/1538-4357/835/2/151

Peer reviewed

ALL-SKY SEARCH FOR TIME-INTEGRATED NEUTRINO EMISSION FROM ASTROPHYSICAL SOURCES WITH 7 YR OF ICECUBE DATA

ICECUBE COLLABORATION: M. G. AARTSEN¹, K. ABRAHAM², M. ACKERMANN³, J. ADAMS⁴, J. A. AGUILAR⁵, M. AHLERS⁶, M. AHRENS⁷, D. ALTMANN⁸, K. ANDEEN⁹, T. ANDERSON¹⁰, I. ANSSEAU⁵, G. ANTON⁸, M. ARCHINGER¹¹, C. ARGÜELLES¹², J. AUFFENBERG¹³, S. AXANI¹², X. BAI¹⁴, S. W. BARWICK¹⁵, V. BAUM¹¹, R. BAY¹⁶, J. J. BEATTY^{17,18}, J. BECKER TJUS¹⁹, K.-H. BECKER²⁰, S. BENZVI²¹, D. BERLEY²², E. BERNARDINI³, A. BERNHARD², D. Z. BESSON²³, G. BINDER^{24,16}, D. BINDIG²⁰, M. BISSOK¹³, E. BLAUFUSS²², S. BLOT³, C. BOHM⁷, M. BÖRNER²⁵, F. BOS¹⁹, D. BOSE²⁶, S. BÖSER¹¹, O. BOTNER²⁷, J. BRAUN⁶, L. BRAYEUR²⁸, H.-P. BRETZ³, S. BRON²⁹, A. BURGMAN²⁷, T. CARVER²⁹, M. CASIER²⁸, E. CHEUNG²², D. CHIRKIN⁶, A. CHRISTOV²⁹, K. CLARK³⁰, L. CLASSEN³¹, S. COENDERS², G. H. COLLIN¹², J. M. CONRAD¹², D. F. COWEN^{10,32}, R. CROSS²¹, M. DAY⁶, J. P. A. M. DE ANDRÉ³³, C. DE CLERCQ²⁸, E. DEL PINO ROSENDO¹¹, H. DEMBINSKI³⁴, S. DE RIDDER³⁵, P. DESIATI⁶, K. D. DE VRIES²⁸, G. DE WASSEIGE²⁸, M. DE WITH³⁶, T. DEYOUNG³³, J. C. DÍAZ-VÉLEZ⁶, V. DI LORENZO¹¹, H. DUJMOVIC²⁶, J. P. DUMM⁷, M. DUNKMAN¹⁰, B. EBERHARDT¹¹, T. EHRHARDT¹¹, B. EICHMANN¹⁹, P. ELLER¹⁰, S. EULER²⁷, P. A. EVENSON³⁴, S. FAHEY⁶, A. R. FAZELY³⁷, J. FEINTZEIG⁶, J. FELDE²², K. FILIMONOV¹⁶, C. FINLEY⁷, S. FLIS⁷, C.-C. FÖSIG¹¹, A. FRANCKOWIAK³, E. FRIEDMAN²², T. FUCHS²⁵, T. K. GAISSER³⁴, J. GALLAGHER³⁸, L. GERHARDT^{24,16}, K. GHORBANI⁶, W. GIANG³⁹, L. GLADSTONE⁶, T. GLAUCH¹³, T. GLÜSENKAMP³, A. GOLDSCHMIDT²⁴, G. GOLUP²⁸, J. G. GONZALEZ³⁴, D. GRANT³⁹, Z. GRIFFITH⁶, C. HAACK¹³, A. HAJ ISMAIL³⁵, A. HALLGREN²⁷, F. HALZEN⁶, E. HANSEN⁴⁰, T. HANSMANN¹³, K. HANSON⁶, D. HEBECKER³⁶, D. HEEREMAN⁵, K. HELBING²⁰, R. HELLAUER²², S. HICKFORD²⁰, J. HIGNIGHT³³, G. C. HILL¹, K. D. HOFFMAN²², R. HOFFMANN²⁰, K. HOLZAPFEL², K. HOSHINA^{6,53}, F. HUANG¹⁰, M. HUBER², K. HULTQVIST⁷, S. IN²⁶, A. ISHIHARA⁴¹, E. JACOBI³, G. S. JAPARIDZE⁴², M. JEONG²⁶, K. JERO⁶, B. J. P. JONES¹², M. JURKOVIC², A. KAPPES³¹, T. KARG³, A. KARLE⁶, U. KATZ⁸, M. KAUER⁶, A. KEIVANI¹⁰, J. L. KELLEY⁶, A. KHEIRANDISH⁶, M. KIM²⁶, T. KINTSCHER³, J. KIRYLUK⁴³, T. KITTLER⁸, S. R. KLEIN^{24,16}, G. KOHNEN⁴⁴, R. KOIRALA³⁴, H. KOLANOSKI³⁶, R. KONIETZ¹³, L. KÖPKE¹¹, C. KOPPER³⁹, S. KOPPER²⁰, D. J. KOSKINEN⁴⁰, M. KOWALSKI^{36,3}, K. KRINGS², M. KROLL¹⁹, G. KRÜCKL¹¹, C. KRÜGER⁶, J. KUNNEN²⁸, S. KUNWAR³, N. KURAHASHI⁴⁵, T. KUWABARA⁴¹, M. LABARE³⁵, J. L. LANFRANCHI¹⁰, M. J. LARSON⁴⁰, F. LAUBER²⁰, D. LENNARZ³³, M. LESIAK-BZDAK⁴³, M. LEUERMANN¹³, L. LU⁴¹, J. LÜNEMANN²⁸, J. MADSEN⁴⁶, G. MAGGI²⁸, K. B. M. MAHN³³, S. M. MANDELARTZ¹⁹, R. MARUYAMA⁴⁷, K. MASE⁴¹, R. MAUNU²², F. MCNALLY⁶, K. MEAGHER⁵, M. MEDICI⁴⁰, M. MEIER²⁵, A. MELI³⁵, T. MENNE²⁵, G. MERINO⁶, T. MEURES⁵, S. MIARECKI^{24,16}, L. MOHRMANN³, T. MONTARULI²⁹, M. MOULAI¹², R. NAHNHAUER³, U. NAUMANN²⁰, G. NEER³³, H. NIEDERHAUSEN⁴³, S. C. NOWICKI³⁹, D. R. NYGREN²⁴, A. OBERTACKER POLLMANN²⁰, A. OLIVAS²², A. O'MURCHADHA⁵, T. PALCZEWSKI⁴⁸, H. PANDYA³⁴, D. V. PANKOVA¹⁰, P. PEIFFER¹¹, Ö. PENEK¹³, J. A. PEPPER⁴⁸, C. PÉREZ DE LOS HEROS²⁷, D. PIELOTH²⁵, E. PINAT⁵, P. B. PRICE¹⁶, G. T. PRZYBYLSKI²⁴, M. QUINNAN¹⁰, C. RAAB⁵, L. RÄDEL¹³, M. RAMEEZ⁴⁰, K. RAWLINS⁴⁹, R. REIMANN¹³, B. RELETHFORD⁴⁵, M. RELICH⁴¹, E. RESCONI², W. RHODE²⁵, M. RICHMAN⁴⁵, B. RIEDEL³⁹, S. ROBERTSON¹, M. RONGEN¹³, C. ROTT²⁶, T. RUHE²⁵, D. RYCKBOSCH³⁵, D. RYSEWYK³³, L. SABBATINI⁶, S. E. SANCHEZ HERRERA³⁹, A. SANDROCK²⁵, J. SANDROOS¹¹, S. SARKAR^{40,50}, K. SATALECKA³, P. SCHLUNDER²⁵, T. SCHMIDT²², S. SCHOENEN¹³, S. SCHÖNEBERG¹⁹, L. SCHUMACHER¹³, D. SECKEL³⁴, S. SEUNARINE⁴⁶, D. SOLDIN²⁰, M. SONG²², G. M. SPICZAK⁴⁶, C. SPIERING³, T. STANEV³⁴, A. STASIK³, J. STETTNER¹³, A. STEUER¹¹, T. STEZELBERGER²⁴, R. G. STOKSTAD²⁴, A. STÖSSL³, R. STRÖM²⁷, N. L. STROTJOHANN³, G. W. SULLIVAN²², M. SUTHERLAND¹⁷, H. TAAVOLA²⁷, I. TABOADA⁵¹, J. TATAR^{24,16}, F. TENHOLT¹⁹, S. TER-ANTONYAN³⁷, A. TERLIUK³, G. TESIĆ¹⁰, S. TILAV³⁴, P. A. TOALE⁴⁸, M. N. TOBIN⁶, S. TOSCANO²⁸, D. TOSI⁶, M. TSELENGIDOU⁸, A. TURCATI², E. UNGER²⁷, M. USNER³, J. VANDENBROUCKE⁶, N. VAN EIJNDHOVEN²⁸, S. VANHEULE³⁵, M. VAN ROSSEM⁶, J. VAN SANTEN³, J. VEENKAMP², M. VEHRING¹³, M. VOGEL⁵², E. VOGEL¹³, M. VRAEGHE³⁵, C. WALCK⁷, A. WALLACE¹, M. WALLRAFF¹³, N. WANDKOWSKY⁶, CH. WEAVER³⁹, M. J. WEISS¹⁰, C. WENDT⁶, S. WESTERHOFF⁶, B. J. WHELAN¹, S. WICKMANN¹³, K. WIEBE¹¹, C. H. WIEBUSCH¹³, L. WILLE⁶, D. R. WILLIAMS⁴⁸, L. WILLS⁴⁵, M. WOLF⁷, T. R. WOOD³⁹, E. WOOLSEY³⁹, K. WOSCHNAGG¹⁶, D. L. XU⁶, X. W. XU³⁷, Y. XU⁴³, J. P. YANEZ³, G. YODH¹⁵, S. YOSHIDA⁴¹, AND M. ZOLL⁷

(Accepted December 6, 2016)

¹Department of Physics, University of Adelaide, Adelaide, 5005, Australia

²Physik-department, Technische Universität München, D-85748 Garching, Germany

³DESY, D-15735 Zeuthen, Germany

⁴Dept. of Physics and Astronomy, University of Canterbury, Private Bag 4800, Christchurch, New Zealand

⁵Université Libre de Bruxelles, Science Faculty CP230, B-1050 Brussels, Belgium

⁶Dept. of Physics and Wisconsin IceCube Particle Astrophysics Center, University of Wisconsin, Madison, WI 53706, USA

⁷Oskar Klein Centre and Dept. of Physics, Stockholm University, SE-10691 Stockholm, Sweden

⁸Erlangen Centre for Astroparticle Physics, Friedrich-Alexander-Universität Erlangen-Nürnberg, D-91058 Erlangen, Germany

⁹Department of Physics, Marquette University, Milwaukee, WI 53201, USA

¹⁰Dept. of Physics, Pennsylvania State University, University Park, PA 16802, USA

¹¹Institute of Physics, University of Mainz, Staudinger Weg 7, D-55099 Mainz, Germany

¹²Dept. of Physics, Massachusetts Institute of Technology, Cambridge, MA 02139, USA

¹³III. Physikalisches Institut, RWTH Aachen University, D-52056 Aachen, Germany

¹⁴Physics Department, South Dakota School of Mines and Technology, Rapid City, SD 57701, USA

¹⁵Dept. of Physics and Astronomy, University of California, Irvine, CA 92697, USA

¹⁶Dept. of Physics, University of California, Berkeley, CA 94720, USA

- ¹⁷Dept. of Physics and Center for Cosmology and Astro-Particle Physics, Ohio State University, Columbus, OH 43210, USA
- ¹⁸Dept. of Astronomy, Ohio State University, Columbus, OH 43210, USA
- ¹⁹Fakultät für Physik & Astronomie, Ruhr-Universität Bochum, D-44780 Bochum, Germany
- ²⁰Dept. of Physics, University of Wuppertal, D-42119 Wuppertal, Germany
- ²¹Dept. of Physics and Astronomy, University of Rochester, Rochester, NY 14627, USA
- ²²Dept. of Physics, University of Maryland, College Park, MD 20742, USA
- ²³Dept. of Physics and Astronomy, University of Kansas, Lawrence, KS 66045, USA
- ²⁴Lawrence Berkeley National Laboratory, Berkeley, CA 94720, USA
- ²⁵Dept. of Physics, TU Dortmund University, D-44221 Dortmund, Germany
- ²⁶Dept. of Physics, Sungkyunkwan University, Suwon 440-746, Korea
- ²⁷Dept. of Physics and Astronomy, Uppsala University, Box 516, SE-75120 Uppsala, Sweden
- ²⁸Vrije Universiteit Brussel, Dienst ELEM, B-1050 Brussels, Belgium
- ²⁹Département de physique nucléaire et corpusculaire, Université de Genève, CH-1211 Genève, Switzerland
- ³⁰Dept. of Physics, University of Toronto, Toronto, ON M5S 1A7, Canada
- ³¹Institut für Kernphysik, Westfälische Wilhelms-Universität Münster, D-48149 Münster, Germany
- ³²Dept. of Astronomy and Astrophysics, Pennsylvania State University, University Park, PA 16802, USA
- ³³Dept. of Physics and Astronomy, Michigan State University, East Lansing, MI 48824, USA
- ³⁴Bartol Research Institute and Dept. of Physics and Astronomy, University of Delaware, Newark, DE 19716, USA
- ³⁵Dept. of Physics and Astronomy, University of Gent, B-9000 Gent, Belgium
- ³⁶Institut für Physik, Humboldt-Universität zu Berlin, D-12489 Berlin, Germany
- ³⁷Dept. of Physics, Southern University, Baton Rouge, LA 70813, USA
- ³⁸Dept. of Astronomy, University of Wisconsin, Madison, WI 53706, USA
- ³⁹Dept. of Physics, University of Alberta, Edmonton, AB T6G 2E1, Canada
- ⁴⁰Niels Bohr Institute, University of Copenhagen, DK-2100 Copenhagen, Denmark
- ⁴¹Dept. of Physics and Institute for Global Prominent Research, Chiba University, Chiba 263-8522, Japan
- ⁴²CTSPS, Clark-Atlanta University, Atlanta, GA 30314, USA
- ⁴³Dept. of Physics and Astronomy, Stony Brook University, Stony Brook, NY 11794-3800, USA
- ⁴⁴Université de Mons, B-7000 Mons, Belgium
- ⁴⁵Dept. of Physics, Drexel University, 3141 Chestnut Street, Philadelphia, PA 19104, USA
- ⁴⁶Dept. of Physics, University of Wisconsin, River Falls, WI 54022, USA
- ⁴⁷Dept. of Physics, Yale University, New Haven, CT 06520, USA
- ⁴⁸Dept. of Physics and Astronomy, University of Alabama, Tuscaloosa, AL 35487, USA
- ⁴⁹Dept. of Physics and Astronomy, University of Alaska Anchorage, 3211 Providence Dr., Anchorage, AK 99508, USA
- ⁵⁰Dept. of Physics, University of Oxford, 1 Keble Road, Oxford OX1 3NP, UK
- ⁵¹School of Physics and Center for Relativistic Astrophysics, Georgia Institute of Technology, Atlanta, GA 30332, USA
- ⁵²Physikalisches Institut, Universität Bonn, Nussallee 12, D-53115 Bonn, Germany
- ⁵³Earthquake Research Institute, University of Tokyo, Bunkyo, Tokyo 113-0032, Japan

ABSTRACT

Since the recent detection of an astrophysical flux of high-energy neutrinos, the question of its origin has not yet fully been answered. Much of what is known about this flux comes from a small event sample of high neutrino purity, good energy resolution, but large angular uncertainties. In searches for point-like sources, on the other hand, the best performance is given by using large statistics and good angular reconstructions. Track-like muon events produced in neutrino interactions satisfy these requirements. We present here the results of searches for point-like sources with neutrinos using data acquired by the IceCube detector over 7 yr from 2008 to 2015. The discovery potential of the analysis in the northern sky is now significantly below $E_\nu^2 d\phi/dE_\nu = 10^{-12} \text{ TeV cm}^{-2} \text{ s}^{-1}$, on average 38% lower than the sensitivity of the previously published analysis of 4 yr exposure. No significant clustering of neutrinos above background expectation was observed, and implications for prominent neutrino source candidates are discussed.

Keywords: astroparticle physics — galaxies: active — neutrinos

1. INTRODUCTION

One outstanding question in astroparticle physics is the origin of ultra-high-energy cosmic rays (UHECRs). In the paradigm of multi-messenger astronomy, both photons and neutrinos can help resolve the sources of UHECRs (Beatty & Westerhoff 2009; Kotera & Olinto 2011). Photons and neutrinos are believed to be produced in the astrophysical beam dump of cosmic-ray particles interacting with matter at the source location. Due to lack of electric charge, they point back to their origin, whereas cosmic rays are deflected by tangled magnetic fields in the universe. Sources of high-energy γ -rays in our Galaxy and extragalactic objects are detected over a wide range of energies (Hinton & Hofmann 2009), but both hadronic and leptonic processes can produce γ -rays. Neutrinos, on the other hand, trace hadronic interactions and therefore are a smoking-gun signature of cosmic-ray acceleration (Learned & Mannheim 2000; Halzen & Hooper 2002; Anchordoqui & Montaruli 2010; Anchordoqui et al. 2014). Other possible types of sources could be hidden in γ -rays and only identified using neutrinos (Murase et al. 2016; Senno et al. 2016).

IceCube recently reported the first observation of high-energy astrophysical neutrinos with more than 5σ significance (Aartsen et al. 2013a, 2014c). Neutrinos with energies up to and exceeding 1 PeV are observed in events starting inside the detector. Since then, this result is confirmed in other detection channels of interactions of $\nu_\mu + \bar{\nu}_\mu$ in the northern sky (Aartsen et al. 2015c, 2016b). The overall flux observed is consistent so far with an isotropic emission over the full sky and all neutrino flavors (Aartsen et al. 2015d,a).

This paper presents the most recent results of searches for point-like steady emission of neutrinos using track-like events traversing the IceCube detector. The statistics are increased by adding 3 yr of exposure to the previous analyses (Abbasi et al. 2011; Aartsen et al. 2013c, 2014e). A sample of 712,830 events is obtained during 7 yr of data recording through 2015 June. In addition, starting tracks are used in a separate sample to help access lower energies in the southern sky (Aartsen et al. 2016a).

In Section 2, the IceCube Neutrino Observatory is introduced and the two samples of through-going and starting tracks are characterized. In Section 3, the statistical method of unbinned likelihood maximization for clustering searches is discussed. Section 4 presents the results and their implications regarding neutrino sources, and in Section 5 conclusions are drawn.

2. THE ICECUBE NEUTRINO OBSERVATORY

Interactions of neutrinos in IceCube are detected using Cerenkov light emitted by relativistic charged secondary particles. To serve this purpose, one cubic kilometer of Antarctic ice was instrumented at the South Pole (Achterberg et al. 2006). A total of 5160 digital optical modules (DOMs) detect light emission in the ice at a depth ranging from 1450 to 2450m. The DOMs consist of a photomultiplier tube, electronics for digitization, and LEDs for detector calibration (Abbasi et al. 2009, 2010). DOMs are attached to 86 cables (called *strings*) in groups of 60 with vertical spacing of 17 m; the mean distance between neighboring strings is ~ 125 m. The eight innermost strings form a denser sub-array called *DeepCore* which targets lower energies (Abbasi et al. 2012). The South Pole IceCube Neutrino Observatory includes the surface array called *IceTop* (Abbasi et al. 2013) that detects and reconstructs air showers above 300 TeV using 82 ice tanks. In the analysis presented here, IceTop’s capabilities are used to veto cosmic-ray-induced backgrounds.

2.1. Neutrino detection channels

Three event topologies are taken into account when considering neutrino interactions in IceCube. *Tracks* are induced by muons traversing the detector. Below 700 GeV, muons lose energy mainly due to ionization; above 700 GeV, stochastic energy losses due to radiative emission become the dominant component. At TeV energies, muons travel long distances, larger than several kilometers in the Antarctic ice (Chirkin & Rhode 2004). Light is constantly emitted along the track. The resulting long lever arm gives a good reconstruction performance with median angular resolution $\Delta\Psi < 1^\circ$.¹ Moreover, the event rate greatly increases because neutrinos can interact far outside the detector prior to the detection of the secondary muon with IceCube. Charged-current interactions of electron or tau neutrinos, as well as neutral current interactions of any neutrino type, produce *shower-* or *cascade-*like events. These types of interactions produce almost spherically symmetric light emission, giving a median angular resolution of $\sim 10^\circ$. Another topology is induced by very high energy charged-current $\nu_\tau + \bar{\nu}_\tau$ interactions with the tau lepton decaying to hadrons after traveling a distance $\sim 50 \text{ m PeV}^{-1} \times E_\tau$, resulting in two cascades separated

¹ Evaluated using dedicated Monte Carlo simulation, verified using experimental data of cosmic-ray shadowing of the Moon (Aartsen et al. 2014d).

distinctly. Such a *double-bang* has not been observed so far (Aartsen et al. 2016c).

Track-like events are more suited than cascades to search for very localized (point-like), faint sources using high statistics and good angular resolution. Such events originate primarily in neutrino charged-current interactions of muon (anti)neutrinos with nucleons, but also in similar interactions of tau neutrinos with the tau lepton decaying to muons and neutrinos, or interactions of electron antineutrinos with electrons by resonant s -channel W^- exchange (Glashow Resonance; Glashow 1960). The energy of a muon track is restricted to the fraction of the track visible in the detector, limiting the energy estimation to the *deposited energy* in the detector or the muon energy upon entering the detector. Independent of the details of neutrino production, the incident flux of astrophysical neutrinos at Earth will consist of an approximately equally shared flavor ratio for all neutrinos due to very long baseline neutrino oscillations (Athar et al. 2006). In the following, only the component of tracks created in muon neutrino interactions is taken into account as the signal of astrophysical neutrinos, likewise to Aartsen et al. (2014e) and Adrian-Martinez et al. (2014). The impact of track events originating from other neutrino interactions is discussed in Section 3.5.

Construction of IceCube finished in December 2010 after 6 yr of deployment. During construction, partial configurations of the detector were successfully taking data, commonly indicated by ICXY, with XY denoting the number of active strings. The first three years of the event sample are in partial detector configurations IC40, IC59, and IC79. The previously published analysis (Aartsen et al. 2014e) included the first year of data-taking with the completed detector IC86 and 3 yr in partial configuration (Abbasi et al. 2011; Aartsen et al. 2013c). These samples focus on through-going track-like events, yielding high statistics over broad energy ranges. In the southern sky, an additional selection of starting tracks, which has a greatly reduced background rate, is performed access lower energies (Aartsen et al. 2016a). This uses completely independent events to the ones selected in the aforementioned through-going track channel.

Table 1. IceCube samples used in this analysis. For each sample, characteristic features are quoted, separated in the two halves of the sky. Previously published results used four years of data: IC40 (Abbasi et al. 2011), IC59+IC79 (Aartsen et al. 2013c), and the first year of IC86 (Aartsen et al. 2014e). Data taken in the seasons from 2012 to 2015 are added in this paper. The separation in the two regions is done at a declination of $\delta = -5^\circ$. In addition, starting tracks are used in a separate sample (Aartsen et al. 2016a) with two additional years of available data.

Sample	Livetime (days)	atm. ν (day^{-1})	Up-going events	Down-going events
IC40	376	40	16323	20577
IC59	348	120	48105	58906
IC79	316	180	54823	38310
IC86	333	210	67938	68302
2012-2015	1058	220	235602	102983
Σ	2431	\dots	422791	289078
starting tracks	1715	< 0.03	0	961

The details of all samples are listed in Tab. 1, including the exposure time and sample size. In this work, the detector livetime is increased by adding data from June 2012 to June 2015 to the analysis, thus increasing the livetime by 1058 days, to a total of 2431 days. The sample of starting tracks (Aartsen et al. 2016a) has an increased livetime of 2 yr to a total of 1715 days that coincide with the five most recent years of the through-going track event sample, starting with IC79.

In the following, the modeling of signal and background for both the through-going and starting track samples is described. The event selection is briefly discussed and the performance of the event sample highlighted. For more detailed information, refer to Abbasi et al. (2011), Aartsen et al. (2013c), and Aartsen et al. (2014e) for through-going muons and Aartsen et al. (2016a) for starting tracks.

In this work, muon tracks induced by astrophysical neutrino interactions are the main signal category in the search for point-like sources of neutrinos. Detailed Monte Carlo simulation is used to evaluate the response of IceCube to such events and distinguish them from atmospheric backgrounds. The median angular resolution $\Delta\Psi$ and the event rate expectation

$$\dot{N}_\nu = \int d\Omega \int_0^\infty dE A_{\text{eff}}(E, \Omega) \times F_\nu(E_\nu, \Omega) \quad (1)$$

given by the detector *effective area* A_{eff} and the incident neutrino flux F_ν can be derived from the simulation (for more information about the details of the Monte Carlo




Figure 1. IceCube effective area as defined in Eq. 1 versus neutrino energy for a flux of $\nu_\mu + \bar{\nu}_\mu$ calculated using simulation of neutrino events for the selection of IC86 (seasons 2012–2015) described in Section 2.2. The effective area for through-going muons is averaged over the solid angle in the declination range (δ) indicated in the legend. Additionally, the effective area for starting tracks in the southern sky ($\delta < -5^\circ$) is shown in black (cf. Section 2.3).

simulation used here, see Aartsen et al. (2016b)).

2.2. Through-going tracks

The main background regarding neutrino searches in IceCube consists of atmospheric muons that are created in extensive air showers and reach IceCube at a depth of ~ 2 km. Almost all events triggering the detector at a rate of ~ 2.8 kHz belong to this component. Similar to Aartsen et al. (2014e), the selection is split into two regions divided at the horizon (declination $\delta = -5^\circ$) due to different background characteristics, as explained in the following.

In the northern sky (up-going region, $\delta \geq -5^\circ$), the main background consists of atmospheric muon events that are mis-reconstructed as up-going; truly up-going muons can only originate from prior neutrino interactions, since all other muons are shielded by the Earth. Hence, the atmospheric muon background is rejected by identifying poorly reconstructed events. In order to achieve this, multivariate selection techniques (boosted decision tree, BDT) are used to discriminate well-reconstructed tracks from neutrino interactions against mis-reconstructed background. The variables used in the BDT are connected to the event quality and a clear track-like topology. Similar to Aartsen et al. (2014e), BDTs are trained for two signal energy spectra of E^{-2} and $E^{-2.7}$ and the cut on the linear combination of BDT scores is optimized to yield the best sensitivity and discovery potential over a wide range of energies with the final cut on BDT output. These spectra are chosen to be sensitive to both hard energy spectra as well as soft or cutoff spectra. A neutrino-dominated sample is obtained in the northern sky. The remaining background




Figure 2. IceCube median angular resolution versus neutrino energy for $\nu_\mu + \bar{\nu}_\mu$ calculated from Monte Carlo simulation for the IC86 sample described in Section 2.2. Through-going tracks (solid black) are shown together with starting tracks (dashed black; see Section 2.3). Moreover, the median kinematic angle of the secondary muon in CC neutrino interactions is shown (dotted black line).

consists of atmospheric neutrinos produced in the northern sky. These neutrinos are an irreducible background, but follow a softer energy spectrum ($\sim E^{-3.7}$ – $E^{-4.0}$ or $\sim E^{-2.7}$ – $E^{-3.0}$ for conventional or prompt neutrinos, respectively) than the expected signal ($\sim E^{-2}$ consistent with diffuse muon neutrino signal at 250 TeV and above; see Aartsen et al. (2016b)). Figure 1 shows the effective area calculated using muon neutrino Monte Carlo simulation of the final event sample for different declination regions. In the northern sky (red, blue), a low energy threshold is achieved, while for near-vertically up-going events absorption becomes an important effect above 100 TeV. Figure 2 shows the median angular resolution of the track reconstruction (solid) with respect to the primary neutrino direction against neutrino energy. Above TeV energies, the kinematic angle (dotted) of muon and parent neutrino becomes negligible and the angular resolution is below 1° . The energy resolution of the track’s energy proxy is $\sim 30\%$ in $\log_{10} E$ (Aartsen et al. 2014a). This is only a lower limit on the energy of a muon that enters the detector from outside and loses energy prior to detection, as well as on energy of the primary neutrino that produced the muon in an interaction with a nucleus. For more information about the neutrino and muon energy estimation, refer to Aartsen et al. (2015c, 2016b).

In the southern sky (down-going, $\delta < -5^\circ$), the picture changes because of large backgrounds of well-reconstructed down-going atmospheric muons. Moreover, muons are produced at high multiplicity in cosmic-ray showers, resulting in bundles of muons; such bundles produce large amounts of light in the detector, thus imitating the signature of a single muon of much higher

energy. Similar to the northern sky, BDTs are used to select only the best-reconstructed events at the highest energies. Following the development of [Aartsen et al. \(2014e\)](#), in addition to event quality and track topology parameters, four variables are used to further discriminate atmospheric muon bundles from single muons. These variables use the deposited energy along the track, as well as the light-arrival time of photons at the DOMs. The Cerenkov light yield for high-energy muons is dominated by stochastic cascades from energy losses along the track superimposed to the Cerenkov cone of the muon track. Bundles of muons at lower energies show less frequent losses by stochastic cascades. The result is a smoother light yield along the track. Furthermore, muon bundles consist of a superposition of many Cherenkov-cones, resulting in many photons arriving earlier than under the assumption of one single Cherenkov cone. For a signal-spectrum E^{-2} , a BDT is trained for final selection of events. The large backgrounds require harsh cuts to reduce their rate significantly, resulting in an effective selection of only very high energy events, as shown in [Fig. 1](#) (yellow and purple). It is evident that the energy threshold in the down-going region increases to ~ 100 TeV and even further for more vertically down-going events. The IceTop surface array is used as an active veto against coincident air-shower events for vertically down-going events. For high energies, this vetoes 90% of the events for vertically down-going events and less for inclined events, with random coincidences in less than 0.1% of the cases ([Aartsen et al. 2013c](#)); compare [Fig. 3](#). The final event rate in the southern sky is optimized to yield the best sensitivity and discovery potential for an E^{-2} spectrum.

The distribution of the cosine zenith $\cos\theta$ (equivalent to the negative sine of declination $-\sin\delta$ in equatorial coordinates) is shown in [Fig. 3](#). The event rate for experimental data of all 3 yr is compared to the expectations of muon neutrinos and atmospheric muons estimated from Monte Carlo simulation. In the northern sky ($\theta > 85^\circ$), the sample is dominated by atmospheric neutrinos produced in the decays of kaons and pions in cosmic-ray air showers ([Honda et al. 2007](#)) and is well described by Monte Carlo simulation. In the southern sky, the neutrino event rate reduces drastically due to the higher energy threshold mentioned before. Instead, atmospheric muons are the dominant component. Astrophysical neutrinos with hard energy spectra (as, e.g. shown with E^{-2}) do not suffer a rate loss as severe as for the soft energy spectrum of atmospheric neutrinos. For very vertically down-going events, the IceTop surface array vetoes atmospheric muon and neutrino events because of their coincident air shower. In this region, the observed event rate is kept constant with the rest of the southern sky using looser cuts on the BDT score, which




Figure 3. Zenith ($\cos\theta$) or declination ($-\sin\delta$) distribution of the through-going track sample after event selection (2012–2015 data). Values of -1 correspond to vertically up-going events. Shown is the experimental data (black), compared to the atmospheric $\nu_\mu + \bar{\nu}_\mu$ expectation of conventional atmospheric (solid gray) and astrophysical neutrinos from Monte Carlo simulation. For simulated atmospheric muons, the plot shows the distribution without the IceTop veto applied.

allows more neutrinos to be detected especially at the lower energy end. Hence, the corresponding event rate in [Fig. 3](#) does not decrease above $\cos\theta > 0.7$.

2.3. Starting tracks

In the southern sky, the large background of atmospheric muons reduces the efficiency to select through-going tracks induced by neutrinos below the PeV regime. A very large fraction of the aforementioned background can be rejected by imposing an active veto at the detector boundary, as, for example, used in [Aartsen et al. \(2013a\)](#). This reduces the detector volume to a smaller fraction of the instrumented volume sacrificing statistics for signal purity. Furthermore, the more clearly an event is identified as a starting track, the more probable it is to be an astrophysical rather than atmospheric background. Down-going atmospheric neutrino events at high energy are likely to be accompanied by muons produced in the same cosmic-ray shower that triggers the veto and reduces the atmospheric neutrino background ([Schönert et al. 2009](#); [Gaisser et al. 2014](#)). In analyses using veto techniques ([Aartsen et al. 2013a, 2015b](#)), the selection is usually more efficient for cascade-like events than tracks, and high astrophysical neutrino purity demands neglecting energies below 60 TeV where backgrounds are more abundant. In searches for point-like sources of astrophysical neutrinos, track-like events are of great importance given their good angular resolution compared to cascade-like events. Furthermore, the purity demands are lower since the signal of a point-like source is restricted to a small por-




Figure 4. Probability distribution of starting distance versus energy proxy (logarithmic, in arbitrary units, a.u.) for both atmospheric background (blue) and neutrinos (red) with E^{-2} spectrum. Different contours depict regions of 20% coverage each. The gray shaded areas shows a region of no coverage resulting from a cut in the selection of Aartsen et al. (2016a).

tion of the sky, hence reducing the background significantly. Consequently, the minimum required total charge deposited in the PMTs of the IceCube detector by an event is lowered to 1500 p.e. compared to 6000 p.e. (Aartsen et al. 2013a), resulting in a higher signal efficiency at lower energies. In addition, only down-going tracks are used, and cuts are imposed that select well-reconstructed track-like events (Aartsen et al. 2016a). For $\nu_\mu + \bar{\nu}_\mu$ events at energies smaller than 200 TeV, the effective area of the analysis is bigger than for vertically through-going tracks ($\delta < -30^\circ$; Fig. 1). For energies up to 1 PeV, the effective area is smaller, but a higher purity is achieved. The angular resolution for starting tracks is shown in Fig. 2 (dashed) and is $\sim 1^\circ$ in the interesting energy region; the reconstruction is worse than for through-going events (solid), due to a smaller lever arm for tracks starting within the fiducial volume of the detector.

In 5 yr, 961 events were recorded in the southern sky starting track sample. The overlap of events in the starting track sample and the through-going events is very small, and overlapping events are removed from the through-going sample because of its higher background rate.

3. METHODS

In order to look for clustering in the sky, the analysis uses an unbinned likelihood maximization, similar to the previous analyses (Aartsen et al. 2014e). The unbinned likelihood is defined as

$$\mathcal{L}(n_S, \gamma) = \prod_i \left(\frac{n_S}{N} \mathcal{S}(|\mathbf{x}_S - \mathbf{x}_i|, E_i; \gamma) + \left(1 - \frac{n_S}{N}\right) \mathcal{B}(\sin \delta_i, E_i) \right) \quad (2)$$

using multiple observables that are introduced in the following. The signal hypothesis used in this work assumes time-integrated emission of neutrinos. Hence, the signature reduces to spatial clustering modeled with a two dimensional Gaussian $\exp(-|\mathbf{x}_S - \mathbf{x}_i|^2 / 2\sigma_i^2) / (2\pi\sigma_i^2)$ using the reconstruction uncertainty σ_i estimated on an event-by-event basis (Neuhöffer 2006; Abbasi et al. 2011). The probability distribution function for the spatial distribution of background is estimated using experimental data and depends only on the event’s declination δ_i , the probability in right ascension is distributed uniformly $1/2\pi$. This yields $\mathcal{P}_B(\sin \delta_i) / 2\pi$ for the spatial probability of background.

In addition, energy information is used to distinguish background with soft spectra ($E^{-3.7}$) from signal with harder spectra of typically E^{-2} . Hence, for each event, probability distributions $\mathcal{E}_{S/B}(E_i)$ for signal and background are evaluated using the event’s energy proxy E_i . For signal, an unbroken power-law with variable index γ , $d\phi/dE_\nu \propto E^{-\gamma}$ is used, the background is estimated from experimental data. The estimation is done declination-dependent, because of the energy-dependence of the effective area (Figure 1). This yields the final modeling of probabilities for signal and background

$$\begin{aligned} \mathcal{S} &= \frac{1}{2\pi\sigma_i^2} e^{-\frac{|\mathbf{x}_S - \mathbf{x}_i|^2}{2\sigma_i^2}} \times \mathcal{E}_S(E_i, \sin \delta_i; \gamma) \\ \mathcal{B} &= \frac{\mathcal{P}_B(\sin \delta_i)}{2\pi} \times \mathcal{E}_B(E_i, \sin \delta_i) \end{aligned} \quad (3)$$

entering the unbinned likelihood calculation in Eq. 2. Two parameters are fit in the likelihood, the number of source events $n_S \geq 0$ and the source spectral index $\gamma \in [1, 4]$.

In contrast to through-going tracks (Section 2.2), no declination dependence of the energy distribution is observed for starting tracks (Section 2.3). This is due to two reasons: the sample only uses down-going events so Earth absorption does not occur, and a uniform charge threshold over all declinations is applied, yielding a uniform effective area (Aartsen et al. 2016a). Meanwhile, starting tracks carry more information than direction and energy; for starting tracks, the vertex of a neutrino interaction can be reconstructed from the first visible light. A background of atmospheric muons can sneak past the veto by not emitting enough light prior to detection. Nevertheless, the higher the energy of the reconstructed track gets, the more likely it is for the vertex to be reconstructed close to the detector boundary due to constant light emission along the track, as shown in Fig. 4. A clear anti-correlation of event en-




Figure 5. Discovery potential (5σ) for this analysis in different bins of neutrino energy E_ν with half-decade width. Within this energy range, an E_ν^{-2} spectrum is used. Three different declinations are shown: Up-going (red, $\delta = 60^\circ$), horizontal (blue, $\delta = 0^\circ$), and down-going (yellow, $\delta = -60^\circ$) events. For down-going events, the dashed line shows the discovery potential *not using* the starting track sample described in Section 2.3.

energy and *starting distance*² is observed for background events (blue). Truly starting signal neutrinos (red) do not show this correlation because the entering neutrino does not emit light. Consequently, the starting distance d_i can be used in addition to the event energy, to disentangle signal and background, modifying the energy likelihood $\mathcal{E}(E_i) \rightarrow \mathcal{E}(E_i, d_i)$, resulting in an additional discrimination power at lower energies.

As in the previous analysis (Aartsen et al. 2014e), the different samples listed in Tab. 1 consist of different detector configurations including partial detector configurations, plus samples using only starting tracks. The total likelihood of all combined samples is the product of all individual likelihoods, or the sum of the logarithms, $\log \mathcal{L}(n_S, \gamma) = \sum_j \log \mathcal{L}(n_S^j, \gamma)$ for all samples j . In the scenario of steady emission, the total number of signal events n_S is split proportionally among the samples given their exposure time and expected signal statistics derived from the effective area (Figure 1 and Eq. 1) and the value of the spectral index fitting parameter γ :

$$n_S^j = n_S \times \frac{\int_0^\infty dE A_{\text{eff}}^j(E, \sin \delta) E^{-\gamma}}{\sum_i \int_0^\infty dE A_{\text{eff}}^i(E, \sin \delta) E^{-\gamma}} \quad (4)$$

Thus, the unbinned likelihood in Eq. 2 is maximized using two parameters only for all samples, that is the number of signal-like events n_S and the spectral in-

dex γ . The null hypothesis is the observation of no signal-like events $n_S = 0$ and defines the test statistic of best-fitting hypothesis ($\hat{n}_S, \hat{\gamma}$) over null hypothesis, $\mathcal{TS} = 2 \log(\mathcal{L}(\hat{n}_S, \hat{\gamma}) / \mathcal{L}(n_S = 0))$. In the maximization of the test statistic, only over-fluctuations $n_S \geq 0$ are taken into account; negative n_S are not part of the physics scenario of neutrino sources (Braun et al. 2008) and not considered here. Thus, the test-statistic of the null hypothesis is expected to split in two fractions, one bound at $n_S = 0$ and over-fluctuations $n_S > 0$. The latter are distributed according to $\chi_{n_{\text{dof}}}^2$ -statistics with $n_{\text{dof}} \sim 1.5$, less than the number of free parameters (2). This is due to n_S and γ being partly degenerate; moreover γ is only defined for $n_S > 0$ as can be seen in Eq. 2. The fraction η of over-fluctuations ranges from 50% to 30% in the northern and southern sky, respectively. From the estimation of the test statistic distribution, the p-value $p = \eta \times \int_{\mathcal{TS}}^\infty dX \chi^2(X | n_{\text{dof}})$ of an observation being consistent with background can be calculated. The p-value will mostly be quoted as $-\log_{10} p$ in the following.

3.1. Neutrino point source sensitivity

Figure 5 shows the 5σ discovery-potential of the unbinned likelihood analysis versus neutrino energy for point sources at various declinations.³ A $\nu_\mu + \bar{\nu}_\mu$ neutrino signal with half-decade width is used for signal injection, using an E^{-2} spectrum within the energy range indicated by the step function. The discovery potential shows a strong variation with declination. In the up-going region ($\delta \geq -5^\circ$), atmospheric muon background is efficiently removed and a large effective area with good angular resolution is achieved above TeV energies, compare Fig. 1 and Fig. 2. This yields a discovery potential reaching from TeV to EeV energies at the horizon ($\delta = 0^\circ$, blue). For vertically up-going events ($\delta = 60^\circ$, red), neutrinos at energies above 100 TeV begin to be absorbed in the Earth, hence reducing the discovery potential compared to the horizon.

In the down-going region (southern sky, $\delta < -5^\circ$), large backgrounds of atmospheric muons result in a higher energy threshold of ~ 100 TeV. Moreover, muon bundles imitate single muons at very high energies resulting in a high energy background. This diminishes the performance compared to the northern sky. At $\delta = -60^\circ$ (yellow), the discovery potential is most effective above energies of 100 TeV; in fact, in between 100 TeV and PeV energies, starting tracks described in Section 2.3 dominate the sensitivity compared with

² Distance along the track pointed back from the reconstructed vertex to the entry point in the detector, see Aartsen et al. (2016a) for more information.

³ The discovery potential is defined as a *false-positive* rate of 5σ or 2.87×10^{-7} with *false-negative* of 50%. The sensitivity is defined as a *false-positive* rate of 50% with *false-negative* of 10%.

through-going muons (Section 2.2). Even though the effective area is of the same order in this energy regime, starting tracks have significantly less background and thus a $\sim 170\times$ higher purity ($0.09 \text{ d}^{-1} \text{ sr}^{-1}$ background events for starting tracks compared to $15.5 \text{ d}^{-1} \text{ sr}^{-1}$ for through-going tracks). Including starting tracks gives a factor of ~ 8 improvement in discovery potential at 100 TeV compared to only using through-going events (dashed yellow in Fig. 5). In the southern sky, similar searches of Adrian-Martinez et al. (2014) test much lower energies, resulting in complementary results combined in Adrian-Martinez et al. (2015).

3.2. Full-sky search

To find the most significant clustering in the sky, the unbinned likelihood maximization is performed on the entire sky. This is done iteratively using a grid with isotropically spaced points (Gorski et al. 2005) finer than the typical event reconstruction uncertainty that enters the likelihood description in Eq. 3.

Thus, for any point in the sky, the best fitting \hat{n}_S , $\hat{\gamma}$ and the test statistic \mathcal{TS} are obtained. The direction with the smallest p-value defines a *hot-spot* showing the biggest deviation from background expectation. This is done for northern and southern sky separately, as they differ in atmospheric backgrounds and energy reach. Thus, two positions in the sky will be reported in the full-sky scan. The significance is trial corrected accounting for the chance of background fluctuations occurring at any position in the sky. The probability to observe no pre-trial $\log_{10} p$ that is smaller than the one at the *hot-spot* for N independent trials is given by

$$d\mathcal{P} = N (1 - p)^{N-1} dp. \quad (5)$$

For both northern and southern sky, the effective number of independent trials N in the sky is fitted to $\sim 190,000$ by repeating the analysis on scrambled data maps. Regions close to the poles (5°) are excluded from the scan because no large off-source regions are available for scrambling. Accounting for the trial factor a pre-trial significance of 5.67σ (p-value 7.13×10^{-9}) is needed for a *hot-spot* to be detected at 3σ significance in the scan of the full sky.

3.3. Hotspot population analysis

The large trial factor of the full-sky scan requires very strong sources that overcome the trial factor. Thus, in addition to looking at only the most significant spot in the sky, the entire sky can be tested for an accumulation of multiple spots at intermediate significance that exceeds the number expected by background.

From the scan of the full sky, the positions of all the local maxima exceeding $-\log_{10} p \geq 3$ are selected. The number of spots observed above a thresh-

old of $-\log_{10} p_{\min}$ is compared to background expectation from repeating the analysis on scrambled data maps. The threshold value is optimized to give the most significant excess over mean background expectation $\lambda(-\log_{10} p_{\min})$ with Poisson statistics. The p-value of the observation of at least n spots, given this expectation, is

$$\mathcal{P} = e^{-\lambda} \sum_{m=n}^{\infty} \frac{\lambda^m}{m!} \quad (6)$$

and defines the test statistic of this test. Due to the optimization of the threshold $-\log_{10} p_{\min}$ to minimize \mathcal{P} , the final result is trial corrected using scrambled experimental data. This is done separately for northern and southern sky. In addition, parts of the sky coinciding with the Galactic plane $\pm 15^\circ$ are analyzed as well for a Galactic contribution.

3.4. Source list searches

In addition to the previously explained searches that did not make any prior assumptions regarding directions in the sky, sources of high-energy γ -rays can be used to search for neutrino emission. Thus, the trial factor of the unbiased full-sky scan can be effectively reduced by probing 74 promising sources selected *a-priori*. The sources are collected in two lists as used in Aartsen et al. (2014e) and Adrian-Martinez et al. (2014); Aartsen et al. (2016a), respectively. The first one (44 sources) probes mainly the northern sky (34 candidates) and the largest fraction of sources are extragalactic objects. The list can be found in Tab. 2. The second list (Table 3) of 30 candidates focuses on the southern sky, especially Galactic sources.

The most significant source in each list will be trial corrected given the number of sources. For the first list, this procedure will be done separately for northern and southern sky. Hence, three post-trial p-values are reported.

3.5. Systematic uncertainties

All analyses described in the previous sections are robust against systematic uncertainties. Background is estimated using experimental data that is scrambled in right ascension and does not require dedicated Monte Carlo simulation. For the calculation of neutrino fluxes however, Monte Carlo simulation is needed to insert neutrino sources into background maps. The resulting neutrino fluxes are affected by systematic uncertainties. Especially, the evaluation of the effective area and reconstruction performance is affected by systematic effects in Monte Carlo, and thus the sensitivity to neutrino fluxes (Equation 1). Main systematic uncertainties include the optical properties (scattering, absorption) of the South Pole ice (Aartsen et al.

2013b), the optical efficiency of Cherenkov light production yield and detection in the DOMs (Abbasi et al. 2010), and different photo-nuclear interaction models (Bugaev & Shlepin 2003a,b; Abramowicz et al. 1991; Abramowicz & Levy 1997). All systematic effects are propagated through the entire likelihood analysis described in Section 3 to obtain the uncertainties on the fluxes using $d\phi/dE_\nu \propto E^{-2}$ spectra. The biggest impact on the fluxes comes from varying the optical efficiency by $\pm 10\%$, resulting in a flux uncertainty of 7.5%. Increasing the absorption or scattering of photons in ice by 10% affects the flux by 5.6%. Uncertainties in the photo-nuclear cross-sections (Bugaev & Shlepin 2003a,b) result in an flux uncertainty of similar size with 5.9%. Adding these values in quadrature yields a total systematic uncertainty of 11% on $\nu_\mu + \bar{\nu}_\mu$ fluxes quoted in the following.

For all locations tested, only the maximal likelihood values of \hat{n}_S and $\hat{\gamma}$ are reported. Because of small event statistics at the position of the likelihood maximization and limited energy resolution of the neutrino energy (compare Section 2.2), uncertainties on the spectral index are of the order ± 1 and reduce to ± 0.5 for values of n_S of ~ 15 and ~ 50 , respectively (Braun et al. 2008). Hence, the impact of systematic uncertainties in the energy reconstruction is small compared to the statistical limitations.

Albeit not a systematic uncertainty per se, so far only fluxes of $\nu_\mu + \bar{\nu}_\mu$ were considered. This is a conservative estimate, because track-like events can also originate in other cases that are discussed in the following. Firstly, tau-leptons created in charged-current $\nu_\tau + \bar{\nu}_\tau$ interactions decay into muons with 17% branching ratio (Jeong & Reno 2010; Olive et al. 2014), resulting in a muon track with lower energy due to the three-body decay $\tau \rightarrow \mu\nu_\mu\nu_\tau$. This decay is important for up-going events, because secondary neutrinos are produced in τ -neutrino regeneration during propagation. Secondly, interactions of $\bar{\nu}_e + e^- \rightarrow W^-$ at the Glashow-resonance (Glashow 1960) at 6.3 PeV produce tracks ($\bar{\nu}_e + e^- \rightarrow \bar{\nu}_\mu + \mu^-$) at 10.6% branching ratio (Olive et al. 2014). Lastly, at the highest energies above PeV, τ -neutrino induced double bangs are well-reconstructable and further increase the number of τ -flavored events in the sample. Accounting for these fluxes assuming an equal flavor ratio at Earth reduces the per-flavor flux necessary for detection by 5% assuming an unbroken E^{-2} spectrum. For harder spectra, the sensitivity gain due to regeneration effects in the northern sky becomes stronger. For example, a spectrum of $d\phi/dE_\nu \propto E^{-1}$ has an 30% improved sensitivity compared to only considering muon neutrinos. This greatly increases the sensitivity with respect to models that predict very hard neutrino energy spectra peaking above




Figure 6. All-sky result of the unbinned likelihood maximization shown in equatorial coordinates (J2000). Shown is the negative logarithm of the pre-trial p-value, $-\log_{10} p$, assuming no clustering as null-hypothesis. The Galactic Plane is shown as black line.

PeV energies (Petropoulou et al. 2015; Reimer 2015).

4. RESULTS AND IMPLICATIONS

In the unbinned likelihood analysis using seven years of IceCube livetime, no significant excess of astrophysical neutrino sources was found. In the following, the results of the three tests introduced in the previous sections are discussed and 90% upper-limits on neutrino source fluxes are calculated. Finally, implications with respect to neutrino models of γ -ray sources and the observed diffuse neutrino flux are presented.

4.1. All sky scan

Figure 6 depicts the pre-trial p-value $-\log_{10} p$ of all points in the sky in equatorial coordinates (J2000) with respect to the null-hypothesis of no observed clustering.

In the northern sky, the most significant position was at $\alpha = 32.2^\circ$, $\delta = 62.1^\circ$ at an accuracy of 0.35° (0.5°) for 1σ (90%) contours using Wilks' theorem with two degrees of freedom. The best fit parameters at the location are $\hat{n}_S = 32.6$ and $\hat{\gamma} = 2.8$, yielding a pre-trial p-value of 1.82×10^{-6} . Looking at each of the combined seasons individually reveals that for each season clustering is observed, providing no indication of time-dependence that could suggest additional evidence for an astrophysical origin.

In the southern sky, the most significant point is at $\alpha = 174.6^\circ$, $\delta = -39.3^\circ$. The best fit point is at $\hat{n}_S = 15.4$, with spectral index $\hat{\gamma} = 2.9$. The uncertainty of the location amounts to 0.22° (0.32°) for 1σ (90%). The pre-trial p-value is 0.93×10^{-6} ; most of the significance at this location is shared by the newly added data of through-going and starting tracks. Indeed, one starting track is within 0.9° distance to the location which is within 1σ of its reconstruction uncertainty.




Figure 7. Trial correction of the most significant spots in the sky that were observed in the seven year search. Solid vertical lines indicate the pre-trial p-value of the most significant spots in each half of the sky; crosses show the distribution of spots similarly obtained in scrambled data trials. The trials are modeled by an analytic parameterization of the trial correction (Equation 5, black dashed line) that corresponds to 1.9×10^5 independent trials per half of the sky.

Due to the large number of tested locations in the sky, the two most significant locations in the sky have to be trial corrected with the trial correction in Eq. 5 that is estimated by repeating the full-sky scan on scrambled data trials, as shown in Fig. 7. This yields post-trial p-values of 29%, 17% for northern and southern sky, respectively. Hence, the full-sky results are in agreement with a pure background assumption, and no significant clustering is observed. For an unbroken E^{-2} power-law spectrum, the 90% upper-limits of the two most significant positions are $E_\nu^2 d\phi/dE_\nu = 4.49 \times 10^{-12}$ TeV cm $^{-2}$ s $^{-1}$ in the northern sky, and $E_\nu^2 d\phi/dE_\nu = 2.92 \times 10^{-11}$ TeV cm $^{-2}$ s $^{-1}$ in the southern sky. For softer spectra of E^{-3} , the 90% upper-limits yield $E^3 d\phi/dE_\nu = 5.08 \times 10^{-11}$ TeV 2 cm $^{-2}$ s $^{-1}$ and $E^3 d\phi/dE_\nu = 1.29 \times 10^{-8}$ TeV 2 cm $^{-2}$ s $^{-1}$ for the northern and southern spot, respectively. In Fig. 8, the solid blue line indicates the 90% upper-limit established by the hottest spot results. A neutrino source at any declination δ that would emit a steady flux higher than this curve, would be detected 90% of the time as having a greater significance than that actually observed for the hottest spots found in the analysis (whose 90% upper-limits are highlighted as stars on the blue line).

Besides the results of the full-sky scan, there are two neutrino events detected with IceCube that are worth commenting on here. The first one is the highest energetic neutrino event detected (4.5 ± 1.2 PeV) so far with IceCube (Schoenen & Rädcl 2015; Aartsen et al. 2016b), a neutrino-induced up-going muon track with very precise angular resolution. This neutrino event is part of the through-going track sample (Section 2.2). At




Figure 8. Discovery potential (5σ , solid red) and sensitivity (dashed red) for a $\nu_\mu + \bar{\nu}_\mu$ unbroken $E_\nu^2 d\phi/dE_\nu$ flux shown against declination δ . The gray line shows the results of (Adrian-Martinez et al. 2014) in the south. Upper limits of source candidates in Tab. 2 and Tab. 3 are depicted by red crosses. The blue line represents the upper limit for the observed most significant spots in each half of the sky for all declinations, the actual declination position of the spots is indicated by a star.

its position ($\alpha = 110^\circ$, $\delta = 11.5^\circ$), no significant clustering is observed (pre-trial 5.2%). A slight excess is indeed observed, but originates from the PeV event alone. The second interesting event is a straight down-going starting track at 430 TeV deposited energy (Aartsen et al. 2015f). Not only does it start inside of the IceCube detector, but the reconstructed track points back to the IceTop surface detector and no atmospheric shower is observed in coincidence with the event. This event is part of the starting track sample (Section 2.3), but no clustering of events apart from the track itself is observed at the location in the sky ($\alpha = 218^\circ$, $\delta = -86^\circ$) and the pre-trial p-value is 0.6%.

4.2. Hotspot population

The search for populations of weak sources in the full-sky in Fig. 6 did not reveal any significant outcome above background expectation. In Fig. 9, the number of spots versus pre-trial p-value $-\log_{10} p_{\min}$ threshold is shown for northern (left) and southern sky (right). The observed number of spots is shown versus background expectation with shaded areas indicating 1σ , 2σ , and 3σ intervals. This is then converted to a local p-value \mathcal{P} according to Eq. 6.

In the northern sky, the most significant excess is observed above a threshold of $-\log_{10} p_{\min} \geq 3.35$ with 72 spots above a scrambled data expectation of 56.7. The local p-value of such an excess is $\mathcal{P} = 2.8\%$ and increases to 25% after trial correction. For the southern sky, 7 spots above an expectation of 2.1 spots at $-\log_{10} p_{\min} \geq 4.66$ are reported. The probability of this




Figure 9. Number of associated spots observed with a minimum pre-trial p-value of $-\log_{10} P_{\min}$ of the unbinned likelihood fit in the northern (left) and southern sky (right), respectively. The top plots show the number of spots (solid black) in comparison to background expectation (dashed black) with confidence intervals of one, two, and three standard deviations shown as shaded blue areas. The points on the far right end of the x-axis correspond to the hottest spots observed in the northern and southern sky. The bottom plots show the local p-value \mathcal{P} (Equation 6) of observing X events given the background mean expectation of Y using Poissonian statistics.




Figure 10. Discovery potential (5σ , solid thin line), sensitivity (dashed line), and 90% upper-limit (solid thick line) for an unbroken E^{-2} spectrum in the population analysis for northern (left) and southern (right) sky, respectively. The flux is shown per source for increasing number of sources distributed uniformly on half of the sky. The black dashed line shows the IceCube-measured astrophysical neutrino flux per source if it was distributed homogeneously among N_{Sources} sources in this half of the sky.

happening by fluctuations of background is 0.62%. After correcting this for trials by scanning in $-\log_{10} P_{\min}$ for the biggest excess, the p-value increases to 8.2%. Restricting the analysis only to regions within $\pm 15^\circ$ of the Galactic Plane the biggest excess is observed at $-\log_{10} P_{\min} \geq 5.68$, corresponding to a single spot, which is also the same spot that was most significant in the northern hemisphere full sky scan. The background expectation is 0.04 giving a local p-value of 3.8% that increases to 26% after trial correction.

Consequently, in both hemispheres and the Galactic Plane, no significant population of sources over background expectation was found. Fig. 10 shows the sensitivity, 5σ discovery potential, and 90% upper-limit for northern (left) and southern sky (right). This is shown against an increasing number of sources, where each source is assumed to be of equal luminosity with an

unbroken E^{-2} spectrum. The flux value is averaged over all declinations of the corresponding half of the sky. By comparing this to a scenario where all of the observed diffuse astrophysical neutrino flux (Aartsen et al. 2014c) $E_\nu^2 d\phi/dE_\nu \sim 10^{-11} \text{ TeV cm}^{-2} \text{ s}^{-1} \text{ sr}^{-1}$ is shared equally among N_{Sources} in the sky, in the northern sky, the analysis result excludes populations of 1000 or fewer equal-strength sources, whereas in the south, the result only excludes populations of 40 or fewer equal-strength sources of the astrophysical flux.

4.3. Source list candidates

In Tab. 2 and Tab. 3, the fit results of the two source lists are quoted. For each source, the fit parameters \hat{n}_S and $\hat{\gamma}$ are quoted. Furthermore, the pre-trial p-value and the mean number of background events B_{1° within a one-degree circle around the source is listed. For each source, 90% upper-limits are calculated using an unbro-

ken E^{-2} spectrum. The upper-limits are shown as red crosses in Fig. 8 at the corresponding declination of the source.

In the first source list, the sources most significant are the blazar 1ES 1959+650 and flat spectrum radio quasar PKS 1406-076 in the northern, southern sky, respectively. At the position of 1ES 1959+650 the pre-trial p-value is 1.8% with best fit-parameters $\hat{n}_S = 15.4$ and $\hat{\gamma} = 3.1$. The resulting 90% upper-limit for an unbroken E^{-2} $\nu_\mu + \bar{\nu}_\mu$ flux of the observed p-value at declination $\delta = 65.15^\circ$ of 1ES 1959+650 yields $E_\nu^2 d\phi/dE_\nu = 2.36 \times 10^{-12} \text{ TeV cm}^{-2} \text{ s}^{-1}$. For PKS 1406-076 at $\delta = -7.87^\circ$, the pre-trial p-value of 5.3% with $\hat{n}_S = 7.3$ and $\hat{\gamma} = 2.6$ results in an 90% upper-limit of $E_\nu^2 d\phi/dE_\nu = 1.65 \times 10^{-12} \text{ TeV cm}^{-2} \text{ s}^{-1}$. For trial correction, the source list is split in a northern and southern part, with the division at $\delta = -5^\circ$. The size of the source lists is then 34 (10) and yields a trial corrected p-value of 54% (37%) for the northern sky (southern sky). Hence, the results of the first source list are in agreement with background expectation.

In the second source list, the most significant source is HESS J1616-508. The fit values $\hat{n}_S = 2.4$ and $\hat{\gamma} = 4.0$ result in a pre-trial p-value of 0.22%. The 90% upper-limit is $E_\nu^2 d\phi/dE_\nu = 1.94 \times 10^{-11} \text{ TeV cm}^{-2} \text{ s}^{-1}$. Trial correction with all 30 sources of the list yields a post-trial p-value of 9.3%. Most of the significance at the position of HESS J1616-508 comes from one starting track only 0.34° away, while no significant clustering of high-energy events is observed in the through-going event samples. Starting tracks access lower energies in the southern sky (cf. Fig. 1, Fig. 5). As explained in Eq. 4, the number of source-like neutrinos n_S is distributed among the different samples according to their signal expectation for a spectral index γ . Consequently, if the clustering is only observed in starting tracks, soft spectral indices at the boundary $\hat{\gamma} = 4$ are preferred as they give more weight to the starting track sample that is most efficient for soft spectra compared to through-going track samples in the southern sky.

In conclusion, both of the two source lists show no significant evidence for clustering of astrophysical neutrinos, and all results are consistent with background.

4.4. Multi-wavelength model constraints

Above, upper limits on neutrino emission from sources were made using unbroken $d\phi/dE_\nu \propto E^{-2}$ fluxes as benchmark. However, more specific estimates for neutrino fluxes can be made using multi-wavelength data of γ -ray sources. In decays of pions, both γ -rays as well as neutrinos are produced (Anchordoqui et al. 2014). Due to long-baseline oscillations, any flavor composition at the source will result in a sizable fraction of muon neutrinos at Earth (Athar et al. 2006). Due to no significant




Figure 11. Differential $\nu_\mu + \bar{\nu}_\mu$ energy spectra versus neutrino energy for the Crab nebula. The figure shows the conversion of the observed gamma-ray flux of the Crab nebula to neutrinos (Kappes et al. 2007) (blue), and a simulation of inelastic p-p scattering at the source (Amato et al. 2003) (red). Thick lines correspond to the 90% upper-limit of this search, thin lines are represent the model. The sensitivity of this analysis is shown as dashed line. 90% upper-limit and sensitivity are shown for the energy interval, where 90% of the events originate that are most signal-like, cf. Fig. 5.

observation of clustering, upper-limits on specific models are set by injecting signal events at the corresponding source declination according to the energy spectrum $E_\nu^2 d\phi/dE_\nu$ given by the model.

The first source considered is the Crab Nebula, a pulsar wind nebula, and the strongest steady TeV γ -ray source in the sky. At a declination of $\delta = 22^\circ$, it is in the region where IceCube covers a wide range of energies efficiently, compare Fig. 5. Two scenarios of neutrino emission from this source are considered. Figure 11 shows the neutrino emission (thin line) with respect to the 90% upper-limit (thick line) of IceCube. At the position of the Crab Nebula, an over-fluctuation with p-value 34% is observed which results in an upper-limit higher than IceCube’s potential sensitivity (dashed). By convoluting the differential discovery potential (Figure 5) at the source position and the model neutrino spectrum, the energy-region, where 90% of the constraining power of IceCube originates, for a specific model is calculated. This is indicated by the energy region where the limits and sensitivities are drawn in Fig. 11 for each of the following models.

Regarding the Crab Nebula, the first model taken into account (red in Fig. 11) is by Amato et al. (2003) and uses inelastic p-p scattering at the source to model the neutrino emission where 60% of the wind luminosity L_{tot} is carried by protons. The model shown is for a Lorentz factor of the wind of $\Gamma = 10^7$ where the energy density peaks at the ~ 500 TeV and assumes a target density of

$$n_t = 10\mu (M_{N_\odot}/R_{\text{pc}}^3) \text{ cm}^{-3} \quad (7)$$

with $\mu = 20$ (shown in Fig. 11) as defined in Eq. 9 in




Figure 12. Sensitivity and 90% upper-limit (dashed and solid lines, respectively) for the proton wind luminosity L_p and target density in Eq. 7 of the Crab Nebula in Amato et al. (2003) for different Lorentz factors Γ . Values of $\mu = 20$ and $\mu = 5$ ($L_p = 60\% L_{\text{tot}}$) are indicated as horizontal lines that correspond to upper limits of the target density and a uniform mass distribution in the nebula, respectively. The dotted line indicates upper-limits from gamma-ray observations compared to $\pi^0 \rightarrow \gamma\gamma$ decays (Amato et al. 2003; de Jager et al. 1996; Aharonian et al. 2000).

Amato et al. (2003). $M_{N\odot}$ and R_{pc} are the mass of the supernova ejecta contained in filamentary structures within the nebula and its radius, respectively. With the increased statistics compared to the previous analysis (Aartsen et al. 2014e), the 90% upper-limit now surpasses the prediction of $\mu = 20$, hence, constraining the target density $\mu^{90\%} < 12$. Upper limits on μ and the proton wind luminosity L_p in units of the total wind luminosity L_{tot} for lower values of the Lorentz wind Γ are shown in Fig. 12. Lower values of Γ shift the neutrino energy spectrum to lower energies into the TeV region where IceCube is most sensitive in the Northern hemisphere. Hence, target densities $\mu < 20$ as mentioned in Atoyan & Aharonian (1996) are partially constrained by IceCube for a proton luminosity fraction of 60%. For $\Gamma = 10^5$, the sensitivity is at the level of $\mu = 5$ corresponding to a uniform mass distribution in the nebula as pointed out by Amato et al. (2003). For Lorentz wind factors $\Gamma < 10^6$, the strongest constraint on the proton luminosity is given by gamma-ray observations via neutral pion decay $\pi^0 \rightarrow \gamma\gamma$ (Amato et al. 2003; de Jager et al. 1996; Aharonian et al. 2000). For $\Gamma \geq 10^6$, neutrino observations give additional information and large values of $\Gamma = 10^7$ are exclusively observable by neutrinos due to the high energy spectrum. Di Palma et al. (2016) revise the predictions for PWNe given new TeV γ -ray observations (Aliu et al. 2008, 2013), predicting ~ 13.09 neutrino events from the Crab nebula per year in the energy range from 1 to 100 TeV. IceCube’s 90% upper-limit at the position of the Crab nebula (see Tab. 2) constrains the number of




Figure 13. Same as Fig. 11 for blazars of type BL Lac modeled in Petropoulou et al. (2015).

neutrinos from the source to not more than 7.0 or 21.2 for an unbroken spectrum of E^{-2} and E^{-3} respectively from a total of seven years of data-taking. This is much lower than the predicted number of neutrino events during the same time range.

The second neutrino spectrum tested for the Crab Nebula is the one by (Kappes et al. 2007) (blue). The neutrino spectrum is connected to γ -rays assuming that both spectra originate from the same pion-component. Thus, the resulting neutrino spectrum is fitted to an exponentially cut-off power-law $d\phi/dE_\nu \propto E^{-2.15} \exp(-\sqrt{E_\nu/1.72 \text{ TeV}})$. Due to the observed over-fluctuation at the position of the Crab Nebula, the 90% upper-limit regarding this model exceeds the benchmark model of 100% pion contribution by a factor of 1.18 and is thus not constraining the amount of hadronic acceleration. Nevertheless, IceCube’s sensitivity (blue dashed line in Fig. 11) is 15% below the model. Hence, with seven years exposure, IceCube is now also sensitive to neutrino fluxes comparable to that of the Crab nebula in gamma-rays, i.e. bright but falling off at relatively low energies of a few TeV.

Another promising category of neutrino emitters are active galactic nuclei (AGN), especially blazars of type BL Lac, as for example highlighted in (Padovani & Resconi 2014). In Glüsenskamp (2015), an analysis of IceCube neutrino events searched for correlations with blazars detected by Fermi-LAT to probe their possible connection to the observed diffuse astrophysical neutrino signal. It was found that such AGN cannot be the dominant contribution to the diffuse neutrino flux over the entire probed energy range, but BL Lacs could possibly explain the high-energy part of the flux (Padovani et al. 2015). In Petropoulou et al. (2015), BL Lacs that were found to be in spatial and energetic agreement with IceCube high-energy starting events were modeled using lepto-hadronic fitting




Figure 14. Same as Fig. 11 for blazars modeled in Reimer (2015).

of multi-wavelength data. Of the six modeled blazars, three are far from being constrained by IceCube due to neutrino absorption in the Earth (1ES 1011+496) or large atmospheric background deep in the southern sky (H2356-309, 1RXS J05435-5532). Figure 13 shows the expected neutrino energy spectra for the three remaining BL Lacs. The 90% upper-limit for 1H1914-194, also in the southern sky, is a factor 3.9 above the prediction. For PG1553+113, the contribution of proton-pion interactions in the model is small (total neutrino flux relative to TeV γ -ray flux $Y_{\nu\gamma} = 0.1$, Eq. 13 in Petropoulou et al. (2015)) resulting in a lower neutrino luminosity and a 90% upper-limit of a factor of three above the prediction.

The third model source considered here is Mrk 421 (blue) and shows a different picture. Mrk 421 is one of the closest blazars with redshift $z \sim 0.03$ and with $\delta = 38.2^\circ$, it is located in the up-going region where IceCube is most sensitive. Moreover, in Petropoulou et al. (2015), the multi-wavelength data observed can be explained with a very high hadronic component ($Y_{\nu\gamma} = 0.7$), thus, realizing a high neutrino luminosity. For such a model, the 90% upper-limit obtained by IceCube is 2/3 of the predicted neutrino flux. Hence, using the results at the position of Mrk 421, the hadronic acceleration cannot be bigger than $Y_{\nu\gamma} \leq 0.47$ assuming a steady emission over the seven years of analyzed data, and thus constrains proton luminosity of the source for steady emission or emission taking into account variability of the source (Petropoulou et al. 2015, 2016).

Other models for active galactic nuclei of type flat spectrum radio quasar or BL Lac were modeled in Reimer (2015). In Fig. 14, 90% upper-limits for 3C273 (red), 3C279 (blue), and 3C454.3 (yellow) are shown. The models in this reference are very characteristic due to their hard spectra extending beyond 10 PeV before cutting off. Hence, IceCube cannot constrain the predic-




Figure 15. Time evolution in discovery potential (5σ) with increasing detector exposure. Different declinations are shown in different colors. The current status of this analysis corresponds to seven years or the year 2015 (indicated by vertical black line). Dashed lines indicate the sensitivity growth assuming a dependence with $1/\sqrt{T}$, dotted with $1/T$.

tion for FSRQ 3C454.3 ($\delta = 16.1^\circ$) because most of the flux is absorbed in the Earth and the 90% upper-limit is a factor of five higher than the prediction. The two other FSRQ on the other hand are close to the horizon where no absorption is present. Therefore, the 90% upper-limit placed by IceCube is lower than the prediction, that is, 13% and 40% for 3C273 and 3C279, respectively. Similar to Mrk 421 in the previous paragraph, assuming that the emission is constant over the livetime of IceCube, this results constrains the amount of hadronic acceleration possible due to a non-observation of neutrinos by IceCube with respect to the model prediction. The other AGN modeled in Reimer (2015) are in the northern sky and due to absorption, the 90% upper-limit by IceCube is at least a factor of 60 above the prediction.

Other models that were tested include galactic source like supernovae remnants (Mandelartz & Becker Tjus 2015) and unidentified TeV sources in the galaxy (Fox et al. 2013) that are mainly in the southern sky which coincides with most of the Galactic Plane. There, IceCube has to cope with large atmospheric muon backgrounds and has reduced sensitivity with high energy-thresholds, compare Fig. 5 and Fig. 8. Consequently, the current 90% upper-limits are at least a factor of five (G40.5-05, Mandelartz & Becker Tjus (2015)) up to more than a hundred (Vela X, Kappes et al. (2007)) above the prediction. At these energies in the southern sky region, neutrino telescopes located in the northern hemisphere can place stronger constraints (Adrian-Martinez et al. 2015; Trovato 2016).

5. CONCLUSIONS

Analyzing the full sky for clustering of astrophysical neutrinos using seven years of through-going muon data

from 2008 to 2015 and five years of starting tracks (2010-2015), we did not find any significant steady point-like emission over background expectation.

An unbiased scan of the full-sky was performed, as well as dedicated searches using promising γ -ray candidates and searches for populations of weak sources. Results are consistent with background and 90% upper-limits on steady neutrino emission are calculated from the observations. In the northern sky ($\delta \geq 5^\circ$), the best limits are set by IceCube and the sensitivity is below $E_\nu^2 d\phi/dE_\nu < 10^{-12} \text{ TeV cm}^{-2} \text{ s}^{-1}$ over a wide declination for the first time, compare Fig. 8. Moreover, the declination dependent flux limit is a factor of ~ 100 (~ 10) below the integrated diffuse astrophysical neutrino flux in the northern (southern) sky. With increased statistics of three additional years compared to the previous analysis, the upper-limits in the southern sky are of the same level as for Adrian-Martinez et al. (2014), however, testing a complementary energy region above PeV neutrino energies. Model-specific 90% upper-limits were calculated for representative cases from the literature, and previously unconstrained neutrino emission scenarios for sources in the northern sky are now disfavored by IceCube. With the newly available data presented here, improved results for other tests are anticipated, for example searches for extended sources or stacking and time-dependent analyses (Aartsen et al. 2014e, 2015e).

With increasing exposure, IceCube continues to improve the sensitivity to steady neutrino fluxes. Figure 15 shows the progression of the 5σ discovery potential with increasing time for three declinations. A continuous gain in performance is observed with time T that is faster than $1/\sqrt{T}$, a scenario assuming statistical limitation by background, and much closer to $1/T$ indicating limitation by signal statistics. This can increase even

further with anticipated improvements in background rejection, angular reconstruction, or detector improvements (Aartsen et al. 2014b).

We acknowledge the support from the following agencies: U.S. National Science Foundation-Office of Polar Programs, U.S. National Science Foundation-Physics Division, University of Wisconsin Alumni Research Foundation, the Grid Laboratory Of Wisconsin (GLOW) grid infrastructure at the University of Wisconsin - Madison, the Open Science Grid (OSG) grid infrastructure; U.S. Department of Energy, and National Energy Research Scientific Computing Center, the Louisiana Optical Network Initiative (LONI) grid computing resources; Natural Sciences and Engineering Research Council of Canada, WestGrid and Compute/Calcul Canada; Swedish Research Council, Swedish Polar Research Secretariat, Swedish National Infrastructure for Computing (SNIC), and Knut and Alice Wallenberg Foundation, Sweden; German Ministry for Education and Research (BMBF), Deutsche Forschungsgemeinschaft (DFG), Helmholtz Alliance for Astroparticle Physics (HAP), Research Department of Plasmas with Complex Interactions (Bochum), Germany; Fund for Scientific Research (FNRS-FWO), FWO Odysseus programme, Flanders Institute to encourage scientific and technological research in industry (IWT), Belgian Federal Science Policy Office (Belspo); University of Oxford, United Kingdom; Marsden Fund, New Zealand; Australian Research Council; Japan Society for Promotion of Science (JSPS); the Swiss National Science Foundation (SNSF), Switzerland; National Research Foundation of Korea (NRF); Villum Fonden, Danish National Research Foundation (DNRF), Denmark

REFERENCES

- Aartsen, M. G., et al. 2013a, *Science*, 342, 1242856
 —. 2013b, *Nucl. Instrum. Meth.*, A711, 73
 —. 2013c, *Astrophys. J.*, 779, 132
 —. 2014a, *JINST*, 9, P03009
 —. 2014b, arXiv:1412.5106
 —. 2014c, *Phys. Rev. Lett.*, 113, 101101
 —. 2014d, *Phys. Rev.*, D89, 102004
 —. 2014e, *Astrophys. J.*, 796, 109
 —. 2015a, *Astrophys. J.*, 809, 98
 —. 2015b, *Phys. Rev.*, D91, 022001
 —. 2015c, *Phys. Rev. Lett.*, 115, 081102
 —. 2015d, *Phys. Rev. Lett.*, 114, 171102
 —. 2015e, *Astrophys. J.*, 807, 46
 —. 2015f, arXiv:1510.05223
 —. 2016a, *Astrophys. J.*, 824, L28
 —. 2016b, arXiv:1607.08006
 —. 2016c, *Phys. Rev.*, D93, 022001
 Abbasi, R., et al. 2009, *Nucl. Instrum. Meth.*, A601, 294
 —. 2010, *Nucl. Instrum. Meth.*, A618, 139
 —. 2011, *Astrophys. J.*, 732, 18
 —. 2012, *Astropart. Phys.*, 35, 615
 —. 2013, *Nucl. Instrum. Meth.*, A700, 188
 Abramowicz, H., Levin, E. M., Levy, A., & Maor, U. 1991, *Phys. Lett.*, B269, 465
 Abramowicz, H., & Levy, A. 1997, arXiv:hep-ph/9712415
 Achterberg, A., et al. 2006, *Astropart. Phys.*, 26, 155
 Adrian-Martinez, S., et al. 2014, *Astrophys. J.*, 786, L5
 —. 2015, arXiv:1511.02149
 Aharonian, F. A., et al. 2000, *Astrophys. J.*, 539, 317
 Aliu, E., et al. 2008, *Science*, 322, 1221
 —. 2013, *Astrophys. J.*, 764, 38
 Amato, E., Guetta, D., & Blasi, P. 2003, *Astron. Astrophys.*, 402, 827
 Anchordoqui, L. A., & Montaruli, T. 2010, *Annual Review of Nuclear and Particle Science*, 60, 129
 Anchordoqui, L. A., et al. 2014, *JHEAp*, 1-2, 1

- Athar, H., Kim, C. S., & Lee, J. 2006, *Mod. Phys. Lett.*, A21, 1049
- Atoyan, A. M., & Aharonian, F. A. 1996, *MNRAS*, 278, 525
- Beatty, J. J., & Westerhoff, S. 2009, *Ann. Rev. Nucl. Part. Sci.*, 59, 319
- Braun, J., Dumm, J., De Palma, F., et al. 2008, *Astropart. Phys.*, 29, 299
- Bugaev, E. V., & Shlepin, Yu. V. 2003a, *Phys. Rev.*, D67, 034027
- . 2003b, *Nucl. Phys. Proc. Suppl.*, 122, 341
- Chirkin, D., & Rhode, W. 2004, arXiv:hep-ph/0407075
- de Jager, O. C., Harding, A. K., Michelson, P. F., et al. 1996, *ApJ*, 457, 253
- Di Palma, I., Guetta, D., & Amato, E. 2016, arXiv:1605.01205
- Fox, D. B., Kashiyama, K., & Mészáros, P. 2013, *Astrophys. J.*, 774, 74
- Gaisser, T. K., Jero, K., Karle, A., & van Santen, J. 2014, *Phys. Rev.*, D90, 023009
- Glashow, S. L. 1960, *Phys. Rev.*, 118, 316
- Glüsenkamp, T. 2015, in 5th Roma International Conference on Astro-Particle physics (RICAP 14) Noto, Sicily, Italy, September 30-October 3, 2014
- Gorski, K. M., Hivon, E., Banday, A. J., et al. 2005, *Astrophys. J.*, 622, 759
- Halzen, F., & Hooper, D. 2002, *Reports on Progress in Physics*, 65, 1025
- Hinton, J. A., & Hofmann, W. 2009, *Ann. Rev. Astron. Astrophys.*, 47, 523
- Honda, M., Kajita, T., Kasahara, K., Midorikawa, S., & Sanuki, T. 2007, *Phys. Rev.*, D75, 043006
- Jeong, Y. S., & Reno, M. H. 2010, *Phys. Rev.*, D82, 033010
- Kappes, A., Hinton, J., Stegmann, C., & Aharonian, F. A. 2007, *Astrophys. J.*, 656, 870, [Erratum: *Astrophys. J.* 661, 1348(2007)]
- Kotera, K., & Olinto, A. V. 2011, *Ann. Rev. Astron. Astrophys.*, 49, 119
- Learned, J. G., & Mannheim, K. 2000, *Annual Review of Nuclear and Particle Science*, 50, 679
- Mandelartz, M., & Becker Tjus, J. 2015, *Astropart. Phys.*, 65, 80
- Murase, K., Guetta, D., & Ahlers, M. 2016, *Phys. Rev. Lett.*, 116, 071101
- Neuhöffer, T. 2006, *Astropart. Phys.*, 25, 220
- Olive, K. A., et al. 2014, *Chin. Phys.*, C38, 090001
- Padovani, P., Petropoulou, M., Giommi, P., & Resconi, E. 2015, *Mon. Not. Roy. Astron. Soc.*, 452, 1877
- Padovani, P., & Resconi, E. 2014, *Mon. Not. Roy. Astron. Soc.*, 443, 474
- Petropoulou, M., Coenders, S., & Dimitrakoudis, S. 2016, *Astropart. Phys.*, 80, 115
- Petropoulou, M., Dimitrakoudis, S., Padovani, P., Mastichiadis, A., & Resconi, E. 2015, *Mon. Not. Roy. Astron. Soc.*, 448, 2412
- Reimer, A. 2015, *Proceedings, 34th International Cosmic Ray Conference (ICRC 2015)*
- Schoenen, S., & Rädcl, L. 2015, *The Astronomer's Telegram*, 7856
- Schönert, S., Gaisser, T. K., Resconi, E., & Schulz, O. 2009, *Phys. Rev.*, D79, 043009
- Senno, N., Murase, K., & Meszaros, P. 2016, *Phys. Rev.*, D93, 083003
- Trovato, A. 2016, *EPJ Web Conf.*, 121, 05011

Table 2. Sources contained in the first source list. In addition to its type, common name, and coordinates (Equatorial (J2000)), the best-fit of signal events n_S and spectral index γ with the pre-trial p-value $-\log_{10} p$ are quoted. For null observations $n_S = 0$, no p-value and spectral index are quoted. A 90% upper-limit for an E^{-2} unbroken power-law is calculated for all sources that showed clustering of neutrinos. The upper limits are shown in Fig. 8, for null-fits the limit equals the sensitivity of this analysis at the corresponding declination.

Type	Source	α 1°	δ 1°	p-value	n_S	γ	B_{1°	$\Phi_{\nu_\mu+\bar{\nu}_\mu}^{90\%}$ ζ^a	
BL Lac	PKS 0537-441	84.71	-44.09	...	0.0	...	45.7	9.79	
	PKS 2155-304	329.72	-30.23	...	0.0	...	52.6	6.07	
	PKS 0235+164	39.66	16.62	0.12	16.2	3.4	72.0	0.94	
	1ES 0229+200	38.20	20.29	0.20	9.5	3.5	66.6	0.84	
	W Comae	185.38	28.23	...	0.0	...	60.8	0.62	
	Mrk 421	166.11	38.21	0.32	2.7	1.8	53.1	0.94	
	Mrk 501	253.47	39.76	0.18	10.5	4.0	52.0	1.15	
	BL Lac	330.68	42.28	...	0.0	...	50.4	0.63	
	H1426+428	217.14	42.67	...	0.0	...	50.9	0.70	
	3C 66A	35.67	43.04	...	0.0	...	50.9	0.70	
	1ES 2344+514	356.77	51.70	...	0.0	...	46.3	0.81	
	1ES 1959+650	300.00	65.15	0.018 ^b	15.4	3.1	42.8	2.36	
	S5 0716+71	110.47	71.34	...	0.0	...	38.4	1.34	
	Flat-spectrum radio quasar	PKS 1454-354	224.36	-35.65	...	0.0	...	49.1	7.99
		PKS 1622-297	246.53	-29.86	0.11	3.8	2.3	52.7	8.20
QSO 1730-130		263.26	-13.08	...	0.0	...	49.8	2.18	
PKS 1406-076		212.24	-7.87	0.053 ^c	7.3	2.6	50.5	1.65	
QSO 2022-077		306.42	-7.64	...	0.0	...	50.5	0.99	
3C 279		194.05	-5.79	0.42	0.5	2.0	54.3	0.63	
3C 273		187.28	2.05	0.25	7.7	3.2	76.4	0.59	
PKS 1502+106		226.10	10.49	0.38	3.1	2.7	73.7	0.59	
PKS 0528+134		82.73	13.53	0.44	2.7	4.0	73.0	0.60	
3C 454.3		343.49	16.15	0.12	4.1	2.0	72.3	0.93	
4C 38.41		248.81	38.13	0.12	6.3	2.4	53.2	1.31	
Galactic center	Sgr A*	266.42	-29.01	...	0.0	...	52.2	6.08	
Not identified	MGRO J1908+06	286.98	6.27	0.025	4.5	2.0	74.9	0.99	
Pulsar wind nebula	Geminga	98.48	17.77	...	0.0	...	69.3	0.49	
	Crab Nebula	83.63	22.01	0.34	6.1	3.8	67.0	0.68	
	MGRO J2019+37	305.22	36.83	0.23	7.8	4.0	53.2	1.04	
Star formation region	Cyg OB2	308.30	41.32	0.26	5.9	4.0	50.6	0.99	
Supernova remnant	IC443	94.21	22.50	0.22	8.1	4.0	66.0	0.83	
	Cas A	350.81	58.81	0.14	8.1	4.0	44.5	1.49	
	TYCHO	6.36	64.18	0.27	4.6	3.4	42.4	1.23	
Starburst/radio galaxy	Cen A	201.37	-43.02	0.21	0.5	1.2	46.2	10.41	
	M87	187.71	12.39	...	0.0	...	73.2	0.48	
	3C 123.0	69.27	29.67	...	0.0	...	59.5	0.57	

Table 2 continued

Table 2 (*continued*)

Type	Source	α 1°	δ 1°	p-value	n_S	γ	B_{1°	$\Phi_{\nu_\mu+\bar{\nu}_\mu}^{90\%}$ ζ^a
HMXB/mqso	Cyg A	299.87	40.73	0.068	2.1	1.4	51.1	1.50
	NGC 1275	49.95	41.51	...	0.0	...	50.6	0.71
	M82	148.97	69.68	...	0.0	...	39.7	1.09
	SS 433	287.96	4.98	0.40	4.1	4.0	75.8	0.47
	HESS J0632+057	98.24	5.81	0.10	13.6	3.6	75.4	0.77
	Cyg X-1	299.59	35.20	0.31	4.5	4.0	54.4	0.90
	Cyg X-3	308.11	40.96	0.077	12.8	4.0	51.3	1.53
	LSI 303	40.13	61.23	...	0.0	...	43.8	0.79

NOTE—Sources of the list are grouped by their classification and ordered in ascending declination δ . High-mass X-ray binaries and micro-quasars are abbreviated by *HMXB/mqso*.

Spectral indices quoted as $\gamma = 4.0$ are at the boundary of the parameter space for minimization. This can happen when there is an over-fluctuation of low-energy events close to the source location. It was ensured that for all events, the minimization converged successfully to the minimum within the parameter-space.

^aUpper limits are in units of $\zeta = 10^{-12} \text{ TeV cm}^{-2} \text{ s}^{-1}$

^bMost significant source in northern sky in this table. The trial-corrected p-value is 54%.

^cMost significant source in southern sky in this table. The trial-corrected p-value is 37%.

Table 3. Sources contained in the second source list, using only sources in the southern sky and focusing on galactic objects. The information listed is the same as in [Tab. 2](#).

Type	Source	α 1°	δ 1°	p-value	n_S	γ	B_{1°	$\Phi_{\nu_\mu+\bar{\nu}_\mu}^{90\%}$ ζ^a
BL Lac	PKS 2005-489	302.37	-48.82	0.071	0.9	1.0	44.7	13.45
	PKS 0426-380	67.17	-37.93	...	0.0	...	47.2	8.93
	PKS 0548-322	87.67	-32.27	...	0.0	...	51.2	6.79
	H2356-309	359.78	-30.63	...	0.0	...	52.1	6.18
	1ES 1101-232	165.91	-23.49	...	0.0	...	52.6	4.64
	1ES 0347-121	57.35	-11.99	0.21	1.4	2.4	52.2	2.16
Flat-spectrum radio quasar	PKS 0454-234	74.27	-23.43	...	0.0	...	52.8	4.58
	PKS 0727-11	112.58	-11.70	0.20	2.7	3.7	52.0	2.30
Not identified	HESS J1507-622	226.72	-62.34	...	0.0	...	43.4	11.02
	HESS J1503-582	226.46	-58.74	...	0.0	...	45.5	11.79
	HESS J1741-302	265.25	-30.20	0.19	2.1	4.0	52.6	7.33
	HESS J1834-087	278.69	-8.76	0.21	1.2	3.7	49.5	1.47
	HESS J1356-645	209.00	-64.50	...	0.0	...	42.4	10.90
Pulsar wind nebula	PSR B1259-63	197.55	-63.52	0.21	1.3	2.0	42.7	11.43
	HESS J1303-631	195.74	-63.20	0.076	4.5	2.3	42.3	13.61
	MSH 15-52	228.53	-59.16	...	0.0	...	44.9	11.28
	HESS J1023-575	155.83	-57.76	...	0.0	...	46.4	11.79

Table 3 *continued*

Table 3 (*continued*)

Type	Source	α 1°	δ 1°	p-value	n_S	γ	B_{1°	$\Phi_{\nu_\mu+\bar{\nu}_\mu}^{90\%}$ ζ^a
	HESS J1616-508	243.78	-51.40	0.0022 ^b	2.4	4.0	45.0	19.37
	HESS J1632-478	248.04	-47.82	0.16	1.5	2.7	44.7	10.79
	Vela X	128.75	-45.60	0.13	2.7	2.4	45.8	10.79
	HESS J1837-069	279.41	-6.95	...	0.0	...	48.1	0.89
Supernova remnant	RCW 86	220.68	-62.48	...	0.0	...	43.1	11.02
	RX J0852.0-4622	133.00	-46.37	...	0.0	...	45.6	10.40
	RX J1713.7-3946	258.25	-39.75	...	0.0	...	45.5	9.22
	W28	270.43	-23.34	...	0.0	...	52.9	4.58
Seyfert galaxy	ESO 139-G12	264.41	-59.94	...	0.0	...	44.0	11.28
HMXB/mqso	Cir X-1	230.17	-57.17	...	0.0	...	46.3	11.03
	GX 339-4	255.70	-48.79	0.15	2.6	2.2	44.8	11.29
	LS 5039	276.56	-14.83	0.26	2.1	4.0	52.3	2.72
Massive star cluster	HESS J1614-518	243.58	-51.82	0.0058	2.2	4.0	45.4	18.33

NOTE—Please refer to [Tab. 2](#) for comments regarding this table.

^aUpper limits are in units of $\zeta = 10^{-12} \text{ TeV cm}^{-2} \text{ s}^{-1}$

^bMost significant source in this table (South only). The trial-corrected p-value is 9.3%.