

# Lawrence Berkeley National Laboratory

## Recent Work

### Title

THE RADIATION- INDUCED OXIDATION OF PEPTIDES IN AQUEOUS SOLUTION

### Permalink

<https://escholarship.org/uc/item/97939199>

### Authors

Atkins, Harriette L.  
Bennett-Corniea, Winifred  
Garrison, Warren M.

### Publication Date

1966-07-01

**University of California**  
**Ernest O. Lawrence**  
**Radiation Laboratory**

**TWO-WEEK LOAN COPY**

*This is a Library Circulating Copy  
which may be borrowed for two weeks.  
For a personal retention copy, call  
Tech. Info. Division, Ext. 5545*

THE RADIATION-INDUCED OXIDATION OF PEPTIDES IN AQUEOUS SOLUTION

Berkeley, California

## **DISCLAIMER**

This document was prepared as an account of work sponsored by the United States Government. While this document is believed to contain correct information, neither the United States Government nor any agency thereof, nor the Regents of the University of California, nor any of their employees, makes any warranty, express or implied, or assumes any legal responsibility for the accuracy, completeness, or usefulness of any information, apparatus, product, or process disclosed, or represents that its use would not infringe privately owned rights. Reference herein to any specific commercial product, process, or service by its trade name, trademark, manufacturer, or otherwise, does not necessarily constitute or imply its endorsement, recommendation, or favoring by the United States Government or any agency thereof, or the Regents of the University of California. The views and opinions of authors expressed herein do not necessarily state or reflect those of the United States Government or any agency thereof or the Regents of the University of California.

Phys. Chem.

UCRL-17004

UNIVERSITY OF CALIFORNIA

Lawrence Radiation Laboratory  
Berkeley, California

AEC Contract No. W-7405-eng-48

THE RADIATION-INDUCED OXIDATION OF PEPTIDES IN AQUEOUS SOLUTION

Harriette L. Atkins, Winifred Bennett-Corniea and Warren M. Garrison

July 1966


THE RADIATION-INDUCED OXIDATION OF PEPTIDES IN AQUEOUS SOLUTION<sup>1</sup>

Harriette L. Atkins, Winifred Bennett-Corniea and Warren M. Garrison



Lawrence Radiation Laboratory  
University of California  
Berkeley, California

July 1966



Radiolysis of compounds of the type  $\text{RCONHCHR}_2$  in dilute aqueous solution containing  $\text{O}_2$  leads to formation of labile peptide derivatives which are readily degraded on mild hydrolysis to give ammonia and carbonyl compounds as characteristic products.<sup>2</sup> It has been proposed<sup>2,3</sup> that such oxidation in the case of peptides derived from the simpler amino acids, glycine and alanine, is initiated by a preferential attack of the OH radical at the carbon-hydrogen position  $\alpha$  to the nitrogen function. The over-all reaction scheme includes the radiation-induced step<sup>4</sup>





followed by



The reducing species  $\text{e}_{\text{aq}}^-$  and  $\text{H}$  are scavenged quantitatively by  $\text{O}_2$  to form  $\text{O}_2^-$  and  $\text{HO}_2$  which are related by the equilibrium<sup>5</sup>  $\text{HO}_2 \rightleftharpoons \text{O}_2^- + \text{H}^+$ . Proposed reaction modes for removal of the intermediate peroxy radicals include





where  $\text{RCON=CR}_2$  represents a dehydropeptide and  $\text{RCONHC(OH)R}_2$  the corresponding hydrate. Such compounds readily decompose on hydrolysis<sup>6</sup>



The above reaction scheme requires that the ammonia and carbonyl yields be in the relationship  $G(\text{NH}_3) = G(\text{R}_2\text{CO}) = G_{\text{OH}}$  where the latter term represents the 100 eV yield for OH production in the radiation induced step, 1. We have measured ammonia and carbonyl yields in the  $\gamma$ -ray induced oxidation of N-acetyl-glycine,<sup>3c</sup> glycine anhydride, N-acetylalanine and alanine anhydride in oxygenated 0.1 M solution and we find for each system,  $G(\text{NH}_3) \approx 3$  which value is consistent with recent measurements<sup>7</sup> of the maximal yield for the production of OH radicals in water under  $\gamma$ -rays. On the other hand, we also find that carbonyl production in these simple peptide systems is not in accord with the quantitative requirements of the proposed oxidation scheme; the initial carbonyl yields are uniformly low with  $G(\text{R}_2\text{CO}) \leq 0.8$ . There is then the question as to whether this apparent discrepancy arises from a) an incorrect formulation of the locus of initial OH attack or from b) unspecified complexities in the chemistry of removal of the peroxy radicals  $\text{RCONHC(O}_2\text{)R}_2$ .

To obtain specific information on this point, we have employed FeIII instead of O<sub>2</sub> as the scavenger of intermediate radicals formed in the radiolysis of N-acetylalanine and N-acetylglycine. Heavy metal ions such as FeIII and CuII oxidize organic free radicals in aqueous solution by electron transfer and by ligand transfer.<sup>8</sup> Such reactions in the case of the peptide radical RCONHCR<sub>2</sub> would correspond to:



respectively. We note that the organic products of reactions 9, 10 are identical with the postulated products of reactions 4, 6.

### Experimental

Solutions containing the acetylamino acid (Nutritional Biochemicals, twice recrystallized) plus ferric sulfate (C.P.) were adjusted to the desired pH with sulfuric acid, evacuated, and then irradiated with Co<sup>60</sup> γ-rays to a dose of not more than  $\sim 5 \times 10^{18}$  eV/ml. Above this value the dose-yield curves were no longer linear. Dose was determined with the Fricke dosimeter,  $\epsilon_{\text{FeIII}} = 2130$  at 305 mμ and 22°C. Immediately after irradiation, an aliquot was passed through a Dowex 50 column (1 cm × 10 cm), acid form, to remove iron. The eluant was made 2 N in hydrochloric acid, heated to 100°C for 60 minutes, cooled, and assayed for α-keto acid and aldehyde through use of the 2,4-dinitrophenylhydrazine reagent.<sup>9</sup> A second aliquot was made 2 N in sodium hydroxide and allowed to stand 24 hours in the outer compartment of a Conway diffusion

cell to liberate ammonia which was collected in 0.1 N sulfuric acid in the inner compartment, and assayed by the standard Tollen procedure. Control runs established that product hydrolysis and ammonia transfer were quantitative under these experimental conditions.


### Results and Discussion

Ammonia production (Fig. 1) in 0.1 M acetylalanine increases abruptly from  $G(\text{NH}_3) = 0.7$  to  $G(\text{NH}_3) = 4.3$  with increasing concentrations of FeIII up to  $\sim 10^{-3}$  M. The ammonia yield then falls gradually to a limiting value of  $G(\text{NH}_3) = 3.3$  at the higher FeIII concentrations. We also find under these conditions that pyruvic acid and ammonia are formed in equal molar yields. Yields of glyoxylic acid and ammonia from acetyl-glycine also show this same quantitative relationship. Aldehyde yields from these systems are low,  $G \approx 0.1$ .

At the higher (FeIII)/(peptide) ratios, the reducing species  $e_{\text{aq}}^-$  and  $\text{H}$  are preferentially scavenged by FeIII and the yield for peptide oxidation through OH attack is in accord with


$$-G(\text{peptide}) = G(\text{NH}_3) = G(\text{RCOCOOH}) \approx 3.2 \approx G_{\text{OH}} + G_{\text{H}_2\text{O}_2}$$

Hydrogen peroxide formed in the radiation-induced step, 1, reacts rapidly with FeII to give an additional yield of OH radicals<sup>10</sup>






The maximum in the yield curve shown in Fig. 1 is attributed to the onset of the reaction



in competition with



at the lower (FeIII)/(peptide) ratios. The  $\text{RCONH}\dot{\text{C}}\text{R}_2$  radicals from both reactions 2 and 12 are then available for oxidation by FeIII. This effect is shown more clearly in Fig. 2 which gives ammonia and pyruvic acid yields as a function of acetylalanine concentration over the range  $10^{-3}$  M to 1.5 M, in the presence of 0.05 M FeIII. The limiting value for peptide oxidation at the higher acetylalanine concentrations is given by

$$-G(\text{peptide}) = G(\text{NH}_3) = G(\text{R}_2\text{CO}) \approx 4.0 \approx G_{\text{OH}} + G_{\text{H}_2\text{O}_2} + G_{\text{H}}$$

We conclude that the reaction of OH and H radicals with these peptide derivatives of the simpler amino acids, glycine and alanine, occurs essentially quantitatively at the  $\alpha$ -position as formulated in reactions 2,12.<sup>11</sup> The present data also establish the quantitative oxidation of  $\text{RCONH}\dot{\text{C}}\text{R}_2$  radicals by FeIII via reactions 9, 10. In the case of acetylalanine such oxidation appears to occur almost exclusively through ligand transfer (reaction 10) since measurements of the optical absorption of the irradiated solutions (after removal of FeIII) reveal negligible absorption above 230 m $\mu$  when read differentially against

unirradiated control solution. Absorption by control solutions containing authentic acetyldehydroalanine<sup>12</sup> ( $\epsilon_{240} = 6050$ ) show that G values of  $>0.1$  for reaction 9 would be detectable. To our knowledge the optical properties of acetyldehydroglycine have not been described.

The low carbonyl yields obtained when  $O_2$  is used in place of FeIII as the radical scavenger are interpreted here as evidence that other more complex degradation reactions occur in parallel with the dehydrogenation and hydroxylation reactions, 4-6. The nature of these more complex branching reactions of the peroxy radical  $RCONHC(O_2)R_2$  is presently under study.


References

- (1) This work was performed under the auspices of the U. S. Atomic Energy Commission.
- (2) M. E. Jayko, B. M. Weeks, and W. M. Garrison, Proc. 2nd Intern. Conf. Peaceful Uses Atomic Energy, Geneva, 29, 30 (1958).
- (3a) W. M. Garrison, M. E. Jayko, and Winifred Bennett-Corniea, Radiation Res. 16, 483 (1962);
- (3b) W. M. Garrison and B. M. Weeks, *ibid* 17, 341 (1962);
- (3c) W. M. Garrison, *ibid*, Supplement 4, 148 (1964).
- (4a) A. O. Allen, Radiation Res. Supplement 4, 54 (1964);
- (4b) E. J. Hart, Science 146, 19 (1964).
- (5) G. Czapski and H. J. Bielski, J. Phys. Chem. 67, 2180 (1963).
- (6) J. P. Greenstein and M. Winitz, "Chemistry of the Amino Acids" (John Wiley and Sons, Inc., New York, N. Y., 1961) p. 843.
- (7) E. Hayon, Trans. Faraday Soc. 61, 723 (1965).
- (8) J. H. Baxendale and D. H. Smithies, Z. Physik Chem. 7, 242 (1956); H. E. DeLaMare, J. K. Kochi, and F. F. Rust, J. Am. Chem. Soc. 85, 1437 (1963).
- (9) G. R. A. Johnson and G. Scholes, Analyst. 79, 217 (1954); T. E. Friedemann and G. E. Haugen, J. Biol. Chem. 147, 415 (1943).
- (10) F. Haber and J. Weiss, Proc. Roy. Soc. A147, 333 (1934).
- (11) We refer to radical attack at the C-H linkage  $\alpha$  to the nitrogen function as the characteristic peptide reaction. With the more complicated amino acid residues, attack may occur in parallel or exclusively at side-chain loci. Ref. 3, also H. A. Sokol, W. Bennett-Corniea and W. M. Garrison, J. Am. Chem. Soc. 87, 1391 (1965).
- (12) We are indebted to Dr. B. M. Weeks for the sample of authentic acetyldehydroalanine which was prepared after the method of V. E. Price and J. P. Greenstein. Arch. Biochem. 10, 383 (1948).

Legends for Figures


Fig. 1. Effect of FeIII concentration on the yields of ammonia (●) and pyruvic acid (▲) from 0.1 M acetylalanine and of ammonia (○) and glyoxylic acid (Δ) from 0.1 M acetyl glycine.

Fig. 2. Effect of acetylalanine concentration on yields of ammonia (●) and of pyruvic acid (▲) from solutions containing 0.05 M FeIII.



MUB-12106

Fig. 1



MUB-12107

Fig. 2

This report was prepared as an account of Government sponsored work. Neither the United States, nor the Commission, nor any person acting on behalf of the Commission:

- A. Makes any warranty or representation, expressed or implied, with respect to the accuracy, completeness, or usefulness of the information contained in this report, or that the use of any information, apparatus, method, or process disclosed in this report may not infringe privately owned rights; or
- B. Assumes any liabilities with respect to the use of, or for damages resulting from the use of any information, apparatus, method, or process disclosed in this report.

As used in the above, "person acting on behalf of the Commission" includes any employee or contractor of the Commission, or employee of such contractor, to the extent that such employee or contractor of the Commission, or employee of such contractor prepares, disseminates, or provides access to, any information pursuant to his employment or contract with the Commission, or his employment with such contractor.

