

UCSF

UC San Francisco Previously Published Works

Title

Molecular Determinants for the Layering and Coarsening of Biological Condensates.

Permalink

<https://escholarship.org/uc/item/9807x5m8>

Journal

Aggregate, 3(6)

Authors

Latham, Andrew

Zhang, Bin

Publication Date

2022-12-01

DOI

10.1002/agt2.306

Copyright Information

This work is made available under the terms of a Creative Commons Attribution License, available at <https://creativecommons.org/licenses/by/4.0/>

Peer reviewed



HHS Public Access

Author manuscript

Aggregate (Hoboken). Author manuscript; available in PMC 2023 April 13.

Published in final edited form as:

Aggregate (Hoboken). 2022 December ; 3(6): . doi:10.1002/agt2.306.

Molecular Determinants for the Layering and Coarsening of Biological Condensates

Andrew P. Latham^{†,‡}, Bin Zhang[†]

[†]Department of Chemistry, Massachusetts Institute of Technology, Cambridge, MA, 02139

[‡]Department of Bioengineering and Therapeutic Sciences, University of California at San Francisco, San Francisco, CA 94143

Abstract


Many membraneless organelles, or biological condensates, form through phase separation, and play key roles in signal sensing and transcriptional regulation. While the functional importance of these condensates has inspired many studies to characterize their stability and spatial organization, the underlying principles that dictate these emergent properties are still being uncovered. In this review, we examine recent work on biological condensates, especially multicomponent systems. We focus on connecting molecular factors such as binding energy, valency, and stoichiometry with the interfacial tension, explaining the nontrivial interior organization in many condensates. We further discuss mechanisms that arrest condensate coalescence by lowering the surface tension or introducing kinetic barriers to stabilize the multidroplet state.

Graphical Abstract

binz@mit.edu .

Conflict of interest

The authors declare no potential conflict of interests.



Introduction

Cells constantly perform myriads biochemical reactions in parallel. To avoid crosstalks and interference among different pathways, cells compartmentalize into membrane-bound vesicles that enclose specific sets of molecules. The recently discovered membraneless organelles, also known as biological condensates, could offer similar specificity by concentrating functionally related molecules via self-assembly, providing additional levels of structure and organization within the cell.^{1–5} Many biological condensates are dynamic and exhibit liquid-like properties. They allow fast molecular exchange with the cellular environment and can assemble/dissolve on demand in response to external signals that modulate the concentration and chemical state of its components.^{6–8}

Biological condensates are involved in a variety of processes throughout the cell. For instance, stress granules form in the cytosol as a mechanism of cellular stress response to protect RNAs from harmful chemicals.⁹ TIS granules form at the interface with the endoplasmic reticulum to assist in the translation of mRNA and the formation of membrane protein complexes.^{10,11} Inside the nucleus, membraneless organelles contribute to genome organization, partitioning chromosomes into active and silent domains.^{1–3,12} They may assist DNA compaction in heterochromatin^{13,14} or promote transcription as in the case of condensates formed by super-enhancers.¹⁵

Significant progress has been made in understanding the mechanisms of biomolecular condensate formation in vitro. In particular, intrinsically disordered proteins (IDPs) are found in many membraneless organelles. They undergo phase separation at cellular concentrations to form dynamic droplets.^{16–19} The propensity of IDPs to phase separate

arises mainly from their ability to associate simultaneously with multiple partners, i.e., high valency. IDPs often harbor several disordered segments that promote electrostatic, cation- π , π - π , hydrogen bonding, or hydrophobic interactions.²⁰ In the meantime, specific interactions between ordered regions with well-defined protein-protein interfaces also contribute to the multivalency.^{21–23} Nucleic acids can form condensates as well.^{24,25} Similar to motifs in proteins, multivalent interactions between nucleic acids contribute to their phase separation propensity.^{26,27} It is also becoming increasingly appreciated that these diverse types of multivalent interactions among biomolecules could produce complex phase behaviors, including phase separation coupled with a percolation transition, which results in gel-like condensates with viscoelastic properties.^{19,28–30}

While the mechanisms of single-component systems are beginning to emerge, biological condensates in the cell contain many components, including proteins, DNA, and RNA. The complex molecular composition of these condensates gives rise to more elaborate mechanisms. For instance, higher valency proteins, frequently referred to as scaffolds, can drive the formation of condensates and incorporate lower valency proteins that are not capable of phase separation on their own, which are often called clients.^{21,35,36} Additionally, many biomolecules are highly charged, and Coulombic interactions can drive their condensation through a mechanism known as complex coacervation.^{37,38} Complex coacervates exemplify a broader class of condensates stabilized by cross-interactions between components. In such cases, condensate stability is often dependent on stoichiometry, resulting in reentrant phase separation.^{39,40} Bridging-induced polymer collapse has been proposed as a mechanism to explain the formation of chromatin bodies.^{41,42} For example, protein molecules may bind with chromatin in multiple locations to introduce physical crosslinks or bridges, the accumulation of which induces polymer collapse to form condensates. For more information on the mechanisms of biomolecular phase separation, we refer the reader to several of many great existing reviews.^{17–20,43}

The complexity in molecular composition also produces novel condensate internal organizations and coarsening behaviors deviating from homogeneous mixtures.^{20,44,45} Order-disorder transitions could occur to form substructures via microphase separation as in block-copolymer systems, producing layering in the condensates.^{46,47} The presence of substructures may facilitate the mini-factories to further streamline the processing and synthesis of biomolecules with dedicated platforms by division of labor. Novel behaviors in phase separation kinetics have also been observed in biological condensates. Defying expectations from the classical nucleation theory, condensates inside cells are often arrested in a multi-droplet state.⁴⁸ Further coarsening is rarely observed over timescales spanning the entire cell cycle (~ 24 hours).⁴⁹

In this review, we examine molecular mechanisms giving rise to emergent behaviors of biological condensates. We begin by exploring examples of condensates with non-trivial internal organizations. Next, we argue that the sub-structures in condensates can be understood and predicted from the interfacial tension among immiscible liquids. Further connecting molecular interactions with interfacial tension provides a conceptual framework to interpret various experimental observations on condensate stability and organization. We

conclude the review by examining the physical characteristics that prevent droplet fusion and thus limit the size of biomolecular condensates.

Layered Internal Organization of Biological Condensates

Many biological condensates display substructures to provide additional spatial and physical control necessary for complex functions. The nucleolus, essential for ribosome biogenesis, presents an excellent example highlighting the advantages of a layered interior organization.^{50,51} It is composed of three layers, the fibrillar center, the dense fibrillar component, and the granular component. This organization allows for each part of the nucleolus to have different protein compositions and perform distinct biological functions: rDNA is transcribed in the fibrillar center, the resulting rRNA is processed in the dense fibrillar center, and ribosome assembly occurs in the granular component (Figure 1A).^{31,52} Such a division of labor helps streamline ribosome production and assembly by assigning steps to different regions of the nucleolus.

The nuclear speckle is another example that displays a layered organization, and the functional significance of maintaining such intricate internal structures is still being revealed. Nuclear speckles are composed of both RNA and protein components and are essential for gene transcription and splicing.⁵³ Recent studies have shown that many RNA components, including snRNA like U2B", localize toward the exterior of the condensate, and scaffold proteins, such as SON, localize toward the interior (Figure 1B).^{32,54,55} This organization may serve as a mechanism to control condensate size, as the accumulation of pre-mRNA at the periphery can recruit more speckle components.³² Further, the localization of RNA to the exterior of the condensate may aid in biological function, for the interface between speckles and the nucleoplasm is likely the location of RNA splicing.⁵⁶

In addition to their differences in molecular composition, the various layers could exhibit distinct material properties as well. For example, the two layers in P granules are found to be either liquid-like or gel-like. P granules are the first proposed droplets formed through liquid-liquid phase separation and localize in the posterior half of *C. elegans* embryos.⁵⁷ The phase separation is driven by the protein, MEG-3,⁵⁸ which was later shown to form gel-like assemblies closer to the exterior of the P granules (Figure 1C).^{33,59} Meanwhile, the core of the P-body contains RNA-binding proteins such as PGL-3 and remains liquid-like.

Similar to P granules, stress granules were found to have liquid and gel-like compartments.⁶¹ Using stochastic optical reconstruction microscopy (STORM), Jain et al. demonstrated that stress granules contain core structures consisting of both proteins and mRNA (Figure 1D).³⁴ Later work demonstrated that protein dynamics within these core structures is slower than that within the surrounding medium.⁶² Time course analysis further showed that these cores form as precursors to the assembly of the liquid shell.⁶³

Both molecular specificity and differential physical properties could contribute to substrate selectivity among the substructures, as elegantly shown in a recent in vitro study. Choi et al. characterized a layered condensate formed by arginine repeats (polyR) and lysine repeats (polyK) with aspartic acid repeats (polyD).⁶⁴ They observed the layering of polyR

dominant and polyK dominant phases and tested the affinity of different phases for nucleic acid substrates, specifically dsRNA and ssRNA. As expected, ssRNA partitioned into the inner, polyR layer, likely due to increased opportunities for cation- π interactions. However, dsRNA partitioned into the polyK layer as a result of their increased stiffness and lack of cation- π interactions. Finally, its differential preference of ssRNA led to significant dehybridization of RNA duplexes within the polyR condensate layer.

Besides the above examples, other biological condensates, including paraspeckles,⁶⁵ anisosomes,⁶⁶ Cajal bodies,^{67,68} and mitochondrial transcriptional condensates,^{69,70} have been known to display complex organization as well. While the physical and biological implications of these structures have been discussed and remain of interest for future studies,^{44,49,52,71} how these organizations emerge from molecular level descriptions of the system has not been adequately reviewed. In the next section, we focus on the physical principles that dictate the internal organization of biological condensates, with a particular emphasis on connecting interactions among components to macroscopic condensate behaviors.

Molecular Determinants of Condensate Organization

Insights into the internal organization of biological condensates can be gained from studies of immiscible liquids. In particular, the geometric outcome for mixing two immiscible droplets is predictable with thermodynamic arguments in terms of their relative surface tensions.^{31,44,52,60,71,72} Two immiscible droplets (dropletA and dropletB) can form four unique configurations (Figure 2A): a layered organization with dropletA on the inside, a layered organization with dropletB on the inside, complete nonwetting to form two separate droplets, and partial wetting into two droplets that share an interface. Therefore, interfacial tensions provide a complete phenomenological description of condensate organization. However, they are emergent properties, and connecting interfacial tensions with the molecular composition is nontrivial.

Microscopic theories help connect surface tension with molecular interactions. The Flory-Huggins theory based on lattice models has proven successful at understanding the thermodynamics of polymer phase separation.⁷³ An essential parameter in the theory is

$$\chi = \frac{z}{2k_B T} [\epsilon_{pp} + \epsilon_{ss} - \epsilon_{ps}] = \frac{z\Delta\epsilon}{2k_B T}, \quad (1)$$

which accounts for the preferences of homotypic interactions, namely polymer-polymer (ϵ_{pp}) and solvent-solvent (ϵ_{ss}), over heterotypic solvent-polymer interactions (ϵ_{ps}). The coordination number (z), temperature (T), and Boltzmann constant (k_B) also determine χ . The theory correctly predicts the dependence of the critical temperature for phase separation on the interaction strength and length of polymers. Generalization of the theory that accounts for both enthalpic and entropic contributions to χ explains the presence of upper liquid critical temperature for certain polymers. Furthermore, it has been shown that the interfacial tension is directly related to χ^α .^{74,75} While the exact value of the exponent, α is subject to debate, a positive correlation between the two is clear. Therefore, increasing

the propensity for homotypic polymer interactions over heterotypic polymer interactions will also increase the surface tension between the polymer and solution phases (Figure 2B).

The above arguments provide a powerful conceptual framework for interpreting experimental observations. For example, poly(proline-arginine) is known to form layered droplets with RNA, where poly-C RNA localized at the solvent interface and poly-A RNA comprised the core.⁷⁶ Modeling demonstrated that this result may occur due to a stronger affinity of poly(proline-arginine) for poly-A RNA than poly-C RNA, a stronger self-affinity of poly-A relative to poly-C, or less favorable interactions for poly-A. Each of these interaction schemes will produce higher χ values and surface tension for the droplet formed by poly-A than those formed by poly-C. Similar results were seen when studying mixtures of arginine repeats (polyR) and lysine repeats (polyK) with uridine-5'-triphosphate trisodium salt (UTP), where polyR formed a core and polyK formed a shell. The differences between the layers were attributed to stronger R-UTP interactions than K-UTP interactions due to the ability to form pi-pi interactions (Figure 3A–C). These stronger interactions with polyR explain the higher surface tension of the corresponding droplet.⁷⁷ Elastin-like polypeptides (ELPs) are another model IDP system that supports similar behaviors. Notably, ELPs undergo a liquid-liquid phase separation above a lower critical solution temperature driven primarily by hydrophobic interactions.^{78–80} For condensates with multiple ELPs, a layered organization has been observed where more hydrophobic ELP sequences with higher values of χ populate the interior.⁸¹

While the Flory-Huggins theory was derived for homopolymers, it can be applied to heteropolymers and complex systems by averaging polymer-polymer and polymer-solvent interactions over the sequence to produce an effective χ (Figure 2C). For example, polymers with more solvated linkers will, on average, have smaller χ values and be pushed to the shell of the droplet, while linkers with less solvation volume will localize toward the interior.⁸⁵ Higher valency has been seen to correlate with the interior of the condensate, as a result of stronger effective interactions.⁸⁶ In a recent study, Latham and Zhang simulated the phase separation of a mixture of two chromatin regulators, HP1 α and histone H1, with DNA with the MOFF force field.^{87–90} They observed that H1 localizes toward the droplet-solvent interface and HP1 α is located toward the middle of the droplet. The observed layering can be explained by stronger HP1 α -HP1 α interactions relative to H1-H1 interactions (Figure 3D–E).⁸²

The effective χ value depends not only on the chemical composition of the molecule but also on the precise arrangement of the chemical groups. For instance, Regy et al. shuffled the sequence of the RGG domain of LAF1 and explored its phase behavior with RNA, specifically a 15 nucleotide adenosine repeat (A₁₅).⁸⁴ They found that a shuffled sequence with more charge blockiness (RGG_{Cshuf}) led to stronger protein-protein and protein-RNA interactions (Figure 3F). Further, the minimum of the protein-RNA potential of mean force moved such that it is minimized when the two biomolecules are held approximately 20 Å apart instead of at a mean distance of zero (Figure 3G). Because of these changes, RGG_{Cshuf} and RNA formed a layered condensate, with RNA located on the exterior of the condensate (Figure 3H,I). Other computational work on polyampholytes has found similar results. Using sequence-specific modeling of K/E mixtures of the same overall composition, Pal

et al. showed that differences in the blockiness of charge distribution can drive condensate layering, likely through changes to the shape or depth of the pairwise PMF for the two chains.⁹¹ These studies typify previous theories on charge patterning^{83,92–95} to demonstrate that higher-order features in protein sequences beyond the amino acid composition can impact the conformation of individual molecules and the collective behavior of condensates.

The effective averaging procedure is inherently a mean-field assumption, which breaks down when functional groups with strong interactions stick together to form clusters. This clustering and even microphase separation can lead to layered organizations in single component systems, as seen in condensates formed with the Velo1 N-terminal prion-like domain.⁹⁶ This protein contains a variety of aromatic residues, which have a strong affinity for each other. The association of these aromatic residues slows down protein dynamics and renders the corresponding condensate more solid-like. Similar results were seen in a computational work on the RNA binding protein Fused in Sarcoma (FUS).⁹⁷ The authors modeled a disordered-to-ordered transition in the prion-like domain that is expected upon condensate aging. After the transition, the higher self-affinity among prion-like domains drove a layered organization and their interior localization.^{98,99} For these systems, by dividing the molecules into multiple groups with individual effective χ values, their organization can be analyzed similarly to the multi-component systems mentioned above (Figure 2D).

The energetics-focused perspective outlined above may miss certain features of phase separation. For instance, experiments have demonstrated that arginine-rich polypeptides can form layered droplets with RNA, with the component in excess preferentially located on the exterior.¹⁰⁰ Modeling has since demonstrated that the polymer in excess localizes to the interface due to the entropic gain of providing multiple binding partners. Further, as the stoichiometry becomes more unbalanced, the amount of unbound polymer at the interface increases, which decreases the surface tension of the condensate.¹⁰¹ In addition, non-equilibrium processes can also play a role in condensate organization. Using a model of the human genome,^{102,103} Jiang et al. examined the role of active forces, such as those that arise from transcription or chromatin remodeling,^{104,105} on phase separation, genome structure, and genome dynamics.¹⁰⁶ They found that applying active forces to euchromatin pushes heterochromatin to the nuclear periphery. This organization competes with strong attractive interactions, which pull heterochromatin toward the center of the nucleus. Similar active processes may play a role in the organization of other systems, including transcriptional condensates.¹⁵

Limitations on Condensate Coarsening

While thermodynamic arguments regarding surface tension have proven successful at explaining many experimental results, some observations appear to defy predictions. In particular, a single condensate is expected at equilibrium to minimize the surface energy.^{107,108} However, multiple nucleoli (~2–5) can stably coexist in the same nucleus,¹⁰⁹ as can paraspeckles¹¹⁰ and nuclear speckles.¹¹¹ Mechanisms that prevent the coarsening and fusion of biological condensates are only beginning to emerge.

The complexity in the molecular interactions that drive condensate formation and their internal organization may produce micelle-like structures with low surface tension, reducing the driving force for coarsening. Similar to surfactants that harbor both hydrophobic and hydrophilic groups, molecules found in condensates differ in their valency. High valency molecules prefer polymer-polymer over polymer-solvent interactions and are thus more “hydrophobic” than low valency ones. The accumulation of low-valency molecules at the condensate exterior could lower the interfacial tension. To demonstrate this principle, Sanchez-Burgos et al. used a minimal scaffold-surfactant model, where scaffolds and surfactants are particles with high and low valency, respectively. They showed that the surface tension, and hence the number of droplets, was tunable by the surfactant-scaffold ratio (Figure 4A).¹¹² This computational framework has been seen in biological contexts. For example, high valency complexes of G3BP and UBAP2L cause the formation of stress granules, while the low valency of USP10 allows it to act as a cap to inhibit stress granule growth.^{115,116} Similar behaviors were also found in a designed ArtiGranule (ArtiG) system, where the presence of modified human ferritin protein with RNA-binding domains (PUM.HD-FFm) reduced the condensate size compared to the protein without RNA-binding domains (mCherry-FFm) (Figure 4B).¹¹³ The authors proposed that RNA binding limits the valency of surface protein molecules, suppressing condensate fusion. The role of valency is also supported by a combination of in-vitro experiments and molecular simulation, where decreasing RNA length, which also lowers RNA valency, was found to reduce the stability, surface tension, and density of protein-RNA condensates.¹¹⁷

Localization of disordered proteins to the interface of condensates can also control the size of biological condensates through other mechanisms. Recent work on P granules suggests that the disordered protein MEG-3 can act as a Pickering agent.¹¹⁸ Folkmann et al. observed that MEG-3 proteins form low dynamic assemblies resembling solid particles, which coat the surface of P granules to reduce the surface tension.³³ Depleting MEG-3 causes P granules to increase dramatically in size.

In addition to lower thermodynamic driving forces, condensate fusion may be slowed down due to kinetic barriers. For example, net charges can induce an accumulation of counterions near the condensate surface, giving rise to the so-called zeta potential. Higher zeta potentials slow or prevent condensate fusion due to stronger electrostatic repulsion.¹¹⁹ Additionally, fusion barriers could arise from breaking existing structures inside the droplets. Using a stickers and spacers model, Ranganathan and Shakhnovich demonstrated that strong interactions with low valency produce finite-sized droplets with saturated interaction sites.²² Existing interactions must be abolished for these droplets to grow. Similarly, TIS granules, which are biological condensates that form near the endoplasmic reticulum,^{10,11} were found to have mesh-like shapes originating from the underlying network of cross-linked, disordered RNA. Fusion of such condensates would require a breakdown of the mesh that faces an enormous energetic penalty.

Just as the structural organization of condensates can hinder their fusion, structures within the solvent phase can also arrest condensate coalescence. Through explicit simulations of the phase separation process for nucleoli formation with the presence of a chromatin network,^{102,103} Qi and Zhang showed that the reorganization of chromatin creates an

entropic barrier to nucleoli fusion,¹¹⁴ and accounts for the observation that many separate nucleoli are observed in the cell (Figure 4C–E).¹²⁰ Other studies have also considered droplet growth in the nucleus by modeling the nucleus as an elastic network.^{121–124} In these cases, the mechanical stress of the permanently cross-linked network was attributed to arresting the coarsening dynamics of condensates. As the dynamical processes of chromatin are known to be timescale dependent,¹²⁵ both the entropic barrier of chromatin reorganization and the mechanical stress of disrupting the chromatin network may limit the coarsening of condensates within the nucleus.

Finally, active processes can also contribute to the arrest of condensate coalescence. Wurtz and Lee introduced a theoretical model in which phase-separating molecules can be converted into soluble molecules through chemical reactions. For certain reaction rates, the steady state of the system corresponds to many monodisperse droplets.¹²⁶ This model may be applicable to a variety of membraneless organelles in the cell, where changes such as post-translational modifications could serve as a control mechanism to inhibit condensate growth.¹²⁷ To demonstrate the role of active processes in a biological context, Guilhas et al. studied ParABS, a system responsible for chromosome and plasmid segregation in bacteria.¹²⁸ While the ParB protein drove phase separation, they found that ParA, specifically ParA's ATPase activity, was necessary to control the size and location of ParABS condensates. Similar mechanisms may also play a role in other condensates, including P granules¹²⁹ and stress granules,^{130,131} where enzymatic activity has been shown to dissolve condensates.¹²⁷

Conclusions and Future Outlook

In this review, we have discussed the molecular factors that dictate the interfacial tension and, thus, the size and structure of biological condensates. Such factors include interaction energy between components, valency, stoichiometry, topological constraints, etc. Continued advances in experimental, computational, and theoretical methods should lead to the discovery of many new biological condensates and improved characterization of existing ones. Such discoveries may allow scientists to perturb and engineer biological condensates for therapeutic purposes.

Acknowledgments

This work was supported by the National Institutes of Health (Grant R35GM133580) and the National Science Foundation (Grant MCB-2042362). A.L. further acknowledges support by the National Science Foundation Graduate Research Fellowship Program.

Biographies



Andrew P. Latham received his B.S. from the University of Notre Dame. He completed a Ph.D. in Chemistry from the Massachusetts Institute of Technology under the supervision of

Dr. Bin Zhang in May 2022, and is now a postdoctoral scholar in Andrej Sali's group at the University of California, San Francisco. His research interests focus on leveraging existing experimental data to build high resolution computational models of complex biological systems, including membraneless organelles.



Bin Zhang obtained his Ph.D. from the California Institute of Technology, where he worked with Thomas F. Miller on Sec-facilitated protein translocation and membrane integration. Upon graduation, Bin accepted a position as a postdoctoral scholar with Peter G. Wolynes at the Center for Theoretical Biological Physics at Rice University. He joined Massachusetts Institute of Technology in 2016, where he is the Pfizer-Laubach Career Development Associate Professor of Chemistry. His group develops multiscale coarse-graining approaches to simulate the organization of biocondensates and the human genome.

References

- (1). Sabari BR; Dall'Agnes A; Young RA Biomolecular Condensates in the Nucleus. Trends Biochem. Sci 2020, 45, 961–977. [PubMed: 32684431]
- (2). Sanulli S; Narlikar J, Liquid-like G interactions in heterochromatin: Implications for mechanism and regulation. Curr. Opin. Cell Biol 2020, 64, 90–96. [PubMed: 32434105]
- (3). Banani SF; Lee HO; Hyman AA; Rosen MK Biomolecular condensates: Organizers of cellular biochemistry. Nat. Rev. Mol. Cell Biol 2017, 18, 285–298. [PubMed: 28225081]
- (4). Uversky VN Intrinsically disordered proteins in overcrowded milieu: Membrane-less organelles, phase separation, and intrinsic disorder. Curr. Opin. Struct. Biol 2017, 44, 18–30. [PubMed: 27838525]
- (5). Woodruff JB; Hyman AA; Boke E Organization and Function of Non-dynamic Biomolecular Condensates. Trends Biochem. Sci 2018, 43, 81–94. [PubMed: 29258725]
- (6). Hnisz D; Shrinivas K; Young RA; Chakraborty AK; Sharp PA A phase separation model predicts key features of transcriptional control. Cell 2017, 169, 13–23. [PubMed: 28340338]
- (7). Riback JA; Zhu L; Ferrolino MC; Tolbert M; Mitrea DM; Sanders DW; Wei MT; Kriwacki RW; Brangwynne CP Composition-dependent thermodynamics of intracellular phase separation. Nature 2020, 581, 209–214. [PubMed: 32405004]
- (8). Klein IA et al. Partitioning of cancer therapeutics in nuclear condensates. Science 2020, 368, 1386–1392. [PubMed: 32554597]
- (9). Riback JA; Katanski CD; Kear-Scott JL; Pilipenko EV; Rojek AE; Sosnick TR; Drummond DA Stress-Triggered Phase Separation Is an Adaptive, Evolutionarily Tuned Response. Cell 2017, 168, 1028–1040.e19. [PubMed: 28283059]
- (10). Ma W; Mayr C A Membraneless Organelle Associated with the Endoplasmic Reticulum Enables 3'UTR-Mediated Protein-Protein Interactions. Cell 2018, 175, 1492–1506.e19. [PubMed: 30449617]
- (11). Ma W; Zheng G; Xie W; Mayr C In vivo reconstitution finds multivalent RNA–RNA interactions as drivers of mesh-like condensates. eLife 2021, 10, 1–32.
- (12). Lin X; Qi Y; Latham AP; Zhang B Multiscale Modeling of Genome Organization with Maximum Entropy Optimization. J. Chem. Phys 2021, 155, 010901. [PubMed: 34241389]
- (13). Larson AG; Elnatan D; Keenen MM; Trnka MJ; Johnston JB; Burlingame AL; Agard DA; Redding S; Narlikar GJ Liquid droplet formation by HP1 α suggests a role for phase separation in heterochromatin. Nature 2017, 547, 236–240. [PubMed: 28636604]

- (14). Strom AR; Emelyanov AV; Mir M; Fyodorov DV; Darzacq X; Karpen GH Phase separation drives heterochromatin domain formation. *Nature* 2017, 547, 241–245. [PubMed: 28636597]
- (15). Sabari BR et al. Coactivator condensation at super-enhancers links phase separation and gene control. *Science* 2018, 361, eaar3958. [PubMed: 29930091]
- (16). Borchers W; Bremer A; Borgia MB; Mittag T How do intrinsically disordered protein regions encode a driving force for liquid-liquid phase separation? *Curr. Opin. Struct. Biol* 2021, 67, 41–50. [PubMed: 33069007]
- (17). Hyman AA; Weber CA; Jülicher F Liquid-Liquid Phase Separation in Biology. *Annu. Rev. Cell Dev. Biol* 2014, 30, 39–58. [PubMed: 25288112]
- (18). Boeynaems S; Alberti S; Fawzi NL; Mittag T; Polymenidou M; Rousseau F; Schymkowitz J; Shorter J; Wolozin B; Van Den Bosch L; Tompa P; Fuxreiter M Protein Phase Separation: A New Phase in Cell Biology. *Trends Cell Biol* 2018, 28, 420–435. [PubMed: 29602697]
- (19). Mittag T; Pappu RV A conceptual framework for understanding phase separation and addressing open questions and challenges. *Mol. Cell* 2022, 82, 2201–2214. [PubMed: 35675815]
- (20). Dignon GL; Best RB; Mittal J Biomolecular Phase Separation : From Molecular Driving Forces to Macroscopic Properties. *Annu. Rev. Phys. Chem* 2020, 71, 53–75. [PubMed: 32312191]
- (21). Banani SF; Rice AM; Peeples WB; Lin Y; Jain S; Parker R; Rosen MK Compositional Control of Phase-Separated Cellular Bodies. *Cell* 2016, 166, 651–663. [PubMed: 27374333]
- (22). Ranganathan S; Shakhnovich EI Dynamic metastable long-living droplets formed by sticker-spacer proteins. *eLife* 2020, 9, e56159. [PubMed: 32484438]
- (23). Harmon TS; Holehouse AS; Rosen MK; Pappu RV Intrinsically disordered linkers determine the interplay between phase separation and gelation in multivalent proteins. *eLife* 2017, 6, 1–31.
- (24). Zhu H et al. RNAPhase: A resource of RNAs undergoing phase separation. *Nucleic Acids Res* 2022, 50, D340–D346. [PubMed: 34718740]
- (25). Jain A; Vale RD RNA phase transitions in repeat expansion disorders. *Nature* 2017, 546, 243–247. [PubMed: 28562589]
- (26). Ries RJ; Zaccara S; Klein P; Olarerin-George A; Namkoong S; Pickering BF; Patil DP; Kwak H; Lee JH; Jaffrey SR m6A enhances the phase separation potential of mRNA. *Nature* 2019, 571, 424–428. [PubMed: 31292544]
- (27). Langdon EM; Qiu Y; Niaki AG; Mclaughlin GA; Weidmann CA; Gerbich TM; Smith JA; Crutchley JM; Termini CM; Weeks KM; Myong S; Gladfelter AS mRNA structure determines specificity of a polyQ-driven phase separation. *Science* 2018, 1, 922–927.
- (28). Majumdar A; Krainer G Phase-separating RNA-binding proteins form heterogeneous distributions of clusters in subsaturated solutions. *Proc. Natl. Acad. Sci. U.S.A* 2022, 119, e2202222119. [PubMed: 35787038]
- (29). Choi JM; Hyman AA; Pappu RV Generalized models for bond percolation transitions of associative polymers. *Phys. Rev. E* 2020, 102, 1–6.
- (30). Das S; Muthukumar M Microstructural Organization in α -Synuclein Solutions. *Macromolecules* 2022, 55, 4228–4236.
- (31). Feric M; Vaidya N; Harmon TS; Mitrea DM; Zhu L; Richardson TM; Kriwacki RW; Pappu RV; Brangwynne CP Coexisting Liquid Phases Underlie Nucleolar Subcompartments. *Cell* 2016, 165, 1686–1697. [PubMed: 27212236]
- (32). Fei J; Jadhavi M; Harmon TS; Li IT; Hua B; Hao Q; Holehouse AS; Reyer M; Sun Q; Freier SM; Pappu RV; Prasanth KV; Ha T Quantitative analysis of multilayer organization of proteins and RNA in nuclear speckles at super resolution. *J. Cell Sci* 2017, 130, 4180–4192. [PubMed: 29133588]
- (33). Folkmann AW; Putnam A; Lee CF; Seydoux G Regulation of biomolecular condensates by interfacial protein clusters. *Science* 2021, 373, 1218–1224. [PubMed: 34516789]
- (34). Jain S; Wheeler JR; Walters RW; Agrawal A; Barsic A; Parker R ATPase-Modulated Stress Granules Contain a Diverse Proteome and Substructure. *Cell* 2016, 164, 487–498. [PubMed: 26777405]
- (35). Elbaum-Garfinkle S; Kim Y; Szczepaniak K; Chen CCH; Eckmann CR; Myong S; Brangwynne CP The disordered P granule protein LAF-1 drives phase separation into droplets with tunable

- viscosity and dynamics. *Proc. Natl. Acad. Sci. U.S.A* 2015, 112, 7189–7194. [PubMed: 26015579]
- (36). Wang J; Choi JM; Holehouse AS; Lee HO; Zhang X; Jahnel M; Maharana S; Lemaitre R; Pozniakovskiy A; Drechsel D; Poser I; Pappu RV; Alberti S; Hyman AA A Molecular Grammar Governing the Driving Forces for Phase Separation of Prion-like RNA Binding Proteins. *Cell* 2018, 174, 688–699.e16. [PubMed: 29961577]
- (37). Sing CE; Perry SL Recent progress in the science of complex coacervation. *Soft Matter* 2020, 16, 2885–2914. [PubMed: 32134099]
- (38). Rumyantsev AM; Jackson NE; De Pablo JJ Polyelectrolyte Complex Coacervates: Recent Developments and New Frontiers. *Annu. Rev. Condens. Matter Phys* 2021, 12, 155–176.
- (39). Banerjee PR; Milin AN; Moosa MM; Onuchic PL; Deniz AA Reentrant Phase Transition Drives Dynamic Substructure Formation in Ribonucleoprotein Droplets. *Angew. Chem* 2017, 129, 11512–11517.
- (40). Henninger JE; Oksuz O; Shrinivas K; Sagi I; LeRoy G; Zheng MM; Andrews JO; Zamudio AV; Lazaris C; Hannett NM; Lee TI; Sharp PA; Cissé II; Chakraborty AK; Young RA RNA-Mediated Feedback Control of Transcriptional Condensates. *Cell* 2021, 184, 207–225.e24. [PubMed: 33333019]
- (41). Erdel F; Rippe K Formation of Chromatin Subcompartments by Phase Separation. *Biophys. J* 2018, 114, 2262–2270. [PubMed: 29628210]
- (42). Ryu JK; Bouchoux C; Liu HW; Kim E; Minamino M; de Groot R; Katan AJ; Bonato A; Marenduzzo D; Michieletto D; Uhlmann F; Dekker C Bridging-induced phase separation induced by cohesin SMC protein complexes. *Sci. Adv* 2021, 7.
- (43). Brangwynne CP; Tompa P; Pappu RV Polymer physics of intracellular phase transitions. *Nat. Phys* 2015, 11, 899–904.
- (44). Fare CM; Villani A; Drake LE; Shorter J Higher-order organization of biomolecular condensates. *Open Biol* 2021, 11.
- (45). Choi JM; Holehouse AS; Pappu RV Physical Principles Underlying the Complex Biology of Intracellular Phase Transitions. *Annu. Rev. Biophys* 2020, 49, 107–133. [PubMed: 32004090]
- (46). Leibler L Theory of Microphase Separation in Block Copolymers. *Macromolecules* 1980, 13, 1602–1617.
- (47). Bates FS; Fredrickson GH Block Copolymer Thermodynamics: Theory and Experiment. *Annu. Rev. Phys. Chem* 1990, 41, 525–557. [PubMed: 20462355]
- (48). Shimobayashi SF; Ronceray P; Sanders DW; Haataja MP; Brangwynne CP Nucleation landscape of biomolecular condensates. *Nature* 2021, 3.
- (49). Gouveia B; Kim Y; Shaevitz JW; Petry S; Stone HA; Brangwynne CP Capillary forces generated by biomolecular condensates. *Nature* 2022, 609, 255–264. [PubMed: 36071192]
- (50). Tschochner H; Hurt E Pre-ribosomes on the road from the nucleolus to the cytoplasm. *Trends Cell Biol* 2003, 13, 255–263. [PubMed: 12742169]
- (51). Boisvert FM; Van Koningsbruggen S; Navascués J; Lamond AI The multifunctional nucleolus. *Nat. Rev. Mol. Cell Biol* 2007, 8, 574–585. [PubMed: 17519961]
- (52). Lafontaine DL; Riback JA; Bascetin R; Brangwynne CP The nucleolus as a multiphase liquid condensate. *Nat. Rev. Mol. Cell Biol* 2021, 22, 165–182. [PubMed: 32873929]
- (53). Alexander KA et al. p53 mediates target gene association with nuclear speckles for amplified RNA expression. *Mol. Cell* 2021, 81, 1666–1681.e6. [PubMed: 33823140]
- (54). Xu S; Lai S-K; Sim DY; Ang W; Li HY; Roca X SRRM2 organizes splicing condensates to regulate alternative splicing. *Nucleic Acids Res* 2022, 50, 8599–8614. [PubMed: 35929045]
- (55). Ilik IA; Malszycki M; Lübke AK; Schade C; Meierhofer D; Akta T Son and srrm2 are essential for nuclear speckle formation. *eLife* 2020, 9, 1–48.
- (56). Liao SE; Regev O Splicing at the phase-separated nuclear speckle interface: A model. *Nucleic Acids Res* 2021, 49, 636–645. [PubMed: 33337476]
- (57). Brangwynne CP; Eckmann CR; Courson DS; Rybarska A; Hoegge C; Gharakhani J; Julicher F; Hyman AA Germline P Granules Are Liquid Droplets That Localize by Controlled Dissolution/Condensation. *Science* 2009, 324, 1729–1732. [PubMed: 19460965]


- (58). Smith J; Calidas D; Schmidt H; Lu T; Rasoloson D; Seydoux G Spatial patterning of P granules by RNA-induced phase separation of the intrinsically-disordered protein MEG-3. *eLife* 2016, 5, 1–18.
- (59). Putnam A; Cassani M; Smith J; Seydoux G granules in *Caenorhabditis elegans* embryos. *Nat. Struct. Mol. Biol* 2019, 26.
- (60). Lu T; Spruijt E Multiphase Complex Coacervate Droplets. *J. Am. Chem. Soc* 2020, 142, 2905–2914. [PubMed: 31958956]
- (61). Hoffmann S; Kedersha N; Anderson P; Ivanov P Molecular mechanisms of stress granule assembly and disassembly. *Biochim. Biophys. Acta Mol. Cell Res* 2021, 1868, 118876. [PubMed: 33007331]
- (62). Niewidok B; Igaev M; da Graca AP; Strassner A; Lenzen C; Richter CP; Piehler J; Kurre R; Brandt R Single-molecule imaging reveals dynamic biphasic partition of RNA-binding proteins in stress granules. *J. Cell Biol* 2018, 217, 1303–1318. [PubMed: 29463567]
- (63). Wheeler JR; Matheny T; Jain S; Abrisch R; Parker R Distinct stages in stress granule assembly and disassembly. *eLife* 2016, 5, 1–25.
- (64). Choi S; Meyer MC; Bevilacqua PC; Keating CD Phase-specific RNA accumulation and duplex thermodynamics in multiphase coacervate models for membraneless organelles. *Nat. Chem* 2022, 14.
- (65). West JA; Mito M; Kurosaka S; Takumi T; Tanegashima C; Chujo T; Yanaka K; Kingston RE; Hirose T; Bond C; Fox A; Nakagawa S Structural, super-resolution microscopy analysis of paraspeckle nuclear body organization. *J. Cell Biol* 2016, 214, 1–14.
- (66). Yu H; Lu S; Gasior K; Singh D; Vazquez-Sanchez S; Tapia O; Toprani D; Beccari MS; Yates JR; Da Cruz S; Newby JM; Lafarga M; Gladfelter AS; Villa E; Cleveland DW HSP70 chaperones RNA-free TDP-43 into anisotropic intranuclear liquid spherical shells. *Science* 2021, 371.
- (67). Gall JG; Bellini M; Wu Z; Murphy C Assembly of the nuclear transcription and processing machinery: Cajal bodies (coiled bodies) and transcriptosomes. *Mol. Biol. Cell* 1999, 10, 4385–4402. [PubMed: 10588665]
- (68). Suzuki H et al. The 3' Pol II pausing at replication-dependent histone genes is regulated by Mediator through Cajal bodies' association with histone locus bodies. *Nat. Commun* 2022, 13, 2905. [PubMed: 35614107]
- (69). Feric M; Sarfallah A; Dar F; Temiakov D; Pappu RV; Misteli T Mesoscale structure – function relationships in mitochondrial transcriptional condensates. *Proc. Natl. Acad. Sci. U.S.A* 2022, 119, e2207303119. [PubMed: 36191226]
- (70). Feric M; Demarest TG; Tian J; Croteau DL; Bohr VA; Misteli T Selfassembly of multi-component mitochondrial nucleoids via phase separation. *EMBO J* 2021, 40, e107165. [PubMed: 33619770]
- (71). Shin Y; Brangwynne CP Liquid phase condensation in cell physiology and disease. *Science* 2017, 357.
- (72). Mountain GA; Keating CD Formation of Multiphase Complex Coacervates and Partitioning of Biomolecules within them. *Biomacromolecules* 2020, 21, 630–640. [PubMed: 31743027]
- (73). Flory PJ Thermodynamics of high polymer solutions. *J. Chem. Phys* 1942, 10, 51.
- (74). Helfand E; Tagami Y Theory of the interface between immiscible polymers. *J. Chem. Phys* 1972, 56, 3592.
- (75). Roe RJ Theory of the interface between polymers or polymer solutions. I. Two components system. *J. Chem. Phys* 1975, 62, 490–499.
- (76). Boeynaems S; Holehouse AS; Weinhardt V; Kovacs D; Van Lindt J; Larabell C; Bosch LVD; Das R; Tompa PS; Pappu RV; Gitler AD Spontaneous driving forces give rise to protein-RNA condensates with coexisting phases and complex material properties. *Proc. Natl. Acad. Sci. U.S.A* 2019, 116, 7889–7898. [PubMed: 30926670]
- (77). Fisher RS; Elbaum-Garfinkle S Tunable multiphase dynamics of arginine and lysine liquid condensates. *Nat. Commun* 2020, 11.
- (78). Urry DW Physical chemistry of biological free energy transduction as demonstrated by elastic protein-based polymers. *J. Phys. Chem. B* 1997, 101, 11007–11028.
- (79). Rauscher S; Pomès R The liquid structure of elastin. *eLife* 2017, 6, e26526. [PubMed: 29120326]

- (80). Reichheld SE; Muiznieks LD; Keeley FW; Sharpe S Direct observation of structure and dynamics during phase separation of an elastomeric protein. *Proc. Natl. Acad. Sci. U.S.A* 2017, 114, E4408–E4415. [PubMed: 28507126]
- (81). Simon JR; Carroll NJ; Rubinstein M; Chilkoti A; López GP Programming molecular self-assembly of intrinsically disordered proteins containing sequences of low complexity. *Nat. Chem* 2017, 9, 509–515. [PubMed: 28537592]
- (82). Latham AP; Zhang B On the stability and layered organization of protein-DNA condensates. *Biophys. J* 2022, 121, 1727–1737. [PubMed: 35364104]
- (83). Sawle L; Ghosh K A theoretical method to compute sequence dependent configurational properties in charged polymers and proteins. *J. Chem. Phys* 2015, 143.
- (84). Regy RM; Dignon GL; Zheng W; Kim YC; Mittal J Sequence dependent phase separation of protein-polynucleotide mixtures elucidated using molecular simulations. *Nucleic Acids Res* 2020, 48, 12593–12603. [PubMed: 33264400]
- (85). Harmon TS; Holehouse AS; Pappu RV Differential solvation of intrinsically disordered linkers drives the formation of spatially organized droplets in ternary systems of linear multivalent proteins. *New J. Phys* 2018, 20.
- (86). Sanchez-Burgos I; Espinosa JR; Joseph JA; Collepardo-Guevara R Valency and binding affinity variations can regulate the multilayered organization of protein condensates with many components. *Biomolecules* 2021, 11, 1–19.
- (87). Latham AP; Zhang B Maximum Entropy Optimized Force Field for Intrinsically Disordered Proteins. *J. Chem. Theory Comput* 2020, 16, 773–781. [PubMed: 31756104]
- (88). Latham AP; Zhang B Consistent Force Field Captures Homologue-Resolved HP1 Phase Separation. *J. Chem. Theory Comput* 2021, 17, 3134–3144. [PubMed: 33826337]
- (89). Latham AP; Zhang B Unifying coarse-grained force fields for folded and disordered proteins. *Curr. Opin. Struct. Biol* 2022, 72, 63–70. [PubMed: 34536913]
- (90). Leicher R; Osunsade A; Chua GNL; Faulkner SC; Latham AP; Watters JW; Nguyen T; Beckwitt EC; Christodoulou-rubalcava S; Young PG; Zhang B; David Y; Liu S Single-stranded nucleic acid binding and coacervation by linker histone H1. *Nat. Struct. Mol. Biol* 2022, 29, 463–471. [PubMed: 35484234]
- (91). Pal T; Wessén J; Das S; Chan HS Subcompartmentalization of polyampholyte species in organelle-like condensates is promoted by charge-pattern mismatch and strong excluded-volume interaction. *Phys. Rev. E* 2021, 103, 042406. [PubMed: 34005864]
- (92). Das RK; Pappu RV Conformations of intrinsically disordered proteins are influenced by linear sequence distributions of oppositely charged residues. *Proc. Natl. Acad. Sci. U.S.A* 2013, 110, 13392–13397. [PubMed: 23901099]
- (93). González-Mozuelos P; Olvera De La Cruz M Random phase approximation for complex charged systems: Application to copolyelectrolytes (polyampholytes). *J. Chem. Phys* 1994, 100, 507–517.
- (94). Amin AN; Lin YH; Das S; Chan HS Analytical Theory for Sequence-Specific Binary Fuzzy Complexes of Charged Intrinsically Disordered Proteins. *J. Phys. Chem. B* 2020, 124, 6709–6720. [PubMed: 32639157]
- (95). Das S; Eisen A; Lin YH; Chan HS A Lattice Model of Charge-Pattern-Dependent Polyampholyte Phase Separation. *J. Phys. Chem. B* 2018, 122, 5418–5431. [PubMed: 29397728]
- (96). Holehouse AS; Ginell GM; Griffith D; Böke E Clustering of Aromatic Residues in Prion-like Domains Can Tune the Formation, State, and Organization of Biomolecular Condensates. *Biochemistry* 2021,
- (97). Patel A et al. A Liquid-to-Solid Phase Transition of the ALS Protein FUS Accelerated by Disease Mutation. *Cell* 2015, 162, 1066–1077. [PubMed: 26317470]
- (98). Garaizar A; Espinosa JR; Joseph JA; Collepardo-Guevara R Kinetic interplay between droplet maturation and coalescence modulates shape of aged protein condensates. *Sci. Rep* 2022, 12, 4390. [PubMed: 35293386]
- (99). Garaizar A; Espinosa JR; Joseph JA; Krainer G; Shen Y; Knowles TP; Collepardo-Guevara R Aging can transform single-component protein condensates into multiphase architectures. *Proc. Natl. Acad. Sci. U.S.A* 2022, 119, 1–11.


- (100). Kaur T; Raju M; Alshareedah I; Davis RB; Potoyan DA; Banerjee PR Sequence-encoded and composition-dependent protein-RNA interactions control multiphasic condensate morphologies. *Nat. Commun* 2021, 12.
- (101). Pyo AG; Zhang Y; Wingreen NS Surface tension and super-stoichiometric surface enrichment in two-component biomolecular condensates. *iScience* 2022, 25, 103852. [PubMed: 35198903]
- (102). Qi Y; Reyes A; Johnstone SE; Aryee MJ; Bernstein BE; Zhang B Data-driven Polymer Model for Mechanistic Exploration of Diploid Genome Organization. *Biophys. J* 2020, 1–28.
- (103). Kamat K; Qi Y; Wang Y; Ma J; Zhang B Genome Compartmentalization with Nuclear Landmarks: Random yet Precise. *bioRxiv* 2021, 2021.11.12.468401.
- (104). Jiang Z; Zhang B Theory of Active Chromatin Remodeling. *Phys. Rev. Lett* 2019, 123, 208102. [PubMed: 31809105]
- (105). Jiang Z; Zhang B On the role of transcription in positioning nucleosomes. *PLOS Comput. Biol* 2021, 17, e1008556. [PubMed: 33417594]
- (106). Jiang Z; Qi Y; Kamat K; Zhang B Phase Separation and Correlated Motions in Motorized Genome. *J. Phys. Chem. B* 2022,
- (107). Kashchiev D *Nucleation*; Elsevier, 2000.
- (108). McGaughey A; Ward C Droplet stability in a finite system: Consideration of the solid–vapor interface. *J. Appl. Phys* 2003, 93, 3619–3626.
- (109). Brangwynne CP; Mitchison TJ; Hyman AA Active liquid-like behavior of nucleoli determines their size and shape in *Xenopus laevis* oocytes. *Proc. Natl. Acad. Sci. U.S.A* 2011, 108, 4334–4339. [PubMed: 21368180]
- (110). Grosch M; Ittermann S; Shaposhnikov D; Drukker M Chromatin-Associated Membraneless Organelles in Regulation of Cellular Differentiation. *Stem Cell Rep* 2020,
- (111). Spector DL; Lamond AI Nuclear speckles. *Cold Spring Harb. Perspect. Biol* 2011, 3, a000646. [PubMed: 20926517]
- (112). Sanchez-Burgos I; Joseph JA; Collepardo-Guevara R; Espinosa JR Size conservation emerges spontaneously in biomolecular condensates formed by scaffolds and surfactant clients. *Sci. Rep* 2021, 11, 1–10. [PubMed: 33414495]
- (113). Garcia-Jove Navarro M; Kashida S; Chouaib R; Souquere S; Pierron G; Weil D; Gueroui Z RNA is a critical element for the sizing and the composition of phase-separated RNA–protein condensates. *Nat. Commun* 2019, 10, 1–13. [PubMed: 30602773]
- (114). Qi Y; Zhang B Chromatin Network Retards Droplet Coalescence. *Nat. Commun* 2021, 6824. [PubMed: 34819511]
- (115). Sanders DW et al. Competing Protein-RNA Interaction Networks Control Multiphase Intracellular Organization. *Cell* 2020, 181, 306–324.e28. [PubMed: 32302570]
- (116). Yang P; Mathieu C; Kolaitis RM; Zhang P; Messing J; Yurtsever U; Yang Z; Wu J; Li Y; Pan Q; Yu J; Erik W; Mittag T; Kim HJ; Taylor JP G3BP1 is a tunable switch that triggers phase separation to assemble stress granules. *Cell* 2021, 181, 325–345.
- (117). Laghmach R; Alshareedah I; Pham M; Raju M; Banerjee PR; Potoyan DA RNA chain length and stoichiometry govern surface tension and stability of protein-RNA condensates. *iScience* 2022, 25, 104105. [PubMed: 35378855]
- (118). Yang Y; Fang Z; Chen X; Zhang W; Xie Y; Chen Y; Liu Z; Yuan W An overview of pickering emulsions: Solid-particle materials, classification, morphology, and applications. *Front. Pharmacol* 2017, 8, 1–20. [PubMed: 28149278]
- (119). Welsh TJ; Krainer G; Espinosa JR; Joseph JA; Sridhar A; Jahnel M; Arter WE; Saar KL; Alberti S; Collepardo-Guevara R; Knowles TP Surface Electrostatics Govern the Emulsion Stability of Biomolecular Condensates. *Nano Lett* 2022, 22, 612–621. [PubMed: 35001622]
- (120). Brangwynne CP; Mitchison TJ; Hyman AA Active liquid-like behavior of nucleoli determines their size and shape in *Xenopus laevis* oocytes. *Proc. Natl. Acad. Sci. U.S.A* 2011, 108, 4334–4339. [PubMed: 21368180]
- (121). Zhang Y; Lee DS; Meir Y; Brangwynne CP; Wingreen NS Mechanical frustration of phase separation in the cell nucleus by chromatin. *Phys. Rev. Lett* 2021, 126, 258102. [PubMed: 34241518]

- (122). Style RW; Sai T; Fanelli N; Ijavi M; Smith-Mannschott K; Xu Q; Wilen LA; Dufresne ER Liquid-Liquid Phase Separation in an Elastic Network. *Phys. Rev. X* 2018, 8, 11028.
- (123). Rosowski KA; Sai T; Vidal-Henriquez E; Zwicker D; Style RW; Dufresne ER Elastic ripening and inhibition of liquid–liquid phase separation. *Nat. Phys* 2020, 16, 422–425. [PubMed: 32273899]
- (124). Lee DS; Wingreen NS; Brangwynne CP Chromatin mechanics dictates sub-diffusion and coarsening dynamics of embedded condensates. *Nat. Phys* 2021, 17, 531–538.
- (125). Zidovska A Chromatin: Liquid or Solid? *Cell* 2020, 183, 1737–1739. [PubMed: 33357397]
- (126). Wurtz JD; Lee CF Chemical-Reaction-Controlled Phase Separated Drops: Formation, Size Selection, and Coarsening. *Phys. Rev. Lett* 2018, 120, 78102.
- (127). Söding J; Zwicker D; Sohrabi-Jahromi S; Boehning M; Kirschbaum J Mechanisms for Active Regulation of Biomolecular Condensates. *Trends Cell Biol* 2020, 30, 4–14. [PubMed: 31753533]
- (128). Guilhas B; Walter JC; Rech J; David G; Walliser NO; Palmeri J; Mathieu-Demaziere C; Parmeggiani A; Bouet JY; Le Gall A; Nollmann M ATP-Driven Separation of Liquid Phase Condensates in Bacteria. *Mol. Cell* 2020, 79, 293–303.e4. [PubMed: 32679076]
- (129). Wang JT; Smith J; Chen BC; Schmidt H; Rasoloson D; Paix A; Lambrus BG; Calidas D; Betzig E; Seydoux G Regulation of RNA granule dynamics by phosphorylation of serine-rich, intrinsically disordered proteins in *C. elegans*. *eLife* 2014, 3, 1–23.
- (130). Wurtz JD; Lee CF Stress granule formation via ATP depletion-Triggered phase separation. *New J. Phys* 2018, 20.
- (131). Wippich F; Bodenmiller B; Trajkovska MG; Wanka S; Aebersold R; Pelkmans L Dual specificity kinase DYRK3 couples stress granule condensation/dissolution to mTORC1 signaling. *Cell* 2013, 152, 791–805. [PubMed: 23415227]


A) Nucleolous




B) Nuclear speckles




D) Stress granule



C) P granule

**Figure 1:**

Examples of biological condensates with a layered organization. A, left) Schematic diagram of subcompartments within the nucleolus. A, right) Nucleoli within an untreated *X. laevis* nucleus. NPM1 (red), FIB1 (green) and POLR1E (blue) are tagged. Scale bar, 20 μm . Image modified from³¹ with permission from Elsevier. B) Organization of two components within nuclear speckles, the protein SON and the snRNA U2B''. Scale bar, 1 μm . Adapted with permission from the Journal of Cell Science.³² C) Photomicrographs examining the *in vivo* assembly of MEG-3:mGFP and PGL-3:mCherry, two main components of P granules. Scale bar, 500 nm. Image minimally modified from.³³ Reprinted with permission from AAAS. D) Stochastic optical reconstruction microscopy (STORM) image of a stress granule (gray), highlighting poly(A)⁺ RNA cores (yellow). Scale bar, 500 nm. Image modified from³⁴ with permission from Elsevier.

**Figure 2:**

Relation between interfacial tension (τ), effective interaction parameter (χ), and condensate organization. A) Mixing two immiscible droplets can lead to four possible organizations: (1) a layered droplet with dropletA on the inside, (2) partially wetted droplets that share an interface (3) complete nonwetting to form two separate droplets, and (4) a layered droplet with dropletB on the inside. The most stable organization is determined by the interfacial tension between dropletA and dropletB (τ_{AB}), between dropletA and the solvent phase (τ_{As}), and between dropletB and the solvent phase (τ_{Bs}). The droplet size (α_A and α_B) can also contribute to the condensate organization, as demonstrated by Lu and Spruijt.⁶⁰ B) For immiscible homopolymers, the Flory-Huggins interaction parameter (χ_A, χ_B) can provide a way to approximate differences in surface tension. C) Flory-Huggins theory can be generalized to heteropolymers by assuming an effective interaction parameter that averages over differences in the sequence ($\chi_A^{eff}, \chi_B^{eff}$). D) Heteropolymers can be divided into segments


with different physical properties, resulting in effective parameters for different portions of the single chain (χ_A^1, χ_B^1).

Author Manuscript

Author Manuscript

Author Manuscript

Author Manuscript

**Figure 3:**

Molecular factors that drive the formation of layered condensates. (A) Hypothetical interaction patterns of polyR, polyK, and UTP that explain the observed condensate organization. (B) Confocal fluorescence images of 50:50 polyK(green):polyR(purple) mixtures at different ratios of UTP. Scale bar, 20 μm . (C) Confocal fluorescence images of fusion of layered coacervates. polyK is labeled in green, and polyR is unlabeled. Images modified from ref.⁷⁷ CC BY 4.0. (D) Approximate interaction patterns of HP1 α , H1, and DNA, which also results in a layered condensate. (E) The slab density profiles support a layered organization for mixtures of HP1 α (blue), H1 (green), and DNA (red). HP1 α coalesces toward the center of the droplet, with H1 to the outside. Image modified from⁸² with permission from Elsevier. (F) Sequence diagram of the native LAF1-RGG sequence (RGG) and the shuffled RGG sequence (RGG_{Cshuf}). Anionic (red) and cationic (blue) amino acids are highlighted. SCD is a measure of charge patterning, where larger, negative values indicate segregated regions of the same charge.⁸³ (G) Potential of mean force (PMF) for protein-protein and protein-RNA interactions with RGG and RGG_{Cshuf}. (H) Simulation


snapshots indicating the preference of A_{15} to localize to the condensate exterior with $RG\bar{G}_{Cshuf}$ but not $RG\bar{G}$. (I) Density profiles of A_{15} in $RG\bar{G}$ and $RG\bar{G}_{Cshuf}$ condensates. Reprinted from ⁸⁴ with permission from Oxford University Press.

Author Manuscript

Author Manuscript

Author Manuscript

Author Manuscript

**Figure 4:**

Molecular factors that limit condensate growth. (A) Computational evidence that lower valency surfactants can limit droplet growth. Surface tension (ρ) depends on the ratio of surfactants (red) to scaffold (blue). Vertical dashed lines indicate the maximum surfactant concentration that allows for phase separation for a given number-droplet regime. Note that the maximum droplet size varies continuously with surfactant concentration even within the same number-droplet regime. Snapshots of simulations in each droplet regime are included. Images modified from ref.¹¹² CC BY 4.0. (B) RNA binding modifies ArtiG size. Confocal images of ArtiG in HeLa cells, 24 hours after transfection of mCherry-FFm and PUM.HD-FFm constructs at ratios of 1:1 (i), 5:1 (ii), 10:1 (iii), and 1:0 (iv). ArtiG^{mCh} indicates ArtiG comprised of mCherry-FFm, while ArtiG^{mCh/PUM} indicates ArtiG comprised of both mCherry-FFm and PUM.HD-FFm. Images modified from ref.¹¹³ CC BY 4.0. (C) Computational modeling shows the presence of an entropic barrier that stabilizes the two-

droplet state for nucleoli. Free energy profile as a function of the radius of gyration, which effectively measures the distance between the two droplets. The free energy is broken into entropic (red) and energetic (black) components before (D) and after (E) the barrier for droplet fusion. Images modified from ref. ¹¹⁴ CC BY 4.0.

Author Manuscript

Author Manuscript

Author Manuscript

Author Manuscript