

UCSF

UC San Francisco Previously Published Works

Title

Chapter Two Residual Immune Dysregulation Syndrome in Treated HIV infection

Permalink

<https://escholarship.org/uc/item/9cj1x4nq>

Authors

Lederman, Michael M
Funderburg, Nicholas T
Sekaly, Rafick P
[et al.](#)

Publication Date

2013

DOI

10.1016/b978-0-12-407707-2.00002-3

Peer reviewed



Published in final edited form as:

Adv Immunol. 2013 ; 119: 51–83. doi:10.1016/B978-0-12-407707-2.00002-3.

Residual Immune Dysregulation Syndrome in Treated HIV infection

Michael M. Lederman^{*,1}, Nicholas T. Funderburg^{*}, Rafick P. Sekaly[†], Nichole R. Klatt[‡], and Peter W. Hunt[§]

^{*}Division of Infectious Diseases and Center for AIDS Research, Case Western Reserve University and University Hospitals/Case Medical Center, Cleveland, Ohio, USA

[†]Vaccine and Gene Therapy Institute, Port St. Lucie, Florida, USA

[‡]Department of Pharmaceutics, WaNPRC, University of Washington, Seattle, Washington, USA

[§]University of California, San Francisco, California, USA

Abstract

Antiretroviral therapy has revolutionized the course of HIV infection, improving immune function and decreasing dramatically the mortality and morbidity due to the opportunistic complications of the disease. Nonetheless, even with sustained suppression of HIV replication, many HIV-infected persons experience a syndrome characterized by increased T cell activation and evidence of heightened inflammation and coagulation. This residual immune dysregulation syndrome or RIDS is more common in persons who fail to increase circulating CD4+ T cells to normal levels and in several epidemiologic studies it has been associated with increased morbidity and mortality. These morbid and fatal events are not the typical opportunistic infections and malignancies seen in the early AIDS era but rather comprise a spectrum of cardiovascular events, liver disease, metabolic disorders, kidney disease, bone disease, and a spectrum of malignant complications distinguishable from the opportunistic malignancies that characterized the earlier days of the AIDS epidemic.

While immune activation, inflammation, and coagulopathy are characteristic of untreated HIV infection and improve with drug-induced control of HIV replication, the drivers of RIDS in treated HIV infection are incompletely understood. And while inflammation, immune activation, and coagulopathy are more common in treated persons who fail to restore circulating CD4+ T cells, it is not entirely clear how these two phenomena are linked.

1. THE ROLE OF ACTIVATION AND INFLAMMATION IN THE NATURAL HISTORY OF INFECTION

The earliest reports of AIDS were prescient in recognizing that profound depletion of circulating CD4+ T cells was central to the immune deficiency that defined the syndrome (Gottlieb et al., 1981; Masur et al., 1981). These investigators also recognized that despite

profound immune deficiency, activation of T cells (Gottlieb et al., 1981) and B cells (Lane et al., 1983) was also characteristic of this syndrome. A number of reports linked phenotypic indices of T cell activation, primarily defined by the expression of CD38 or CD38 and HLA-DR to the risk of disease progression in natural history studies (Giorgi et al., 1999, 1993, 2002). The importance of immune activation in the pathogenesis of disease is also underscored by the recognition that African nonhuman primates that have coexisted with endemic infection with related simian immunodeficiency viruses generally tolerate high-level SIV replication without progressive immune deficiency or opportunistic infections (Silvestri, Paiardini, Pandrea, Lederman, & Sodora, 2007). In these animals, immune activation is transient after acute infection and is typically brought under control (Harris et al., 2010). In contrast, Asian rhesus macaques infected experimentally with SIV that is nonpathogenic in African animals experience a pathogenic process that is characterized by both progressive immune deficiency and sustained high levels of systemic immune activation (Silvestri et al., 2007).

Early events in both pathogenic SIV infection and HIV infection are typically associated with high levels of virus in plasma that in time diminish to a lower steady state level. This is associated with a modest decrease in numbers of circulating CD4+ T cells that in most persons progressively decline in the absence of effective antiretroviral therapy (ART). But the effects of acute HIV infection on circulating CD4+ T cell numbers are very poor reflections of pathologic events elsewhere in the body. In both pathogenic SIV infection and HIV infection of humans, acute infection is associated with “catastrophic” infection and depletion of mucosal CD4+ T cells that are typically enriched with cells coexpressing the virus chemokine receptor CCR5 that is used by the vast majority of all SIV and HIV viruses that are transmitted. This gut mucosal CD4+ T cell depletion may have a profound impact on the determinants of pathogenesis and persistent inflammation as discussed in the sections below.

As noted above, broad systemic immune activation is characteristic of untreated HIV infection. A robust activation and inflammatory state are seen during acute infection (Stacey et al., 2009), which is often reflected clinically as a “viral syndrome” with fever, rash, lymphadenopathy, and occasionally an aseptic meningitis (Schacker, Collier, Hughes, Shea, & Corey, 1996; Tindall et al., 1988). These clinical manifestations typically resolve and markers of inflammation attenuate but still remain relatively and persistently elevated. Thus, laboratory indices of T cell activation and B cell activation, activation of monocytes and dendritic cells (DCs), and natural killer cells are recognized concomitants of untreated infection. Likewise, inflammatory and coagulation indices are typically and persistently elevated during untreated infection (Funderburg et al., 2010, 2012). With application of antiretroviral therapies, and as circulating CD4+ T cell numbers rise, these markers improve; yet in many, and especially among those who fail to restore circulating CD4+ T cell numbers, persistent immune activation, inflammation, and coagulation abnormalities persist (Lederman et al., 2011) and are the subject of this review. With ART-induced suppression of HIV replication, circulating CD4+ T cell numbers typically increase in a biphasic pattern. An early first-phase increase that includes all lymphocyte subpopulations is likely the consequence of rapid redistribution of cells from lymph nodes (LNs) where they were likely sequestered in the inflammatory environment that accompanies untreated infection (Autran

et al., 1997; Biancotto et al., 2007; Bucy et al., 1999). It is plausible to attribute much of the activation/inflammation and coagulation abnormalities in untreated infection directly to HIV replication, especially as therapy-induced control of HIV replication improves (but typically does not normalize) these activation and coagulation abnormalities. Viral elements are recognized by innate immune sensors (Beignon et al., 2005; Heil et al., 2004) and *in vitro*, ligation of these sensors by viral elements can drive innate immune cell activation directly and T cell activation indirectly (Funderburg et al., 2008; Meier et al., 2007). How much of this activation is mediated by cognate peptide recognition is less clear. On the one hand, chronic HIV infection is characterized by profound and persistent expansion of circulating CD8 T cell numbers, yet few of these cells can be shown to be HIV reactive (Betts et al., 2001). Proportionally more are demonstrably reactive with peptides derived from cytomegalovirus (Naeger et al., 2010) and it can be argued that sustained replication of this and other intracellular pathogens promoted by a failure of host defenses may underlie the dramatic expansion of CD8 T cells seen in chronic HIV infection. On the other hand, the relative expansion of these cells persists long after ART controls HIV replication and risk for opportunistic infections subsides. The mechanisms underlying this sustained CD8 T cell expansion are not understood.

2. INFLAMMATORY AND COAGULATION INDICES PREDICT MORBIDITY IN TREATED HIV INFECTION

Despite dramatic improvements in the modern treatment era, HIV-infected persons maintaining ART-mediated viral suppression continue to have increased mortality compared to the general population, particularly if ART is delayed until late in the disease course (Lewden et al., 2012, 2007; The Antiretroviral Therapy Cohort Collaboration, 2008; Lohse et al., 2007; van Sighem, Gras, Reiss, Brinkman, & de Wolf, 2010). These morbidities include cardiovascular disease (Freiberg et al., 2013; Triant, Lee, Hadigan, & Grinspoon, 2007; Tseng et al., 2012), cancer (Deeken et al., 2012; Grulich, van Leeuwen, Falster, & Vajdic, 2007; Patel et al., 2008; Silverberg et al., 2009), osteoporosis (Brown & Qaqish, 2006; Triant, Brown, Lee, & Grinspoon, 2008), neurocognitive dysfunction (Heaton et al., 2010), type II diabetes (Brown et al., 2005), thromboembolic disease (Copur, Smith, Gomez, Bergman, & Homel, 2002; Fultz, McGinnis, Skanderson, Ragni, & Justice, 2004; Sullivan, Dworkin, Jones, & Hooper, 2000), and even frailty (Desquilbet et al., 2007, 2009), a syndrome of multimorbidity typically only seen in geriatric populations. While health-related behaviors and toxicities of antiretroviral drugs may both contribute to these risks, several recent studies highlight the contribution of immune dysfunction and inflammation to morbidity and mortality in this setting.

3. PERSISTENT CD41+ T CELL LYMPHOPENIA PREDICTS CLINICAL OUTCOMES DURING ART

As discussed above, many HIV-infected individuals fail to recover normal CD4+ T cell counts (i.e., >500 cells/mm³) despite over a decade of sustained ART-mediated viral suppression (Kelley et al., 2009). These persons appear to be at highest risk for subsequent mortality in several large multicenter cohort studies (Lewden et al., 2012; The Antiretroviral

Therapy Cohort Collaboration, 2008). Persistently low CD4+ T cell counts despite ART has been consistently associated with an increased risk for combined non-AIDS morbidity and mortality in several large cohort studies and clinical trials in North America and Europe (Achhra et al., 2010; Baker et al., 2008; Marin et al., 2009; Mocroft et al., 2010; Smurzynski et al., 2010). Interestingly, there continues to be an incremental decrease in mortality risk with increasing CD4+ T cell counts even among those with CD4+ T cell counts >500 cells/mm³, suggesting that continued CD4+ T cell recovery—even after recovering “normal” levels—may continue to confer clinical benefit (Young et al., 2012). Several studies have also linked persistent CD4+ T cell lymphopenia to an increased risk of specific non-AIDS morbidities including cardiovascular disease (Lichtenstein et al., 2010; Triant et al., 2010), osteoporosis, and fracture risk (Yong, Elliott, Woolley, & Hoy, 2011). Persistently low CD4+ T cell counts despite ART also appear to predict several infection-related cancers (Grulich et al., 2007; Silverberg et al., 2009). While all of these chronic morbidities appear to be linked to poor ART-mediated CD4+ T cell recovery, HIV-associated neurocognitive disorders and thromboembolic disease appear to be predicted by lower pre-ART nadir—but not current—CD4+ T cell counts, suggesting that the degree or pre-ART immunodeficiency might be a more important determinant of certain morbidities than the degree of ART-mediated CD4+ T cell restoration (Ellis et al., 2011; Musselwhite et al., 2011). Collectively, these studies suggest that the degree of pre-ART immunodeficiency and the extent of CD4+ T cell recovery during ART are important predictors of morbidity and mortality in treated HIV disease.

4. IMMUNE ACTIVATION/INFLAMMATION PREDICTS MORBIDITY AND MORTALITY DURING ART

While the degree of CD4+ T cell lymphopenia is an important predictor of morbidity and mortality in treated HIV infection, the extent of persistent immune activation and inflammation have emerged as perhaps even stronger—and independent—predictors of morbidity and mortality in this setting. For example, in a recent nested case-control study within the Strategies for Management of Antiretroviral Therapy (SMART) trial, higher plasma levels of the inflammatory cytokines IL-6 and C-reactive protein (CRP) and the coagulation marker D-dimer strongly predicted higher overall mortality and cardiovascular events (Duprez et al., 2009; Kuller et al., 2008). A key finding from this study was that the degree to which these inflammatory markers predicted mortality in treated HIV infection was much stronger than that observed in older HIV-uninfected individuals (Harris et al., 1999; Reuben et al., 2002; Volpato et al., 2001; Wikby et al., 2006), consistent with the hypothesis that chronic inflammation is a much more important determinant of mortality in the context of HIV infection than it is for the general population.

Several more recent studies have also demonstrated strong associations between inflammatory and coagulation markers and subsequent morbidity and mortality in treated HIV infection. For example, higher fibrinogen and CRP levels predicted increased mortality after adjustment for plasma HIV RNA levels and CD4+ T cell counts in a largely treated group of HIV-infected individuals in the Fat Redistribution and Metabolic Change in HIV Infection (FRAM) cohort (Tien et al., 2010). Importantly, these bio-markers continued to

predict mortality even when restricting to HIV-infected individuals with CD4+ T cell counts >500 cells/mm³, suggesting that inflammation may be an important driver of morbidity and mortality even for those with optimal CD4+ T cell recovery. Similarly, a nested case-control study from two AIDS Clinical Trials Group (ACTG) trials of HIV-infected individuals starting ART assessed the impact of pre-ART immune activation and inflammatory markers on subsequent AIDS and death (Kalayjian et al., 2010). Similarly, a nested case-control study showed that higher pre-ART levels of IL-6, CRP, D-dimer, and the tissue fibrosis biomarker hyaluronic acid predicted subsequent mortality and AIDS events in individuals with advanced disease prior to ART (Boulware et al., 2011). A consistent finding from all of these published studies—as well as two other recently presented studies (Hunt et al., 2012; Tenorio et al., 2013)—was that inflammatory markers continued to predict subsequent mortality during treated HIV infection even after adjustment for concurrent and pre-ART nadir CD4+ T cell count, suggesting an independent role of the inflammatory state in driving mortality in this setting.

In more mechanistic work from the SMART study, Sandler and colleagues showed that higher plasma levels of the soluble monocyte activation marker sCD14 predict earlier mortality (Sandler et al., 2011), findings later replicated by several other groups (Hunt et al., 2012; Justice et al., 2012; Tenorio et al., 2013). Since CD14 is cleaved from the monocyte cell surface upon lipopolysaccharide (LPS)-mediated monocyte activation, this study provided the first evidence potentially linking microbial translocation to mortality in treated HIV infection. While some have argued that sCD14 elevations may not always reflect LPS-induced activation of monocytes, our own group has recently demonstrated that specific markers of gut epithelial barrier integrity including intestinal fatty acid-binding protein and zonulin-1 levels during ART-mediated viral suppression also independently predict subsequent mortality in individuals who initiated ART with a diagnosis of AIDS (Hunt et al., 2012). These studies support microbial translocation as an important target of interventions in subsequent clinical trials.

Persistent inflammation, monocyte activation, and coagulation during treated HIV infection have also been strongly linked to several specific morbidities in this setting. For example, higher plasma CRP, IL-6, and D-dimer levels predict cardiovascular events in ART-suppressed HIV-infected individuals (Duprez et al., 2009; Ford et al., 2010; Triant, Meigs, & Grinspoon, 2009). As has been established in HIV-uninfected persons, soluble markers of inflammation also appear to predict incident type II diabetes in the context of treated HIV infection, independent of other established risk factors (Brown, Tassiopoulos, Bosch, Shikuma, & McComsey, 2010). Similarly, in a recent nested case-control study from the Multicenter AIDS Cohort Study (MACS), higher levels of inflammatory (including IL-6 and CRP) and B cell stimulatory biomarkers predicted subsequent development of non-central nervous system Non-Hodgkin lymphoma years later, even after adjustment for CD4+ T cell count (Breen et al., 2011). Increased peripheral blood markers of monocyte activation (i.e., sCD14 and sCD163) and inflammation are also associated with HIV-associated neurocognitive dysfunction during ART-mediated viral suppression (Burdo et al., 2013; Letendre et al., 2012; Lyons et al., 2011). Interesting recent work from the Veterans Aging Cohort Study cohort also suggests that higher plasma levels of sCD14, IL-6, and D-dimer were not only independently associated with mortality but also strongly associated with

anemia as well as with greater abnormalities in surrogate markers of liver fibrosis and renal insufficiency (Justice et al., 2012). These data support a model whereby chronic inflammation causes damage to multiple organ systems, all contributing to an increased risk of morbidity and mortality. The concept that inflammation may contribute to the multiple morbidities commonly associated with the aging process was further underscored by a recent study linking inflammatory markers to frailty in treated HIV infection (Erlandson et al., 2013).

While markers of innate immune activation and inflammation strongly predict morbidity and mortality in treated HIV infection, markers of T cell activation are less strongly and consistently associated with clinical outcomes in this setting. While higher frequencies of activated (CD38+ HLA-DR+) CD8+ T cells during early ART-mediated viral suppression predicted subsequent mortality in HIV-infected Ugandans independent of CD4+ T cell count (Hunt, Cao, et al., 2011), and among HIV-infected North Americans in the Study of the Ocular Complications of AIDS (Hunt et al., 2012), these associations were much weaker than those observed for inflammatory, coagulation, and monocyte activation markers. Furthermore, other recent studies failed to identify an independent association between T cell activation and subsequent clinical outcomes during treated HIV infection (Ford et al., 2010; Tenorio et al., 2013).

5. FAILURE TO RESTORE CIRCULATING CD4+ T CELLS IN HIV INFECTION

While most persons treated with ART experience substantial and progressive gains in circulating CD4+ T cell numbers, a significant proportion of treated patients fails to achieve normalization of these counts (Benveniste et al., 2005; Gazzola, Tincati, Bellistri, Monforte, & Marchetti, 2009; Kelley et al., 2009). Not surprisingly, this “immunologic failure” is more common in persons who fail to effectively suppress HIV replication (Gandhi et al., 2006). Yet, even among persons whose treatment regimens decrease HIV levels in plasma below the limits of assay detection, immunologic failure occurs and as expected, is seen more often in persons who start therapy with lower CD4+ T cell numbers (Lederman et al., 2011). Immunologic failure also appears to be more common in older subjects and among men (Gandhi et al., 2006; Lederman et al., 2011). With time and sustained suppression of HIV replication, more immune failure subjects tend to normalize circulating CD4+ T cell counts (Hunt et al., 2003; Kelley et al., 2009), yet a substantial minority of treated persons (in our clinic at University Hospitals Case Medical Center as many as one in four with suppressed viral replication under treatment) will sustain circulating CD4+ T cell numbers below the lower limits of normal (B. Rodriguez and H. Myerson unpublished). In these immune failure subjects, immune activation, inflammation, and coagulation abnormalities persist and are demonstrable at levels higher than those seen in healthy subjects and among HIV-infected subjects who experience “normalization” of blood CD4 T cell numbers (Lederman et al., 2011). In these subjects, both CD4+ and CD8+ T cells coexpress activation markers such as CD38 and HLA-DR, yet even though both CD4+ and CD8+ T cells are activated, only CD4+ T cells are more often demonstrable in cell cycle and these are primarily memory-phenotype cells. An apparent failure of T cell maturation is suggested in this setting by the diminished numbers of circulating CD4+ and CD8+ naïve T cells (Lederman et al., 2011). This is likely mediated in part by diminished T cell expression of the IL-7 receptor alpha

chain CD127 (Colle et al., 2006; Marziali et al., 2006) as well as by a suggested failure of reliable T cell access to IL-7 in lymphoid tissues as a result of fibrosis and architectural distortion of the fibroblast reticular network (Zeng et al., 2011). At the same time, immune failure patients sustain high systemic levels of inflammation and coagulation with increased expression of interferon-stimulated genes (Fernandez et al., 2011) and increased plasma levels of IL-6, sCD14, and D-dimers (Lederman et al., 2011), and not surprisingly, immune failure subjects appear at increased risk for the non-AIDS morbidities now seen in treated HIV infection as discussed above. Thus, immune failure and inflammation are linked and plausibly both are the consequence of concurrent LN fibrosis and sustained damage to the gut as outlined directly below.

6. THE LN IN TREATED HIV INFECTION

LNs were identified as major sites of viral replication in HIV-infected subjects (Pantaleo et al., 1991, 1993). A number of studies in humans and nonhuman primates have reported higher frequencies of HIV/SIV-infected cells as well as higher copy numbers of viral transcripts in CD4⁺ T cells isolated from lymphoid tissues (GALT, LN) when compared to the peripheral blood (Chun et al., 1997; Fox, Kotler, Tierney, Wilson, & Fauci, 1989; Kiviat et al., 1998; Pantaleo et al., 1991; Zuckerman et al., 2004). Several features of lymphoid tissues provide a favorable environment for HIV replication. They include (1) a privileged tissue architecture that favors close cellular contact between immune cells, thereby promoting cell-to-cell transmission of HIV and ensuring viral dissemination; (2) a significant enrichment in the frequency of cells that are highly permissive to HIV infection, such as activated CD4⁺ T cells that can produce large numbers of viral particles; and (3) a proinflammatory environment (Andersson et al., 1998; Biancotto et al., 2007) that enhances viral production from infected cells and promotes new infections. These factors contribute to the high levels of HIV replication observed in lymphoid organs from HIV-infected subjects and provide an explanation for the major role played by these compartments in the pathophysiology of HIV infection. Early in infection, there is evidence that this inflammatory environment promotes the accumulation of T regulatory (Treg) cells and the induction of collagen synthesis as a result of TGF β expression (Estes et al., 2007). This increase in collagen deposition and resultant fibrosis is thought to disrupt the normal architecture of the fibroblastic reticular network that is responsible for trafficking of cells cytokines and antigen through the LN T cell zone and perhaps limiting T cell access to the homeostatic cytokine IL-7 (Zeng et al., 2011). These structural abnormalities are typically not repaired with ART-induced suppression of HIV replication and are related inversely both to the numbers of circulating CD4⁺ T cells and their restoration with ART (Schacker, Nguyen, Beilman, et al., 2002; Schacker, Nguyen, Martinez, et al., 2002). A biphasic restoration of circulating CD4⁺ T cells is typically seen with application of ART (Autran et al., 1997; Lederman et al., 1998; Pakker et al., 1998). The first rapid phase seen in the first few weeks of therapy is thought to be a consequence of systemic redistribution of sequestered lymphocytes from inflamed and activated lymphoid tissues as viral replication is halted (Bucy et al., 1999) while the second slower phase is thought to represent homeostatic CD4⁺ T cell restoration. We have recently found that LN T cells in untreated HIV infection have an impaired responsiveness to sphingosine-1 phosphate, the phospholipid that mediates

chemotactic egress of lymphocytes from lymphoid tissues (Mudd et al., 2013). This defect can be induced in vitro by both T cell receptor engagement and by exposure to type 1 interferons as well as to a variety of microbial toll-like receptor agonists (Mudd et al., 2013). Treatment with antiretroviral drugs largely (but not completely) corrects this defect thereby providing a mechanistic explanation for the rapid systemic first-phase CD4+ T cell restoration with ART.

7. THE GUT IN TREATED HIV INFECTION

The mucosal immune system is dramatically affected during HIV infection. Massive damage to the tight epithelial barrier of the GI tract is linked to microbial translocation and severe immunological dysfunction (Brenchley & Douek, 2008; Klatt, Funderburg, & Brenchley, 2013). Long-term ART partially restores this mucosal damage and dysfunction; however, complete restoration does not often occur. One of the most notable immune abnormalities during acute HIV infection is a rapid and severe depletion of CD4+ T cells from mucosal tissues (Brenchley, Price, & Douek, 2006; Brenchley et al., 2004; Mattapallil et al., 2005; Veazey et al., 1998). While long-term therapy usually induces reconstitution of CD4+ T cells in systemic circulation, depletion is typically maintained in mucosal tissues throughout treatment (Marchetti et al., 2008; Mavigner et al., 2012; Mehandru et al., 2006; Tincati et al., 2009). Initially after infection, CD4+ T cells are profoundly depleted in mucosal tissues, likely due to direct infection by HIV (Brenchley, Price, & Douek, 2006; Brenchley et al., 2004; Mattapallil et al., 2005; Picker, 2006). The selective targeting of the GI tract is likely because the majority of mucosal CD4+ T cells express the HIV coreceptor CCR5 and activated memory cells that the virus infects preferentially (Douek et al., 2002; Mattapallil et al., 2005). There is also evidence that the gut homing integrin $\alpha 4\beta 7$ is also used by HIV to enhance cellular susceptibility to infection (Arthos et al., 2008). Nonetheless, and perhaps in part because of this, complete CD4+ T cell restoration in the gut rarely occurs despite clinically effective suppression of HIV replication by ART (Brenchley & Douek, 2008; Macal et al., 2008). This also may be due, in part, to incomplete suppression of virus replication in gastrointestinal tissues, permitting residual HIV replication and continued CD4+ T cell infection. Persistent damage to the gut mucosal barrier is also a likely important contributor to the failure of CD4+ T cell restoration in the GI tract during ART.

During HIV infection, there is significant damage to the tight epithelial barrier of the GI tract, leading to breaches that allow microbial products to penetrate the lamina propria (Estes et al., 2010; Klatt et al., 2010; Nazli et al., 2010). This damage not only contributes to microbial translocation but also is associated with loss of essential immunoregulatory cells such as CD103+ DCs and impaired production of immunoregulatory cytokines such as IL-17 and IL-22 by T cells and innate lymphocytes (Klatt, Estes et al., 2012). While damage to the structural barrier has not been well studied in the context of ART, it is clear that the associated immune dysfunctions persist despite treatment-induced virus suppression. Indeed, loss of IL-17 producing CD4+ T cells (Th17 cells) in the GI tract has been associated with increased immune activation and decreased CD4+ T cell restoration, despite virologic control on ART (Gordon et al., 2010; Klatt & Brenchley, 2010; Mavigner et al., 2012). Furthermore, the ratio of Th17 to Treg cells remains decreased after treatment, an imbalance

that may drive elevated indoleamine 2,3-dioxygenase (IDO) production by DCs, further potentiating immune activation, decreasing T cell proliferation, and impairing production of IL-17 and IL-22 (Favre et al., 2010; Reeves et al., 2011). Overall, antigen presenting cell dysfunction persists despite virus suppression on therapy. DCs may contribute to immune dysfunction via IDO production, via altered cytokine production, and by altered mucosal homing (Chehimi et al., 2002; Favre et al., 2010; Jasny et al., 2012; Miller et al., 2012; Reeves et al., 2012). And despite growing evidence that treated HIV infection is characterized by an increased inflammatory signature, the role of the innate immune system in mucosal immune dysfunction during ART is incompletely studied. For example, while monocytes and mucosal macrophages are clearly dysfunctional even in treated HIV infection (Klatt, Funderburg, et al., 2013; Lichtfuss et al., 2011), their role in HIV persistence and in immune activation and microbial translocation remains unclear.

Chronic mucosal dysfunction and immune activation may be a result of persistent microbial translocation despite ART. During HIV infection, mucosal damage allows microbial products to translocate across the tight epithelial barrier of the gut into the lamina propria and eventually into circulation (Brenchley, Price, Schacker, et al., 2006; Klatt, Funderburg, et al., 2013). Although most work to date is correlative, we suspect that persistent microbial translocation is an important driver of chronic, pathological immune activation in HIV infection (Brenchley & Douek, 2008, 2012; Brenchley, Price, Schacker, et al., 2006; Klatt, Funderburg, et al., 2013). Recent work in the nonhuman primate model indicates that during acute SIV infection, blockade of microbial translocation with the LPS-binding resin sevelamer decreases both systemic microbial translocation as well as systemic indices of both T cell activation and inflammation (Pandrea, 2013). And while microbial translocation is reduced with suppressive ART, most studies find elevated levels of these products in circulation in these subjects when compared to levels found in uninfected persons (Chege et al., 2011; Klatt, Funderburg, et al., 2013; Lederman et al., 2011; Lichtfuss et al., 2011; Merlini et al., 2011; Pilakka-Kanthikeel et al., 2012). This enduring microbial translocation may well underlie the persistent immune activation and inflammatory sequelae that occur despite treatment and are associated with increased morbidity and mortality in treated patients (Kuller et al., 2008; Lichtfuss et al., 2011; Rodger et al., 2009; Sandler et al., 2011).

The mechanisms that underlie mucosal immune barrier dysfunction are still poorly understood. While virus replication likely plays a role, particularly early in infection, the persistent mucosal dysfunction during ART when virus replication is effectively suppressed suggests that factors distinguishable from viral replication underlie the persistence of gut lesions (Klatt, Funderburg, et al., 2013). Furthermore, lack of mucosal dysfunction, including an absence of microbial translocation or loss of Th17 cells, in non-progressive SIV infection of naturally adapted nonhuman primates, which have high virus replication but do not progress to AIDS, further indicates that mucosal damage is not solely due to the virus (Brenchley, Price, Schacker, et al., 2006; Brenchley et al., 2008; Klatt, Silvestri, & Hirsch, 2012). We suspect that virus replication and CD4+ T cell depletion results in immune activation and inflammation, damaging the gut epithelial barrier permitting microbial translocation, that may perpetuate mucosal and systemic immune activation ultimately contributing to CD4+ T cell restoration failure. As is seen in systemic lymphoid tissues during pathogenic infection, and as a consequence of inflammation (Estes, Baker, et

al., 2008), there is also collagen deposition and fibrosis of lymphoid follicles of the GI tract that alters their architecture and appears to limit CD4+ T cell restoration at these sites (Estes, Haase, & Schacker, 2008). Interplay between gut mucosal immune defenses and the microbiota is likely as in ART-treated SIV-infected macaques, prebiotic/probiotic treatment enhanced CD4+ T cell restoration in the gut possibly by decreasing fibrosis (Klatt, Canary, et al., 2013).

8. OTHER POSSIBLE DRIVERS OF RESIDUAL IMMUNE DYSREGULATION IN TREATED HIV INFECTION

As noted earlier, several recent studies have identified markers of immune activation, inflammation, and coagulation as robust predictors of morbidity and mortality in treated HIV infection (Gray et al., 1999; Kalayjian et al., 2010; Kuller et al., 2008; Sandler et al., 2011); it remains unclear, however, which of these elements are directly driving pathogenesis and which are just markers of the persistent inflammatory environment that characterizes this setting. It is likely that morbidities are multifactorial in origin, with risks driven in part by age, in part by exposures such as cigarettes, and in part by this inflammatory environment. And if, as suspected, multiple factors acting in concert are driving pathogenesis and disease course, unraveling the contributions of each will be challenging. What is more, the underlying drivers of sustained inflammation and coagulation in the setting of treated virologically controlled infection are not clearly defined and may include any of the drivers below alone or in combination.

8.1. Microbial translocation

As noted earlier, microbial translocation has been linked strongly to both immune activation and the pathogenesis of disease in treated HIV infection (Klatt, Funderburg, et al., 2013). And while ART reduces plasma levels of microbial products, plasma levels most typically remain elevated when compared to levels in uninfected controls (Brenchley, Price, & Douek, 2006; Jiang et al., 2009; Lederman et al., 2011; Marchetti et al., 2008). Bacterial products can drive monocyte expression of inflammatory cytokines and coagulation factors (Funderburg et al., 2010, 2012; Pasare & Medzhitov, 2004) that have both been linked to morbidity and mortality in treated infection (Duprez et al., 2009; Kuller et al., 2008). Moreover, in vitro exposure of peripheral blood cells to these products can drive activation and cycling of memory CD4+ T cells (but not so much for CD8+ T cells) (Funderburg et al., 2008), inducing an activation signature that is similar to that seen among treated patients with immune failure (Lederman et al., 2011).

8.2. Homeostatic proliferation

Among persons with profound lymphopenia, circulating levels of the homeostatic cytokine interleukin-7 are elevated (Fry et al., 2001; Napolitano et al., 2001). And while T cell responses to IL-7 may be impaired in HIV infection (Bazdar & Sieg, 2007; Colle et al., 2006; Kalinowska, Bazdar, Lederman, Funderburg, & Sieg, 2013; Marziali et al., 2006), the role of this cytokine in driving T cell activation and cycling in patients with immune failure and profound lymphopenia is plausible as this has been linked to cellular activation in persons with idiopathic CD4 lymphopenia (ICL; Zonios et al., 2008). On the other hand, in

ICL, plasma levels of the inflammatory cytokines IL-6 and the receptors for tumor necrosis factor that have been linked to morbidity and mortality in treated HIV infection are not elevated (I. Sereti, personal communication) suggesting that factors other than cytopenia are likely important in driving inflammation and activation in treated HIV infection. Yet after administration of IL-7 to HIV-infected patients with immune failure, selected indices of immune activation are attenuated (Sereti et al., 2012), but this could be related to IL-7-driven repopulation of the damaged gut mucosal immune system (Cimbro et al., 2012; Sereti et al., 2012). Likewise in nonhuman primates that are naturally infected with the simian immune deficiency virus, immune activation is attenuated after acute infection and CD4 lymphopenia is uncommon (Brenchley, Price, & Douek, 2006; Estes, Gordon, et al., 2008). Whether the benign course of infection is a consequence or a cause of low-level cellular activation remains an unanswered question.

8.3. HIV replication

While HIV replication is attenuated dramatically in most ART-treated patients, with sensitive enough assays, low levels of virus often can be found in plasma (Hatano et al., 2010; Maldarelli et al., 2007; Palmer et al., 2008) and HIV RNA can be found in LN and gut mucosal tissues (Cory, Schacker, Stevenson, & Fletcher, 2013; Yukl et al., 2010). Viral products can drive activation of both adaptive and innate immune responses (Beignon et al., 2005; Heil et al., 2004; Meier et al., 2007) and also can activate the procoagulant activity of circulating monocytes (Funderburg et al., 2010, 2012), potentially contributing to thrombotic risk. Yet, results of ART intensification studies have provided mixed results, with some showing no effect on T cell activation in the blood or GALT (Gandhi et al., 2011; Hatano et al., 2011) and another reporting a decrease in CD8+ T cell activation in a subset of patients (Buzon et al., 2010).

8.4. Copathogens

Coinfection with pathogens that cause chronic infection such as hepatitis C virus, cytomegalovirus, and other human herpesviruses is common in HIV-infected persons and in many regions of epidemic infection, infestation with long-lived parasites such as helminthes is also prevalent. It is reasonable to propose that these persistent infections could contribute to immune activation in treated HIV infection (Borkow, Teicher, & Bentwich, 2007; Modjarrad & Vermund, 2010) as they may serve as persistent sources of antigen that can drive T cell activation as well as other microbial elements that can activate innate immune receptors. Recently, an 8-week trial of valganciclovir, a drug used for the treatment of cytomegalovirus infection in patients dually infected with HIV and CMV reduced indices of CD8 T cell activation (Hunt et al., 2011). CMV coinfection also has been linked to the increased risk of cardiovascular disease in HIV infection (Triant et al., 2007) as those patients with more active CMV-specific T cell responses are more likely to have thickening of the carotid artery intima (Sacre et al., 2012)—a radiographic finding that is associated with increased risk of vascular morbidity, and recent data suggest that CMV-specific T cells may play a role in the endovascular inflammation of HIV infection (Sacre et al., 2012).

8.5. Inflammatory lipids

Patients with chronic HIV infection have altered lipid and metabolic profiles (Grinspoon & Carr, 2005; Rose et al., 2008). These altered lipid profiles may contribute to inflammation in HIV disease, as there is a growing appreciation that lipid subclasses, including oxidized low-density lipoprotein, may activate innate immune receptors (Stewart et al., 2010) and induce proinflammatory cytokine production through activation of the inflammasome (Erridge, 2009). Recently, in a cross-sectional study, metabolic factors (low-density LDL, apolipoprotein A) were better correlated with cardiovascular risk than were inflammatory markers (CRP, IL-6, TNF- α) in HIV-infected patients with ART-controlled viremia (Piconi et al., 2013). There is reason to suspect that certain aspects of the inflammatory lipid profile may be related to certain classes of antiretroviral therapies as the recent randomized SPIRAL study demonstrated that among patients with virologic control on ART, switching treatment from a protease inhibitor-based regimen to an integrase inhibitor-based regimen improved both the lipid profile of these subjects and diminished levels of inflammatory markers in plasma (Martinez et al., 2012) although the lipid changes and inflammatory marker changes were only weakly correlated.

9. THERAPEUTIC APPROACHES

Our evolving understanding of the possible underlying causes of persistent immune activation/inflammation/coagulation during treated HIV infection has given rise to several therapeutic approaches to decrease immune activation, inflammation, and coagulation in this setting.

9.1. Targeting residual viral replication

While most HIV-infected patients with access to ART are able to maintain plasma HIV RNA levels below the limits of detection of current clinical assays, HIV RNA continues to be readily detectable in tissues (Anton et al., 2003), and even in plasma, using ultrasensitive methods (Maldarelli et al., 2007; Palmer et al., 2008). There has been extensive controversy over whether the detected HIV RNA in these studies represents ongoing continued productive viral replication or simply release of HIV from infected cells in the absence of new infection of target cells. While there is little evidence for viral evolution in these individuals, and most treatment-intensification studies have failed to demonstrate a reduction in low-level viremia or immune activation (Dinoso et al., 2009; Gandhi et al., 2010; Hatano et al., 2011; Hunt et al., 2013), addition of raltegravir to an apparently suppressive ART treatment regimen appeared to decrease new infection of CD4⁺ T cells and reduce T cell activation or D-dimer levels in two independent randomized controlled trials (Hatano et al., 2013; Massanella et al., 2012). These apparent immunologic and virologic benefits appeared to be most apparent in individuals receiving protease inhibitor-based therapy in both of these studies, and while unproven, there is some speculation that incomplete drug penetration into lymphoid tissues for some antiretroviral drugs and/or classes (Cory et al., 2013) may allow for at least some aborted rounds of viral replication—insufficient to result in viral evolution—but sufficient to contribute to immune activation. That said, the immunologic benefit of treatment intensification is marginal at best and will

likely be an insufficient strategy for managing chronic immune activation in treated HIV infection.

9.2. Targeting chronic viral coinfections

Most HIV-infected individuals are infected chronically with many other viruses including cytomegalovirus. Since CMV is responsible for at least 10% of the entire circulating memory T cell repertoire in healthy HIV-uninfected CMV-seropositive individuals, and presumably even higher levels in treated HIV infection (Naeger et al., 2010), our group recently tested the hypothesis that asymptomatic CMV (and/or other herpesvirus) replication contributes to persistent immune activation in treated HIV infection. Indeed, we found that treating asymptomatic CMV coinfection with valganciclovir significantly reduces CD8+ T cell activation in the context of treated HIV infection (Hunt, Martin, et al., 2011). While drug-related toxicities preclude the long-term use of valganciclovir, other drugs targeting CMV are in clinical development and may hold promise for future therapeutic approaches to reduce T cell activation in treated HIV infection. The contribution of other chronic viral infections to systemic immune activation in treated HIV infection is less clear but is an area of active study.

9.3. Targeting microbial translocation

Given the strong link between markers of gut epithelial barrier dysfunction, monocyte activation, and mortality, several studies are now targeting microbial translocation to reduce immune activation in treated HIV infection. While an early trial of hyperimmune bovine colostrum showed no benefit (and failed to reduce microbial translocation; Byakwaga et al., 2011), other recent pilot studies of prebiotic interventions show some promise in reducing systemic markers of monocyte activation and/or microbial translocation, though these studies need to be replicated in larger trials (Gori et al., 2011). Other ongoing studies in the NIH-funded ACTG are assessing the impact of rifaximin (an antibiotic that is not systemically absorbed) and sevelamer (which binds bacterial LPS in the gut lumen) on microbial translocation and immune activation in HIV infection.

9.4. Interventions to improve CD41 T cell recovery

While the relationship between inflammation and clinical outcomes appears independent of CD4+ T cell count, certain interventions designed to improve CD4+ T cell recovery may well help decrease microbial translocation and systemic immune activation, particularly if they restore important CD4+ T cell subsets in the gut mucosa. While IL-2 therapy failed to prevent morbidity and mortality despite raising CD4+ T cell counts (Abrams et al., 2009), IL-7 administration may prove more promising since the phenotype of the expanded cell populations appears more favorable (i.e., without Treg characteristics) and it appears to restore CD4+ T cells in the gut-associated lymphoid tissue and decrease D-dimer levels in peripheral blood (I Sereti et al., 2012).

9.5. Targeting innate immune responses

As there is substantial evidence linking innate immune activation to the pathogenesis of disease in treated HIV infection, some groups have conducted trials of weak to moderate

inhibitors of innate immune responses. For example, a small pilot trial of chloroquine appeared to have favorable immunologic effects in untreated HIV infection (Murray et al., 2010), but hydroxychloroquine appeared to increase plasma HIV RNA levels while having very little effect on T cell activation in a much larger randomized controlled trial (Paton et al., 2012). Interestingly, when hydroxychloroquine was assessed in ART-suppressed patients, it appeared to reduce T cell activation significantly (Piconi et al., 2011), suggesting that modulating the innate immune response might have qualitatively different effects when interferons are no longer needed for the control of HIV replication.

Several commonly used medications with anti-inflammatory properties decrease monocyte activation and may hold promise in treated HIV infection. For example, a recent randomized controlled trial of rosuvastatin in ART-suppressed HIV-infected individuals demonstrated significant reductions in plasma sCD14 levels and cellular markers of monocyte activation (McComsey et al., 2013). Similarly, an uncontrolled trial of aspirin in HIV-infected individuals suggested a reduction in sCD14 levels after 1 week (O'Brien et al., 2013) though this observation needs to be confirmed in a randomized controlled trial. Whether these interventions are associated with clinical benefits can only be confirmed in much larger clinical event trials.

10. SUMMARY

Despite effective control of HIV replication with combination antiretroviral therapies, there is increasing evidence that selected morbidities and mortalities are increased in treated HIV-infected persons. These outcomes appear linked to incomplete systemic CD4+ T cell restoration with therapy and also with an increased immune activation, inflammation, and coagulation profile. And while activation, inflammation, and coagulation abnormalities are greater in those with incomplete systemic immune restoration, both the determinants of these processes and their relationships to each other are incompletely understood. Defining the pathogenesis of this RIDS will help to identify targets for therapeutic intervention in treated HIV infection and may also highlight determinants of morbidities that affect the general population.

Acknowledgments

The authors would like to thank the members of the Cleveland Immunopathogenesis Consortium (CLIC) funded by the National Institutes of Health (AI 076174) for the discussions and guidance that have helped to shape this chapter.

References

- Abrams D, Levy Y, Losso MH, Babiker A, Collins G, Cooper DA, et al. Interleukin-2 therapy in patients with HIV infection. *The New England Journal of Medicine*. 2009; 361(16):1548–1559. <http://dx.doi.org/10.1056/NEJMoa0903175>. [PubMed: 19828532]
- Achhra AC, Amin J, Law MG, Emery S, Gerstoft J, Gordin FM, et al. Immunodeficiency and the risk of serious clinical endpoints in a well studied cohort of treated HIV-infected patients. *AIDS*. 2010; 24(12):1877–1886. <http://dx.doi.org/QA1.0b013e31823fd1f2/QAD.0b013e32833b1b2>. [PubMed: 20588170]

- Andersson J, Fehniger TE, Patterson BK, Pottage J, Agnoli M, Jones P, et al. Early reduction of immune activation in lymphoid tissue following highly active HIV therapy. *AIDS*. 1998; 12(11):F123–F129. [PubMed: 9708402]
- Anton PA, Mitsuyasu RT, Deeks SG, Scadden DT, Wagner B, Huang C, et al. Multiple measures of HIV burden in blood and tissue are correlated with each other but not with clinical parameters in aviremic subjects. *AIDS*. 2003; 17(1):53–63. [PubMed: 12478069]
- Arthos J, Cicala C, Martinelli E, Macleod K, Van Ryk D, Wei D, et al. HIV-1 envelope protein binds to and signals through integrin alpha4beta7, the gut mucosal homing receptor for peripheral T cells. *Nature Immunology*. 2008; 9(3):301–309. <http://dx.doi.org/10.1038/ni1566>. [PubMed: 18264102]
- Autran B, Carcelain G, Li TS, Blanc C, Mathez D, Tubiana R, et al. Positive effects of combined antiretroviral therapy on CD4+ T cell homeostasis and function in advanced HIV disease. *Science*. 1997; 277(5322):112–116. [see comments]. [PubMed: 9204894]
- Baker JV, Peng G, Rapkin J, Abrams DI, Silverberg MJ, MacArthur RD, et al. CD4+ count and risk of non-AIDS diseases following initial treatment for HIV infection. *AIDS*. 2008; 22(7):841–848. <http://dx.doi.org/QAI.0b013e31823fd1f2/QAD.0b013e3282f7cb76>. [PubMed: 18427202]
- Bazdar DA, Sieg SF. Interleukin-7 enhances proliferation responses to T-cell receptor stimulation in naive CD4+ T cells from human immunodeficiency virus-infected persons. *Journal of Virology*. 2007; 81(22):12670–12674. <http://dx.doi.org/10.1128/JVI.00476-07>. [PubMed: 17855550]
- Beignon AS, McKenna K, Skoberne M, Manches O, DaSilva I, Kavanagh DG, et al. Endocytosis of HIV-1 activates plasmacytoid dendritic cells via Toll-like receptor-viral RNA interactions. *The Journal of Clinical Investigation*. 2005; 115(11):3265–3275. [PubMed: 16224540]
- Benveniste O, Flahault A, Rollot F, Elbim C, Estaquier J, Pedron B, et al. Mechanisms involved in the low-level regeneration of CD4+ cells in HIV-1-infected patients receiving highly active antiretroviral therapy who have prolonged undetectable plasma viral loads. *Journal of Infectious Diseases*. 2005; 191(10):1670–1679. <http://dx.doi.org/10.1086/429670>. [PubMed: 15838794]
- Betts MR, Ambrozak DR, Douek DC, Bonhoeffer S, Brenchley JM, Casazza JP, et al. Analysis of total human immunodeficiency virus (HIV)-specific CD4(+) and CD8(+) T-cell responses: Relationship to viral load in untreated HIV infection. *Journal of Virology*. 2001; 75(24):11983–11991. [PubMed: 11711588]
- Biancotto A, Grivel JC, Iglehart SJ, Vanpouille C, Lisco A, Sieg SF, et al. Abnormal activation and cytokine spectra in lymph nodes of people chronically infected with HIV-1. *Blood*. 2007; 109(10):4272–4279. [PubMed: 17289812]
- Borkow G, Teicher C, Bentwich Z. Helminth-HIV coinfection: Should we deworm? *PLoS Neglected Tropical Diseases*. 2007; 1(3):e160. <http://dx.doi.org/10.1371/journal.pntd.0000160>. [PubMed: 18167516]
- Boulware DR, Hullsiek KH, Puroton CE, Rupert A, Baker JV, French MA, et al. Higher levels of CRP, D-dimer, IL-6, and hyaluronic acid before initiation of antiretroviral therapy (ART) are associated with increased risk of AIDS or death. *Journal of Infectious Diseases*. 2011; 203(11):1637–1646. <http://dx.doi.org/10.1093/infdis/jir134>. [PubMed: 21592994]
- Breen EC, Hussain SK, Magpantay L, Jacobson LP, Detels R, Rabkin CS, et al. B-cell stimulatory cytokines and markers of immune activation are elevated several years prior to the diagnosis of systemic AIDS-associated non-Hodgkin B-cell lymphoma. *Cancer Epidemiology, Biomarkers & Prevention*. 2011; 20(7):1303–1314. <http://dx.doi.org/10.1158/1055-9965.EPI-11-0037>.
- Brenchley JM, Douek DC. HIV infection and the gastrointestinal immune system. *Mucosal Immunology*. 2008; 1(1):23–30. <http://dx.doi.org/10.1038/mi.2007.1>. [PubMed: 19079157]
- Brenchley JM, Douek DC. Microbial translocation across the GI tract. *Annual Review of Immunology*. 2012; 30:149–173. <http://dx.doi.org/10.1146/annurev-immunol-020711-075001>.
- Brenchley JM, Paiardini M, Knox KS, Asher AI, Cervasi B, Asher TE, et al. Differential Th17 CD4 T-cell depletion in pathogenic and nonpathogenic lentiviral infections. *Blood*. 2008; 112(7):2826–2835. <http://dx.doi.org/10.1182/blood-2008-05-159301>. [PubMed: 18664624]
- Brenchley JM, Price DA, Douek DC. HIV disease: Fallout from a mucosal catastrophe? *Nature Immunology*. 2006; 7(3):235–239. [PubMed: 16482171]

- Brenchley JM, Price DA, Schacker TW, Asher TE, Silvestri G, Rao S, et al. Microbial translocation is a cause of systemic immune activation in chronic HIV infection. *Nature Medicine*. 2006; 12(12): 1365–1371.
- Brenchley JM, Schacker TW, Ruff LE, Price DA, Taylor JH, Beilman GJ, et al. CD4+ T cell depletion during all stages of HIV disease occurs predominantly in the gastrointestinal tract. *The Journal of Experimental Medicine*. 2004; 200(6):749–759. <http://dx.doi.org/10.1084/jem.20040874>. [PubMed: 15365096]
- Brown TT, Cole SR, Li X, Kingsley LA, Palella FJ, Riddler SA, et al. Antiretroviral therapy and the prevalence and incidence of diabetes mellitus in the multicenter AIDS cohort study. *Archives of Internal Medicine*. 2005; 165(10):1179–1184. <http://dx.doi.org/10.1001/archinte.165.10.1179>. [PubMed: 15911733]
- Brown TT, Qaqish RB. Antiretroviral therapy and the prevalence of osteopenia and osteoporosis: A meta-analytic review. *AIDS*. 2006; 20(17):2165–2174. [PubMed: 17086056]
- Brown TT, Tassiopoulos K, Bosch RJ, Shikuma C, McComsey GA. Association between systemic inflammation and incident diabetes in HIV-infected patients after initiation of antiretroviral therapy. *Diabetes Care*. 2010; 33(10):2244–2249. <http://dx.doi.org/10.2337/dc10-0633>. [PubMed: 20664016]
- Bucy RP, Hockett RD, Derdeyn CA, Saag MS, Squires K, Sillers M, et al. Initial increase in blood CD4(+) lymphocytes after HIV antiretroviral therapy reflects redistribution from lymphoid tissues. *The Journal of Clinical Investigation*. 1999; 103(10):1391–1398. [PubMed: 10330421]
- Burdo, TH.; Weiffenbach, A.; Woods, SP.; Letendre, S.; Ellis, RJ.; Williams, KC. Elevated sCD163 is a marker of neurocognitive impairment in HIV infection. *AIDS*. 2013. <http://dx.doi.org/10.1097/QAD.0b013e32836010bd>. Epub ahead of print
- Buzon MJ, Massanella M, Llibre JM, Esteve A, Dahl V, Puertas MC, et al. HIV-1 replication and immune dynamics are affected by raltegravir intensification of HAART-suppressed subjects. *Nature Medicine*. 2010; 16(4):460–465. <http://dx.doi.org/10.1038/nm.2111>.
- Byakwaga, H.; Kelly, M.; Purcell, DF.; French, MA.; Amin, J.; Lewin, SR., et al. Intensification of antiretroviral therapy with raltegravir or addition of hyperimmune bovine colostrum in HIV-infected patients with suboptimal CD4+ T-cell response: A randomized controlled trial. *Journal of Infectious Diseases*. 2011. http://dx.doi.org/10.1093/infdis/jir559***
- Chege D, Sheth PM, Kain T, Kim CJ, Kovacs C, Loutfy M, et al. Sigmoid Th17 populations, the HIV latent reservoir, and microbial translocation in men on long-term antiretroviral therapy. *AIDS*. 2011; 25(6):741–749. <http://dx.doi.org/QAI.0b013e31823fd1f2/QAD.0b013e328344cefb>. [PubMed: 21378536]
- Chehimi J, Campbell DE, Azzoni L, Bacheller D, Papasavvas E, Jerandi G, et al. Persistent decreases in blood plasmacytoid dendritic cell number and function despite effective highly active antiretroviral therapy and increased blood myeloid dendritic cells in HIV-infected individuals. *The Journal of Immunology*. 2002; 168(9):4796–4801. [PubMed: 11971031]
- Chun TW, Carruth L, Finzi D, Shen X, DiGiuseppe JA, Taylor H, et al. Quantification of latent tissue reservoirs and total body viral load in HIV-1 infection. *Nature*. 1997; 387(6629):183–188. [PubMed: 9144289]
- Cimbro R, Vassena L, Arthos J, Cicala C, Kehrl JH, Park C, et al. IL-7 induces expression and activation of integrin alpha4beta7 promoting naive T-cell homing to the intestinal mucosa. *Blood*. 2012; 120(13):2610–2619. <http://dx.doi.org/10.1182/blood-2012-06-434779>. [PubMed: 22896005]
- Colle JH, Moreau JL, Fontanet A, Lambotte O, Joussemet M, Jacod S, et al. Regulatory dysfunction of the interleukin-7 receptor in CD4 and CD8 lymphocytes from HIV-infected patients—Effects of antiretroviral therapy. *Journal of Acquired Immune Deficiency Syndromes*. 2006; 42(3):277–285. <http://dx.doi.org/QAI.0b013e31823fd1f2/01.qai.0000214823.11034.4e>. [PubMed: 16810123]
- Copur AS, Smith PR, Gomez V, Bergman M, Homel P. HIV infection is a risk factor for venous thromboembolism. *AIDS Patient Care and STDs*. 2002; 16(5):205–209. <http://dx.doi.org/10.1089/10872910252972258>. [PubMed: 12055028]
- Cory TJ, Schacker TW, Stevenson M, Fletcher CV. Overcoming pharmacologic sanctuaries. *Current Opinion in HIV and AIDS*. 2013; 8(3):190–195. <http://dx.doi.org/QAI.0b013e31823fd1f2/COH.0b013e32835fc68a>. [PubMed: 23454865]

- Deeken JF, Tjen ALA, Rudek MA, Okuliar C, Young M, Little RF, et al. The rising challenge of non-AIDS-defining cancers in HIV-infected patients. *Clinical Infectious Diseases: An Official Publication of the Infectious Diseases Society of America*. 2012; 55(9):1228–1235. <http://dx.doi.org/10.1093/cid/cis613>. [PubMed: 22776851]
- Desquilbet L, Jacobson LP, Fried LP, Phair JP, Jamieson BD, Holloway M, et al. HIV-1 infection is associated with an earlier occurrence of a phenotype related to frailty. *The Journals of Gerontology Series A, Biological Sciences and Medical Sciences*. 2007; 62(11):1279–1286.
- Desquilbet L, Margolick JB, Fried LP, Phair JP, Jamieson BD, Holloway M, et al. Relationship between a frailty-related phenotype and progressive deterioration of the immune system in HIV-infected men. *Journal of Acquired Immune Deficiency Syndromes*. 2009; 50(3):299–306. <http://dx.doi.org/QAI.0b013e31823fd1f2/QAI.0b013e3181945eb0>. [PubMed: 19194312]
- Dinso JB, Kim SY, Wiegand AM, Palmer SE, Gange SJ, Cranmer L, et al. Treatment intensification does not reduce residual HIV-1 viremia in patients on highly active antiretroviral therapy. *Proceedings of the National Academy of Sciences of the United States of America*. 2009; 106(23):9403–9408. <http://dx.doi.org/10.1073/pnas.0903107106>. [PubMed: 19470482]
- Douek DC, Brenchley JM, Betts MR, Ambrozak DR, Hill BJ, Okamoto Y, et al. HIV preferentially infects HIV-specific CD4+ T cells. *Nature*. 2002; 417(6884):95–98. [PubMed: 11986671]
- Duprez DA, Kuller LH, Tracy R, Otvos J, Cooper DA, Hoy J, et al. Lipoprotein particle subclasses, cardiovascular disease and HIV infection. *Atherosclerosis*. 2009; 207(2):524–529. <http://dx.doi.org/10.1016/j.atherosclerosis.2009.05.001>. [PubMed: 19515371]
- Ellis RJ, Badiee J, Vaida F, Letendre S, Heaton RK, Clifford D, et al. CD4 nadir is a predictor of HIV neurocognitive impairment in the era of combination antiretroviral therapy. *AIDS*. 2011; 25(14):1747–1751. <http://dx.doi.org/QAI.0b013e31823fd1f2/QAD.0b013e32834a40cd>. [PubMed: 21750419]
- Erlanson, KM.; Allshouse, AA.; Jankowski, CM.; Lee, EJ.; Rufner, KM.; Palmer, BE., et al. Association of functional impairment with inflammation and immune activation in HIV-1-infected adults on effective antiretroviral therapy. *The Journal of Infectious Diseases*. 2013. <http://dx.doi.org/10.1093/infdis/jit147>. Epub ahead of print
- Erridge C. The roles of toll-like receptors in atherosclerosis. *Journal of Innate Immunity*. 2009; 1(4):340–349. <http://dx.doi.org/10.1159/000191413>. [PubMed: 20375591]
- Estes J, Baker JV, Brenchley JM, Khoruts A, Barthold JL, Bantle A, et al. Collagen deposition limits immune reconstitution in the gut. *Journal of Infectious Diseases*. 2008; 198(4):456–464. <http://dx.doi.org/10.1086/590112>. [PubMed: 18598193]
- Estes JD, Gordon SN, Zeng M, Chahroudi AM, Dunham RM, Staprans SI, et al. Early resolution of acute immune activation and induction of PD-1 in SIV-infected sooty mangabeys distinguishes nonpathogenic from pathogenic infection in rhesus macaques. *The Journal of Immunology*. 2008; 180(10):6798–6807. [PubMed: 18453600]
- Estes JD, Haase AT, Schacker TW. The role of collagen deposition in depleting CD4+ T cells and limiting reconstitution in HIV-1 and SIV infections through damage to the secondary lymphoid organ niche. *Seminars in Immunology*. 2008; 20(3):181–186. <http://dx.doi.org/10.1016/j.smim.2008.04.002>. [PubMed: 18595731]
- Estes JD, Harris LD, Klatt NR, Tabb B, Pittaluga S, Paiardini M, et al. Damaged intestinal epithelial integrity linked to microbial translocation in pathogenic simian immunodeficiency virus infections. *PLoS Pathogens*. 2010; 6(8):e1001052. [PubMed: 20808901]
- Estes JD, Wietgreffe S, Schacker T, Southern P, Beilman G, Reilly C, et al. Simian immunodeficiency virus-induced lymphatic tissue fibrosis is mediated by transforming growth factor beta 1-positive regulatory T cells and begins in early infection. *Journal of Infectious Diseases*. 2007; 195(4):551–561. <http://dx.doi.org/10.1086/510852>. [PubMed: 17230415]
- Favre D, Mold J, Hunt PW, Kanwar B, Loke P, Seu L, et al. Tryptophan catabolism by indoleamine 2,3-dioxygenase 1 alters the balance of TH17 to regulatory T cells in HIV disease. *Science Translational Medicine*. 2010; 2(32):32ra36. <http://dx.doi.org/10.1126/scitranslmed.3000632>.
- Fernandez S, Tanaskovic S, Helbig K, Rajasuriar R, Kramski M, Murray JM, et al. CD4+ T-cell deficiency in HIV patients responding to antiretroviral therapy is associated with increased expression of interferon-stimulated genes in CD4+ T cells. *Journal of Infectious Diseases*. 2011; 204(12):1927–1935. <http://dx.doi.org/10.1093/infdis/jir659>. [PubMed: 22006994]

- Ford ES, Greenwald JH, Richterman AG, Rupert A, Dutcher L, Badralmaa Y, et al. Traditional risk factors and D-dimer predict incident cardiovascular disease events in chronic HIV infection. *AIDS*. 2010; 24(10):1509–1517. <http://dx.doi.org/QAI.0b013e31823fd1f2/QAD.0b013e32833ad914>. [PubMed: 20505494]
- Fox CH, Kotler D, Tierney A, Wilson CS, Fauci AS. Detection of HIV-1 RNA in the lamina propria of patients with AIDS and gastrointestinal disease. *Journal of Infectious Diseases*. 1989; 159(3):467–471. [PubMed: 2915167]
- Freiberg MS, Chang CC, Kuller LH, Skanderson M, Lowy E, Kraemer KL, et al. HIV infection and the risk of acute myocardial infarction. *JAMA Internal Medicine*, 1–9. 2013; 173(8):614–622. <http://dx.doi.org/10.1001/jamainternmed.2013.3728>.
- Fry TJ, Connick E, Falloon J, Lederman MM, Liewehr DJ, Spritzler J, et al. A potential role for interleukin-7 in T-cell homeostasis. *Blood*. 2001; 97(10):2983–2990. [PubMed: 11342421]
- Fultz SL, McGinnis KA, Skanderson M, Ragni MV, Justice AC. Association of venous thromboembolism with human immunodeficiency virus and mortality in veterans. *The American Journal of Medicine*. 2004; 116(6):420–423. <http://dx.doi.org/10.1016/j.amjmed.2003.10.011>. [PubMed: 15006592]
- Funderburg N, Luciano AA, Jiang W, Rodriguez B, Sieg SF, Lederman MM. Toll-like receptor ligands induce human T cell activation and death, a model for HIV pathogenesis. *PLoS One*. 2008; 3(4):e1915. [PubMed: 18382686]
- Funderburg NT, Mayne E, Sieg SF, Asaad R, Jiang W, Kalinowska M, et al. Increased tissue factor expression on circulating monocytes in chronic HIV infection: Relationship to in vivo coagulation and immune activation. *Blood*. 2010; 115(2):161–167. <http://dx.doi.org/10.1182/blood-2009-03-210179>. [PubMed: 19828697]
- Funderburg NT, Zidar DA, Shive C, Lioi A, Mudd J, Musselwhite LW, et al. Shared monocyte subset phenotypes in HIV-1 infection and in uninfected subjects with acute coronary syndrome. *Blood*. 2012; 120(23):4599–4608. <http://dx.doi.org/10.1182/blood-2012-05-433946>. [PubMed: 23065151]
- Gandhi RT, Coombs RW, Chan ES, Bosch RJ, Zheng L, Margolis DM, et al. No effect of raltegravir intensification on viral replication markers in the blood of HIV-1-infected patients receiving antiretroviral therapy. *Journal of Acquired Immune Deficiency Syndromes*. 2011; 59(3):229–235. <http://dx.doi.org/QAI.0b013e31823fd1f2/QAI.0b013e31823fd1f2>. [PubMed: 22083073]
- Gandhi RT, Spritzler J, Chan E, Asmuth DM, Rodriguez B, Merigan TC, et al. Effect of baseline- and treatment-related factors on immunologic recovery after initiation of antiretroviral therapy in HIV-1-positive subjects: Results from ACTG 384. *Journal of Acquired Immune Deficiency Syndromes*. 2006; 42(4):426–434. <http://dx.doi.org/10.1097/01.qai.0000226789.51992.3f>. [PubMed: 16810109]
- Gandhi RT, Zheng L, Bosch RJ, Chan ES, Margolis DM, Read S, et al. The effect of raltegravir intensification on low-level residual viremia in HIV-infected patients on antiretroviral therapy: A randomized controlled trial. *PLoS Medicine*. 2010; 7(8) <http://dx.doi.org/10.1371/journal.pmed.1000321>.
- Gazzola L, Tincati C, Bellistri GM, Monforte A, Marchetti G. The absence of CD4+ T cell count recovery despite receipt of virologically suppressive highly active antiretroviral therapy: Clinical risk, immunological gaps, and therapeutic options. *Clinical Infectious Diseases*. 2009; 48(3):328–337. <http://dx.doi.org/10.1086/595851>. [PubMed: 19123868]
- Giorgi JV, Hultin LE, McKeating JA, Johnson TD, Owens B, Jacobson LP, et al. Shorter survival in advanced human immunodeficiency virus type 1 infection is more closely associated with T lymphocyte activation than with plasma virus burden or virus chemokine coreceptor usage. *Journal of Infectious Diseases*. 1999; 179(4):859–870. [PubMed: 10068581]
- Giorgi JV, Liu Z, Hultin LE, Cumberland WG, Hennessey K, Detels R. Elevated levels of CD38+ CD8+ T cells in HIV infection add to the prognostic value of low CD4+ T cell levels: Results of 6 years of follow-up. The Los Angeles Center, Multicenter AIDS Cohort Study. *Journal of Acquired Immune Deficiency Syndromes*. 1993; 6(8):904–912. [PubMed: 7686224]
- Giorgi JV, Lyles RH, Matud JL, Yamashita TE, Mellors JW, Hultin LE, et al. Predictive value of immunologic and virologic markers after long or short duration of HIV-1 infection. *Journal of Acquired Immune Deficiency Syndromes*. 2002; 29(4):346–355. [PubMed: 11917238]

- Gordon SN, Cervasi B, Odorizzi P, Silverman R, Abera F, Ginsberg G, et al. Disruption of intestinal CD4+T cell homeostasis is a key marker of systemic CD4+T cell activation in HIV-infected individuals. *The Journal of Immunology*. 2010; 185(9):5169–5179. <http://dx.doi.org/10.4049/jimmunol.1001801>. [PubMed: 20889546]
- Gori A, Rizzardini G, Van't Land B, Amor KB, van Schaik J, Torti C, et al. Specific prebiotics modulate gut microbiota and immune activation in HAART-naïve HIV-infected adults: Results of the “COPA” pilot randomized trial. *Mucosal Immunology*. 2011; 4(5):554–563. <http://dx.doi.org/10.1038/mi.2011.15>. [PubMed: 21525866]
- Gottlieb MS, Schroff R, Schanker HM, Weisman JD, Fan PT, Wolf RA, et al. Pneumocystis carinii pneumonia and mucosal candidiasis in previously healthy homosexual men: Evidence of a new acquired cellular immunodeficiency. *The New England Journal of Medicine*. 1981; 305(24):1425–1431. [PubMed: 6272109]
- Gray CM, Lawrence J, Schapiro JM, Altman JD, Winters MA, Crompton M, et al. Frequency of class I HLA-restricted anti-HIV CD8+ T cells in individuals receiving highly active antiretroviral therapy (HAART). *The Journal of Immunology*. 1999; 162(3):1780–1788. [PubMed: 9973442]
- Grinspoon S, Carr A. Cardiovascular risk and body-fat abnormalities in HIV-infected adults. *The New England Journal of Medicine*. 2005; 352(1):48–62. <http://dx.doi.org/10.1056/NEJMra041811>. [PubMed: 15635112]
- Gulich AE, van Leeuwen MT, Falster MO, Vajdic CM. Incidence of cancers in people with HIV/AIDS compared with immunosuppressed transplant recipients: A meta-analysis. *The Lancet*. 2007; 370(9581):59–67. [http://dx.doi.org/10.1016/S0140-6736\(07\)61050-2](http://dx.doi.org/10.1016/S0140-6736(07)61050-2).
- Harris TB, Ferrucci L, Tracy RP, Corti MC, Wacholder S, Ettinger WH Jr, et al. Associations of elevated interleukin-6 and C-reactive protein levels with mortality in the elderly. *The American Journal of Medicine*. 1999; 106(5):506–512. [PubMed: 10335721]
- Harris LD, Tabb B, Sodora DL, Paiardini M, Klatt NR, Douek DC, et al. Downregulation of robust acute type I interferon responses distinguishes non-pathogenic simian immunodeficiency virus (SIV) infection of natural hosts from pathogenic SIV infection of rhesus macaques. *Journal of Virology*. 2010; 84(15):7886–7891. <http://dx.doi.org/10.1128/JVI.02612-09>. [PubMed: 20484518]
- Hatano H, Delwart EL, Norris PJ, Lee TH, Neilands TB, Kelley CF, et al. Evidence of persistent low-level viremia in long-term HAART-suppressed, HIV-infected individuals. *AIDS*. 2010; 24(16):2535–2539. <http://dx.doi.org/10.1097/QAD.0b013e32833dba03>. [PubMed: 20651585]
- Hatano H, Hayes TL, Dahl V, Sinclair E, Lee TH, Hoh R, et al. A randomized, controlled trial of raltegravir intensification in antiretroviral-treated, HIV-infected patients with a suboptimal CD4+ T cell response. *Journal of Infectious Diseases*. 2011; 203(7):960–968. <http://dx.doi.org/10.1093/infdis/jiq138>. [PubMed: 21402547]
- Hatano, H.; Strain, M.; Scherzer, R.; Sinclair, E.; Palmer, S.; Busch, M., et al. Increase in 2-LTR circles after raltegravir intensification in HAART-suppressed patients with high CD4+T Cell counts: A randomized, controlled trial. Paper presented at the in the Program and Abstracts from the 20th Conference on Retroviruses and Opportunistic Infections; Atlanta, GA. 2013. Abstract # 42
- Heaton RK, Clifford DB, Franklin DR Jr, Woods SP, Ake C, Vaida F, et al. HIV-associated neurocognitive disorders persist in the era of potent antiretroviral therapy: CHARTER Study. *Neurology*. 2010; 75(23):2087–2096. <http://dx.doi.org/10.1212/WNL.0b013e318200d727>. [PubMed: 21135382]
- Heil F, Hemmi H, Hochrein H, Ampenberger F, Kirschning C, Akira S, et al. Species-specific recognition of single-stranded RNA via toll-like receptor 7 and 8. *Science*. 2004; 303(5663):1526–1529. [PubMed: 14976262]
- Hunt PW, Cao HL, Muzoora C, Ssewanyana I, Bennett J, Emenyonu N, et al. Impact of CD8+ T-cell activation on CD4+ T-cell recovery and mortality in HIV-infected Ugandans initiating antiretroviral therapy. *AIDS*. 2011; 25(17):2123–2131. <http://dx.doi.org/10.1097/QAD.0b013e32834c4ac1>. [PubMed: 21881481]
- Hunt PW, Deeks SG, Rodriguez B, Valdez H, Shade SB, Abrams DI, et al. Continued CD4 cell count increases in HIV-infected adults experiencing 4 years of viral suppression on antiretroviral therapy. *AIDS*. 2003; 17(13):1907–1915. [PubMed: 12960823]

- Hunt PW, Martin JN, Sinclair E, Epling L, Teague J, Jacobson MA, et al. Valganciclovir reduces T cell activation in HIV-infected individuals with incomplete CD4+ T cell recovery on antiretroviral therapy. *Journal of Infectious Diseases*. 2011; 203(10):1474–1483. <http://dx.doi.org/10.1093/infdis/jir060>. [PubMed: 21502083]
- Hunt, P.; Rodriguez, B.; Shive, C.; Clagett, B.; Funderburg, N.; Natta, MV., et al. Gut epithelial barrier dysfunction, inflammation, and coagulation predict higher mortality during treated HIV/AIDS. Paper presented at the in the Program and Abstracts from the 19th Conference on Retroviruses and Opportunistic Infections; Seattle, WA. 2012. Abstract #278
- Hunt PW, Shulman N, Hayes TL, Dahl V, Somsouk M, Funderburg NT, et al. The immunologic effects of maraviroc intensification in treated HIV-infected individuals with incomplete CD4+ T cell recovery: A randomized trial. *Blood*. 2013; 121(23):4635–4646. <http://dx.doi.org/10.1182/blood-2012-06-436345>. [PubMed: 23589670]
- Jasny E, Geer S, Frank I, Vagenas P, Aravantinou M, Salazar AM, et al. Characterization of peripheral and mucosal immune responses in rhesus macaques on long-term tenofovir and emtricitabine combination antiretroviral therapy. *Journal of Acquired Immune Deficiency Syndromes*. 2012; 61(4):425–435. <http://dx.doi.org/10.1097/QAI.0b013e318266be53>. [PubMed: 22820802]
- Jiang W, Lederman MM, Hunt P, Sieg SF, Haley K, Rodriguez B, et al. Plasma levels of bacterial DNA correlate with immune activation and the magnitude of immune restoration in persons with antiretroviral-treated HIV infection. *Journal of Infectious Diseases*. 2009; 199(8):1177–1185. <http://dx.doi.org/10.1086/597476>. [PubMed: 19265479]
- Justice AC, Freiberg MS, Tracy R, Kuller L, Tate JP, Goetz MB, et al. Does an index composed of clinical data reflect effects of inflammation, coagulation, and monocyte activation on mortality among those aging with HIV? *Clinical Infectious Diseases: An Official Publication of the Infectious Diseases Society of America*. 2012; 54(7):984–994. <http://dx.doi.org/10.1093/cid/cir989>. [PubMed: 22337823]
- Kalayjian RC, Machekano RN, Rizk N, Robbins GK, Gandhi RT, Rodriguez BA, et al. Pretreatment levels of soluble cellular receptors and interleukin-6 are associated with HIV disease progression in subjects treated with highly active antiretroviral therapy. *Journal of Infectious Diseases*. 2010; 201(12):1796–1805. <http://dx.doi.org/10.1086/652750>. [PubMed: 20446847]
- Kalinowska M, Bazdar DA, Lederman MM, Funderburg N, Sieg SF. Decreased IL-7 responsiveness is related to oxidative stress in HIV disease. *PLoS One*. 2013; 8(3):e58764. <http://dx.doi.org/10.1371/journal.pone.0058764>. [PubMed: 23505558]
- Kelley CF, Kitchen CM, Hunt PW, Rodriguez B, Hecht FM, Kitahata M, et al. Incomplete peripheral CD4+ cell count restoration in HIV-infected patients receiving long-term antiretroviral treatment. *Clinical Infectious Diseases*. 2009; 48(6):787–794. <http://dx.doi.org/10.1086/597093>. [PubMed: 19193107]
- Kiviat NB, Critchlow CW, Hawes SE, Kuypers J, Surawicz C, Goldbaum G, et al. Determinants of human immunodeficiency virus DNA and RNA shedding in the anal-rectal canal of homosexual men. *Journal of Infectious Diseases*. 1998; 177(3):571–578. [PubMed: 9498434]
- Klatt NR, Brechley JM. Th17 cell dynamics in HIV infection. *Current Opinion in HIV and AIDS*. 2010; 5(2):135–140. <http://dx.doi.org/10.1097/COH.0b013e31823364846>. [PubMed: 20543590]
- Klatt NR, Canary LA, Sun X, Vinton CL, Funderburg NT, Morcock DR, et al. Probiotic/prebiotic supplementation of antiretrovirals improves gastrointestinal immunity in SIV-infected macaques. *The Journal of Clinical Investigation*. 2013; 123(2):903–907. <http://dx.doi.org/10.1172/JCI66227>. [PubMed: 23321668]
- Klatt NR, Estes JD, Sun X, Ortiz AM, Barber JS, Harris LD, et al. Loss of mucosal CD103+ DCs and IL-17+ and IL-22+ lymphocytes is associated with mucosal damage in SIV infection. *Mucosal Immunology*. 2012; 5(6):646–657. <http://dx.doi.org/10.1038/mi.2012.38>. [PubMed: 22643849]
- Klatt NR, Funderburg NT, Brechley JM. Microbial translocation, immune activation, and HIV disease. *Trends in Microbiology*. 2013; 21(1):6–13. <http://dx.doi.org/10.1016/j.tim.2012.09.001>. [PubMed: 23062765]
- Klatt NR, Harris LD, Vinton CL, Sung H, Briant JA, Tabb B, et al. Compromised gastrointestinal integrity in pigtail macaques is associated with increased microbial translocation, immune activation, and IL-17 production in the absence of SIV infection. *Mucosal Immunology*. 2010; 3(4):387–398. <http://dx.doi.org/10.1038/mi.2010.14>. [PubMed: 20357762]

- Klatt NR, Silvestri G, Hirsch V. Nonpathogenic simian immunodeficiency virus infections. *Cold Spring Harbor Perspectives in Medicine*. 2012; 2(1):a007153. <http://dx.doi.org/10.1101/cshperspect.a007153>. [PubMed: 22315718]
- Kuller LH, Tracy R, Belloso W, De Wit S, Drummond F, Lane HC, et al. Inflammatory and coagulation biomarkers and mortality in patients with HIV infection. *PLoS Medicine*. 2008; 5(10):e203. <http://dx.doi.org/10.1371/journal.pmed.0050203>. [PubMed: 18942885]
- Lane HC, Masur H, Edgar LC, Whalen G, Rook AH, Fauci AS. Abnormalities of B-cell activation and immunoregulation in patients with the acquired immunodeficiency syndrome. *The New England Journal of Medicine*. 1983; 309(8):453–458. [PubMed: 6224088]
- Lederman MM, Calabrese L, Funderburg NT, Clagett B, Medvik K, Bonilla H, et al. Immunologic failure despite suppressive antiretroviral therapy is related to activation and turnover of memory CD4 cells. *Journal of Infectious Diseases*. 2011; 204(8):1217–1226. <http://dx.doi.org/10.1093/infdis/jir507>. [PubMed: 21917895]
- Lederman MM, Connick E, Landay A, Kuritzkes DR, Spritzler J, St Clair M, et al. Immunologic responses associated with 12 weeks of combination antiretroviral therapy consisting of zidovudine, lamivudine, and ritonavir: Results of AIDS Clinical Trials Group Protocol 315. *Journal of Infectious Diseases*. 1998; 178(1):70–79. [PubMed: 9652425]
- Letendre, S.; Marcotte, T.; Deutsch, R.; Franklin, D.; Rosario, D.; Alexander, T., et al. A concise panel of biomarkers diagnoses and predicts neurocognitive status in HIV-infected individuals. Paper presented at the in the Program and Abstracts from the 19th Conference on Retroviruses and Opportunistic Infections; Seattle, WA. 2012. Abstract #82
- Lewden C, Bouteloup V, De Wit S, Sabin C, Mocroft A, Wasmuth JC, et al. All-cause mortality in treated HIV-infected adults with CD4 $500/\text{mm}^3$ compared with the general population: Evidence from a large European observational cohort collaboration. *International Journal of Epidemiology*. 2012; 41(2):433–445. <http://dx.doi.org/10.1093/ije/dyr164>. [PubMed: 22493325]
- Lewden C, Chene G, Morlat P, Raffi F, Dupon M, Dellamonica P, et al. HIV-infected adults with a CD4 cell count greater than $500 \text{ cells}/\text{mm}^3$ on long-term combination antiretroviral therapy reach same mortality rates as the general population. *Journal of Acquired Immune Deficiency Syndromes*. 2007; 46(1):72–77. <http://dx.doi.org/10.1097/QAI.0b013e318134257a>. [PubMed: 17621240]
- Lichtenstein KA, Armon C, Buchacz K, Chmiel JS, Buckner K, Tedaldi EM, et al. Low CD4+ T cell count is a risk factor for cardiovascular disease events in the HIV outpatient study. *Clinical Infectious Diseases*. 2010; 51(4):435–447. <http://dx.doi.org/10.1086/655144>. [PubMed: 20597691]
- Lichtfuss GF, Hoy J, Rajasuriar R, Kramski M, Crowe SM, Lewin SR. Biomarkers of immune dysfunction following combination antiretroviral therapy for HIV infection. *Biomarkers in Medicine*. 2011; 5(2):171–186. <http://dx.doi.org/10.2217/bmm.11.15>. [PubMed: 21473720]
- Lohse N, Hansen AB, Pedersen G, Kronborg G, Gerstoft J, Sorensen HT, et al. Survival of persons with and without HIV infection in Denmark, 1995–2005. *Annals of Internal Medicine*. 2007; 146(2):87–95. [PubMed: 17227932]
- Lyons JL, Uno H, Ancuta P, Kamat A, Moore DJ, Singer EJ, et al. Plasma sCD14 is a biomarker associated with impaired neurocognitive test performance in attention and learning domains in HIV infection. *Journal of Acquired Immune Deficiency Syndromes*. 2011; 57(5):371–379. <http://dx.doi.org/10.1097/QAI.0b013e3182237e54>. [PubMed: 21646912]
- Macal M, Sankaran S, Chun TW, Reay E, Flamm J, Prindiville TJ, et al. Effective CD4+ T-cell restoration in gut-associated lymphoid tissue of HIV-infected patients is associated with enhanced Th17 cells and polyfunctional HIV-specific T-cell responses. *Mucosal Immunology*. 2008; 1(6):475–488. <http://dx.doi.org/10.1038/mi.2008.35>. [PubMed: 19079215]
- Maldarelli F, Palmer S, King MS, Wiegand A, Polis MA, Mican J, et al. ART suppresses plasma HIV-1 RNA to a stable set point predicted by pretherapy viremia. *PLoS Pathogens*. 2007; 3(4):e46. [PubMed: 17411338]
- Marchetti G, Bellistri GM, Borghi E, Tincati C, Ferramosca S, La Francesca M, et al. Microbial translocation is associated with sustained failure in CD4+ T-cell reconstitution in HIV-infected patients on long-term highly active antiretroviral therapy. *AIDS*. 2008; 22(15):2035–2038. <http://dx.doi.org/10.1097/QAD.0b013e3283112d29>. [PubMed: 18784466]

- Marin B, Thiebaut R, Bucher HC, Rondeau V, Costagliola D, Dorrucchi M, et al. Non-AIDS-defining deaths and immunodeficiency in the era of combination antiretroviral therapy. *AIDS*. 2009; 23(13):1743–1753. <http://dx.doi.org/10.1097/QAD.0b013e32832e9b78>. [PubMed: 19571723]
- Martinez E, D'Albuquerque PM, Llibre JM, Gutierrez F, Podzamczar D, Antela A, et al. Changes in cardiovascular biomarkers in HIV-infected patients switching from ritonavir-boosted protease inhibitors to raltegravir. *AIDS*. 2012; 26(18):2315–2326. <http://dx.doi.org/10.1097/QAD.0b013e328359f29c>. [PubMed: 23018438]
- Marziali M, De Santis W, Carello R, Leti W, Esposito A, Isgro A, et al. T-cell homeostasis alteration in HIV-1 infected subjects with low CD4 T-cell count despite undetectable virus load during HAART. *AIDS*. 2006; 20(16):2033–2041. <http://dx.doi.org/10.1097/01.aids.0000247588.69438.fd>. [PubMed: 17053349]
- Massanella M, Negredo E, Puig J, Puertas MC, Buzon MJ, Perez-Alvarez N, et al. Raltegravir intensification shows differing effects on CD8 and CD4 T cells in HIV-infected HAART-suppressed individuals with poor CD4 T-cell recovery. *AIDS*. 2012; 26(18):2285–2293. <http://dx.doi.org/10.1097/QAD.0b013e328359f20f>. [PubMed: 23018435]
- Masur H, Michelis MA, Greene JB, Onorato I, Stouwe RA, Holzman RS, et al. An outbreak of community-acquired *Pneumocystis carinii* pneumonia: Initial manifestation of cellular immune dysfunction. *The New England Journal of Medicine*. 1981; 305(24):1431–1438. [PubMed: 6975437]
- Mattapallil JJ, Douek DC, Hill B, Nishimura Y, Martin M, Roederer M. Massive infection and loss of memory CD4+ T cells in multiple tissues during acute SIV infection. *Nature*. 2005; 7037; 434:1093–1097. [PubMed: 15793563]
- Mavigner M, Cazabat M, Dubois M, L'Faqihi FE, Requena M, Pasquier C, et al. Altered CD4+ T cell homing to the gut impairs mucosal immune reconstitution in treated HIV-infected individuals. *The Journal of Clinical Investigation*. 2012; 122(1):62–69. <http://dx.doi.org/10.1172/JCI59011>. [PubMed: 22156200]
- McComsey, G.; Jiang, Y.; Debanne, S.; Clagett, B.; Robinson, J.; Labbato, D., et al. Effect of statins on immune activation and inflammation in HIV+subjects on ART: A randomized placebo controlled trial. Paper presented at the In the Program and Abstracts from the 20th Conference on Retroviruses and Opportunistic Infections; Atlanta, GA. 2013. Abstract #186LB
- Mehandru S, Poles MA, Tenner-Racz K, Jean-Pierre P, Manuelli V, Lopez P, et al. Lack of mucosal immune reconstitution during prolonged treatment of acute and early HIV-1 infection. *PLoS Medicine*. 2006; 3(12):e484. [PubMed: 17147468]
- Meier A, Alter G, Frahm N, Sidhu H, Li B, Bagchi A, et al. MyD88-dependent immune activation mediated by human immunodeficiency virus type 1-encoded toll-like receptor ligands. *Journal of Virology*. 2007; 81(15):8180–8191. [PubMed: 17507480]
- Merlini E, Bai F, Bellistri GM, Tincati C, d'Arminio Monforte A, Marchetti G. Evidence for polymicrobial flora translocating in peripheral blood of HIV-infected patients with poor immune response to antiretroviral therapy. *PLoS One*. 2011; 6(4):e18580. <http://dx.doi.org/10.1371/journal.pone.0018580>. [PubMed: 21494598]
- Miller EA, Spadaccia MR, O'Brien MP, Rolnitzky L, Sabado R, Manches O, et al. Plasma factors during chronic HIV-1 infection impair IL-12 secretion by myeloid dendritic cells via a virus-independent pathway. *Journal of Acquired Immune Deficiency Syndromes*. 2012; 61(5):535–544. <http://dx.doi.org/10.1097/QAI.0b013e31826afbce>. [PubMed: 22902724]
- Mocroft A, Reiss P, Gasiorowski J, Ledergerber B, Kowalska J, Chiesi A, et al. Serious fatal and nonfatal non-AIDS-defining illnesses in Europe. *Journal of Acquired Immune Deficiency Syndromes*. 2010; 55(2):262–270. <http://dx.doi.org/10.1097/QAI.0b013e3181e9be6b>. [PubMed: 20700060]
- Modjarrad K, Vermund SH. Effect of treating co-infections on HIV-1 viral load: A systematic review. *The Lancet Infectious Diseases*. 2010; 10(7):455–463. [http://dx.doi.org/10.1016/S1473-3099\(10\)70093-1](http://dx.doi.org/10.1016/S1473-3099(10)70093-1). [PubMed: 20610327]
- Mudd JC, Murphy P, Manion M, Debernardo R, Hardacre J, Ammori J, et al. Impaired T cell responses to sphingosine-1-phosphate in HIV-1 infected lymph nodes. *Blood*. 2013; 121(15): 2914–2922. <http://dx.doi.org/10.1182/blood-2012-07-445783>. [PubMed: 23422746]

- Murray SM, Down CM, Boulware DR, Stauffer WM, Cavert WP, Schacker TW, et al. Reduction of immune activation with chloroquine therapy during chronic HIV infection. *Journal of Virology*. 2010; 84(22):12082–12086. <http://dx.doi.org/10.1128/JVI.01466-10>. [PubMed: 20844049]
- Musselwhite LW, Sheikh V, Norton TD, Rupert A, Porter BO, Penzak SR, et al. Markers of endothelial dysfunction, coagulation and tissue fibrosis independently predict venous thromboembolism in HIV. *AIDS*. 2011; 25(6):787–795. <http://dx.doi.org/10.1097/QAD.0b013e3283453fcb>. [PubMed: 21412059]
- Naeger DM, Martin JN, Sinclair E, Hunt PW, Bangsberg DR, Hecht F, et al. Cytomegalovirus-specific T cells persist at very high levels during long-term antiretroviral treatment of HIV disease. *PLoS One*. 2010; 5(1):e8886. <http://dx.doi.org/10.1371/journal.pone.0008886>. [PubMed: 20126452]
- Napolitano LA, Grant RM, Deeks SG, Schmidt D, De Rosa SC, Herzenberg LA, et al. Increased production of IL-7 accompanies HIV-1-mediated T-cell depletion: Implications for T-cell homeostasis. *Nature Medicine*. 2001; 7(1):73–79.
- Nazli A, Chan O, Dobson-Belaire WN, Ouellet M, Tremblay MJ, Gray-Owen SD, et al. Exposure to HIV-1 directly impairs mucosal epithelial barrier integrity allowing microbial translocation. *PLoS Pathogens*. 2010; 6(4):e1000852. [PubMed: 20386714]
- O'Brien M, Montenont E, Hu L, Nardi MA, Valdes V, Merolla M, et al. Aspirin attenuates platelet activation and immune activation in HIV-infected subjects on antiretroviral therapy: A pilot study. *Journal of Acquired Immune Deficiency Syndromes*. 2013; 63(3):280–288. <http://dx.doi.org/10.1097/QAI.0b013e31828a292c>. [PubMed: 23406976]
- Pakker NG, Notermans DW, de Boer RJ, Roos MT, de Wolf F, Hill A, et al. Biphasic kinetics of peripheral blood T cells after triple combination therapy in HIV-1 infection: A composite of redistribution and proliferation. *Nature Medicine*. 1998; 4(2):208–214.
- Palmer S, Maldarelli F, Wiegand A, Bernstein B, Hanna GJ, Brun SC, et al. Low-level viremia persists for at least 7 years in patients on suppressive antiretroviral therapy. *Proceedings of the National Academy of Sciences of the United States of America*. 2008; 105(10):3879–3884. [PubMed: 18332425]
- Pandrea I. Therapeutic interventions to modulate microbial translocation, immune activation and cardiovascular comorbidities. Lessons from nonhuman primate models. Paper presented at the Keystone Symposia on Molecular and Cellular Biology. *Immune Activation in HIV infection: Basic mechanisms and clinical implications*; Breckenridge, Colorado. 2013.
- Pantaleo G, Graziosi C, Butini L, Pizzo PA, Schnittman SM, Kotler DP, et al. Lymphoid organs function as major reservoirs for human immunodeficiency virus. *Proceedings of the National Academy of Sciences of the United States of America*. 1991; 88(21):9838–9842. [PubMed: 1682922]
- Pantaleo G, Graziosi C, Demarest JF, Butini L, Montroni M, Fox CH, et al. HIV infection is active and progressive in lymphoid tissue during the clinically latent stage of disease. *Nature*. 1993; 362(6418):355–358. <http://dx.doi.org/10.1038/362355a0>. [PubMed: 8455722]
- Pasare C, Medzhitov R. Toll-dependent control mechanisms of CD4 T cell activation. *Immunity*. 2004; 21(5):733–741. [PubMed: 15539158]
- Patel P, Hanson DL, Sullivan PS, Novak RM, Moorman AC, Tong TC, et al. Incidence of types of cancer among HIV-infected persons compared with the general population in the United States, 1992–2003. *Annals of Internal Medicine*. 2008; 148(10):728–736. [PubMed: 18490686]
- Paton NI, Goodall RL, Dunn DT, Franzen S, Collaco-Moraes Y, Gazzard BG, et al. Effects of hydroxychloroquine on immune activation and disease progression among HIV-infected patients not receiving antiretroviral therapy: A randomized controlled trial. *JAMA: The Journal of the American Medical Association*. 2012; 308(4):353–361. <http://dx.doi.org/10.1001/jama.2012.6936>. [PubMed: 22820788]
- Picker LJ. Immunopathogenesis of acute AIDS virus infection. *Current Opinion in Immunology*. 2006; 18(4):399–405. [PubMed: 16753288]
- Piconi S, Parisotto S, Rizzardini G, Passerini S, Meraviglia P, Schiavini M, et al. Atherosclerosis is associated with multiple pathogenic mechanisms in HIV-infected antiretroviral-naïve or treated individuals. *AIDS*. 2013; 27(3):381–389. <http://dx.doi.org/10.1097/QAD.0b013e32835abcc9>. [PubMed: 23079800]

- Piconi S, Parisotto S, Rizzardini G, Passerini S, Terzi R, Argentero B, et al. Hydroxychloroquine drastically reduces immune activation in HIV-infected, antiretroviral therapy-treated immunologic nonresponders. *Blood*. 2011; 118(12):3263–3272. <http://dx.doi.org/10.1182/blood-2011-01-329060>. [PubMed: 21576701]
- Pilakka-Kanthikeel S, Huang S, Fenton T, Borkowsky W, Cunningham CK, Pahwa S. Increased Gut microbial translocation in HIV-infected children persists in virologic responders and virologic failures after antiretroviral therapy. *The Pediatric Infectious Disease Journal*. 2012; 31(6):583–591. <http://dx.doi.org/10.1097/INF.0b013e31824da0f5>. [PubMed: 22333700]
- Reeves RK, Evans TI, Gillis J, Wong FE, Kang G, Li Q, et al. SIV infection induces accumulation of plasmacytoid dendritic cells in the gut mucosa. *Journal of Infectious Diseases*. 2012; 206(9):1462–1468. <http://dx.doi.org/10.1093/infdis/jis408>. [PubMed: 22711907]
- Reeves RK, Rajakumar PA, Evans TI, Connole M, Gillis J, Wong FE, et al. Gut inflammation and indoleamine deoxygenase inhibit IL-17 production and promote cytotoxic potential in NKp44+ mucosal NK cells during SIV infection. *Blood*. 2011; 118(12):3321–3330. <http://dx.doi.org/10.1182/blood-2011-04-347260>. [PubMed: 21791421]
- Reuben DB, Cheh AI, Harris TB, Ferrucci L, Rowe JW, Tracy RP, et al. Peripheral blood markers of inflammation predict mortality and functional decline in high-functioning community-dwelling older persons. *Journal of the American Geriatrics Society*. 2002; 50(4):638–644. [PubMed: 11982663]
- Rodger AJ, Fox Z, Lundgren JD, Kuller LH, Boesecke C, Gey D, et al. Activation and coagulation biomarkers are independent predictors of the development of opportunistic disease in patients with HIV infection. *Journal of Infectious Diseases*. 2009; 200(6):973–983. <http://dx.doi.org/10.1086/605447>. [PubMed: 19678756]
- Rose H, Hoy J, Woolley I, Tchoua U, Bukrinsky M, Dart A, et al. HIV infection and high density lipoprotein metabolism. *Atherosclerosis*. 2008; 199(1):79–86. <http://dx.doi.org/10.1016/j.atherosclerosis.2007.10.018>. [PubMed: 18054941]
- Sacre K, Hunt PW, Hsue PY, Maidji E, Martin JN, Deeks SG, et al. A role for cytomegalovirus-specific CD4+CX3CR1+ T cells and cytomegalovirus-induced T-cell immunopathology in HIV-associated atherosclerosis. *AIDS*. 2012; 26(7):805–814. <http://dx.doi.org/10.1097/QAD.0b013e328351f780>. [PubMed: 22313962]
- Sandler NG, Wand H, Roque A, Law M, Nason MC, Nixon DE, et al. Plasma levels of soluble CD14 independently predict mortality in HIV infection. *Journal of Infectious Diseases*. 2011; 203(6):780–790. <http://dx.doi.org/10.1093/infdis/jiq118>. [PubMed: 21252259]
- Schacker T, Collier AC, Hughes J, Shea T, Corey L. Clinical and epidemiologic features of primary HIV infection. *Annals of Internal Medicine*. 1996; 125(4):257–264. [PubMed: 8678387]
- Schacker TW, Nguyen PL, Beilman GJ, Wolinsky S, Larson M, Reilly C, et al. Collagen deposition in HIV-1 infected lymphatic tissues and T cell homeostasis. *The Journal of Clinical Investigation*. 2002; 110(8):1133–1139. [PubMed: 12393849]
- Schacker TW, Nguyen PL, Martinez E, Reilly C, Gatell JM, Horban A, et al. Persistent abnormalities in lymphoid tissues of human immunodeficiency virus-infected patients successfully treated with highly active antiretroviral therapy. *Journal of Infectious Diseases*. 2002; 186(8):1092–1097. [PubMed: 12355359]
- Sereti I, Estes J, Thompson W, Fischl M, Croughs T, Beq S, et al. Gut mucosa T lymphocyte restoration in chronically HIV+patients treated with recombinant interleukin-7. Paper presented at the in the Program and Abstracts from the 19th Conference on Retroviruses and Opportunistic Infections; Seattle, WA. 2012. Abstract # 94
- Silverberg MJ, Chao C, Leyden WA, Xu L, Tang B, Horberg MA, et al. HIV infection and the risk of cancers with and without a known infectious cause. *AIDS*. 2009; 23(17):2337–2345. <http://dx.doi.org/10.1097/QAD.0b013e3283319184>. [PubMed: 19741479]
- Silvestri G, Paiardini M, Pandrea I, Lederman MM, Sodora DL. Understanding the benign nature of SIV infection in natural hosts. *The Journal of Clinical Investigation*. 2007; 117(11):3148–3154. [PubMed: 17975656]
- Smurzynski M, Wu K, Benson CA, Bosch RJ, Collier AC, Koletar SL. Relationship between CD4+ T-cell counts/HIV-1 RNA plasma viral load and AIDS-defining events among persons followed in the ACTG longitudinal linked randomized trials study. *Journal of Acquired Immune Deficiency*

- Syndromes. 2010; 55(1):117–127. <http://dx.doi.org/10.1097/QAI.0b013e3181e8c129>. [PubMed: 20622677]
- Stacey AR, Norris PJ, Qin L, Haygreen EA, Taylor E, Heitman J, et al. Induction of a striking systemic cytokine cascade prior to peak viremia in acute human immunodeficiency virus type 1 infection, in contrast to more modest and delayed responses in acute hepatitis B and C virus infections. *Journal of Virology*. 2009; 83(8):3719–3733. <http://dx.doi.org/10.1128/JVI.01844-08>. [PubMed: 19176632]
- Stewart CR, Stuart LM, Wilkinson K, van Gils JM, Deng J, Halle A, et al. CD36 ligands promote sterile inflammation through assembly of a toll-like receptor 4 and 6 heterodimer. *Nature Immunology*. 2010; 11(2):155–161. <http://dx.doi.org/10.1038/ni.1836>. [PubMed: 20037584]
- Sullivan PS, Dworkin MS, Jones JL, Hooper WC. Epidemiology of thrombosis in HIV-infected individuals. The adult/adolescent spectrum of HIV disease project. *AIDS*. 2000; 14(3):321–324. [PubMed: 10716509]
- Tenorio, A.; Zheng, E.; Bosch, R.; Deeks, SG.; Rodriguez, B.; Krishnan, S., et al. Soluble markers of inflammation & coagulation, but not T-cell activation, predict non-AIDS-defining events during suppressive antiretroviral therapy. Paper presented at the in the Program and Abstracts of the 20th Conference on Retroviruses and Opportunistic Infections; 2013; Atlanta, GA. 2013. Abstract #790
- The Antiretroviral Therapy Cohort Collaboration. Life expectancy of individuals on combination antiretroviral therapy in high-income countries: A collaborative analysis of 14 cohort studies. *The Lancet*. 2008; 372(9635):293–299. [http://dx.doi.org/10.1016/S0140-6736\(08\)61113-7](http://dx.doi.org/10.1016/S0140-6736(08)61113-7).
- Tien PC, Choi AI, Zolopa AR, Benson C, Tracy R, Scherzer R, et al. Inflammation and mortality in HIV-infected adults: Analysis of the FRAM study cohort. *Journal of Acquired Immune Deficiency Syndromes*. 2010; 55(3):316. [PubMed: 20581689]
- Tincati C, Biasin M, Bandera A, Violin M, Marchetti G, Piacentini L, et al. Early initiation of highly active antiretroviral therapy fails to reverse immunovirological abnormalities in gut-associated lymphoid tissue induced by acute HIV infection. *Antiviral Therapy*. 2009; 14(3):321–330. [PubMed: 19474466]
- Tindall B, Barker S, Donovan B, Barnes T, Roberts J, Kronenberg C, et al. Characterization of the acute clinical illness associated with human immunodeficiency virus infection. *Archives of Internal Medicine*. 1988; 148(4):945–949. [PubMed: 3258508]
- Triant VA, Brown TT, Lee H, Grinspoon SK. Fracture prevalence among human immunodeficiency virus (HIV)-infected versus non-HIV-infected patients in a large U.S. healthcare system. *The Journal of Clinical Endocrinology and Metabolism*. 2008; 93(9):3499–3504. [PubMed: 18593764]
- Triant VA, Lee H, Hadigan C, Grinspoon SK. Increased acute myocardial infarction rates and cardiovascular risk factors among patients with human immunodeficiency virus disease. *The Journal of Clinical Endocrinology and Metabolism*. 2007; 92(7):2506–2512. <http://dx.doi.org/10.1210/jc.2006-2190>. [PubMed: 17456578]
- Triant VA, Meigs JB, Grinspoon SK. Association of C-reactive protein and HIV infection with acute myocardial infarction. *Journal of Acquired Immune Deficiency Syndromes*. 2009; 51(3):268. [PubMed: 19387353]
- Triant VA, Regan S, Lee H, Sax PE, Meigs JB, Grinspoon SK. Association of immunologic and virologic factors with myocardial infarction rates in a US healthcare system. *Journal of Acquired Immune Deficiency Syndromes*. 2010; 55(5):615–619. <http://dx.doi.org/10.1097/QAI.0b013e3181f4b752>. [PubMed: 20827215]
- Tseng ZH, Secemsky EA, Dowdy D, Vittinghoff E, Moyers B, Wong JK, et al. Sudden cardiac death in patients with human immunodeficiency virus infection. *Journal of the American College of Cardiology*. 2012; 59(21):1891–1896. <http://dx.doi.org/10.1016/j.jacc.2012.02.024>. [PubMed: 22595409]
- van Sighem AI, Gras LA, Reiss P, Brinkman K, de Wolf F. Life expectancy of recently diagnosed asymptomatic HIV-infected patients approaches that of uninfected individuals. *AIDS*. 2010; 24(10):1527–1535. <http://dx.doi.org/10.1097/QAD.0b013e32833a3946>. [PubMed: 20467289]

- Veazey RS, DeMaria M, Chalifoux LV, Shvetz DE, Pauley DR, Knight HL, et al. Gastrointestinal tract as a major site of CD4+ T cell depletion and viral replication in SIV infection. *Science*. 1998; 280(5362):427–431. <http://dx.doi.org/10.1126/science.280.5362.427>. [PubMed: 9545219]
- Volpato S, Guralnik JM, Ferrucci L, Balfour J, Chaves P, Fried LP, et al. Cardiovascular disease, interleukin-6, and risk of mortality in older women: The women's health and aging study. *Circulation*. 2001; 103(7):947–953. [PubMed: 11181468]
- Wikby A, Nilsson BO, Forsey R, Thompson J, Strindhall J, Lofgren S, et al. The immune risk phenotype is associated with IL-6 in the terminal decline stage: Findings from the Swedish NONA immune longitudinal study of very late life functioning. *Mechanisms of Ageing and Development*. 2006; 127(8):695–704. [PubMed: 16750842]
- Yong MK, Elliott JH, Woolley IJ, Hoy JF. Low CD4 count is associated with an increased risk of fragility fracture in HIV-infected patients. *Journal of Acquired Immune Deficiency Syndromes*. 2011; 57(3):205–210. <http://dx.doi.org/10.1097/QAI.0b013e31821ecf4c>. [PubMed: 21522014]
- Young J, Psychogiou M, Meyer L, Ayayi S, Grabar S, Raffi F, et al. CD4 cell count and the risk of AIDS or death in HIV-Infected adults on combination antiretroviral therapy with a suppressed viral load: A longitudinal cohort study from COHERE. *PLoS Medicine*. 2012; 9(3):e1001194. <http://dx.doi.org/10.1371/journal.pmed.1001194>. [PubMed: 22448150]
- Yukl SA, Gianella S, Sinclair E, Epling L, Li Q, Duan L, et al. Differences in HIV burden and immune activation within the gut of HIV-positive patients receiving suppressive antiretroviral therapy. *Journal of Infectious Diseases*. 2010; 202(10):1553–1561. <http://dx.doi.org/10.1086/656722>. [PubMed: 20939732]
- Zeng M, Smith AJ, Wietgreffe SW, Southern PJ, Schacker TW, Reilly CS, et al. Cumulative mechanisms of lymphoid tissue fibrosis and T cell depletion in HIV-1 and SIV infections. *The Journal of Clinical Investigation*. 2011; 121(3):998–1008. <http://dx.doi.org/10.1172/JCI45157>. [PubMed: 21393864]
- Zonios DI, Falloon J, Bennett JE, Shaw PA, Chaitt D, Baseler MW, et al. Idiopathic CD4+ lymphocytopenia: Natural history and prognostic factors. *Blood*. 2008; 112(2):287–294. <http://dx.doi.org/10.1182/blood-2007-12-127878>. [PubMed: 18456875]
- Zuckerman RA, Whittington WL, Celum CL, Collis TK, Lucchetti AJ, Sanchez JL, et al. Higher concentration of HIV RNA in rectal mucosa secretions than in blood and seminal plasma, among men who have sex with men, independent of antiretroviral therapy. *Journal of Infectious Diseases*. 2004; 190(1):156–161. <http://dx.doi.org/10.1086/421246>. [PubMed: 15195255]