

Lawrence Berkeley National Laboratory

LBL Publications

Title

Consider the Anoxic Microsite: Acknowledging and Appreciating Spatiotemporal Redox Heterogeneity in Soils and Sediments.

Permalink

<https://escholarship.org/uc/item/9cj5w6rj>

Journal

ACS Earth and Space Chemistry, 7(9)

ISSN

2472-3452

Authors

Lacroix, Emily
Aeppli, Meret
Boye, Kristin
[et al.](#)

Publication Date

2023-09-21

DOI

10.1021/acsearthspacechem.3c00032

Peer reviewed

Consider the Anoxic Microsite: Acknowledging and Appreciating Spatiotemporal Redox Heterogeneity in Soils and Sediments

Emily M. Lacroix,* Meret Aeppli, Kristin Boye, Eoin Brodie, Scott Fendorf, Marco Keiluweit, Hannah R. Naughton, Vincent Noël, and Debjani Sihi


Cite This: *ACS Earth Space Chem.* 2023, 7, 1592–1609

Read Online

ACCESS |

Metrics & More

Article Recommendations



ABSTRACT: Reduction–oxidation (redox) reactions underlie essentially all biogeochemical cycles. Like most soil properties and processes, redox is spatiotemporally heterogeneous. However, unlike other soil features, redox heterogeneity has yet to be incorporated into mainstream conceptualizations of soil biogeochemistry. Anoxic microsites, the defining feature of redox heterogeneity in bulk oxic soils and sediments, are zones of oxygen depletion in otherwise oxic environments. In this review, we suggest that anoxic microsites represent a critical component of soil function and that appreciating anoxic microsites promises to advance our understanding of soil and sediment biogeochemistry. In sections 1 and 2, we define anoxic microsites and highlight their dynamic properties, specifically anoxic microsite distribution, redox gradient magnitude, and temporality. In section 3, we describe the influence of anoxic microsites on several key elemental cycles, organic carbon, nitrogen, iron, manganese, and sulfur. In section 4, we evaluate methods for identifying and characterizing anoxic microsites, and in section 5, we highlight past and current approaches to modeling anoxic microsites. Finally, in section 6, we suggest steps for incorporating anoxic microsites and redox heterogeneities more broadly into our understanding of soils and sediments.

KEYWORDS: *anoxic microsite, oxygen, soil, biogeochemistry, redox heterogeneity*

1. INTRODUCTION

Reduction–oxidation (redox) reactions, the biotic or abiotic transfer of electrons, underlie essentially all biogeochemical cycles. Redox constrains the energetic bounds of cellular growth, governs the bioavailability of nutrients and contaminants, and drives chemical weathering of rocks and minerals. In soils, redox reactions are paramount as they connect elemental cycles across the atmosphere, biosphere, hydrosphere, and lithosphere, the mixed media of soils.

Oxygen (O_2) is a central regulator of soil redox cycling. A potent oxidant, O_2 facilitates abiotic redox transformations, particularly at redox interfaces, where strong oxidizers (e.g.,

O_2) and reduced species meet. Oxygen also has a pronounced effect on microbial respiration, the principal biotic driver of soil redox cycling. Respiration represents an essential series of redox reactions for most organisms, providing energy through electron movement from one chemical species (i.e., the

Special Issue: Environmental Redox Processes and Contaminant and Nutrient Dynamics

Received: January 30, 2023

Revised: May 7, 2023

Accepted: July 21, 2023

Published: August 23, 2023



electron donor) to another (i.e., the electron acceptor). Oxygen is the most energetically favorable electron acceptor. Thus, when O₂ is present (i.e., oxic conditions), O₂ is primarily utilized as the terminal electron acceptor and aerobic respiration proceeds. However, when O₂ declines below a physiological threshold (i.e., anoxic conditions), microorganisms can utilize alternative electron acceptors and perform anaerobic respiration or fermentation. Anaerobic respiration yields considerably less energy than aerobic respiration, resulting in slower substrate consumption and organism growth.¹

Soil O₂, and thus redox status, is heterogeneous across both space and time. Redox heterogeneity primarily has been defined at the landscape scale, vis-à-vis wetland and aquifer delineation, with some attention granted to soils undergoing redox fluctuations over time.² Generally, flooded soils, such as those in lowland areas, are assumed to be anoxic,³ while upland soils are assumed to be entirely oxic and, thus, dominated by oxygen-dependent redox processes. However, redox heterogeneity can also occur at finer scales, within bulk oxic soils and over shorter periods of time. Seemingly aberrant biogeochemical behavior is often attributed to redox heterogeneity in soils,^{4–10} yet there are few studies that directly investigate the influence of redox heterogeneity on biogeochemistry.

Redox heterogeneity is not unique: nearly all soil properties and processes are heterogeneous across space and time. However, unlike redox, other aspects of soil heterogeneity are better defined. Small-scale and short-term spatiotemporal variability in soil biological activity is represented by biological “hot spots” and “hot moments”,¹¹ whereas larger scale and longer term variability can be classified by biome and seasonal effect.^{12–14} Soil structural units (e.g., aggregates) are a defining feature of soil pore heterogeneity,^{15–18} and soil textural classes arose from an attempt to classify heterogeneity in the soil particle size distribution across landscapes.¹⁹ At landscape scales, rainfall patterns, macrotopography, and groundwater intrusion dictate when and where soils receive and accumulate moisture,¹⁹ and at smaller scales, macropores and preferential flow paths guide water and associated resources to some portions of soil while leaving other portions without.²⁰

Defining the features of soil heterogeneity and clarifying the relationships between these heterogeneities across scales has advanced our understanding of soils as a complex system. Biological activity in soils depends on water, and thus preferential flow, for microbial habitat space and substrates for metabolism.^{21–25} Preferential flow is dictated by soil structure,²⁰ and in turn, soil structure and preferential flow can be driven by texture and biological activity.^{17,26,27} Environmental forcings (e.g., precipitation, snow melt, temperature) modulate these relationships over days, months, and years. For instance, freeze–thaw cycles alter soil pore size and connectivity as well as water and nutrient availability, which subsequently control soil microbial activity.^{28–30} Alternatively, temperature-driven changes in root architecture can alter the distribution of soil root channels,³¹ which serve as both effective conduits for water^{22,32,33} and specialized habitats for microorganisms.^{11,34} Clarifying how redox heterogeneity emerges from other physical and biological soil heterogeneities promises to advance our understanding of soil biogeochemistry.

Anoxic microsites, nonmajority soil pore volumes that are without O₂, are a defining feature of soil redox heterogeneity in bulk oxic soils and sediments. In the early 1960s, Currie

(1961) postulated that based on the laws of gaseous diffusion, soil aggregates may be anoxic at their cores.¹⁵ Years later, microelectrode measurements confirmed localized anoxia at the interior of both saturated³⁵ and unsaturated³⁶ aggregates. Evidence of anoxic microsites has been observed in soils across multiple ecosystems, such as tropical and temperate forests, agricultural lands, deserts, and floodplains.^{37–41} Increasingly, anoxic microsites are recognized as an important but ill-defined control on soil greenhouse gas emissions^{42–44} and contaminant transport.^{45,46} Yet, despite their prevalence, long-known presence, and presumed importance, our understanding of anoxic microsites remains incomplete. The soil properties and processes that shape anoxic microsite dynamics have only been partially characterized, and the influence of anoxic microsites on various biogeochemical cycles is still undefined.

In this review, we propose that anoxic microsites are a critical component of soil function and that failure to acknowledge soil redox heterogeneity limits our understanding of soil and sediment biology, mineral transformations, nutrient and contaminant dynamics, and biogeochemical cycling more broadly. In section 2, we define anoxic microsites and their mutable properties, specifically anoxic microsite distribution, redox gradient magnitude, and temporality. In section 3, we describe how anoxic microsites may alter cycles of key redox-active elements. In section 4, we review and evaluate methods for identifying and characterizing anoxic microsites, and in section 5, we review existing models that represent anoxic microsites and their environmental significance. Finally, in section 6, we suggest steps to incorporate anoxic microsites, their measure, and influence into the canons of soil biogeochemistry.

2. DEFINING ANOXIC MICROSITES

2.1. Formation. Anoxic conditions establish when O₂ supply is slower than O₂ demand. Oxygen supply in soils is controlled predominantly by water content and soil structure. Oxygen diffuses 10 000 times more slowly through water-filled pores than through air-filled pores,¹⁵ and as a result, most submerged soils are considered bulk anoxic.³ Even in seemingly well-drained forest soils, both long- and short-term precipitation have been associated with O₂ depletion⁴⁷ and redox heterogeneity.⁴⁸ Soil structure and texture also control soil O₂ supply. Clay particles flocculate and, with organic matter, aggregate. The continued expansion/contraction of aggregates—through hydration and thermal cycles and/or the action of plants roots and soil organisms—results in the formation of soil structural units, which can range from microaggregates, tens of micrometers in diameter, to peds, a few centimeters in diameter.^{18,19} The small, tortuous pores characteristic of these units retain water across a wide range of soil moistures and extend the diffusion path length of O₂, impeding O₂ supply.^{15,36,49–51}

Oxygen demand in soils is controlled principally by microbial demand for O₂ during aerobic respiration, though abiotic reactions may also deplete soil O₂.^{10,52} The rate and magnitude of microbial O₂ demand varies with resource availability and temperature.^{53–56} For example, electron-donor availability, most notably organic carbon (OC), partially dictates soil O₂ content. Higher concentrations of bioavailable OC are frequently linked to decreased O₂ concentrations as a result of enhanced aerobic respiration.^{51,57} Soil moisture, in addition to inhibiting O₂ supply, can stimulate respiration and increase soil O₂ demand by increasing the connectivity of soil




Figure 1. Depictions of anoxic microsities: (a) basic conceptualization of anoxic microsities, highlighting the redox interface, the boundary where oxidants and reductants meet; (b) anoxic microsities as redox gradients, hosting a multitude of microbial metabolisms.




Figure 2. Representations of anoxic microsities: (a) distribution, (b) gradient magnitude, and (c) temporality.

pores, allowing mobile, dissolved OC to be transported to decomposer communities.⁵⁸ Temperature can also alter microbial activity, with decomposition rates and O₂ con-

sumption accelerating in warmer temperatures^{59–61} and slowing in cooler temperatures.⁵⁶

Anoxic microsites are emergent features of heterogeneous O_2 supply and demand within bulk oxic soils. Thus, anoxic microsites form in soil volumes with limited O_2 supply, excessive O_2 demand, or a mild imbalance of the two. The interior of aggregates or peds, common locations for anoxic microsites,^{15,36,49} represents zones of limited O_2 supply. Anoxic microsites have also been associated with actively growing plant roots⁶² and preferential flow paths,²⁴ both of which represent zones with ample bioavailable OC and, therefore, high O_2 demand. Anoxic microsites are also frequently observed within or near plant detritus;^{40,63,64} these microsites are attributed to both high biological O_2 demand, as microorganisms break down and respire OC, and impeded O_2 supply, as plant tissues readily absorb water.⁶³

2.2. Expanding and Revising Terminology. The phrase “anoxic microsite” is imperfect. The prefix “micro-” connotes small spatial scales, yet anoxic microsites represent sizes ranging from micrometers to decimeters;^{36,51,65} the definition of microsite is entirely dependent on one’s frame of reference. The term “anoxic” emphasizes O_2 presence and absence (Figure 1a), yet the biogeochemical influence of anoxic microsites is more nuanced. Anoxic microsites represent gradients in redox potential below the anoxic threshold and may host a multitude of chemical transformations at various distances from each other (Figure 1b). Additionally, the exterior of an anoxic microsite represents a critical redox interface where reduced species and strong oxidants meet (Figure 1a). Finally, the suffix “-site” implies a false sense of constancy. The word “site” (e.g., building site, camp site, archeological site) often refers to exact locations of structures that last for days or years, whereas the lifetime, size, and location of an anoxic microsite is likely shorter, variable, and still largely unknown.

Despite its imperfections, the term anoxic microsite has merit. The term anoxic centers the importance of O_2 in regulating biogeochemical cycles, providing a single metric by which we may define a region of interest. The term microsite provides some sense of relative scale: in a soil containing anoxic microsites, the anoxic portion of the sediment represents the minority of soil pore space. Beyond anoxia and minority soil volume, anoxic microsites can exhibit a variety of characteristics. To examine the diversity that exists within the anoxic microsite classification and highlight the biogeochemical influence of anoxic microsites, we define three domains of variability in how anoxic microsites manifest: anoxic microsite distribution, redox gradient magnitude, and temporality (Figure 2).

Anoxic microsite distribution, the size and location of anoxic microsites, ultimately determines the susceptibility of anoxic microsites to disturbance and the absolute volume over which anaerobic processes may occur. Anoxic microsite distribution theoretically ranges from small and sparse (Figure 2a, lower left) to large and highly concentrated (Figure 2a, upper right). Small and sparse distributions result in less anoxic volume over a given space than large and concentrated anoxic microsites. Additionally, anoxic microsites can vary in size within the same soil profile³⁶ and can be unevenly distributed, clustering in specific horizons or near plant roots and preferential flow paths.^{24,62,66}

Redox gradients within anoxic microsites represent vector quantities, both a direction and a magnitude of change. Often, anoxic microsites are depicted as centered on areas of limited diffusion, such as aggregates, containing a singular redox

minimum toward the center of the aggregate (Figure 1b). In reality, an anoxic microsite may contain multiple redox minima, asymmetrical redox gradients,³⁶ or redox minima centered on areas of preferential flow, such as the exterior of an aggregate.^{24,57} The orientation of redox minima within a single anoxic microsite likely alters microsite biogeochemical function. We limit our discussion to the influence of redox gradient magnitude as the principals of redox gradient magnitude can be applied across multiple microsite orientations and scales.

Redox gradient magnitude governs (i) the extent of reducing conditions in anoxic microsites and (ii) the distance reduced and oxidized species must diffuse to react with other oxidizing or reducing species, respectively (examples provided throughout section 3). As a result, redox gradient magnitude controls the processes that occur within and near anoxic microsites. In our usage, large redox decline over a confined distance constitutes a sharp or steep gradient (Figure 2b, lower panel), whereas a smaller redox decline over the same distance constitutes a milder gradient (Figure 2b, upper panel). Gradient magnitude within anoxic microsites depends on the supply of electron donors (e.g., OC) and acceptors, which is constrained by donor/acceptor concentration and their ability to move through soil pore space (i.e., diffusion).⁶⁷ For example, supplying nitrate (NO_3^- ; i.e., an electron acceptor) to an anoxic microsite that supports Fe reduction effectively decreases redox gradient magnitude. Given sufficient time and limiting OC or nutrients, providing an alternative, more favorable electron acceptor to pair with the oxidation of OC can shift the dominant respiration pathway(s) of a system.⁶⁸ Alternatively, supplying excess OC increases redox gradient magnitude by increasing demand for and depletion of electron acceptors over a confined distance.⁶⁹ Due to their influence on diffusive flux, soil structure and texture may attenuate or amplify the response of gradient magnitude to changes in electron-donor and -acceptor availability.^{24,67}

Anoxic microsite temporality or transience represents the variable frequencies with which anoxic microsites are present and the fluctuations of redox gradient magnitude. The temporality of anoxic microsites determines the net flux of a given element over a constrained period. Anoxic microsites can endure for hours or months^{24,51,70} and possibly from seconds to years. Anoxic microsite temporality is driven by changes in soil O_2 supply and demand that result from changes in moisture, OC availability, soil structure, and potentially other drivers (Figure 2c). Climatic (e.g., precipitation, wet–dry, freeze–thaw), biotic (e.g., plant root exudates, earthworm burrows), and anthropogenic (e.g., farming practices like tillage, OM amendments) perturbations and dynamics drive changes in soil moisture, OC availability, and structure and, in turn, likely alter anoxic microsite temporality.^{17,24,71,72}

3. INFLUENCE OF ANOXIC MICROSITES ON BIOGEOCHEMICAL CYCLES

Classic soil dogma asserts that upland soils and bulk oxic aquifers exclusively host aerobic biogeochemical transformations.⁷³ However, there is increasing evidence that anoxic microsites exist within bulk oxic soils and sediments, dictating the fate of several elements; the precise influence of anoxic microsites on various biogeochemical cycles is still being investigated. Here, we explore the various redox processes that may be hosted within and near anoxic microsites using anoxic microsite distribution, gradient magnitude, and temporality to

discuss how different manifestations of anoxic microsites can have varying effects on key redox transformations. Where possible, we cite anoxic microsite-specific literature. Otherwise, we apply knowledge about anaerobic processes to hypothesize how anoxic microsite presence and variability may alter elemental cycles.

3.1. Soil Organic Carbon. Anoxic microsite distribution can alter the fluxes and distribution of soil OC, but evidence is mixed as to whether anoxic microsites prompt soil OC accrual (i.e., decreased oxidation of OC to CO₂) or loss (i.e., enhanced oxidation of OC to CO₂). Anoxic conditions can slow the oxidation of OC to CO₂ by (i) precluding the oxidative depolymerization of large macromolecules^{68,74,75} and (ii) decreasing the energetic yield of microbial respiration.^{1,76,77} Thus, anoxic microsites and proxies for anoxic microsite volume, such as soil moisture or Fe(II) concentration, are often positively correlated with soil OC content in bulk oxic soils.^{78,79} Similarly, soils that support more anaerobic respiration, as indicated by Fe(II) and methanogen abundance, have been shown to lose the most OC upon oxygenation,⁷² and prolonged O₂ limitations within soil structural units can partially explain discrete zones of relatively old OC.⁶⁶ However, anoxic conditions can also prompt the reductive dissolution of Fe-oxides (see section 3.3), which can liberate mineral-bound OC, increasing microbial access to OC.⁸⁰ Under unsaturated, intermediate moisture contents (~75% saturated), Huang and Hall (2017) attributed elevated soil OC loss as CO₂ and CH₄ to anoxic microsites and the reductive dissolution of Fe-oxides.⁸¹ Yet, the balance between OC accrual and loss within anoxic microsites and subsequent impacts on ecosystem-scale OC cycling remain undefined.

Redox gradient magnitude determines the rate of soil OC loss and/or accrual in anoxic microsites as well as the identity of the OC byproducts. Naughton et al. (2021) decreased redox gradient magnitude in model anoxic microsites by adding NO₃⁻, a relatively favorable electron acceptor.⁶⁸ The decrease in gradient magnitude increased respiration (i.e., OC oxidation) by enabling denitrification, which proceeds more quickly than OC oxidation paired with sulfate (SO₄²⁻) or Fe reduction. Redox gradient magnitude also determines whether anoxic microsites can support methanogenesis. Methanogenesis is one of the least thermodynamically favorable respiration pathways. Thus, methanogens are outcompeted for OC and nutrient resources when other electron acceptors and hence more energetically favorable respiration pathways are present.³ However, in the absence of OC limitations, methanogenesis can proceed alongside other anaerobic respiration pathways.⁸² Evidence of methanogenesis has been observed in bulk oxic tropical, desert, arid agricultural, Mediterranean grassland, and temperate forest soils,^{37,38,72,83–85} indicating that these soils host anoxic microsites with redox declines large enough to support methanogenesis. Similarly, redox gradient magnitude determines the fate of specific OC compounds. Soil OC with a nominal oxidation state of C of less than -1.8 or -0.8 is thermodynamically nonrespirable when paired with the reduction of Fe(III) or SO₄²⁻, respectively.⁴³ As a result, reduced (low nominal oxidation state of C) OC compounds, such as lipids, accumulate within sulfidic and/or iron-reducing anoxic microsites^{39,67} but not within denitrifying microsites.

Environmental forces, such as weather events and climate, drive anoxic microsite temporality, influencing net C fluxes over time. In humid tropical soils, dry periods of more than 2

days were associated with concomitant declines in methane (CH₄) emissions, presumably due to the aeration of anoxic microsites.⁷¹ High-latitude upland soils typically serve as net CH₄ sinks, but during the nongrowing season, soils can become a net source of CH₄ depending on soil moisture and temperature.⁸⁶ Similarly, precipitation events and generally rainy seasons can support anaerobic respiration of OC not observed in drier periods¹⁰ or turn upland temperate forest soils into temporary CH₄ sources.⁴ In laboratory systems, relative enrichment of a low nominal oxidation state of C compounds was established in less than 40 days,^{67,68} and net CH₄ production from a bulk oxic biological soil crust was observed in less than 7 days.⁸⁴ However, the intensity of environmental forcings and minimum duration over which microsites must persist to induce observable changes in C flux or speciation remains unknown.

3.2. Nitrogen. Anoxic microsite distribution partially determines N retention and loss from ecosystems. Anoxic soils favor microbial denitrification, the reduction of NO₃⁻ to dinitrogen gas (N₂),⁸⁷ with intermediates nitric oxide (NO) and nitrous oxide (N₂O), potent greenhouse gases that often escape full metabolization and contribute to global warming.⁸⁸ Denitrification is seemingly ubiquitous across upland soils and is regularly observed in agricultural fields, peatlands, and humid tropical forests.^{36,89,90} Net denitrification and N loss from soils is associated with anoxic microsite abundance, with variance in denitrification potential over larger (i.e., meter) scales attributed to key drivers of anoxic microsite formation, such as high water and organic matter contents.^{91–93} However, denitrification is dependent in part on adequate OC and NO₃⁻ supply.^{94,95} As a result, denitrification scales nonlinearly with anoxic volume.^{36,96} The relative positioning of anoxic microsites alters denitrification rates and the proportion of denitrification byproducts. Schlüter et al. (2019) embedded model aggregates in layered and random orientations and found that more evenly spaced anoxic microsites (i.e., random orientation) resulted in greater denitrification rates as well as increased NO and N₂O product ratios.⁹⁷ Finally, the absolute location of anoxic microsites can dictate N availability. For example, anoxic microsites associated with plant roots may limit plant N availability by hosting redox interfaces that promote rapid denitrification and/or nitrification, resulting in gaseous or aqueous N loss.^{98–100}

The relationship between microsite redox gradient magnitude and N cycling is complex. Gradient magnitude controls N response to anoxic microsites. However, influxes of N, particularly NO₃⁻, can alter redox gradient magnitude, with enhanced NO₃⁻ supply decreasing redox gradient magnitude.⁶⁸ At the exterior of an anoxic microsite, coupled denitrification–nitrification results in constant cycling between nitrite (NO₂⁻) and NO₃⁻,^{89,101–103} with nitrification supplying the NO₃⁻ precursors for denitrification and possibly enhancing N₂O production.¹⁰² Microbial anammox functions similarly, with NO₂⁻ or NO₃⁻ supplied at the oxic–anoxic interface fueling anaerobic oxidation of ammonium (NH₄⁺) to N₂ or the intermediate byproduct, N₂O.^{104–106} In addition to being oxidized at the oxic–anoxic interface, NO₂⁻ can act as an oxidant when reacting with other anaerobic byproducts. During chemodenitrification, NO₂⁻ oxidizes Fe(II) to produce Fe(III) and N₂O.^{107,108} During nitrite dependent anaerobic methane oxidation (n-DAMO), NO₂⁻ oxidizes CH₄ to produce CO₂ and N₂, thereby mitigating CH₄ and N₂O emissions.^{105,109–112} Chemodenitrification has been directly

observed in anoxic microsites,¹¹³ but n-DAMO has only been observed in flooded paddy soils thus far.¹¹⁴ Gradient sharpness presumably controls the likelihood that anoxic microsites will host chemodenitrification or n-DAMO. If the zone of Fe reduction within the gradient is relatively close to the nitrite byproduct, chemodenitrification may be more likely to occur. Conversely, if the zone of methanogenesis is physically separated from a suitable NO_2^- supply, n-DAMO may be less likely to occur.

Anoxic microsite temporality drives pulses in N loss from soils. Once anoxic or hypoxic (low-oxygen) conditions are established, denitrifiers respond quickly due to their often facultative nature, high abundance (up to 20% of culturable taxa),¹¹⁵ and rapid production of denitrifying enzymes.^{116,117} As a result, even ephemeral anoxic microsites caused by heavy rains, soil freeze–thaw, or even drought-induced cell lysis can result in significant N_2O emissions.^{113,118,119} Sustained patterns in O_2 availability, such as fluctuating or static redox conditions, can also alter the identity of N byproducts from soils. For example, Pett-Ridge et al. (2013) found that soils under fluctuating redox conditions supported more nitrification than soils under static oxic or anoxic incubations.¹²⁰ Static anoxic conditions are more likely to facilitate complete denitrification, reducing oxidized N species fully to N_2 ¹²¹ and limiting greenhouse gas production. Similarly, static anoxic conditions are more likely to promote dissimilatory nitrate reduction to ammonium,⁸⁷ limiting aqueous N loss.

3.3. Iron and Manganese. Anoxic microsite distribution dictates the volume and precise location of Fe and Mn redox transformations and thereby affects soil development, nutrient acquisition, and contaminant mobilization. Although the reduction of Mn–oxides is thermodynamically more favorable than the reduction of Fe–oxides,³ Fe is more abundant and thus often accounts for the majority of soil anaerobic respiration.⁶⁷ Sustained or repeated Fe reduction within anoxic microsites can result in profile- and feature-scale Fe loss as Fe(II). For example, along a Hawaiian precipitation gradient, wetter bulk soils have lower Fe–oxide and oxyhydroxide contents than drier bulk soils.¹²² Similarly, anoxic microsites formed within and near root channels drive Fe reduction and depletion over pedogenic time scales.¹²³ Reductive dissolution of Fe–oxides and Mn–oxides mobilizes OC,^{81,124,125} P,^{126,127} nutrient cations,¹²⁸ and contaminants¹²⁹ bound to these Fe–oxides and Mn–oxides. For example, in agricultural soils, anoxic microsites in a compacted plough layer drove reduction of Fe–oxides and Mn–oxides and comobilized P into soil water.¹³⁰ Similarly, rhizosphere-associated anoxic microsites have been shown to enhance ferrihydrite dissolution and mobilize adsorbed As and U, facilitating plant uptake of these contaminants.¹³¹

Redox gradients within anoxic microsites control the collocation of Fe, Mn, and other reactants, thereby controlling which Fe and Mn transformations occur within or near anoxic microsites. Several Fe and Mn redox reactions are facilitated at redox interfaces.^{132,133} Microbial Fe oxidizers are often situated in microaerophilic zones at oxic–anoxic interfaces and conserve energy by oxidizing Fe(II) to Fe(III).^{134,135} Fe(II) can also be oxidized to Fe(III) by abiotic reaction with molecular O_2 or hydrogen peroxide (i.e., the Fenton reaction), the byproducts of which include Fe(III) and highly reactive hydroxyl radicals.^{136,137} Similarly, Mn(II) can be oxidized abiotically to Mn(III/IV) by reaction with reactive oxygen species¹³⁸ or O_2 given sufficiently high pH.^{137,139} Sharp redox

gradients shorten the diffusion path length between Fe(II), Mn(II), and the oxic interface, increasing the likelihood of the Fenton reaction and/or Mn oxidation.^{132,136} As a result, anoxic microsites with sharper redox gradients may produce more reactive oxygen species and Mn(III) at their exterior than microsites with milder gradients. Gradient magnitude also controls Fe and Mn transformations strictly within the anoxic microsite. For example, Pallud et al. (2010) imposed a sharp redox gradient within a model aggregate system by increasing the concentration of electron donors.⁶⁹ Concentrations of Fe(II) and inorganic carbon were high enough to promote siderite (FeCO_3) formation, whereas in a shallower gradient system, no siderite was formed. Similarly, gradient sharpness partially controls the likelihood of chemodenitrification, the abiotic oxidation of Fe(II) by NO_2^- (see section 3.2).¹⁴⁰

Finally, anoxic microsite temporality controls the net fluxes and byproducts of Fe and Mn throughout soils. Frequent transitions from anoxic to oxic conditions can increase Fe–oxide crystallinity, decreasing the surface area, sorbent capacity, and reducibility of these oxides.^{122,141–143} If the same volume of soil repeatedly endures redox fluctuations, distinct patches of soil, such as the rhizosphere, may contain more crystalline Fe–oxides over pedogenic time.¹⁴⁴ Changes in anoxic microsite distribution and gradient magnitude are often driven by changes in seasons, influencing the mobility of Fe and Mn from soils. For instance, warm, wet conditions in summer months may be more likely to support anoxic microsite formation and persistence, leading to greater soluble Fe, Mn, and P in summer months than in cool, dry winter months.¹³⁰

3.4. Sulfur. Anoxic microsite distribution partially dictates the mobility of S within soils and sediments. Within anoxic microsites, microorganisms use SO_4^{2-} as an electron acceptor, producing aqueous sulfide ($\text{HS}^-/\text{S}^{2-}$) and/or hydrogen sulfide (H_2S).¹⁴⁵ These sulfide species can react with transition metals to form low-solubility elemental S and/or (di)sulfide minerals, essentially immobilizing S,^{145–148} contaminant metals, and OC from soil and groundwater.^{149–161} Thus, anoxic microsites can export or immobilize S, with S mobility depending on the geochemistry of the anoxic microsite. For example, immobilization of sulfide via reductive dissolution of Fe(III)–(hydroxy)oxides and Fe–sulfide formation requires high ratios of S(II) to Fe(II).^{162–164} However, in systems with inadequate S^{2-} or excess Fe(II), anoxic microsites mobilize S and associated nutrients and contaminants as colloids.¹⁶⁵ The export of these sulfidic colloidal species can extend the boundaries of anoxic microsites by stimulating microbial activity and accumulation of reduced S products downgradient of the “original” anoxic microsite.^{166,167}

Redox gradient magnitude also determines the fate of S in sediments with anoxic microsites. Sulfate reduction yields less energy to microorganisms than other electron acceptors, and therefore, SO_4^{2-} reduction is often limited or inhibited by other electron acceptors of higher redox potential.^{3,168} Thus, anoxic microsites with mild redox gradients are less likely to support SO_4^{2-} reduction and S^{2-} generation than similarly sized anoxic microsites with sharper redox gradients. Gradient magnitude also dictates the distance S^{2-} species must travel to encounter oxidizing conditions. For example, hydrogen sulfide can react with Fe(III)–oxides to produce Fe(II) and elemental S.¹⁶⁹ However, the likelihood of this interaction is dependent on gradient sharpness, with sharper gradients (i.e., shorter distances between sulfidic species and Fe(III) species)

Table 1. Techniques for Identifying and Characterizing Anoxic Microsites

approach	scale(s)	advantages	disadvantages
Bulk and 1D			
oxygen probes	>5 μm	direct measurement of anoxic/oxic conditions in situ deployment	false negatives; limited information on how much Eh declines below anoxic threshold
Eh probes	from 5 μm to multimillimeter	in situ deployment reveals possible terminal electron acceptors	difficult to link to specific redox transformations
mediated amperometry	~100 mg of soil	provides reasonable averages for electron-accepting/donating capacity over small volumes	cannot implement in situ
redox-active species	see below	reveals specific biogeochemical transformations	false negatives
gaseous	from 1 g of soil to field	real-time indicator for anoxic microsites	consumption/production dynamics obscure data
solid/dissolved	~0.25–0.5 g of soil	relatively easy to measure	may not reflect conditions at time of sampling
microbial	~0.25–5g of soil	indicator in real time and/or for the recent past	limited spatial resolution
16S		fairly common technique; enables use of “big data”	phylogeny does not reveal function
SIP/BONCAT		links phylogenetics to function	technically cumbersome
qPCR/ddPCR		reflect microbial function and/or metabolic capacity	difficulties with environmental samples
2D			
planar optodes:	<1 mm	high resolution	unavoidable “wall” effect
O ₂		optodes are reusable	difficult to implement in field
H ₂ S or Fe		optode reactivity is irreversible	optodes are single use (i.e., irreversible)
CARD-FISH	~0.5 cm ³ of soil	links microorganisms to microstructural features	thin sectioning is technically difficult
X-ray imaging	1 μm	high resolution; enables μm to multicentimeter mapping of anoxic features	resin embedding is technically difficult
Simulated			
no-flow/capillary fringe	multicentimeter	represents unsaturated systems	difficult to obtain adequate liquid sample volume
flow-through reactors	multicentimeter	controlled advection dynamics	limited applicability to unsaturated systems
microfluidic systems	micrometer	enables fine-scale measurements	abstracted from environmental conditions

potentially more likely to support the formation of elemental S, metastable FeS, and associated byproducts.¹⁴⁵ Finally, the concept of gradient magnitude provokes interesting questions regarding the transport of sulfidic byproducts into and through oxic environments. For example, metal sulfide clusters generated in sulfidic conditions¹⁷⁰ are remarkably resistant to oxidation and were found to contribute to contaminant transport in oxic rivers.¹⁷¹ By logical extension, anoxic microsites in oxic aquifers could release metal sulfide clusters to oxic environments, but how far these aqueous metal sulfide clusters can migrate away from their generation point, i.e., the interior of an anoxic microsite, remains unknown.

In sulfate-rich soils, long periods of anoxia can promote the formation of Fe–sulfides, which, when oxidized, generate acidity. Although we typically associate acid sulfate soils with flooding and draining wetlands, presumably these acidification dynamics can occur at a smaller spatial scale. For example, if a persistent, sulfidic anoxic microsite were to become oxic, local acidification would transpire, which in turn alters soil weathering and nutrient, contaminant, and OC mobility. Regular redox fluctuations in sulfidic soils can drive pulses of contaminant release with subsequent cycles releasing more contaminant metals depending on the substrate.^{172–174}

4. TECHNIQUES FOR CHARACTERIZING ANOXIC MICROSITES

The net influence of anoxic microsites on soil biogeochemistry is still largely unknown due, in part, to methodological constraints on identifying and characterizing anoxic microsites. Here, we evaluate existing approaches for detecting anoxic microsites and assessing their distribution, redox gradient magnitude, and temporality. We categorize these techniques into (1) bulk approaches that indicate anoxic microsite

presence and contribution to biogeochemical processes, (2) spatially explicit 2D imaging approaches, and (3) simulated systems (Table 1).

4.1. Bulk and 1D Measurements. Many bulk soil measurements can determine whether anoxic microsites are present and reveal some aspects of their distribution and gradient magnitude. In general, bulk and 1D measurements tend to be easier and cheaper to implement than 2D measures. Bulk and 1D measurements commonly used to identify anoxic microsites include (i) soil O₂ and redox state, (ii) concentrations of redox-active chemical species, and (iii) characteristics of soil microbial communities.

4.1.1. Oxygen and Soil Redox State. Oxygen concentrations offer a direct and instantaneous measure of anoxic microsites. Early evidence for the existence of anoxic microsites consisted of O₂ distribution measurements within wet aggregates of a silt loam soil using O₂ microelectrodes,³⁶ and this line of work was later carried out in combination with measurements of soil redox potential (Eh).¹⁷⁵ In more recent work, Hall et al. (2016) used O₂ sensors in the field to measure variations in bulk O₂ concentration semicontinuously for multiple months.¹⁷⁶ However, bulk O₂ content does not always reflect the spatial heterogeneity of biogeochemical conditions. For example, Keiluweit et al. (2018) confirmed the presence of anoxic microsites with 2D planar optodes, but bulk O₂ measurements showed consistently oxic conditions.⁵¹ Similarly, in the field, Teh et al. (2005) consistently observed methanogenesis in soils which, when measured by bulk O₂ probes, were consistently oxic.⁸⁵ As a result, bulk O₂ measurements in a field or laboratory setting can result in “false negatives” in identifying anoxic microsites.

Soil redox potential (Eh) has also been used to characterize anoxic microsites in the field and laboratory. Flessa and Beese

(1995) used redox microelectrodes to characterize changes in redox potential across depth,¹¹⁸ and more recently, redox electrodes were used to identify functionally different microsites in soils and overall redox heterogeneity.⁴⁸ Installing redox electrodes is relatively simple and inexpensive. However, Eh measurements can be difficult to interpret.¹⁷⁷ Like O₂ probes, redox probes make stationary point measurements and, therefore, can miss anoxic microsites in space. Additionally, depending on the size of the redox probe, Eh values reflect an integrated average over some volume, which can obfuscate microsites smaller than the probe resolution. Finally, Eh values need to be interpreted alongside pH in order to deduce the redox conditions.¹⁷⁸

To avoid the issues associated with point measurements, mediated amperometric approaches allow characterization of the overall redox state of a soil on bulk samples, enabling the averaging of redox state over a confined soil volume. Mediated amperometry enables quantification of electron-accepting and -donating capacities of soil. These capacities represent the cumulative response of redox-active compounds in the samples, including solid-phase electron acceptors and their reduced species, quinone/hydroquinone groups in organic matter, and S²⁻. Aeppli et al. (2022) used this approach to characterize spatiotemporal variations in soil redox state in a simulated aquifer system.¹⁷⁹ However, unlike Eh and O₂ measures, mediated amperometry has yet to be successfully deployed in a field setting.

4.1.2. Redox-Active Chemical Species. Gaseous byproducts of anaerobic respiration can also provide evidence for anoxic microsites. For instance, CH₄ has been used as an indicator for methanogenesis in well-oxygenated soils in the field^{85,180,181} and laboratory.^{37,51,85,182,183} Similarly, N₂O emissions may be used as an indicator for anoxic microsites.¹¹⁸ Although N₂O and CH₄ fluxes are useful real-time indicators for anoxic microsites, both can be difficult to interpret on their own. Methane produced in anoxic microsites can be consumed by methanotrophs elsewhere in the soil profile, obfuscating any detectable signal at the soil surface. Nitrous oxide can be produced via multiple pathways, both abiotic (e.g., chemodenitrification)^{107,140} and biotic (e.g., nitrification and denitrification).¹⁸⁴ Thus, N₂O fluxes are not necessarily indicative of denitrification. To address these ambiguities, authors have used isotopic approaches to attribute N₂O to specific (de)nitrification pathways^{107,140,185,186} and to detect CH₄ production even when net CH₄ flux from soils is negative or zero.^{37,181} Alternatively, others have used the co-occurrence of net CH₄ oxidation and N₂O reduction as an indicator for the presence of anoxic microsites.¹⁸⁷

Dissolved and solid-phase measurements can also reveal anoxic microsites. Iron and Mn reduction can be monitored by measuring Fe(II) and Mn(II) in the dissolved, adsorbed, or solid phase.^{51,67,79} Similarly, S²⁻ formed during microbial SO₄²⁻ reduction can be monitored in the dissolved phase or solid phase through precipitation of iron–sulfide minerals.⁶⁵ Whereas measuring solid-phase Fe(II), Mn(II), and S²⁻ often requires more advanced and costly X-ray techniques (see section 4.2.3), monitoring dissolved or extractable species is relatively simple and low cost with several methods performed colorimetrically.^{188–190} Thus, when appropriate, many opt for dissolution protocols (e.g., Fe with weak hydrochloric acid) coupled with subsequent dissolved-phase measurements (e.g., Ferrozine for Fe(II), iodine titration for S²⁻, and/or mass spectrometry when appropriate). Of course, there are

challenges to measuring and interpreting Fe(II), Mn(II), and S²⁻. For example, Fe(II) readily reacts with O₂ to form Fe(III),¹⁹¹ and aqueous sulfides react to form hydrogen sulfide, which escapes soil but can be detected as H₂S gas.¹⁹² Mn(II) is relatively stable under oxic conditions but can be oxidized biotically to Mn(IV). As a result, Fe(II), Mn(II), and S²⁻ can represent both present and near-past anoxic conditions.

Although the products of specific redox reactions can indicate which biogeochemical transformations are occurring in soils and sediments, these chemical species should not be used as sole indicators of anoxic microsites. For example, a soil may host anoxic microsites that support Fe and Mn reduction but fail to support denitrification due to insufficient NO₃⁻ and/or fail to support sulfate reduction or methanogenesis due to insufficient redox decline. In this case, relying on N₂O, CH₄, or S²⁻ as the only indicators for anoxic microsites could conceal anoxic microsite presence.

4.1.3. Microbial Analyses. The presence of facultative or obligate anaerobes can provide strong evidence for present or historical anoxic (micro)environments. Common microbial analyses require isolation of DNA and/or RNA from approximately 0.25 to 5 g of soil. Minimum DNA and RNA extraction mass limits the ability to spatially resolve micrometer- or even millimeter-scale anoxic microsites but does allow for relative comparisons of anoxic microsite distribution. Analyses of DNA and RNA require different interpretations. DNA represents the metabolic potential of the microbial community and is relatively long lived in soils,^{193,194} whereas RNA is ephemeral and represents present microbial function.¹⁹⁵ As a result, the presence of anaerobe DNA indicates metabolic potential for soil anaerobiosis in the past or present, whereas anaerobe RNA indicates active anaerobes at the time of sampling.¹⁹⁶

Amplicon sequencing, typically of 16S rRNA subunits, yields the genetic makeup of the microbial community and their relatedness to known isolates. Through these associations and given the conserved nature of dissimilatory anaerobic respiration pathways within specific microbial taxonomic groups, “community analysis” or “molecular fingerprinting” of soil microorganisms can be used to determine whether a sample contains anaerobes and, therefore, anoxic microsites.^{179,197,198} However, sequencing alone does not specifically target active cells or microbial functions.¹⁹⁹ Sequencing must be used in conjunction with bioorthogonal noncanonical amino acid tagging (BONCAT) coupled with fluorescence-activated cell sorting (FACS)²⁰⁰ or stable isotope probing (SIP) with labeled ¹³C and ¹⁵N substrates or even H₂¹⁸O to link microbial community composition to functional traits (e.g., growth rate, death rate, substrate use efficiency).^{201–203}

Quantification of target genes, typically those associated with anaerobic metabolisms, may be better suited than 16S rRNA sequencing to confirm and compare the presence of anoxic microsites. Quantitative polymerase chain reaction (qPCR) compares the amplification of a gene of interest using specific primers—for instance, primers targeting enzyme components involved in denitrification or methanogenesis—against amplification of a standard to yield gene copy number of that target on a soil mass basis.^{46,113,179} The emerging digital (dPCR) and digital droplet PCR (ddPCR) instruments enhance the sensitivity of the method, improving the detection sensitivity of small microbial populations like methanogens in upland soils.⁷² Notably, ddPCR does not require specimen standards, which can be laborious or even impossible to obtain.²⁰⁴

Depending on the quality of the nucleic acid extracts, resolution in environmental samples can be poor for both qPCR and (d)dPCR.²⁰⁵

Finally, a more exploratory approach to characterize microbial community functional potential and activity is through shotgun sequencing of whole-soil DNA or RNA samples to yield metagenomic and metatranscriptomic data.²⁰⁶ Metagenomic techniques offer a glimpse into the whole soil functional potential, such as respiratory pathways, growth history, transporters, and other traits relevant to microbial lifestyle, and the addition of metatranscriptomics suggests microbial investment in these functions.^{38,180,207} As a diagnostic method of anaerobiosis, however, these techniques suffer from high computational and analytical costs. Microarray chips that use nucleic acid probes attached to fluorophores offer a less comprehensive but functionally similar way to identify specific functional genes or taxa and quantify them via fluorescence upon hybridization of isolated and digested sample DNA or RNA-derived cDNA with the complementary sequence for the gene of interest (e.g., “GeoChip” as in refs 208 and 209).

4.2. Two-Dimensional Measurements. In general, spatially resolved measurements of relatively undisturbed soil allow us to define anoxic microsite distribution and gradient magnitude more clearly. Most of the bulk techniques described above require some destruction of soil structure, undermining the applicability of findings to natural systems. Two- and three-dimensional measures allow researchers to maintain soil structure while revealing patterns about the location and size of microsites and the degree of redox decline within anoxic microsites. To our knowledge, no 3D soil imaging techniques directly reveal the absence of O₂ nor the stable byproducts of anaerobic processes. So, although 3D techniques like computed tomography (CT) and 3D OrbiSIMS can provide useful information on soil properties that lead to or emerge from anoxic microsite formation,^{210–212} we do not discuss 3D techniques here.

4.2.1. Planar Optode Measurements. Planar optode systems have enabled the measurement of multiple chemical species over space and time. Oxygen-sensing planar optodes have successfully revealed anoxic microsite presence and development primarily in laboratory settings.^{51,63,213,214} Hydrogen sulfide and Fe(II) planar optodes are less commonly used, in part, because these sensors tend to be irreversible (i.e., single use), limiting their reusability and ability to provide time-resolved measures.²¹⁵ Practical constraints limit the use of planar optodes in a field setting,²¹⁶ and inserting a “wall” into 3D soil structure alters diffusion dynamics, potentially misrepresenting the in situ behavior of anoxic microsites.²¹⁵

4.2.2. Spatially Resolved Microbial Techniques. Opportunities to identify and characterize anoxic microsites through spatially resolved microbial techniques are limited. Catalyzed reporter deposition fluorescence in situ hybridization (CARD-FISH) is a reliable and flexible technique that allows for the identification of functional genes or microbial biomarkers in 2D. During CARD-FISH, a reporter-labeled oligonucleotide probe binds to a complementary DNA segment in a 2D mounted sample. The binding of the probe activates the fluorescent label, and a confocal laser microscope is used to view and record the signal of the fluorescent probe. Tyramide signal amplification in catalyzed reporter deposition (CARD) significantly improves signal-to-noise fluorescence intensity.²¹⁷ Yet, CARD-FISH still has significant limitations for identifying

and characterizing anoxic microsites. First, 2D CARD-FISH is limited to analyses of small, thin slices of sample, which requires some disruption of soil structure and makes the technique difficult to implement on larger samples. Second, it can be difficult or even impossible to discern fluorescent tags from naturally fluorescent soil particles, even with the additional fluorescence provided by the tyramide signal.²¹⁷

4.2.3. X-ray Fluorescence Imaging Combined with Spectroscopy. Synchrotron X-ray methods may be used to detect solid-phase evidence of anoxic microsites, past or present. Synchrotron-based X-ray fluorescence yield 2D imaging and spectroscopy methods provide high chemical sensitivity (fluorescent X-rays can be detected at very low signal strength) and the ability to map micrometer-scale redox contrasts for a large subset of elements, including S, Fe, Mn, As, U, Cr, and Mo.^{17,18,34–36} X-ray 2D imaging and spectroscopy can reveal clusters of reduced, solid-phase elements which are interpreted to signify past or present anoxic microsites. X-ray 2D imaging and spectroscopy is particularly well suited for the study of anoxic microsites in its ability to bridge data across scales. Anoxic microsites can be mere micrometers in diameter,⁵¹ requiring equally small spatial resolution for detection; yet, it is also necessary to quantify the spatial distribution of these microsites in a larger context, such as along multicentimeter soil cores. X-ray 2D imaging can provide spatial resolution down to $\sim 1 \mu\text{m}$ ²¹⁸ and measurements on samples tens of centimeters long, such as lengthwise soil core slices (up to 100 cm long)²¹⁹ or 30 cm \times 20 cm rhizoboxes.²²⁰

Because X-ray 2D imaging and spectroscopy tracks the byproducts of anoxic microsites and not the lack of O₂ itself, X-ray 2D imaging and spectroscopy is an inherently indirect technique. However, depending on which element(s) are mapped, X-ray 2D imaging and spectroscopy methods can provide information on current, recent, and/or past anoxic conditions in the sample’s history.⁶⁵ For example, detection of Fe(II) clusters would indicate anoxic microsites present at or close to the time of sampling. Alternatively, detection of lepidocrocite, the oxidized, stable product of Fe(II)–minerals and elemental S, would indicate that Fe(II)–sulfides and therefore anoxic microsites were present at some point in time but not at the time of sample collection.

Like other 2D methods, X-ray imaging and spectroscopy is well suited to reveal anoxic microsite distribution but comes with logistical shortcomings. As with planar optodes, proper X-ray imaging requires a relatively flat surface. Thus, intact soil cores must be cut for imaging. Slicing intact soil cores is technically difficult, may result in smearing or loss of particles, and can introduce O₂. Cores can be cast in epoxy to stabilize particles and diminish the risk of oxidation upon cutting, but the penetration of epoxy is limited in fresh (moist/wet) soil, making it difficult to ensure even epoxy casts. Further, synchrotron analyses are time intensive, and accessibility is limited.

4.3. Simulated Experimental Systems. One way to get around the considerable technical difficulties in detecting and characterizing anoxic microsites and their influence is to construct laboratory analogues of varying complexity, such as no-flow/capillary-fringe incubation reactors,^{51,221–223} diffusion-limited flow reactors,^{67,68,224–227} flow-through columns/reactors with dual-flow regimes,^{166,168,179,226,228} and microfluidic systems.^{214,229–231} In these systems, distinct redox zones of various dimensions and composition are created by

permeability contrasts (e.g., by inserting a diffusion-limited aggregate/lens within an advection dominated domain^{166,168,179,226,228}). Within these distinct zones, anoxic microsites are established in response to variation in flow and exchange rates in reactors^{67,68,221,225–227} or microfluidic systems.²²⁹ Biogeochemical factors in these experimental systems are measured over time and space to obtain detailed information about types and rates of reactions inside anoxic microsites and exchange flows and rates across redox boundaries and gradients. Experimental systems are also well suited to investigate system responses to manipulation, such as changes in flow rates, influent water chemistry, or water saturation. Despite limitations arising from creating a simplified system (compared to nature), the knowledge gained from well-designed simulated systems is invaluable as presently it is extremely difficult if not impossible to obtain dependable measures of anoxic microsites in the field. Simulated experiments, therefore, serve as a critical link between field measurements and numerical models by elucidating key mechanisms and reaction rates in systems with anoxic microsites.^{167,229}

5. MODELING ANOXIC MICROSITES

Modeling anoxic microsites remains a challenge. Classical methods for modeling redox processes constrain biogeochemical function through a series of equations drawn from soil physics, chemistry, and biology. These equations are then solved for discrete points in time using singular values for soil moisture content, O₂ concentration, redox potential, and microbial kinetics, among other parameters.^{232–234} These classical methods allow for representations of temporal variability in soil biogeochemistry but assume that soil properties are spatially constant (Figure 3).²³² As a result, classical methods cannot adequately model the spatiotemporal variability of anoxic microsites. Nevertheless, there have been several promising advances in the modeling of anoxic microsites.

Spatially discretizing soil into distinct biogeochemical environments represents a relatively simple approach to modeling anoxic microsites. For example, Currie (1961) discretizes soils into aggregate and interaggregate pores with distinct O₂ diffusion coefficients for each pore category.¹⁵ Parkin (1987) tediously separated soil volumes, measuring the denitrifying enzyme activity and denitrification rates of discretized units to construct a bimodal classification of high denitrification or minimal denitrification soil volumes.⁶⁴ Others have called for the discretization and modeling of anoxic microsites based on aggregates and aggregate size⁴⁴ and/or distinct soil environments, such as rhizosphere or subsoils.²³³ One shortcoming of spatial discretization is the oversimplification of redox conditions; it is unlikely that soil moisture, O₂ concentrations, and/or denitrification capacities are truly bimodal or categorical throughout an entire soil volume (Figure 3).

To overcome some of the shortcomings that result from treating soils like distinct biogeochemical units, models that integrate probability distribution (or density) functions allow for continuous representations of soil spatial heterogeneity. Instead of representing key soil parameters as single values, probability distribution (or density) functions represent soil parameters as a continuous distribution of values, which are informed and constrained by soil measurements (Figure 3). For example, in an extension of the classical dual-Arrhenius




Figure 3. Depiction of different modeling approaches for representing anoxic microsites. Each rectangle represents a confined soil volume with shades of blue representing different oxygen (O₂) concentrations to represent redox heterogeneity. In theory, terminal electron acceptors other than O₂ can also be represented using these same approaches. PDFs = probability distribution (or density) functions.

Michaelis–Menten model, Sihi et al. (2020) represented soil OC content and moisture as skewed distributions with low probability of high-moisture and high-OC contents.¹⁸⁷ Instead of using single values in the network of equations represented in the dual-Arrhenius Michaelis–Menten model, distributions of OC and moisture were used to produce a distribution of outcomes, such as CO₂ efflux, which were used to predict subsequent distributions of other soil variables, such as O₂ concentration, N₂O, and CH₄ fluxes.¹⁸⁷

Finally, other models more explicitly represent soils as porous media through which water and O₂ move while sustaining biogeochemical transformations. Ebrahimi and Or (2015) developed a 3D angular pore network model to characterize anoxic microsite distribution, lifetime, and response to hydration conditions within a single aggregate.²³⁵ The 3D angular pore network model was later expanded to include interactions of aggregates of different sizes and orientations⁵⁰ and upscaled to demonstrate how aggregate size distributions alter anoxic microsite distribution and, thus, biogeochemical gas fluxes from soil profiles.²³⁶ Pore-scale modeling of anoxic microsites is often paired with X-ray microcomputed tomography (μ CT), which can constrain values for diffusivity and water distribution.^{96,237} Larger scale models, based on laboratory experiments, have accurately represented anoxic microsites and their biogeochemical

influence in 2D using the principles of Darcian flow and reactive transport modeling.¹⁶⁷

6. ACKNOWLEDGING AND APPRECIATING ANOXIC MICROSITES

While anoxic microsites often comprise only a small fraction of the total soil volume, they can have disproportionately large effects on the fate and transport of critical elements and compounds. However, decades after their first direct measurement,³⁶ anoxic microsites are not yet integrated into our mainstream conceptualizations of soil biogeochemistry. Water content is typically assumed to be the sole control on soil O₂ content,²³⁸ with unsaturated soils tacitly accepted as entirely aerobic.⁷³ Even the Soil Science Society of America's *Methods of Soil Analysis* portrays anoxic microsites as anomalous, stating that "apparent bad readings" on redox probes "may be the microzone around the electrode [driving]... readings not representative of the bulk soil".²³⁹ By continuing to overlook anoxic microsites, we pre-emptively accept the null hypothesis, H₀: Anoxic microsites have no effect on soil biogeochemistry.

In addition to acknowledging that anoxic microsites exist, we must appreciate the diversity that exists within the anoxic microsite classification, such as their varying distributions, redox gradients, and temporality. Connecting anoxic microsite features to soil properties will help reveal (i) the suite of factors that regulate anoxic microsite dynamics (i.e., the formation and persistence) and (ii) the influence of anoxic microsites on soil biogeochemistry. Filling both knowledge gaps is prerequisite to improve Earth System Models and project soils' responses to change. We envision a basic framework for advancing the study of anoxic microsites.

First, we recommend investing in our ability to measure the distribution and impact of anoxic microsites in soils. Presently, researchers must combine a bevy of techniques (Table 1) to confirm the presence and define the characteristics of anoxic microsites in soils and sediments. Furthermore, there is no reliable way to detect anoxic microsites in intact soils, and there are no unified metrics for anoxic microsite distribution, gradient magnitude, nor temporality. We suggest developing measures that (i) directly or indirectly quantify anoxic volume, (ii) characterize dominant electron-acceptor species across 2D or 3D space, and (iii) provide a sense of spatially resolved redox history, revealing anoxic microsites in real time as well as in the recent past. Ideally, these measures could be employed on intact soils. Developing and using these measures will help elucidate anoxic microsite distribution, forecast redox gradient-dependent reactions, and project fluxes of redox-sensitive elements over time.

Second, to hasten translation of information from measurements to models, robust, easily measurable, and scalable predictors of anoxic microsites should be defined. Repeating newly developed and/or refined measures of anoxic microsite characteristics across a variety of soils and ecosystems promises to reveal edaphic and/or climatic parameters that effectively predict anoxic microsite distribution, gradient magnitude, and temporality. These predictors can help project anoxic microsite dynamics across ecosystems, allowing more research to start linking anoxic microsite presence and traits to larger scale ecosystem function through field, laboratory, or modeling studies.

In brief, it is time to consider the anoxic microsite. Anoxic microsites exist and alter critical biogeochemical cycles. Ignoring anoxic microsites precludes needed technological

advances and studies of redox heterogeneity and biogeochemical cycling. Through identifying anoxic microsites, characterizing their variability, and quantifying their impact on various elemental cycles, we can advance our understanding of soils and improve our ability to project redox-sensitive elemental response to change.

■ AUTHOR INFORMATION

Corresponding Author

Emily M. Lacroix – Institut des Dynamiques de la Surface Terrestre (IDYST), Université de Lausanne, 1015 Lausanne, Switzerland; Department of Earth System Science, Stanford University, Stanford, California 94305, United States; orcid.org/0000-0002-6919-788X; Email: emily.lacroix@unil.ch

Authors

Meret Aeppli – Institut d'ingénierie de l'environnement (IIE), École Polytechnique Fédérale de Lausanne, 1015 Lausanne, Switzerland; orcid.org/0000-0003-3335-3673

Kristin Boye – Environmental Geochemistry Group, SLAC National Accelerator Laboratory, Menlo Park, California 94025, United States; orcid.org/0000-0003-2087-607X

Eoin Brodie – Lawrence Berkeley Laboratory, Earth and Environmental Sciences Area, Berkeley, California 94720, United States; orcid.org/0000-0002-8453-8435

Scott Fendorf – Department of Earth System Science, Stanford University, Stanford, California 94305, United States; orcid.org/0000-0002-9177-1809

Marco Keiluweit – Institut des Dynamiques de la Surface Terrestre (IDYST), Université de Lausanne, 1015 Lausanne, Switzerland; orcid.org/0000-0002-7061-8346

Hannah R. Naughton – Lawrence Berkeley Laboratory, Earth and Environmental Sciences Area, Berkeley, California 94720, United States; orcid.org/0000-0003-0807-0647

Vincent Noël – Environmental Geochemistry Group, SLAC National Accelerator Laboratory, Menlo Park, California 94025, United States

Debjani Sihi – Department of Environmental Sciences, Emory University, Atlanta, Georgia 30322, United States; orcid.org/0000-0002-5513-8862

Complete contact information is available at:

<https://pubs.acs.org/10.1021/acsearthspacechem.3c00032>

Author Contributions

E.M.L., M.A., K.B., M.K., H.R.N., V.N., and D.S. wrote the manuscript. E.M.L. developed figures and tables with input from other authors. All authors contributed to editing and selecting content.

Notes

The authors declare no competing financial interest.

■ ACKNOWLEDGMENTS

Funding was provided by the Stanford Precourt Energy Institute Seed Grant (to E.M.L.) and the NSF EAR (Earth Sciences) Postdoctoral Fellowship, No. 1952802 (to H.R.N.). The U.S. Department of Energy (DOE) also provided support. The office of Biological and Environmental Research (BER), Environmental System Science program supported Contract No. DE-SC0022314 (to D.S.) and Contract No. DE-AC02-76SF00515 (to the SLAC Floodplain Hydro-Biogeochemistry Science Focus Area, SFA). Both SLAC and Lawrence Berkeley

Laboratory are supported by the U.S. Department of Energy, Office of Science, Office of Basic Energy Sciences.

REFERENCES

- (1) Jin, Q.; Bethke, C. M. A New Rate Law Describing Microbial Respiration. *Appl. Environ. Microbiol.* **2003**, *69* (4), 2340–2348.
- (2) Zhang, Z.; Furman, A. Soil Redox Dynamics under Dynamic Hydrologic Regimes - A Review. *Sci. Total Environ.* **2021**, *763*, 143026.
- (3) Ponnampertuma, F. N. The Chemistry of Submerged Soils. *Adv. Agron.* **1972**, *24*, 29–96.
- (4) Megonigal, J. P.; Guenther, A. B. Methane Emissions from Upland Forest Soils and Vegetation. *Tree Physiol.* **2008**, *28* (4), 491–498.
- (5) Young, I. M.; Crawford, J. W. Interactions and Self-Organization in the Soil-Microbe Complex. *Science* **2004**, *304* (5677), 1634–1637.
- (6) Ribas, A.; Mattana, S.; Llorba, R.; Debouk, H.; Sebastià, M. T.; Domene, X. Biochar Application and Summer Temperatures Reduce N₂O and Enhance CH₄ Emissions in a Mediterranean Agroecosystem: Role of Biologically-Induced Anoxic Microsites. *Sci. Total Environ.* **2019**, *685*, 1075–1086.
- (7) Davidson, E. A. Pulses of Nitric Oxide and Nitrous Oxide Flux Following Wetting of Dry Soil: An Assessment of Probable Sources and Importance Relative to Annual Fluxes. *Ecol. Bull.* **1992**, *42*, 149–155.
- (8) Vidal, A.; Hirte, J.; Bender, S. F.; Mayer, J.; Gattinger, A.; Hoschen, C.; Schädler, S.; Iqbal, T. M.; Mueller, C. W. Linking 3D Soil Structure and Plant-Microbe-Soil Carbon Transfer in the Rhizosphere. *Front. Environ. Sci.* **2018**, *6* (FEB), 1–14.
- (9) Heckman, K. A.; Nave, L. E.; Bowman, M.; Gallo, A.; Hatten, J. A.; Matosziuk, L. M.; Possinger, A. R.; SanClements, M.; Strahm, B. D.; Weiglein, T. L.; Rasmussen, C.; Swanston, C. W. Divergent Controls on Carbon Concentration and Persistence between Forests and Grasslands of the Conterminous US. *Biogeochemistry* **2021**, *156*, 41–56.
- (10) Hodges, C.; Kim, H.; Brantley, S. L.; Kaye, J. Soil CO and O Concentrations Illuminate the Relative Importance of Weathering and Respiration to Seasonal Soil Gas Fluctuations. *Soil Sci. Soc. Am. J.* **2019**, *83* (4), 1167.
- (11) Kuzuyakov, Y.; Blagodat'skaya, E. Microbial Hotspots and Hot Moments in Soil: Concept & Review. *Soil Biol. Biochem.* **2015**, *83*, 184–199.
- (12) LaCroix, R. E.; Tfaily, M. M.; McCreight, M.; Jones, M. E.; Spokas, L.; Keiluweit, M. Shifting Mineral and Redox Controls on Carbon Cycling in Seasonally Flooded Mineral Soils. *Biogeosciences* **2019**, *16*, 2573–2589.
- (13) Ren, F.; Yang, X.; Zhou, H.; Zhu, W.; Zhang, Z.; Chen, L.; Cao, G.; He, J. S. Contrasting Effects of Nitrogen and Phosphorus Addition on Soil Respiration in an Alpine Grassland on the Qinghai-Tibetan Plateau. *Sci. Rep.* **2016**, *6*, 34786.
- (14) Zhao, Z.; Peng, C.; Yang, Q.; Meng, F.-R.; Song, X.; Chen, S.; Epule, T. E.; Li, P.; Zhu, Q. Model Prediction of Biome-Specific Global Soil Respiration from 1960 to 2012. *Earths Future* **2017**, *5* (7), 715–729.
- (15) Currie, J. A. Gaseous Diffusion in the Aeration of Aggregated Soils. *Soil Sci.* **1961**, *92* (1), 40–45.
- (16) Tisdall, J. M.; Oades, J. M. Organic Matter and Water-stable Aggregates in Soils. *J. Soil Sci.* **1982**, *33* (2), 141–163.
- (17) Six, J.; Bossuyt, H.; Degryze, S.; Denef, K. A History of Research on the Link between (Micro)Aggregates, Soil Biota, and Soil Organic Matter Dynamics. *Soil Tillage Res.* **2004**, *79* (1), 7–31.
- (18) Totsche, K. U.; Amelung, W.; Gerzabek, M. H.; Guggenberger, G.; Klumpp, E.; Knief, C.; Lehdorff, E.; Mikutta, R.; Peth, S.; Prechtel, A.; Ray, N.; Kögel-Knabner, I. Microaggregates in Soils. *J. Plant Nutr. Soil Sci.* **2018**, *181* (1), 104–136.
- (19) Brady, N. C.; Weil, R. R. *The Nature and Properties of Soils*; Prentice Hall, Inc.: Upper Saddle River, NJ, 2008.
- (20) Nimmo, J. R. Preferential Flow Occurs in Unsaturated Conditions. *Hydrol. Process.* **2012**, *26* (5), 786–789.
- (21) Young, I. M.; Ritz, K. Tillage, Habitat Space and Function of Soil Microbes. *Soil Tillage Res.* **2000**, *53* (3–4), 201–213.
- (22) Bundt, M.; Widmer, F.; Pesaro, M.; Zeyer, J.; Blaser, P. Preferential Flow Paths: Biological “hot Spots” in Soils. *Soil Biol. Biochem.* **2001**, *33* (6), 729–738.
- (23) Wolf, A. B.; Vos, M.; De Boer, W.; Kowalchuk, G. A. Impact of Matric Potential and Pore Size Distribution on Growth Dynamics of Filamentous and Non-Filamentous Soil Bacteria. *PLoS One* **2013**, *8* (12), e83661.
- (24) Franklin, S.; Vasilas, B.; Jin, Y. More than Meets the Dye: Evaluating Preferential Flow Paths as Microbial Hotspots. *Vadose Zone J.* **2019**, *18* (1), 1–8.
- (25) Bak, F.; Nybroe, O.; Zheng, B.; Badawi, N.; Hao, X.; Nicolaisen, M. H.; Aamand, J. Preferential Flow Paths Shape the Structure of Bacterial Communities in a Clayey till Depth Profile. *FEMS Microbiol. Ecol.* **2019**, *95* (3), 1–12.
- (26) Morales, V. L.; Parlange, J. Y.; Steenhuis, T. S. Are Preferential Flow Paths Perpetuated by Microbial Activity in the Soil Matrix? A Review. *J. Hydrol.* **2010**, *393* (1–2), 29–36.
- (27) Crawford, J. W.; Deacon, L.; Grinev, D.; Harris, J. A.; Ritz, K.; Singh, B. K.; Young, I. Microbial Diversity Affects Self-Organization of the Soil-Microbe System with Consequences for Function. *J. R. Soc. Interface* **2012**, *9*, 1302.
- (28) Liptzin, D.; Williams, M. W.; Helmig, D.; Seok, B.; Filippa, G.; Chowanski, K.; Hueber, J. Process-Level Controls on CO₂ Fluxes from a Seasonally Snow-Covered Subalpine Meadow Soil, Niwot Ridge, Colorado. *Biogeochemistry* **2009**, *95* (1), 151–166.
- (29) Rooney, E. C.; Bailey, V. L.; Patel, K. F.; Dragila, M.; Battu, A. K.; Buchko, A. C.; Gallo, A. C.; Hatten, J.; Possinger, A. R.; Qafoku, O.; Reno, L. R.; SanClements, M.; Varga, T.; Lybrand, R. A. Soil Pore Network Response to Freeze-Thaw Cycles in Permafrost Aggregates. *Geoderma* **2022**, *411*, 115674.
- (30) Ma, R.; Jiang, Y.; Liu, B.; Fan, H. Effects of Pore Structure Characterized by Synchrotron-Based Micro-Computed Tomography on Aggregate Stability of Black Soil under Freeze-Thaw Cycles. *Soil Tillage Res.* **2021**, *207*, 104855.
- (31) Calleja-Cabrera, J.; Boter, M.; Oñate-Sánchez, L.; Pernas, M. Root Growth Adaptation to Climate Change in Crops. *Front. Plant Sci.* **2020**, *11*, 544.
- (32) Lin, H. Temporal Stability of Soil Moisture Spatial Pattern and Subsurface Preferential Flow Pathways in the Shale Hills Catchment. *Vadose Zone J.* **2006**, *5*, 317–340.
- (33) Zhang, Y. H.; Niu, J. Z.; Yu, X. X.; Zhu, W. L.; Du, X. Q. Effects of Fine Root Length Density and Root Biomass on Soil Preferential Flow in Forest Ecosystems. *For. Syst.* **2015**, *24* (1), e012.
- (34) Nuccio, E. E.; Starr, E.; Karaoz, U.; Brodie, E. L.; Zhou, J.; Tringe, S. G.; Malmstrom, R. R.; Woyke, T.; Banfield, J. F.; Firestone, M. K.; Pett-Ridge, J. Niche Differentiation Is Spatially and Temporally Regulated in the Rhizosphere. *ISME J.* **2020**, *14* (4), 999–1014.
- (35) Greenwood, D. J.; Goodman, D. Direct Measurements of the Distribution of Oxygen in Soil Aggregates and in Columns of Fine Soil Crumbs. *J. Soil Sci.* **1967**, *18* (1), 182–196.
- (36) Sextone, A. J.; Revsbech, N. P.; Parkin, T. B.; Tiedje, J. M. Direct Measurement of Oxygen Profiles and Denitrification Rates in Soil Aggregates. *Soil Sci. Soc. Am. J.* **1985**, *49* (3), 645–651.
- (37) Brewer, P. E.; Calderón, F.; Vigil, M.; von Fischer, J. C. Impacts of Moisture, Soil Respiration, and Agricultural Practices on Methanogenesis in Upland Soils as Measured with Stable Isotope Pool Dilution. *Soil Biol. Biochem.* **2018**, *127*, 239–251.
- (38) Angel, R.; Claus, P.; Conrad, R. Methanogenic Archaea Are Globally Ubiquitous in Aerated Soils and Become Active under Wet Anoxic Conditions. *ISME J.* **2012**, *6* (4), 847–862.
- (39) Boye, K.; Noël, V.; Tfaily, M. M.; Bone, S. E.; Williams, K. H.; Bargar, J. R.; Fendorf, S. Thermodynamically Controlled Preservation of Organic Carbon in Floodplains. *Nat. Geosci.* **2017**, *10*, 415–419.
- (40) Van Der Lee, G. E. M.; De Winder, B.; Bouten, W.; Tietema, A. Anoxic Microsites in Douglas Fir Litter. *Soil Biol. Biochem.* **1999**, *31* (9), 1295–1301.

- (41) O'Connell, C. S.; Ruan, L.; Silver, W. L. Drought Drives Rapid Shifts in Tropical Rainforest Soil Biogeochemistry and Greenhouse Gas Emissions. *Nat. Commun.* **2018**, *9* (1), 1348.
- (42) Wilmoth, J. L. Redox Heterogeneity Entangles Soil and Climate Interactions. *Sustainability* **2021**, *13*, 10084.
- (43) Keiluweit, M.; Nico, P. S.; Kleber, M.; Fendorf, S. Are Oxygen Limitations under Recognized Regulators of Organic Carbon Turnover in Upland Soils? *Biogeochemistry* **2016**, *127*, 157–171.
- (44) Wang, B.; Brewer, P. E.; Shugart, H. H.; Lerdau, M. T.; Allison, S. D. Soil Aggregates as Biogeochemical Reactors and Implications for Soil-Atmosphere Exchange of Greenhouse Gases—A Concept. *Glob. Change Biol.* **2019**, *25* (2), 373–385.
- (45) Essaid, H. I.; Bekins, B. A.; Cozzarelli, I. M. Organic Contaminant Transport and Fate in the Subsurface: Evolution of Knowledge and Understanding. *Water Resour. Res.* **2015**, *51* (7), 4861–4902.
- (46) Campbell, K. M.; Kukkadapu, R. K.; Qafoku, N. P.; Peacock, A. D.; Leshner, E.; Williams, K. H.; Bargar, J. R.; Wilkins, M. J.; Figueroa, L.; Ranville, J.; Davis, J. A.; Long, P. E. Geochemical, Mineralogical and Microbiological Characteristics of Sediment from a Naturally Reduced Zone in a Uranium-Contaminated Aquifer. *Appl. Geochem.* **2012**, *27* (8), 1499–1511.
- (47) Silver, W. L.; Lugo, A. E.; Keller, M. Soil Oxygen Availability and Biogeochemistry along Rainfall and Topographic Gradients in Upland Wet Tropical Forest Soils. *Biogeochemistry* **1999**, *44* (3), 301–328.
- (48) Wanzek, T.; Keiluweit, M.; Baham, J.; Dragila, M.; Fendorf, S.; Fiedler, S.; Nico, P. S.; Kleber, M. Quantifying Biogeochemical Heterogeneity in Soil Systems. *Geoderma* **2018**, *324*, 89–97.
- (49) Ebrahimi, A.; Or, D. Dynamics of Soil Biogeochemical Gas Emissions Shaped by Remolded Aggregate Sizes and Carbon Configurations under Hydration Cycles. *Glob. Change Biol.* **2018**, *24* (1), e378–e392.
- (50) Ebrahimi, A.; Or, D. Microbial Community Dynamics in Soil Aggregates Shape Biogeochemical Gas Fluxes from Soil Profiles - Upscaling an Aggregate Biophysical Model. *Glob. Change Biol.* **2016**, *22* (9), 3141–3156.
- (51) Keiluweit, M.; Gee, K.; Denney, A.; Fendorf, S. Anoxic Microsites in Upland Soils Dominantly Controlled by Clay Content. *Soil Biol. Biochem.* **2018**, *118*, 42–50.
- (52) Berner, R. A. GEOCARBSULF: A Combined Model for Phanerozoic Atmospheric O₂ and CO₂. *Geochim. Cosmochim. Acta* **2006**, *70*, 5653–5664.
- (53) Fontaine, S.; Barot, S.; Barré, P.; Bdioui, N.; Mary, B.; Rumpel, C. Stability of Organic Carbon in Deep Soil Layers Controlled by Fresh Carbon Supply. *Nature* **2007**, *450*, 277–280.
- (54) Klotzbücher, T.; Kaiser, K.; Guggenberger, G.; Gatzek, C.; Kalbitz, K. A New Conceptual Model for the Fate of Lignin in Decomposing Plant Litter. *Ecology* **2011**, *92* (5), 1052–1062.
- (55) Viscarra Rossel, R. A.; Lee, J.; Behrens, T.; Luo, Z.; Baldock, J.; Richards, A. Continental-Scale Soil Carbon Composition and Vulnerability Modulated by Regional Environmental Controls. *Nat. Geosci.* **2019**, *12*, 547–552.
- (56) Hobbie, S. E.; Schimel, J. P.; Trumbore, S. E.; Randerson, J. R. Controls over Carbon Storage and Turnover in High-Latitude Soils. *Glob. Change Biol.* **2000**, *6*, 196–210.
- (57) Lacroix, E. M.; Rossi, R. J.; Bossio, D.; Fendorf, S. Effects of Moisture and Physical Disturbance on Pore-Scale Oxygen Content and Anaerobic Metabolisms in Upland Soils. *Sci. Total Environ.* **2021**, *780*, 146572.
- (58) Bailey, V. L.; Smith, A. P.; Tfaily, M.; Fansler, S. J.; Bond-Lamberty, B. Differences in Soluble Organic Carbon Chemistry in Pore Waters Sampled from Different Pore Size Domains. *Soil Biol. Biochem.* **2017**, *107*, 133–143.
- (59) Pold, G.; Grandy, A. S.; Melillo, J. M.; DeAngelis, K. M. Changes in Substrate Availability Drive Carbon Cycle Response to Chronic Warming. *Soil Biol. Biochem.* **2017**, *110*, 68–78.
- (60) Sierra, C. A.; Malghani, S.; Loescher, H. W. Interactions among Temperature, Moisture, and Oxygen Concentrations in Controlling Decomposition Rates in a Boreal Forest Soil. *Biogeosciences* **2017**, *14* (3), 703–710.
- (61) Voroney, R.; Heck, R. The Soil Habitat. In *Soil Microbiology, Ecology, and Biochemistry*; Paul, E. A., Ed.; Academic Press: San Francisco, CA, 2015; pp 15–40. DOI: 10.1016/B978-0-12-415955-6.00002-5.
- (62) Fischer, W. R.; Flessa, H.; Schaller, G. PH Values and Redox Potentials in Microsites of the Rhizosphere. *Z. Für Pflanzenernähr. Bodenkd.* **1989**, *152* (2), 191–195.
- (63) Kravchenko, A. N.; Toosi, E. R.; Guber, A. K.; Ostrom, N. E.; Yu, J.; Azeem, K.; Rivers, M. L.; Robertson, G. P. Hotspots of Soil N₂O Emission Enhanced through Water Absorption by Plant Residue. *Nat. Geosci.* **2017**, *10* (7), 496–500.
- (64) Parkin, T. B. Soil Microsites as a Source of Denitrification Variability. *Soil Sci. Soc. Am. J.* **1987**, *51* (5), 1194–1199.
- (65) Noël, V.; Boye, K.; Kukkadapu, R. K.; Bone, S.; Lezama Pacheco, J. S.; Cardarelli, E.; Janot, N.; Fendorf, S.; Williams, K. H.; Bargar, J. R. Understanding Controls on Redox Processes in Floodplain Sediments of the Upper Colorado River Basin. *Sci. Total Environ.* **2017**, *603–604*, 663–675.
- (66) Lacroix, E. M.; Masue-Slowey, Y.; Dlott, G. A.; Keiluweit, M.; Chadwick, O. A.; Fendorf, S. Mineral Protection and Resource Limitations Combine to Explain Profile-Scale Soil Carbon Persistence. *J. Geophys. Res. Biogeosciences* **2022**, *127* (4), 1–14.
- (67) Keiluweit, M.; Wanzek, T.; Kleber, M.; Nico, P.; Fendorf, S. E. Anaerobic Microsites Have an Unaccounted Role in Soil Carbon Stabilization. *Nat. Commun.* **2017**, *8* (1771), 1–8.
- (68) Naughton, H. R.; Keiluweit, M.; Tfaily, M. M.; Dynes, J. J.; Regier, T.; Fendorf, S. Development of Energetic and Enzymatic Limitations on Microbial Carbon Cycling in Soils. *Biogeochemistry* **2021**, *153* (2), 191–213.
- (69) Pallud, C.; Kausch, M.; Fendorf, S.; Meile, C. Spatial Patterns and Modeling of Reductive Ferrihydrite Transformation Observed in Artificial Soil Aggregates. *Environ. Sci. Technol.* **2010**, *44* (1), 74–79.
- (70) Yabusaki, S. B.; Wilkins, M. J.; Fang, Y.; Williams, K. H.; Arora, B.; Bargar, J.; Beller, H. R.; Bouskill, N. J.; Brodie, E. L.; Christensen, J. N.; Conrad, M. E.; Danczak, R. E.; King, E.; Soltanian, M. R.; Spycher, N. F.; Steefel, C. I.; Tokunaga, T. K.; Versteeg, R.; Waichler, S. R.; Wainwright, H. M. Water Table Dynamics and Biogeochemical Cycling in a Shallow, Variably-Saturated Floodplain. *Environ. Sci. Technol.* **2017**, *51* (6), 3307–3317.
- (71) Liptzin, D.; Silver, W. L.; Detto, M. Temporal Dynamics in Soil Oxygen and Greenhouse Gases in Two Humid Tropical Forests. *Ecosystems* **2011**, *14* (2), 171–182.
- (72) Lacroix, E. M.; Mendillo, J.; Gomes, A.; Dekas, A. E.; Fendorf, S. Contributions of Anoxic Microsites to Soil Carbon Protection across Soil Textures. *Geoderma* **2022**, *425*, 116050.
- (73) Kumaragamage, D.; Warren, C. J.; Spiers, G. Soil Chemistry. *Digging into Canadian Soils: An Introduction to Soil Science*; Canadian Society of Soil Science, 2021.
- (74) Freeman, C.; Ostle, N.; Kang, H. An Enzymic 'Latch' on a Global Carbon Store. *Nature* **2001**, *409*, 149–150.
- (75) Sinsabaugh, R. L. Phenol Oxidase, Peroxidase and Organic Matter Dynamics of Soil. *Soil Biol. Biochem.* **2010**, *42* (3), 391–404.
- (76) Jin, Q.; Bethke, C. M. Predicting the Rate of Microbial Respiration in Geochemical Environments. *Geochim. Cosmochim. Acta* **2005**, *69* (5), 1133–1143.
- (77) LaRowe, D. E.; Van Cappellen, P. Degradation of Natural Organic Matter: A Thermodynamic Analysis. *Geochim. Cosmochim. Acta* **2011**, *75* (8), 2030–2042.
- (78) Schuur, E. A.; Chadwick, O. A.; Matson, P. A. Carbon Cycling and Soil Carbon Storage in Mesic to Wet Hawaiian Montane Forests. *Ecology* **2001**, *82* (11), 3182–3196.
- (79) Hall, S. J.; Silver, W. L. Reducing Conditions, Reactive Metals, and Their Interactions Can Explain Spatial Patterns of Surface Soil Carbon in a Humid Tropical Forest. *Biogeochemistry* **2015**, *125* (2), 149–165.

- (80) Chen, C.; Hall, S. J.; Coward, E.; Thompson, A. Iron-Mediated Organic Matter Decomposition in Humid Soils Can Counteract Protection. *Nat. Commun.* **2020**, *11*, 2255.
- (81) Huang, W.; Hall, S. J. Elevated Moisture Stimulates Carbon Loss from Mineral Soils by Releasing Protected Organic Matter. *Nat. Commun.* **2017**, *8* (1774), 1–10.
- (82) Dalcin Martins, P.; Hoyt, D. W.; Bansal, S.; Mills, C. T.; Tfaily, M.; Tangen, B. A.; Finocchiaro, R. G.; Johnston, M. D.; McAdams, B. C.; Solensky, M. J.; Smith, G. J.; Chin, Y.; Wilkins, M. J. Abundant Carbon Substrates Drive Extremely High Sulfate Reduction Rates and Methane Fluxes in Prairie Pothole Wetlands. *Glob. Change Biol.* **2017**, *23* (8), 3107–3120.
- (83) Praeg, N.; Schwinghammer, L.; Illmer, P. Larix Decidua and Additional Light Affect the Methane Balance of Forest Soil and the Abundance of Methanogenic and Methanotrophic Microorganisms. *FEMS Microbiol. Lett.* **2019**, *366* (24), fnz259.
- (84) Angel, R.; Matthies, D.; Conrad, R. Activation of Methanogenesis in Arid Biological Soil Crusts Despite the Presence of Oxygen. *PLoS One* **2011**, *6* (5), No. e20453.
- (85) Teh, Y. A.; Silver, W. L.; Conrad, M. E. Oxygen Effects on Methane Production and Oxidation in Humid Tropical Forest Soils. *Glob. Change Biol.* **2005**, *11* (8), 1283–1297.
- (86) Treat, C. C.; Bloom, A. A.; Marushchak, M. E. Nongrowing Season Methane Emissions—a Significant Component of Annual Emissions across Northern Ecosystems. *Glob. Change Biol.* **2018**, *24* (8), 3331–3343.
- (87) Tiedje, J. M. Ecology of Denitrification and Dissimilatory Nitrate Reduction to Ammonium. In *Environmental Microbiology of Anaerobes*; Zehnder, A. J. B., Ed.; John Wiley and Sons: New York, 1988; pp 179–244.
- (88) Zumft, W. G. Cell Biology and Molecular Basis of Denitrification. *Microbiol. Mol. Biol. Rev. MMBR* **1997**, *61* (4), 533–616.
- (89) Pett-Ridge, J.; Silver, W. L.; Firestone, M. K. Redox Fluctuations Frame Microbial Community Impacts on N-Cycling Rates in a Humid Tropical Forest Soil. *Biogeochemistry* **2006**, *81* (1), 95–110.
- (90) Hayden, M. J.; Ross, D. S. Denitrification as a Nitrogen Removal Mechanism in a Vermont Peatland. *J. Environ. Qual.* **2005**, *34* (6), 2052–2061.
- (91) Robertson, G. P.; Hutson, M. A.; Evans, F. C.; Tiedje, J. M. Spatial Variability in a Successional Plant Community: Patterns of Nitrogen Availability. *Ecology* **1988**, *69* (5), 1517–1524.
- (92) Parkin, T. B. Characterizing the Variability of Soil Denitrification. *Denitrification in soil and sediment*; Springer, 1990; pp 213–228.
- (93) Holmes, W. E.; Zak, D. R. Soil Microbial Control of Nitrogen Loss Following Clear-cut Harvest in Northern Hardwood Ecosystems. *Ecol. Appl.* **1999**, *9* (1), 202–215.
- (94) Højberg, O.; Schnider, U.; Winteler, H. V.; Sørensen, J.; Haas, D. Oxygen-Sensing Reporter Strain of Pseudomonas Fluorescens for Monitoring the Distribution of Low-Oxygen Habitats in Soil. *Appl. Environ. Microbiol.* **1999**, *65* (9), 4085–4093.
- (95) Seech, A. G.; Beauchamp, E. G. Denitrification in Soil Aggregates of Different Sizes. *Soil Sci. Soc. Am. J.* **1988**, *52* (6), 1616–1621.
- (96) Rohe, L.; Apelt, B.; Vogel, H.-J.; Well, R.; Wu, G.-M.; Schlüter, S. Denitrification in Soil as a Function of Oxygen Availability at the Microscale. *Biogeosciences* **2021**, *18* (3), 1185–1201.
- (97) Schlüter, S.; Zawallich, J.; Vogel, H.-J.; Dörsch, P. Physical Constraints for Respiration in Microbial Hotspots in Soil and Their Importance for Denitrification. *Biogeosciences* **2019**, *16* (18), 3665–3678.
- (98) Schimel, J. P.; Bennett, J. Nitrogen Mineralization: Challenges of a Changing Paradigm. *Ecology* **2004**, *85* (3), 591–602.
- (99) Hodge, A.; Stewart, J.; Robinson, D.; Griffiths, B. S.; Fitter, A. H. Competition between Roots and Soil Micro-Organisms for Nutrients from Nitrogen-Rich Patches of Varying Complexity. *J. Ecol.* **2000**, *88* (1), 150–164.
- (100) Davidson, E. A.; Swank, W. T. Nitrous Oxide Dissolved in Soil Solution: An Insignificant Pathway of Nitrogen Loss from a Southeastern Hardwood Forest. *Water Resour. Res.* **1990**, *26* (7), 1687–1690.
- (101) Maier, S.; Kratz, A. M.; Weber, J.; Prass, M.; Liu, F.; Clark, A. T.; Abed, R. M. M.; Su, H.; Cheng, Y.; Eickhorst, T.; Fiedler, S.; Pöschl, U.; Weber, B. Water-Driven Microbial Nitrogen Transformations in Biological Soil Crusts Causing Atmospheric Nitrous Acid and Nitric Oxide Emissions. *ISME J.* **2022**, *16* (4), 1012–1024.
- (102) Friedl, J.; Scheer, C.; De Rosa, D.; Müller, C.; Grace, P. R.; Rowlings, D. W. Sources of Nitrous Oxide from Intensively Managed Pasture Soils: The Hole in the Pipe. *Environ. Res. Lett.* **2021**, *16* (6), 065004.
- (103) Hall, S. J.; Reyes, L.; Huang, W.; Homyak, P. M. Wet Spots as Hotspots: Moisture Responses of Nitric and Nitrous Oxide Emissions From Poorly Drained Agricultural Soils. *J. Geophys. Res. Biogeosciences* **2018**, *123* (12), 3589–3602.
- (104) Kuypers, M. M. M.; Sliemers, A. O.; Lavik, G.; Schmid, M.; Jørgensen, B. B.; Kuenen, J. G.; Sinninghe Damsté, J. S.; Strous, M.; Jetten, M. S. M. Anaerobic Ammonium Oxidation by Anammox Bacteria in the Black Sea. *Nature* **2003**, *422* (6932), 608–611.
- (105) Zhu, G.; Jetten, M. S. M.; Kuschik, P.; Ettwig, K. F.; Yin, C. Potential Roles of Anaerobic Ammonium and Methane Oxidation in the Nitrogen Cycle of Wetland Ecosystems. *Appl. Microbiol. Biotechnol.* **2010**, *86* (4), 1043–1055.
- (106) Humbert, S.; Tarnawski, S.; Fromin, N.; Mallet, M.-P.; Aragno, M.; Zopfi, J. Molecular Detection of Anammox Bacteria in Terrestrial Ecosystems: Distribution and Diversity. *ISME J.* **2010**, *4* (3), 450–454.
- (107) Grabb, K. C.; Buchwald, C.; Hansel, C. M.; Wankel, S. D. A Dual Nitrite Isotopic Investigation of Chemodenitrification by Mineral-Associated Fe(II) and Its Production of Nitrous Oxide. *Geochim. Cosmochim. Acta* **2017**, *196*, 388–402.
- (108) Stevenson, F. J.; Cole, M. A. *Cycles of Soils: Carbon, Nitrogen, Phosphorus, Sulfur, Micronutrients*; John Wiley & Sons, 1999.
- (109) Shen, L.; Liu, S.; Huang, Q.; Lian, X.; He, Z.; Geng, S.; Jin, R.; He, Y.; Lou, L.; Xu, X.; et al. Evidence for the Cooccurrence of Nitrite-Dependent Anaerobic Ammonium and Methane Oxidation Processes in a Flooded Paddy Field. *Appl. Environ. Microbiol.* **2014**, *80* (24), 7611–7619.
- (110) Welte, C. U.; Rasigraf, O.; Vaksmaa, A.; Versantvoort, W.; Arshad, A.; Op den Camp, H. J. M.; Jetten, M. S. M.; Lüke, C.; Reimann, J. Nitrate- and Nitrite-Dependent Anaerobic Oxidation of Methane. *Environ. Microbiol. Rep.* **2016**, *8* (6), 941–955.
- (111) Hu, B.; Shen, L.; Lian, X.; Zhu, Q.; Liu, S.; Huang, Q.; He, Z.; Geng, S.; Cheng, D.; Lou, L.; Xu, X.; Zheng, P.; He, Y. Evidence for Nitrite-Dependent Anaerobic Methane Oxidation as a Previously Overlooked Microbial Methane Sink in Wetlands. *Proc. Natl. Acad. Sci. U. S. A.* **2014**, *111* (12), 4495–4500.
- (112) Cheng, C.; Zhang, J.; He, Q.; Wu, H.; Chen, Y.; Xie, H.; Pavlostathis, S. G. Exploring Simultaneous Nitrous Oxide and Methane Sink in Wetland Sediments under Anoxic Conditions. *Water Res.* **2021**, *194*, 116958.
- (113) Harris, E.; Diaz-Pines, E.; Stoll, E.; Schloter, M.; Schulz, S.; Duffner, C.; Li, K.; Moore, K. L.; Ingrisch, J.; Reinthaler, D.; Zechmeister-Boltenstern, S.; Glatzel, S.; Brüggemann, N.; Bahn, M. Denitrifying Pathways Dominate Nitrous Oxide Emissions from Managed Grassland during Drought and Rewetting. *Sci. Adv.* **2021**, *7* (6), No. eabb7118.
- (114) Fan, L.; Schneider, D.; Dippold, M. A.; Poehlein, A.; Wu, W.; Gui, H.; Ge, T.; Wu, J.; Thiel, V.; Kuzyakov, Y.; et al. Active Metabolic Pathways of Anaerobic Methane Oxidation in Paddy Soils. *Soil Biol. Biochem.* **2021**, *156*, 108215.
- (115) Payne, W. J. *Denitrification*; Wiley: New York, 1981.
- (116) Schepers, J. S.; Raun, W. *Nitrogen in Agricultural Systems*; ASA-CSSA-SSSA, 2008.
- (117) Coyne, M. S. Biological Denitrification. In *Agronomy Monographs*; Schepers, J. S., Raun, W. R., Eds.; American Society of Agronomy, Crop Science Society of America, Soil Science Society of

- America: Madison, WI, USA, 2015; pp 201–253; DOI: 10.2134/agronmonogr49.c7.
- (118) Flessa, H.; Beese, F. Effects of Sugarbeet Residues on Soil Redox Potential and Nitrous Oxide Emission. *Soil Sci. Soc. Am. J.* **1995**, *59* (4), 1044–1051.
- (119) Wagner-Riddle, C.; Congreves, K. A.; Abalos, D.; Berg, A. A.; Brown, S. E.; Ambadan, J. T.; Gao, X.; Tenuta, M. Globally Important Nitrous Oxide Emissions from Croplands Induced by Freeze-Thaw Cycles. *Nat. Geosci.* **2017**, *10* (4), 279–283.
- (120) Pett-Ridge, J.; Petersen, D. G.; Nuccio, E.; Firestone, M. K. Influence of Oxidic/Anoxic Fluctuations along Ammonia Oxidizers and Nitrification Potential in a Wet Tropical Soil. *FEMS Microbiol. Ecol.* **2013**, *85* (1), 179–194.
- (121) Winnick, M. J. Stream Transport and Substrate Controls on Nitrous Oxide Yields From Hyporheic Zone Denitrification. *AGU Adv.* **2021**, *2* (4), No. e2021AV000517.
- (122) Thompson, A.; Rancourt, D. G.; Chadwick, O. A.; Chorover, J. Iron Solid-Phase Differentiation along a Redox Gradient in Basaltic Soils. *Geochim. Cosmochim. Acta* **2011**, *75* (1), 119–133.
- (123) Schulz, M.; Stonestrom, D.; Lawrence, C.; Bullen, T.; Fitzpatrick, J.; Kyker-Snowman, E.; Manning, J.; Mnich, M. Structured Heterogeneity in a Marine Terrace Chronosequence: Upland Mottling. *Vadose Zone J.* **2016**, *15* (2), 1–14.
- (124) Patzner, M. S.; Mueller, C. W.; Malusova, M.; Baur, M.; Nikeleit, V.; Scholten, T.; Hoeschen, C.; Byrne, J. M.; Borch, T.; Kappler, A.; Bryce, C. Iron Mineral Dissolution Releases Iron and Associated Organic Carbon during Permafrost Thaw. *Nat. Commun.* **2020**, *11* (1), 1–11.
- (125) Fritzsche, A.; Bosch, J.; Sander, M.; Schroder, C.; Byrne, J. M.; Ritschel, T.; Joshi, P.; Maisch, M.; Meckenstock, R. U.; Kappler, A.; Totsche, K. U. Organic Matter from Redoximorphic Soils Accelerates and Sustains Microbial Fe(III) Reduction. *Environ. Sci. Technol.* **2021**, *55* (15), 10821–10831.
- (126) Patrick, W. H.; Khalid, R. A. Phosphate Release and Sorption by Soils and Sediments: Effect of Aerobic and Anaerobic Conditions. *Science* **1974**, *186* (4158), 53–55.
- (127) Lin, Y.; Gross, A.; O'Connell, C. S.; Silver, W. L. Anoxic Conditions Maintained High Phosphorus Sorption in Humid Tropical Forest Soils. *Biogeosciences* **2020**, *17* (1), 89–101.
- (128) Hall, S. J.; Huang, W. Iron Reduction: A Mechanism for Dynamic Cycling of Occluded Cations in Tropical Forest Soils? *Biogeochemistry* **2017**, *136* (1), 91–102.
- (129) Cummings, D. E.; Caccavo, F.; Fendorf, S.; Rosenzweig, R. F. Arsenic Mobilization by the Dissimilatory Fe(III)-Reducing Bacterium *Shewanella Alga* BrY. *Environ. Sci. Technol.* **1999**, *33* (5), 723–729.
- (130) Warrinnier, R.; Bossuyt, S.; Resseguier, C.; Cambier, P.; Houot, S.; Gustafsson, J. P.; Diels, J.; Smolders, E. Anaerobic Respiration in the Unsaturated Zone of Agricultural Soil Mobilises Phosphorus and Manganese. *Environ. Sci. Technol.* **2020**, *54*, 4922.
- (131) Malakar, A.; Kaiser, M.; Snow, D. D.; Walia, H.; Panda, B.; Ray, C. Ferrihydrate Reduction Increases Arsenic and Uranium Bioavailability in Unsaturated Soil. *Environ. Sci. Technol.* **2020**, *54* (21), 13839–13848.
- (132) Jones, M. E.; Nico, P. S.; Ying, S.; Regier, T.; Thieme, J.; Keiluweit, M. Manganese-Driven Carbon Oxidation at Oxidic-Anoxic Interfaces. *Environ. Sci. Technol.* **2018**, *52* (21), 12349–12357.
- (133) Keiluweit, M.; Nico, P.; Harmon, M. E.; Mao, J.; Pett-Ridge, J.; Kleber, M. Long-Term Litter Decomposition Controlled by Manganese Redox Cycling. *Proc. Natl. Acad. Sci. U. S. A.* **2015**, *112* (38), No. E5253–E5260.
- (134) Weber, K. A.; Achenbach, L. A.; Coates, J. D. Microorganisms Pumping Iron: Anaerobic Microbial Iron Oxidation and Reduction. *Nat. Rev. Microbiol.* **2006**, *4* (10), 752–764.
- (135) Kappler, A.; Bryce, C.; Mansor, M.; Lueder, U.; Byrne, J. M.; Swanner, E. D. An Evolving View on Biogeochemical Cycling of Iron. *Nat. Rev. Microbiol.* **2021**, *19* (6), 360–374.
- (136) Trusiak, A.; Treibergs, L.; Kling, G.; Cory, R. The Controls of Iron and Oxygen on Hydroxyl Radical ($\bullet\text{OH}$) Production in Soils. *Soil Syst.* **2019**, *3* (1), 1.
- (137) Kleber, M.; Bourg, I. C.; Coward, E. K.; Hansel, C. M.; Myneni, S. C. B.; Nunan, N. Dynamic Interactions at the Mineral–Organic Matter Interface. *Nat. Rev. Earth Environ.* **2021**, *2*, 402–421.
- (138) Nico, P. S.; Anastasio, C.; Zasoski, R. J. Rapid Photo-Oxidation of Mn(II) Mediated by Humic Substances. *Geochim. Cosmochim. Acta* **2002**, *66* (23), 4047–4056.
- (139) Duckworth, O. W.; Sposito, G. Siderophore-Manganese(III) Interactions. I. Air-Oxidation of Manganese(II) Promoted by Desferrioxamine B. *Environ. Sci. Technol.* **2005**, *39* (16), 6037–6044.
- (140) Jones, L. C.; Peters, B.; Lezama Pacheco, J. S.; Casciotti, K. L.; Fendorf, S. Stable Isotopes and Iron Oxide Mineral Products as Markers of Chemodenitrification. *Environ. Sci. Technol.* **2015**, *49* (6), 3444–3452.
- (141) Bhattacharyya, A.; Campbell, A. N.; Tfaily, M. M.; Lin, Y.; Kukkadapu, R. K.; Silver, W. L.; Nico, P. S.; Pett-Ridge, J. Redox Fluctuations Control the Coupled Cycling of Iron and Carbon in Tropical Forest Soils. *Environ. Sci. Technol.* **2018**, *52* (24), 14129–14139.
- (142) Barcellos, D.; O'Connell, C. S.; Silver, W.; Meile, C.; Thompson, A. Hot Spots and Hot Moments of Soil Moisture Explain Fluctuations in Iron and Carbon Cycling in a Humid Tropical Forest Soil. *Soil Syst.* **2018**, *2* (4), 59.
- (143) Chen, C.; Meile, C.; Wilmoth, J.; Barcellos, D.; Thompson, A. Influence of PO₂ on Iron Redox Cycling and Anaerobic Organic Carbon Mineralization in a Humid Tropical Forest Soil. *Environ. Sci. Technol.* **2018**, *52* (14), 7709–7719.
- (144) Garcia Arredondo, M.; Lawrence, C.; Schulz, M.; Tfaily, M. M.; Kukkadapu, R.; Jones, M. E.; Boye, K.; Keiluweit, M. Root-Driven Weathering Impacts on Mineral–Organic Associations in Deep Soils over Pedogenic Time Scales. *Geochim. Cosmochim. Acta* **2019**, *263*, 68–84.
- (145) Rickard, D.; Luther, G. W. Chemistry of Iron Sulfides. *Chem. Rev.* **2007**, *107* (2), 514–562.
- (146) Rickard, D.; Morse, J. W. Acid Volatile Sulfide (AVS). *Mar. Chem.* **2005**, *97* (3), 141–197.
- (147) Rickard, D. T. Kinetics and Mechanism of Pyrite Formation at Low Temperatures. *Am. J. Sci.* **1975**, *275* (6), 636.
- (148) Luther, G. W. Pyrite Synthesis via Polysulfide Compounds. *Geochim. Cosmochim. Acta* **1991**, *55* (10), 2839–2849.
- (149) Morse, J. W.; Arakaki, T. Adsorption and Coprecipitation of Divalent Metals with Mackinawite (FeS). *Geochim. Cosmochim. Acta* **1993**, *57* (15), 3635–3640.
- (150) Patterson, R. R.; Fendorf, S.; Fendorf, M. Reduction of Hexavalent Chromium by Amorphous Iron Sulfide. *Environ. Sci. Technol.* **1997**, *31* (7), 2039–2044.
- (151) Bostick, B. C.; Fendorf, S. Arsenite Sorption on Troilite (FeS) and Pyrite (FeS₂). *Geochim. Cosmochim. Acta* **2003**, *67* (5), 909–921.
- (152) Livens, F. R.; Jones, M. J.; Hynes, A. J.; Charnock, J. M.; Mosselmans, J. F. W.; Hennig, C.; Steele, H.; Collison, D.; Vaughan, D. J.; Patrick, R. A. D.; Reed, W. A.; Moyes, L. N. X-Ray Absorption Spectroscopy Studies of Reactions of Technetium, Uranium and Neptunium with Mackinawite. *J. Environ. Radioact.* **2004**, *74* (1–3), 211–219.
- (153) Vorlicek, T. P.; Kahn, M. D.; Kasuya, Y.; Helz, G. R. Capture of Molybdenum in Pyrite-Forming Sediments: Role of Ligand-Induced Reduction by Polysulfides 1 Associate Editor: M. Goldhaber. *Geochim. Cosmochim. Acta* **2004**, *68* (3), 547–556.
- (154) Jeong, H. Y.; Klaue, B.; Blum, J. D.; Hayes, K. F. Sorption of Mercuric Ion by Synthetic Nanocrystalline Mackinawite (FeS). *Environ. Sci. Technol.* **2007**, *41* (22), 7699–7705.
- (155) Dahl, T. W.; Chappaz, A.; Fitts, J. P.; Lyons, T. W. Molybdenum Reduction in a Sulfidic Lake: Evidence from X-Ray Absorption Fine-Structure Spectroscopy and Implications for the Mo Paleoproxy. *Geochim. Cosmochim. Acta* **2013**, *103*, 213–231.
- (156) Jeong, H. Y.; Han, Y.-S.; Park, S. W.; Hayes, K. F. Aerobic Oxidation of Mackinawite (FeS) and Its Environmental Implication

- for Arsenic Mobilization. *Geochim. Cosmochim. Acta* **2010**, *74* (11), 3182–3198.
- (157) Kulp, T. R.; Miller, L. G.; Braiotta, F.; Webb, S. M.; Kocar, B. D.; Blum, J. S.; Oremland, R. S. Microbiological Reduction of Sb(V) in Anoxic Freshwater Sediments. *Environ. Sci. Technol.* **2014**, *48* (1), 218–226.
- (158) Chang, H. S.; Buettner, S. W.; Seaman, J. C.; Jaffé, P. R.; Koster Van Groos, P. G.; Li, D.; Peacock, A. D.; Scheckel, K. G.; Kaplan, D. I. Uranium Immobilization in an Iron-Rich Rhizosphere of a Native Wetland Plant from the Savannah River Site under Reducing Conditions. *Environ. Sci. Technol.* **2014**, *48* (16), 9270–9278.
- (159) Veeramani, H.; Alessi, D. S.; Suvorova, E. I.; Lezama-Pacheco, J. S.; Stubbs, J. E.; Sharp, J. O.; Dippon, U.; Kappler, A.; Bargar, J. R.; Bernier-Latmani, R. Products of Abiotic U(VI) Reduction by Biogenic Magnetite and Vivianite. *Geochim. Cosmochim. Acta* **2011**, *75* (9), 2512–2528.
- (160) Kyung, D.; Sihm, Y.; Kim, S.; Bae, S.; Amin, M. T.; Alazba, A. A.; Lee, W. Synergistic Effect of Nano-Sized Mackinawite with Cyano-Cobalamin in Cement Slurries for Reductive Dechlorination of Tetrachloroethylene. *J. Hazard. Mater.* **2016**, *311*, 1–10.
- (161) Butler, E. C.; Hayes, K. F. Kinetics of the Transformation of Halogenated Aliphatic Compounds by Iron Sulfide. *Environ. Sci. Technol.* **2000**, *34* (3), 422–429.
- (162) Wan, M.; Shchukarev, A.; Lohmayer, R.; Planer-Friedrich, B.; Peiffer, S. Occurrence of Surface Polysulfides during the Interaction between Ferric (Hydr)Oxides and Aqueous Sulfide. *Environ. Sci. Technol.* **2014**, *48* (9), 5076–5084.
- (163) Peiffer, S.; Behrends, T.; Hellige, K.; Larese-Casanova, P.; Wan, M.; Pollok, K. Pyrite Formation and Mineral Transformation Pathways upon Sulfidation of Ferric Hydroxides Depend on Mineral Type and Sulfide Concentration. *Chem. Geol.* **2015**, *400*, 44–55.
- (164) Wan, M.; Schröder, C.; Peiffer, S. Fe(III):S(-II) Concentration Ratio Controls the Pathway and the Kinetics of Pyrite Formation during Sulfidation of Ferric Hydroxides. *Geochim. Cosmochim. Acta* **2017**, *217*, 334–348.
- (165) Noël, V.; Kumar, N.; Boye, K.; Barragan, L.; Lezama-Pacheco, J. S.; Chu, R.; Tolic, N.; Brown, G. E.; Bargar, J. R. FeS Colloids - Formation and Mobilization Pathways in Natural Waters. *Environ. Sci. Nano* **2020**, *7* (7), 2102–2116.
- (166) Kumar, N.; Noël, V.; Planer-Friedrich, B.; Besold, J.; Lezama-Pacheco, J.; Bargar, J. R.; Brown, G. E.; Fendorf, S.; Boye, K. Redox Heterogeneities Promote Thioarsenate Formation and Release into Groundwater from Low Arsenic Sediments. *Environ. Sci. Technol.* **2020**, *54* (6), 3237–3244.
- (167) Babey, T.; Boye, K.; Tolar, B.; Engel, M.; Noël, V.; Perzan, Z.; Kumar, N.; Francis, C. A.; Bargar, J. R.; Maher, K. Simulation of Anoxic Lenses as Exporters of Reactivity in Alluvial Aquifer Sediments. *Geochim. Cosmochim. Acta* **2022**, *334*, 119–134.
- (168) Engel, M.; Noël, V.; Kukkadapu, R. K.; Boye, K.; Bargar, J. R.; Fendorf, S. Nitrate Controls on the Extent and Type of Metal Retention in Fine-Grained Sediments of a Simulated Aquifer. *Environ. Sci. Technol.* **2022**, *56* (20), 14452–14461.
- (169) Dos Santos Afonso, M.; Stumm, W. Reductive Dissolution of Iron(III) (Hydr)Oxides by Hydrogen Sulfide. *Langmuir* **1992**, *8* (6), 1671–1675.
- (170) Weber, F.-A.; Voegelin, A.; Kaegi, R.; Kretzschmar, R. Contaminant Mobilization by Metallic Copper and Metal Sulphide Colloids in Flooded Soil. *Nat. Geosci.* **2009**, *2* (4), 267–271.
- (171) Rozan, T. F.; Lassman, M. E.; Ridge, D. P.; Luther, G. W. Evidence for Iron, Copper and Zinc Complexation as Multinuclear Sulphide Clusters in Oxidic Rivers. *Nature* **2000**, *406* (6798), 879–882.
- (172) Sharma, N.; Wang, Z.; Catalano, J. G.; Giammar, D. E. Dynamic Responses of Trace Metal Bioaccessibility to Fluctuating Redox Conditions in Wetland Soils and Stream Sediments. *ACS Earth Space Chem.* **2022**, *6* (5), 1331–1344.
- (173) Karimian, N.; Johnston, S. G.; Burton, E. D. Acidity Generation Accompanying Iron and Sulfur Transformations during Drought Simulation of Freshwater Re-Flooded Acid Sulfate Soils. *Geoderma* **2017**, *285*, 117–131.
- (174) Karimian, N.; Johnston, S. G.; Burton, E. D. Iron and Sulfur Cycling in Acid Sulfate Soil Wetlands under Dynamic Redox Conditions: A Review. *Chemosphere* **2018**, *197*, 803–816.
- (175) Zausig, J.; Stepniewski, W.; Horn, R. Oxygen Concentration and Redox Potential Gradients in Unsaturated Model Soil Aggregates. *Soil Sci. Soc. Am. J.* **1993**, *57*, 908–916.
- (176) Hall, S. J.; Liptzin, D.; Buss, H. L.; DeAngelis, K.; Silver, W. L. Drivers and Patterns of Iron Redox Cycling from Surface to Bedrock in a Deep Tropical Forest Soil: A New Conceptual Model. *Biogeochemistry* **2016**, *130* (1–2), 177–190.
- (177) Fiedler, S.; Vepraskas, M. J.; Richardson, J. L. Soil Redox Potential: Importance, Field Measurements, and Observations. In *Advances in Agronomy*; Sparks, D. L., Ed.; Academic Press, 2007; Vol. 94, pp 1–54; DOI: 10.1016/S0065-2113(06)94001-2.
- (178) Husson, O. Redox Potential (Eh) and PH as Drivers of Soil/Plant/Microorganism Systems: A Transdisciplinary Overview Pointing to Integrative Opportunities for Agronomy. *Plant Soil* **2013**, *362* (1–2), 389–417.
- (179) Aeppli, M.; Babey, T.; Engel, M.; Lacroix, E. M.; Tolar, B. B.; Fendorf, S.; Bargar, J. R.; Boye, K. Export of Organic Carbon from Reduced Fine-Grained Zones Governs Biogeochemical Reactivity in a Simulated Aquifer. *Environ. Sci. Technol.* **2022**, *56* (4), 2738–2746.
- (180) Angle, J. C.; Morin, T. H.; Solden, L. M.; Narrowe, A. B.; Smith, G. J.; Borton, M. A.; Rey-Sanchez, C.; Daly, R. A.; Mirfenderesgi, G.; Hoyt, D. W.; Riley, W. J.; Miller, C. S.; Bohrer, G.; Wrighton, K. C. Methanogenesis in Oxygenated Soils Is a Substantial Fraction of Wetland Methane Emissions. *Nat. Commun.* **2017**, *8* (1), 1–9.
- (181) von Fischer, J. C.; Hedin, L. O. Controls on Soil Methane Fluxes: Tests of Biophysical Mechanisms Using Table Isotope Tracers. *Global Biogeochem. Cycles* **2007**, *21* (2), 1–9.
- (182) Roscioli, J. R.; Meredith, L. K.; Shorter, J. H.; Gil-Loaiza, J.; Volkmann, T. H. M. Soil Gas Probes for Monitoring Trace Gas Messengers of Microbial Activity. *Sci. Rep.* **2021**, *11* (1), 1–11.
- (183) Wagner, D. Effect of Varying Soil Water Potentials on Methanogenesis in Aerated Marshland Soils. *Sci. Rep.* **2017**, *7* (1), 14706.
- (184) Khalil, K.; Mary, B.; Renault, P. Nitrous Oxide Production by Nitrification and Denitrification in Soil Aggregates as Affected by O₂ concentration. *Soil Biol. Biochem.* **2004**, *36* (4), 687–699.
- (185) Hall, S. J.; Weintraub, S. R.; Bowling, D. R. Scale-Dependent Linkages between Nitrate Isotopes and Denitrification in Surface Soils: Implications for Isotope Measurements and Models. *Oecologia* **2016**, *181* (4), 1221–1231.
- (186) Ostrom, N. E.; Sutka, R.; Ostrom, P. H.; Grandy, A. S.; Huizinga, K. M.; Gandhi, H.; von Fischer, J. C.; Robertson, G. P. Isotopologue Data Reveal Bacterial Denitrification as the Primary Source of N₂O during a High Flux Event Following Cultivation of a Native Temperate Grassland. *Soil Biol. Biochem.* **2010**, *42* (3), 499–506.
- (187) Sihi, D.; Davidson, E. A.; Savage, K. E.; Liang, D. Simultaneous Numerical Representation of Soil Microsite Production and Consumption of Carbon Dioxide, Methane, and Nitrous Oxide Using Probability Distribution Functions. *Glob. Change Biol.* **2020**, *26* (1), 200–218.
- (188) Viollier, E.; Inglett, P. W.; Hunter, K.; Roychoudhury, A. N.; Van Cappellen, P. The Ferrozine Method Revisited: Fe(II)/Fe(III) Determination in Natural Waters. *Appl. Geochem.* **2000**, *15*, 785–790.
- (189) Fogo, J. K.; Popowsky, M. Spectrophotometric Determination of Hydrogen Sulfide. *Anal. Chem.* **1949**, *21* (6), 732–734.
- (190) Strickland, J.; Parsons, T. A *Practical Handbook of Seawater Analysis*, 2nd ed.; Fisheries Research Board of Canada: Ottawa, 1972.
- (191) Stumm, W.; Morgan, J. J. *Aquatic Chemistry: Chemical Equilibria and Rates in Natural Waters*; Wiley: New York, 1996.
- (192) Yu, Q.; Si, G.; Zong, T.; Mulder, J.; Duan, L. High Hydrogen Sulfide Emissions from Subtropical Forest Soils Based on Field Measurements in South China. *Sci. Total Environ.* **2019**, *651*, 1302–1309.

- (193) Pietramellara, G.; Ascher, J.; Borgogni, F.; Ceccherini, M. T.; Guerri, G.; Nannipieri, P. Extracellular DNA in Soil and Sediment: Fate and Ecological Relevance. *Biol. Fertil. Soils* **2009**, *45* (3), 219–235.
- (194) Carini, P.; Marsden, P. J.; Leff, J. W.; Morgan, E. E.; Strickland, M. S.; Fierer, N. Relic DNA Is Abundant in Soil and Obscures Estimates of Soil Microbial Diversity. *Nat. Microbiol.* **2017**, *2*, 16242.
- (195) Belasco, J. G. mRNA Degradation in Prokaryotic Cells: An Overview. *Control of Messenger RNA Stability*; Elsevier, 1993; pp 3–12; DOI: 10.1016/B978-0-08-091652-1.50005-0.
- (196) Hammerl, V.; Kastl, E.-M.; Schloter, M.; Kublik, S.; Schmidt, H.; Welzl, G.; Jentsch, A.; Beierkuhnlein, C.; Gschwendtner, S. Influence of Rewetting on Microbial Communities Involved in Nitrification and Denitrification in a Grassland Soil after a Prolonged Drought Period. *Sci. Rep.* **2019**, *9* (1), 2280.
- (197) Hansel, C. M.; Fendorf, S.; Jardine, P. M.; Francis, C. A. Changes in Bacterial and Archaeal Community Structure and Functional Diversity along a Geochemically Variable Soil Profile. *Appl. Environ. Microbiol.* **2008**, *74* (5), 1620–1633.
- (198) Nogueira, M. A.; Nehls, U.; Hampp, R.; Poralla, K.; Cardoso, E. J. B. N. Mycorrhiza and Soil Bacteria Influence Extractable Iron and Manganese in Soil and Uptake by Soybean. *Plant Soil* **2007**, *298* (1–2), 273–284.
- (199) Alteio, L. V.; Senecca, J.; Canarini, A.; Angel, R.; Jansa, J.; Guseva, K.; Kaiser, C.; Richter, A.; Schmidt, H. A Critical Perspective on Interpreting Amplicon Sequencing Data in Soil Ecological Research. *Soil Biol. Biochem.* **2021**, *160*, 108357.
- (200) Couradeau, E.; Sasse, J.; Goudeau, D.; Nath, N.; Hazen, T. C.; Bowen, B. P.; Chakraborty, R.; Malmstrom, R. R.; Northen, T. R. Probing the Active Fraction of Soil Microbiomes Using BONCAT-FACS. *Nat. Commun.* **2019**, *10* (1), 2770.
- (201) Coskun, Ö. K.; Özen, V.; Wankel, S. D.; Orsi, W. D. Quantifying Population-Specific Growth in Benthic Bacterial Communities under Low Oxygen Using H218O. *ISME J.* **2019**, *13* (6), 1546–1559.
- (202) Pett-Ridge, J.; Firestone, M. K. Using Stable Isotopes to Explore Root-Microbe-Mineral Interactions in Soil. *Rhizosphere* **2017**, *3* (April), 244–253.
- (203) Zhang, L.; Dumont, M. G.; Bodelier, P. L. E.; Adams, J. M.; He, D.; Chu, H. DNA Stable-Isotope Probing Highlights the Effects of Temperature on Functionally Active Methanotrophs in Natural Wetlands. *Soil Biol. Biochem.* **2020**, *149*, 107954.
- (204) Voegel, T. M.; Larrabee, M. M.; Nelson, L. M. Development of Droplet Digital PCR Assays to Quantify Genes Involved in Nitrification and Denitrification, Comparison with Quantitative Real-Time PCR and Validation of Assays in Vineyard Soil. *Can. J. Microbiol.* **2021**, *67* (2), 174–187.
- (205) Kokkoris, V.; Vukicevich, E.; Richards, A.; Thomsen, C.; Hart, M. M. Challenges Using Droplet Digital PCR for Environmental Samples. *Appl. Microbiol.* **2021**, *1* (1), 74–88.
- (206) Myrold, D. D.; Zeglin, L. H.; Jansson, J. K. The Potential of Metagenomic Approaches for Understanding Soil Microbial Processes. *Soil Sci. Soc. Am. J.* **2014**, *78* (1), 3–10.
- (207) Matheus Carnevali, P. B.; Lavy, A.; Thomas, A. D.; Crits-Christoph, A.; Diamond, S.; Méheust, R.; Olm, M. R.; Sharrar, A.; Lei, S.; Dong, W.; Falco, N.; Bouskill, N.; Newcomer, M. E.; Nico, P.; Wainwright, H.; Dwivedi, D.; Williams, K. H.; Hubbard, S.; Banfield, J. F. Meanders as a Scaling Motif for Understanding of Floodplain Soil Microbiome and Biogeochemical Potential at the Watershed Scale. *Microbiome* **2021**, *9* (1), 121.
- (208) He, Z.; Gentry, T. J.; Schadt, C. W.; Wu, L.; Liebich, J.; Chong, S. C.; Huang, Z.; Wu, W.; Gu, B.; Jardine, P.; Criddle, C.; Zhou, J. GeoChip: A Comprehensive Microarray for Investigating Biogeochemical, Ecological and Environmental Processes. *ISME J.* **2007**, *1* (1), 67–77.
- (209) Van Nostrand, J. D.; Wu, W.-M.; Wu, L.; Deng, Y.; Carley, J.; Carroll, S.; He, Z.; Gu, B.; Luo, J.; Criddle, C. S.; Watson, D. B.; Jardine, P. M.; Marsh, T. L.; Tiedje, J. M.; Hazen, T. C.; Zhou, J. GeoChip-Based Analysis of Functional Microbial Communities during the Reoxidation of a Bioreduced Uranium-Contaminated Aquifer. *Environ. Microbiol.* **2009**, *11* (10), 2611–2626.
- (210) Passarelli, M. K.; Pirkl, A.; Moellers, R.; Grinfeld, D.; Kollmer, F.; Havelund, R.; Newman, C. F.; Marshall, P. S.; Arlinghaus, H.; Alexander, M. R.; West, A.; Horning, S.; Niehuis, E.; Makarov, A.; Dollery, C. T.; Gilmore, I. S. The 3D OrbiSIMS—Label-Free Metabolic Imaging with Subcellular Lateral Resolution and High Mass-Resolving Power. *Nat. Methods* **2017**, *14* (12), 1175–1183.
- (211) Zhang, J.; Brown, J.; Scurr, D. J.; Bullen, A.; MacLellan-Gibson, K.; Williams, P.; Alexander, M. R.; Hardie, K. R.; Gilmore, I. S.; Rakowska, P. D. Cryo-OrbiSIMS for 3D Molecular Imaging of a Bacterial Biofilm in Its Native State. *Anal. Chem.* **2020**, *92* (13), 9008–9015.
- (212) Helliwell, J. R.; Sturrock, C. J.; Grayling, K. M.; Tracy, S. R.; Flavel, R. J.; Young, I. M.; Whalley, W. R.; Mooney, S. J. Applications of X-Ray Computed Tomography for Examining Biophysical Interactions and Structural Development in Soil Systems: A Review. *Eur. J. Soil Sci.* **2013**, *64* (3), 279–297.
- (213) Zhu, K.; Bruun, S.; Larsen, M.; Glud, R. N.; Jensen, L. S. Spatial Oxygen Distribution and Nitrous Oxide Emissions from Soil after Manure Application: A Novel Approach Using Planar Optodes. *J. Environ. Qual.* **2014**, *43* (5), 1809–1812.
- (214) Ceriotti, G.; Borisov, S. M.; Berg, J. S.; de Anna, P. Morphology and Size of Bacterial Colonies Control Anoxic Microenvironment Formation in Porous Media. *Environ. Sci. Technol.* **2022**, *56* (23), 17471–17480.
- (215) Li, C.; Ding, S.; Yang, L.; Zhu, Q.; Chen, M.; Tsang, D. C. W.; Cai, G.; Feng, C.; Wang, Y.; Zhang, C. Planar Optode: A Two-Dimensional Imaging Technique for Studying Spatial-Temporal Dynamics of Solutes in Sediment and Soil. *Earth-Sci. Rev.* **2019**, *197*, 102916.
- (216) Vieweg, M.; Trauth, N.; Fleckenstein, J. H.; Schmidt, C. Robust Optode-Based Method for Measuring in Situ Oxygen Profiles in Gravelly Streambeds. *Environ. Sci. Technol.* **2013**, *47* (17), 9858–9865.
- (217) Eickhorst, T.; Tippkötter, R. Improved Detection of Soil Microorganisms Using Fluorescence in Situ Hybridization (FISH) and Catalyzed Reporter Deposition (CARD-FISH). *Soil Biol. Biochem.* **2008**, *40* (7), 1883–1891.
- (218) Prietzel, J.; Thieme, J.; Salomé, M. Assessment of Sulfur and Iron Speciation in a Soil Aggregate by Combined S and Fe Micro-XANES: Microspatial Patterns and Relationships. *J. Synchrotron Radiat.* **2010**, *17* (2), 166–172.
- (219) O'Rourke, S. M.; Turner, J. N.; Holden, N. M. Estimating Fine Resolution Carbon Concentration in an Intact Soil Profile by X-Ray Fluorescence Scanning. In *Soil Carbon*; Hartemink, A. E., McSweeney, K., Eds.; Springer International Publishing: Cham, 2014; pp 179–187; DOI: 10.1007/978-3-319-04084-4_19.
- (220) Neumann, R. B.; Seyfferth, A. L.; Teshera-Levye, J.; Ellingson, J. Soil Warming Increases Arsenic Availability in the Rice Rhizosphere. *Agric. Environ. Lett.* **2017**, *2* (1), 170006.
- (221) Hansel, C. M.; Benner, S. G.; Neiss, J.; Dohnalkova, A.; Kukkadapu, R. K.; Fendorf, S. Secondary Mineralization Pathways Induced by Dissimilatory Iron Reduction of Ferrihydrite under Advective Flow. *Geochim. Cosmochim. Acta* **2003**, *67* (16), 2977–2992.
- (222) Kurt, Z.; Mack, E. E.; Spain, J. C. Biodegradation of Cis-Dichloroethene and Vinyl Chloride in the Capillary Fringe. *Environ. Sci. Technol.* **2014**, *48* (22), 13350–13357.
- (223) Kurt, Z.; Mack, E. E.; Spain, J. C. Natural Attenuation of Nonvolatile Contaminants in the Capillary Fringe. *Environ. Sci. Technol.* **2016**, *50* (18), 10172–10178.
- (224) Tufano, K. J.; Benner, S. G.; Mayer, K. U.; Marcus, M. A.; Nico, P. S.; Fendorf, S. Aggregate-Scale Heterogeneity in Iron (Hydr)Oxide Reductive Transformations. *Vadose Zone J.* **2009**, *8* (4), 1004–1012.
- (225) Ying, S. C.; Kocar, B. D.; Griffis, S. D.; Fendorf, S. Competitive Microbially and Mn Oxide Mediated Redox Processes

Controlling Arsenic Speciation and Partitioning. *Environ. Sci. Technol.* **2011**, *45* (13), 5572–5579.

(226) Ying, S. C.; Masue-Slowey, Y.; Kocar, B. D.; Griffis, S. D.; Webb, S.; Marcus, M. A.; Francis, C. A.; Fendorf, S. Distributed Microbially- and Chemically-Mediated Redox Processes Controlling Arsenic Dynamics within Mn-/Fe-Oxide Constructed Aggregates. *Geochim. Cosmochim. Acta* **2013**, *104*, 29–41.

(227) Mock, R. P.; Schaefer, M. V.; Pacheco, J. L.; Lake, L.; Lee, I.; Ying, S. C. Influence of Fe(II) on Arsenic(III) Oxidation by Birmessite in Diffusion-Limited Systems. *ACS Earth Space Chem.* **2019**, *3* (4), 550–561.

(228) Engel, M.; Boye, K.; Noël, V.; Babey, T.; Bargar, J. R.; Fendorf, S. Simulated Aquifer Heterogeneity Leads to Enhanced Attenuation and Multiple Retention Processes of Zinc. *Environ. Sci. Technol.* **2021**, *55* (5), 2939–2948.

(229) Wilmoth, J. L.; Doak, P. W.; Timm, A.; Halsted, M.; Anderson, J. D.; Ginovart, M.; Prats, C.; Portell, X.; Retterer, S. T.; Fuentes-Cabrera, M. A. Microfluidics and Agent-Based Modeling Framework for Investigating Spatial Organization in Bacterial Colonies: The Case of *Pseudomonas Aeruginosa* and H1-Type VI Secretion Interactions. *Front. Microbiol.* **2018**, *9*, 33.

(230) Aufrecht, J.; Khalid, M.; Walton, C. L.; Tate, K.; Cahill, J. F.; Retterer, S. T. Hotspots of Root-Exuded Amino Acids Are Created within a Rhizosphere-on-a-Chip. *Lab. Chip* **2022**, *22* (5), 954–963.

(231) Stanley, C. E.; Grossmann, G.; Casadevall i Solvas, X.; deMello, A. J. Soil-on-a-Chip: Microfluidic Platforms for Environmental Organismal Studies. *Lab. Chip* **2016**, *16* (2), 228–241.

(232) Davidson, E. A.; Samanta, S.; Caramori, S. S.; Savage, K. The Dual Arrhenius and Michaelis-Menten Kinetics Model for Decomposition of Soil Organic Matter at Hourly to Seasonal Time Scales. *Glob. Change Biol.* **2012**, *18* (1), 371–384.

(233) Davidson, E. A.; Savage, K. E.; Finzi, A. C. A Big-Microsite Framework for Soil Carbon Modeling. *Glob. Change Biol.* **2014**, *20*, 3610–3620.

(234) Zakem, E. J.; Polz, M. F.; Follows, M. J. Redox-Informed Models of Global Biogeochemical Cycles. *Nat. Commun.* **2020**, *11* (1), 5680.

(235) Ebrahimi, A.; Or, D. Hydration and Diffusion Processes Shape Microbial Community Organization and Function in Model Soil Aggregates. *Water Resour. Res.* **2015**, *51* (12), 9804–9827.

(236) Ebrahimi, A.; Or, D. On Upscaling of Soil Microbial Processes and Biogeochemical Fluxes From Aggregates to Landscapes. *J. Geophys. Res. Biogeosci.* **2018**, *123*, 1526–1547.

(237) Yan, Z.; Wang, T.; Wang, L.; Yang, X.; Smith, P.; Hilpert, M.; Li, S.; Shang, J.; Bailey, V.; Liu, C. Microscale Water Distribution and Its Effects on Organic Carbon Decomposition in Unsaturated Soils. *Sci. Total Environ.* **2018**, *644*, 1036–1043.

(238) Linn, D. M.; Doran, J. W. Effect of Water-Filled Pore Space on Carbon Dioxide and Nitrous Oxide Production in Tilled and Nontilled Soils. *Soil Sci. Soc. Am. J.* **1984**, *48*, 1267–1272.

(239) Patrick, W. H.; Gambrell, R. P.; Faulkner, S. P. Redox Measurements of Soils. In *Methods of Soil Analysis: Part 3 Chemical Methods*; Sparks, D. L., Page, A. L., Helmke, P. A., Loeppert, R. H., Soltanpour, P. N., Tabatabai, M. A., Johnston, C. T., Sumner, M. E., Eds.; Soil Science Society of America, American Society of Agronomy: Madison, WI, USA, 2018; pp 1255–1273; DOI: [10.2136/sssabookser5.3.c42](https://doi.org/10.2136/sssabookser5.3.c42).