

UC San Diego

UC San Diego Previously Published Works

Title

Liberating Chiral Lipid Mediators, Inflammatory Enzymes, and LIPID MAPS from Biological Grease

Permalink

<https://escholarship.org/uc/item/9dm057nn>

Journal

Journal of Biological Chemistry, 291(47)

ISSN

0021-9258

Author

Dennis, Edward A

Publication Date

2016-11-01

DOI

10.1074/jbc.x116.723791

Peer reviewed

Liberating Chiral Lipid Mediators, Inflammatory Enzymes, and LIPID MAPS from Biological Grease

Published, JBC Papers in Press, August 23, 2016, DOI 10.1074/jbc.X116.723791

Edward A. Dennis

From the Department of Chemistry and Biochemistry and Department of Pharmacology, School of Medicine, University of California at San Diego, La Jolla, California 92093-0601

Edited by Herbert Tabor

In 1970, it was well accepted that the central role of lipids was in energy storage and metabolism, and it was assumed that amphipathic lipids simply served a passive structural role as the backbone of biological membranes. As a result, the scientific community was focused on nucleic acids, proteins, and carbohydrates as information-containing molecules. It took considerable effort until scientists accepted that lipids also “encode” specific and unique biological information and play a central role in cell signaling. Along with this realization came the recognition that the enzymes that act on lipid substrates residing in or on membranes and micelles must also have important signaling roles, spurring curiosity into their potentially unique modes of action differing from those acting on water-soluble substrates. This led to the creation of the concept of “surface dilution kinetics” for describing the mechanism of enzymes acting on lipid substrates, as well as the demonstration that lipid enzymes such as phospholipase A₂ (PLA₂) contain allosteric activator sites for specific phospholipids as well as for membranes. As our understanding of phospholipases advanced, so did the understanding that many of the lipids released by these enzymes are chiral information-containing signaling molecules; for example, PLA₂ regulates the generation of precursors for the biosynthesis of eicosanoids and other bioactive lipid mediators of inflammation and resolution underlying disease progression. The creation of the LIPID MAPS initiative in 2003 and the ensuing development of the lipidomics field have revealed that lipid metabolites are central to human metabolism. Today lipids are recognized as key mediators of health and disease as we enter a new era of biomarkers and personalized medicine. This article is my personal “reflection” on these scientific advances.

Early on I did not know I was going to be a scientist. I was aware of science as a possible career as two of my maternal uncles were chemists. I also enjoyed my experiences in science; for example, my teachers in biology and chemistry at Senn High School in Chicago inspired me to successfully compete in the 1958 city science fair with a project on air pollution. However, I was equally inspired by my English teacher who encouraged me to consider organizational leadership directions. Thus, my intent when I matriculated at Yale was to obtain a liberal arts education. But in choosing a major I was influenced by William von Eggers Doering who first exposed me to the beauty of three-dimensional stereochemistry as a sophomore when I took organic chemistry for chemists—although I was not yet a chemistry major, I took the course and I was hooked. Doering’s class stimulated me to major in chemistry and to carry out



FIGURE 1. Frank H. Westheimer (right) in his 90s pictured with R. Stephen Berry (left) who first described the concept of pseudorotation in the trigonal bipyramid structure of PF_5 .

undergraduate research on the stereochemistry of the HCl elimination reaction with a great role model for me, Harry H. Wasserman.

My fascination with stereochemistry continued as I transitioned to graduate school in the Chemistry Department at Harvard University. During my second year, we were given a 1-week cumulative exam in which we were to make an “original” proposal. I thought about how carbon had four bonds and could form two isomers and wondered how many isomers would be possible for phosphorus, which could have five bonds. This led me to learn what inorganic chemists already knew well, namely that phosphorus bonds were arranged in a trigonal bipyramid or square pyramid with both equatorial and apical bonds. I then reasoned that in going from carbon to phosphorus, the number of isomers increases from 2 isomers to 30 isomers, so I proposed synthesizing a pentacoordinate phosphorus with different links so as to make “orbital isomers” of pentacoordinate phosphorus. I received a C (failing) grade with the professor’s comment that orbital isomers could not be made. However, my graduate research with the father of enzyme mechanisms and their stereochemistry, Frank H. Westheimer (1) (Fig. 1), was on the mechanism of phosphate ester hydrolysis. As my thesis work proceeded and I realized that the intermediate or transition state in this process was pentacoordinate phosphorus, I was able to show that orbital isomers could exist and the concept of pseudorotation was key to understanding the mechanism (2). Later, I independently applied pseudorotation to understanding the mechanism of ribonuclease (3).

My graduate work had kindled a strong interest in the enzymology of phosphate groups, so I stayed in Boston for a postdoctoral position in the Department of Biological Chemistry at Harvard Medical School on phospholipid biosynthesis (4) with Eugene P. Kennedy (5) (Fig. 2), who had discovered the metabolic pathways by which phospholipids are made. At that time, the prevailing view was that lipids played a central role in metabolism and especially in the storage and generation of energy; on the other hand, membranes were thought to be just a sea of unimportant lipids or “grease” in which membrane proteins carried out their supposedly important duties in “the viscous phospholipid bilayer solvent” (6). Thus, the scientific community was studying membrane structure but only by looking at the functional role of proteins in membranes. I had joined the lab with the intent of finding the enzyme that makes an unusual P–C bond in *Tetrahymena* phospholipids. As a precursor to this task, I started working with the more typical enzymes in the laboratory. As I began learning the lab’s protocols, I was stunned to realize that scientists in the field were using detergents without having any idea at the chemical level why these lipids were needed! My recognition of this may have been related to the fact that I was an unusual addition to the group because I came from a chemistry background, and so I always wanted to know what was happening at the molecular level. The discovery that lipids were necessary for enzymatic function planted the seed of what was a radical view at the time that these detergents are lipids and they have essential functions.

I started my independent scientific career in 1970 as a new Assistant Professor at the University of California at San Diego (UCSD) intent on pursuing mechanistic questions about these enzymes. I believed that the optimal way to start was with an enzyme that acted in or on the membrane, and particularly one with activity toward membrane phospholipids. This decision started a theme that would continue throughout my research career—from developing the early methodology needed to study membrane-active enzymes, to tracking down the destinies and functions of lipids in the corresponding metabolic pathways, to creating conferences and consortia to harness and build on our increasing technological and conceptual breakthroughs in lipid enzymology, to seeing the specific implications of these scientific advances on understanding and improving human health. It has been very exciting to have contributed to these advances! Throughout these adventures, I have been very fortunate to work with fabulous colleagues, both internal and external to my lab, and I cannot begin to thank each of them appropriately within



FIGURE 2. Symposium on Cellular Regulation and Membranes in honor of Eugene P. Kennedy on his 65th birthday organized by Edward Dennis on April 12–14, 1985 in Woods Hole, MA. A, Albert Lehninger; B, Fritz Lipmann; C, Edward Dennis; D, Marilyn Rumley; E, Adelaide Kennedy; F, Roberta Friedkin; G, Phyllis Elfman; H, Morris Friedkin; and I, Eugene Kennedy. Kennedy and Friedkin were Lehninger's first two graduate students, and Kennedy carried out postdoctoral studies with Lipmann. Rumley was a technician and Elfman was a secretary in the Kennedy laboratory. Both Lehninger and Lipmann passed away the following year.

the space allotted me here; I have done my best to cite their contributions in the reference section, which only hints as to their importance in my scientific travels.

New Methods and Mechanistic Concepts for Membrane-active Enzymes

My first NIH RO1 grant was to study the reconstitution of phosphatidylserine decarboxylase—a holdover from my time in the Kennedy lab—in phosphatidylserine-containing surfactant mixed micelles. This work was designed to confirm that proteins were actually interacting with phospholipids in a productive manner. If the enzyme was active, CO_2 would be released, which could be measured by a novel gas chromatography method (7). However, a serendipitous conversation with Dr. Beatriz M. Braganca, a scientist on sabbatical from India working in a neighboring lab, turned my attention to cobra venom as a commercially available and plentiful source of phospholipase A_2 (PLA_2) (8). This water-soluble enzyme hydrolyzes the fatty acid on the *sn*-2 position of membrane phospholipids to produce a lysophospholipid and a free fatty acid. It is more stable than phosphatidylserine decarboxylase, and its basic characteristics simplify several technical challenges. Finally, it also has many important biological functions including biosynthesis and generating lipid cell signals (9) (Fig. 3). We jumped at the opportunity to switch to this more advantageous model, and our studies on PLA_2 led to my second RO1 grant, now in its 40th year.

With PLA_2 under study, our first challenge was to understand why detergents were necessary to obtain enzy-

matic activity. This led us to pursue NMR techniques to characterize the interaction of phospholipids with surfactants, principally the non-ionic detergent Triton X-100 (10–12). We proposed that one role of the detergent was to solubilize the phospholipid into mixed micelles, which provided a surface to present the substrate to the enzyme (8). Another possible function of the Triton X-100 would be to solubilize a membrane-localized enzyme so that it could be present in aqueous solution and be available to the substrate; however, the fact that PLA_2 is already water-soluble simplified the kinetic problem enormously. This experimental system provided an ideal model for understanding protein-membrane interactions. But it was an uphill battle to convince those studying bilayer membranes that mixed micelles had any biological relevance! It was a battle worth fighting though, as surfactant micelles (and later bicelles) are now a mainstay in NMR studies of membrane proteins (13–16). Indeed, NMR work on mixed micelles of surfactants and phospholipids established the physical chemical principles that would allow the surfactants to be used to study a variety of phospholipids, including sphingolipids. Moreover, this established the generality of the mixed micelle system for studying protein-membrane interactions (17, 18).

A second critical step in our studies was formulating the concept of “surface dilution kinetics” (19, 20). This introduces two pre-catalytic steps into the reaction, first the enzyme binding to the membrane and then the enzyme separately finding and binding to its substrate in the two-dimensional interface (Fig. 4). This framework allowed us




FIGURE 3. Phospholipases A_2 function as degradative enzymes when they produce lysophospholipids and free fatty acids as products, as biosynthetic enzymes when the lysophospholipid product is reacylated with a PUFA to produce important phospholipids, and as signaling enzymes when the products are converted to agonists of GPCRs. Adapted from Ref. 9.




FIGURE 4. The concept of “surface dilution kinetics” requires two distinct steps for enzymes (E) acting on substrates (B) contained in lipid-water interfaces or surfaces (A) of membranes and micelles. Shown is an illustration of the application of “surface dilution kinetics” to the phospholipase A_2 - (blue) catalyzed hydrolysis of phospholipid substrate (red) contained in mixed micelles with nonionic surfactants such as Triton X-100 (yellow). The enzyme first associates with the mixed micelle surface (Bulk Step), and in a subsequent step (Surface Step), the enzyme associated with the micelle binds a phospholipid substrate molecule in its catalytic site and carries out hydrolysis, producing as products a lysophospholipid and a fatty acid, which may be released to solution or may be retained in the micelle surface. Reprinted from Ref. 21, adapted from Refs. 19 and 20.

to translate existing thinking about enzyme mechanism to the study of enzymes acting in/on membranes. I presented this concept at the first California Membrane Conference, March 13–17, 1972, held in Squaw Valley (the first meeting in what became the Keystone Symposia) and also reported our first micelle studies (10). Eventually, the surface dilution concept was generalized (21) to apply to many other enzymes that act on phospholipids (22, 23).

Using the mixed micelle system and the surface dilution kinetic approach, it was now possible to pursue the mechanism of action of PLA_2 (24). We first discovered that the enzyme was active only in the presence of certain phospholipids. This allowed application of the then underappreciated concept of allostery developed by Jean-Pierre Changeux (25), a concept I first learned about in the mid-1960s when Frank Westheimer returned from lecturing at

the Pasteur Institute in Paris. Specifically, we hypothesized that PLA_2 contained an “activator site,” although the enzyme functions as a monomer (a departure from Changeux’s original concept), and that when the water-soluble enzyme associated with the mixed micelle (membrane), it underwent a conformational change (at that time, represented simply as a square changing to a circle) to enable binding of the substrate phospholipid at the catalytic site. Our data showed that the whole surface of the enzyme interacted with the micelle (membrane), perhaps sensing as many as 30 phospholipid head groups at once, and that one of those lipids had to be a phosphocholine-containing lipid that acted at a “specific site” that was distinct from the catalytic site (26, 27). This allosteric activator site was later confirmed by kinetics (28) and X-ray crystallography (29, 30) and verified by mutational studies (31) and gene synthesis (32).

Detailed binding (33) and kinetic studies on PLA_2 (34–38) and the determination of the rate constants for acyl and phosphoryl migration in the lysophospholipid product (39) were critical for elucidating the kinetic analysis of PLA_2 , including its allosteric activation (40, 41) and varying activity on different membrane forms (42–44). The same approaches were also critical in my lab’s study of the related mammalian lysophospholipases (45–49) and indeed many other phospholipases (22, 50–58). Along the way, we’ve also explored PLA_2 ’s evolution from snake venom to human forms (59) and its reported inhibition by lipocortin (annexin 1), which we demonstrated occurred via “surface dilution”; thus it was not true inhibition as claimed in the literature (60–62). We also realized early on that having tools to probe PLA_2 function would be



FIGURE 5. Two key contributors to eicosanoid chemistry. *Top*, Edward Dennis and Bengt Samuelsson celebrating Samuelsson's 80th birthday at the first Lipid Mediators in Health and Disease Symposium held at the Nobel Forum, Stockholm, Sweden on August 27–29, 2014. *Bottom*, Edward Dennis introducing E. J. Corey as Keynote Speaker at the VI International Conference on Eicosanoids and Other Bioactive Lipids in Cancer, Inflammation, and Related Diseases held September 12–15, 1999 in Boston, MA.

useful, and so we focused attention on the discovery, design, and characterization of a large variety of chemical inhibitors of PLA₂ (63–76). Application of these inhibitors—in tandem with experiments probing enzyme aggregation (77)—not only provided insight into the catalytic mechanism, but also played a key role in advancing the understanding of PLA₂ and its regulation in biological systems (78).


A Turning Point for Lipid Signaling

Despite the great progress in understanding PLA₂ mechanisms we felt we were making, the biological implications for these studies remained obscure to the field at large. However, John Vane's discovery that aspirin and other non-steroidal anti-inflammatory drugs (NSAIDs) worked by acting on lipid biosynthesis (79) changed the landscape: suddenly, arachidonic acid metabolites—specifically prostaglandins—were important. Bengt Samuelsson's (Fig. 5, *top*) discovery of the first leukotrienes (80)

further cemented the relevance of lipid mediators. This was recognized with the awarding of the 1982 Nobel Prize in Physiology or Medicine to Sune Bergström, John Vane, and Bengt Samuelsson. The 1990 Nobel Prize in Chemistry was received by E. J. Corey (Fig. 5, *bottom*), who had previously synthesized many of the key chiral eicosanoids. Of course, we were excited to see that the new questions raised by their discoveries pointed back to our favorite enzyme, PLA₂. Specifically, free arachidonic acid is at very low levels in cells and is generated spontaneously as needed for prostaglandin/leukotriene biosynthesis; PLA₂ is responsible for the initiation of this arachidonic acid cascade (9, 78, 81) (Fig. 6).

These conceptual linkages led many investigators, myself included, to pursue the intracellular forms of PLA₂ responsible for arachidonic acid release. The first two "FASEB Summer Conferences on Phospholipase A₂," which I organized with Moseley Waite, included watershed moments. In 1988, two groups announced the sequence and cloning of the human secreted sPLA₂, and in 1992, two groups announced the sequencing and cloning of the human cytosolic cPLA₂. In 1995, at the third FASEB Summer Conference, our lab reported on the identification, complete purification, and characterization of the Ca²⁺-independent iPLA₂ (82). At that time, the explosion of PLA₂s initiated a discussion among the conference attendees that led to the Group numbering system for PLA₂s (83), which with several evolutions (84–86) is universally used today to distinguish the 16 groups and many subgroups of PLA₂ (9). In the meantime, another unique form of mammalian sPLA₂ secreted from macrophages was found, namely the Group V sPLA₂ (87). And on the functional side, iPLA₂ was sequenced and cloned (88, 89), and its remodeling function in macrophages (90–93) and unusual abundance in brain (94) were established.

In the 1980s, my laboratory turned its attention to the function of these enzymes in cellular metabolism and the generation of lipid cell signals—what I viewed as the most exciting new frontiers in cell biology and immunology. To kick start this new *in vivo* direction, I first needed to know more about these complex systems. Thus, in 1983, I received a Guggenheim Fellowship to spend a sabbatical with Manfred Karnovsky at Harvard Medical School, an expert on primary macrophage cells, and with Lawrence "Laurie" Levine at Brandeis University, the developer of radioimmune assays for eicosanoids. When I returned to UCSD, we decided to pursue macrophages as the archetypal mammalian cell—the "*E. coli* of mammalian cells"—in which to explore PLA₂ function and eicosanoid mediator production. In macrophage cell lines, the



Nature Reviews | Immunology

FIGURE 6. Central role of phospholipase A₂ in eicosanoid biosynthesis and receptor signaling. Reprinted from Ref. 81.

numerous types of PLA₂s that existed in a single cell type were characterized (95, 96) and the role of PLA₂ in activating the arachidonic acid cascade by a variety of agonists was explored (97, 98). This led to characterizing the eicosanoids produced downstream of PLA₂ and their autocrine and paracrine function in lipid signaling in the cellular milieu (99–101). This all became considerably more complex as we realized that the activation of eicosanoid synthesis could involve activation of multiple PLA₂s (102–106) stimulated by the products of other PLA₂s from distinct subcellular locations (107, 108).

Once the role of phospholipases in the field of signal transduction was established and the central role of the downstream eicosanoid mediators derived from arachidonic acid released by PLA₂ was shown, our lab's interests naturally shifted to the emerging broader role of lipid mediators in signal transduction. Fred Fox invited me to organize the first Keystone Symposium on Cellular Activation and Signal Initiation in 1988, and we brought together Tony Hunter, Robin Irvine, and Michael Berridge for this conference, which launched great interest in this field. Chairing this conference also reminded me of my early interests in organizational leadership, as I greatly

enjoyed organizing what I hoped would lead to a creative and engaging scientific meeting. It was no surprise that I had gravitated toward this kind of role in the past, and I made it a point to participate in similar opportunities in the future. Throughout the years, I have greatly valued these organizational leadership experiences, whether as president of the board of the Keystone Conferences, chair of the Gordon Research Conferences Board of Directors, chair of my department, or Editor-in-Chief of the Journal of Lipid Research. After the 1988 conference, I organized several more Keystone conferences on lipid signaling including an important one in 1994 on Lipid Second Messengers (109) and the 1989 Gordon Research Conference on Lipid Metabolism where this topical theme continued.

It was during this period that it became recognized that there are numerous lipid second messengers, many generated by phospholipases and lipid kinases/phosphatases, that play critical roles in cell signaling (51). It took a great deal of time and effort before the important roles of lipid cell signals were recognized by the broader biochemical community, which, at the time, was much more focused on signaling and regulation by phosphorylation and dephosphorylation of proteins and more traditional recep-

tors. One challenge the field faced was in communicating the newly discovered roles of lipids in cell signaling. As a result, Ralph Bradshaw and I made it a priority to highlight the central role of lipid cell signaling and integration of lipid signaling with more familiar pathways (110) as we organized and edited the three-volume *Handbook of Cell Signaling*, which appeared in 2003 (111).

Back in my laboratory, lipid cell signaling had always been focused on the metabolites of arachidonic acid—the eicosanoids and related oxidized polyunsaturated fatty acids (PUFA)—that activate a large variety of G-coupled protein receptors (GPCRs) (81). We had already learned through our work on cobra venom sPLA₂ that inhibitors could help us interrogate enzymes, but the inhibitors that worked on sPLA₂ didn't necessarily translate well to the mammalian systems, especially the intracellular PLA₂s that now occupied our attention. We thus expanded our initial inhibitor discovery work first to mammalian sPLA₂s (112), including the use of antisense technology (113), and then to other PLA₂ groups—especially the intracellular cPLA₂ and iPLA₂—by activated ketones (114, 115), fluorophosphonates (116), and other traditional inhibitors (98). In the last decade, this has expanded into a long collaboration with Professor George Kokotos from the University of Athens (9), in which we are designing and testing novel inhibitors including oxoamides (117–120), polyfluoroketones (121, 122), and thiazolyl ketones (123). Specifically, we've leaned on insights from molecular dynamics and simulations to inform rational drug design and medicinal chemistry efforts capped off with extensive biochemical characterization. Some of these inhibitors have shown specificity in important neurological disease models including experimental autoimmune encephalomyelitis, Wallerian degeneration, hyperalgesia, and spinal cord injury (124–127).

The Mechanism Comes Together

While these exciting changes and advances were occurring in the field at large and in my lab, we hadn't forgotten our goal to understand how these enzymes work at the molecular level. Research on enzymes that acted on soluble substrates had reached a high level of sophistication in their description of the catalytic mechanism, and I hoped that we might be able to do the same for our favorite enzyme! We therefore continued our detailed kinetic, mechanistic, crystallographic, NMR, and other structural studies on these enzymes. Many studies were carried out on iPLA₂, especially on its aggregation using radiation inactivation approaches (82, 128) and its additional lysophospholipase and acyl transferase activities (129) and

similar catalytic activities for cPLA₂ (130, 131). We were particularly intrigued by the discovery of the dramatic and specific activation of cPLA₂ by phosphatidylinositol 4,5-bisphosphate (PIP₂) (132–134) at a uniquely defined allosteric activator site as it both generalized our previous discovery regarding allosteric sites but also established a novel mechanism to accomplish the task. In the last decade, the lab has also made use of a new approach utilizing deuterium exchange mass spectrometry (DXMS) (135). This approach has traditionally been used to study protein-protein and protein-ligand interactions, but we applied it to look at the interaction of water-soluble proteins with membranes and demonstrated this approach with PLA₂ (136–138). We have now studied four of the major types of PLA₂, including the secreted (139), cytosolic (140, 141), and Ca²⁺-independent forms (142), as well as lipoprotein-associated/platelet-activating factor (PAF) acetyl hydrolase (143) and the interactions of the latter with lipoproteins such as HDL (144). These DXMS methods also led to a better understanding of the catalytic action of an unusual form with a Cys in the catalytic site (145).

We also launched a collaboration with my UCSD colleague J. Andrew McCammon, leading to the incorporation of molecular dynamics and simulations guided by DXMS into our consideration of how the enzymes work. This technique was used to create structural complexes of each enzyme with a single phospholipid substrate molecule in its catalytic site to visualize substrate extraction. Simulations of the enzyme-substrate-membrane system revealed important information about the mechanisms by which enzymes associate with the membrane (146) and then extract and bind their phospholipid substrate (147) (Fig. 7). This allowed us to extend early observations of allosteric sites on PLA₂s to include membranes as allosteric ligands. We also examined the effects of potent specific inhibitors of these PLA₂s such as pyrophenone and oxoamides for cPLA₂ (148) and various fluoroketones with iPLA₂ (149–151) using these techniques.

These combined results led us to a complete model of the catalytic cycle of two different human PLA₂s, cPLA₂ and iPLA₂ (147). After some forty years of studying PLA₂, a most fulfilling moment for me was when our lab developed detailed simulations (and representative movies) depicting the association of PLA₂ with a membrane during which the membrane acts as an allosteric ligand of the enzyme's interfacial surface by shifting its conformation from a closed (inactive) state in water to an open (active) state. This is followed by the extraction of a substrate




FIGURE 7. Cytosolic phospholipase A₂ extracting, binding, and hydrolyzing phospholipid substrate releasing arachidonic acid for eicosanoid biosynthesis (147). Cover for July, 2015 *Journal of Lipid Research* issue introducing a Thematic Reviews series on Phospholipases (51).

phospholipid from the membrane into the active site of the enzyme, and then achievement of the energetically optimal conformation for catalysis and finally diffusion of the products into the membrane surface (147). My quest to match the quality of information available for “normal” water-soluble enzyme systems, the initial steps of which were recognized 15 years earlier by my receipt of an award from the ASBMB in lipid enzymology (Fig. 8), had been attained.

Even more exciting, this work led to a general hypothesis that “membranes serve as allosteric regulators of enzymes that act in or on membranes” (135, 147) (Fig. 9).

Traditionally, allosteric regulators had been conceived of as small molecules binding to “topographically distinct sites” from the catalytic site, such as phosphatidylcholine for sPLA₂ and PIP₂ for cPLA₂, and generally on oligomeric proteins exhibiting sigmoid kinetics. Here, we have an enormous ligand (when compared with the size of a typical enzyme) binding to the enzyme but effecting a significant conformational change that is essential for catalysis. We anticipate that this new concept will have broad applicability to many systems including membrane proteins and provide a new paradigm for understanding lipid-protein interactions.



FIGURE 8. Walter Shaw, President of Avanti Polar Lipids, presenting the ASBMB Avanti Award in Lipids to Edward Dennis on June 5, 2000 in Boston MA.




FIGURE 9. Membranes are allosteric activators of phospholipases. Reprinted from Ref. 147.

Charting a Course for Lipid Research

Even at the beginning of the 21st century—at the dawn of genomics and with interest in proteomics growing—little was known about lipids themselves, such as how many existed, what signaling roles they had, how they should be measured, and even how lipids should be classified and named in the expanding world of omics. Given my interests in organizational leadership, I recognized an opportunity. In 2002, I proposed to NIH an initiative titled LIPID MAPS, standing for “LIPID Metabolites And Pathways Strategies,” which would develop the field of “lipidomics.” This “omic” can be considered part of metabolomics, which aims at the identification and quantification of all of the metabolites in the cell including its nucleic acids, amino acids, sugars, and fats. The ultimate goal of biochemists is to follow the fluxes of molecules in biological pathways. But by far the largest number of distinct molecular species in cellular metabolism lies in the lipids, where tens of thousands of distinct molecular species exist. Thus, the initiative would undertake the identification, characterization, and quantification of all lipid metabolites. With a \$25,000 “planning grant,” which was used to recruit the top academicians in each lipid type as

well as bioinformatics experts, the eventual group of 12 principal investigators came from eight universities and a commercial company (Fig. 10). The resulting ten-year Large Scale Collaborative “Glue” Grant awarded by the National Institute of General Medical Sciences was charged with developing the field of lipidomics, and it produced over 400 publications (152). At last lipidomics could join the omics evolution (153)!

The success of LIPID MAPS over the years is one of the great highlights of my career. Together, we developed novel liquid chromatographic-mass spectrometric-based lipidomics techniques termed “CLASS” (138) using a quantitative approach based on over 500 LIPID MAPS synthetic standards and specific agonists such as Kdo₂-Lipid A (154). This approach was applied to the overall omics analysis of some 500 lipid species in immunologically activated macrophages integrating transcriptomics, proteomics, and metabolomics of lipid metabolites (155). Novel methods for separating and assessing the purity of subcellular organelles (156) were developed and employed to determine the subcellular localization of many individual lipid molecular species and their changes during macrophage activation (157). A major finding from the integrated LIPID MAPS study of activated macrophages was the importance of desmosterol in linking lipid metabolism with inflammation (158). In another integrated study, a second sterol, 25-hydroxycholesterol, was shown to activate the stress response to reprogram transcription and translation (159).

We also profiled human plasma to quantify some 600 distinct lipid molecular species present across all mammalian lipid categories (160). We have high hopes for the potential implications of these data for clinical diagnostics and disease biomarkers as well as understanding the mechanisms of disease (161). However, these data also made a big impact on lipid research in another way. As we were sorting through all of the information, we realized that we didn’t have sufficient classification, nomenclature, or structural drawing representations for lipids to be able to describe our results properly. So, we developed some (162, 163)! We’ve also gone further—the initiative includes a website (www.lipidmaps.org) and databases where over 40,000 individual molecular species of lipids can be searched; tutorials on lipid metabolism using the structural drawing tool can be found on this site as well. This standardization may be the most impactful and lasting achievement of the investigators of the LIPID MAPS Consortium and their international collaborators, the International Lipid Classification and Nomenclature




FIGURE 10. **LIPID MAPS Consortium leaders at annual meeting December, 2008.** Left to right: Christian R. H. Raetz (Duke University), Joseph L. Witztum (UCSD), Nicholas Winograd (Penn State) who replaced Stephen White (University of California, Irvine), Alfred H. Merrill (Georgia Tech), Robert C. Murphy (University of Colorado), Michael S. VanNieuwenhze (Indiana University), Edward A. Dennis (UCSD), David W. Russell (UT Southwestern University), Walter A. Shaw (Avanti Polar Lipids), Christopher K. Glass (UCSD), Jean Chin (NIH representative), Shankar Subramaniam (UCSD), and H. Alex Brown (Vanderbilt University).

Committee (ILCNC), who continue to work together so productively.

Lipidomics of Fatty Acids and Eicosanoids

Beyond these collaborative projects, my own laboratory's work under the LIPID MAPS initiative has continued its focus on the lipidomics of fatty acyls including fatty acids, eicosanoids and related oxidized fatty acids, and fatty amides. We like to think of this focus as looking to the future of the field (Fig. 11). So far, this focus has led to a robust and comprehensive approach to the lipidomics analysis of hundreds of fatty acids including their numerous metabolites arising from an array of cyclooxygenases, lipoxygenases, cytochrome P450s, and non-enzymatic oxidation reactions (164), such as the production of dihomoprostaglandins in stimulated macrophages resulting from the elongation of arachidonic acid to adrenic acid (165). We also used lipidomics to analyze very long chain

fatty acids that are particularly abundant in the phospholipids of retinal membranes where they are critical for function (166) and demonstrated the essential role of ELOVL-4 in the synthesis of these fatty acids. Today over 200 fatty acyls can routinely be analyzed and quantified with internal standards (167) using the ultrahigh performance liquid chromatograph/mass spectrometric (UPLC/MS) techniques developed over the years for eicosanoids and related oxidized PUFAs (168) as well as free fatty acids (169) and complex lipids including phospholipids (170).

It is fascinating to see how this basic science has so many potential applications in nutrition and disease, and so we've continued to search for new biological areas relevant to lipids. One study demonstrated how a single immune cell can store a pro-resolving lipid precursor and then release it for bioactive maturation and secretion, conceptually similar to the production and inflammasome-dependent maturation of the pro-inflammatory IL-1 family of




FIGURE 11. Edward Dennis in his laboratory with mass spectrometer on the initiation of LIPID MAPS.

cytokines. Specifically, we examined the role of lipoxins as pro-resolution lipid mediators that inhibit phlogistic neutrophil recruitment and promote wound healing by macrophage recruitment via potent and specific signaling through the LXA₄ receptor. Lipoxins are generated as a consequence of sequential activation of the Toll-like receptor 4 (TLR4), a receptor for endotoxin, and P2X₇, a purinergic receptor for extracellular ATP (171). Initial activation of TLR4 results in accumulation of the COX-2-derived lipoxin precursor 15-hydroxyeicosatetraenoic acid (15-HETE) in esterified form within membrane phospholipids. 15-HETE production can be enhanced by aspirin. Subsequent activation of P2X₇ results in efficient release of 15-HETE from membrane phospholipids by cPLA₂ and the conversion of 15-HETE to bioactive lipoxins by 5-lipoxygenase (5-LOX). This study illustrated how aspirin in addition to inhibiting prostaglandin production also promotes pro-resolution eicosanoids. In its broader context, this result illustrates the importance of eicosanoids in the general response to infection, their relationship to inflammasome formation, and the consequent “eicosanoid storm” (81).

Once we had started examining the role of eicosanoids in inflammation, yet more questions surfaced. It was known that endotoxin-stimulated macrophages could serve as models for bacterial inflammation and infection, but the role of lipids in these processes was unclear. Using lipidomics analysis of eicosanoids, we were able to identify synergy in cellular lipid signaling of TLRs and purinergic receptors (172). This same approach was used to determine the course of infection-induced arthritic inflammation and its resolution in Lyme disease (173) and its broader implications for the cyclooxygenase pathway. We’ve also looked at lipids introduced from the environment. For example, we studied the effects of fish oil con-

sumption on Lyme disease progression (174). In this research, novel eicosanoid mediators were found, including anti-inflammatory mediators that differentiate the pathogenicity of influenza strains in mouse and in human lavages (175). Lipidomic analysis of cells supplemented with small amounts of the fish oil-derived ω -3 fatty acids, eicosapentaenoic acid (EPA) or docosahexaenoic acid (DHA), clarified their effects on the inflammatory eicosanoid in inhibiting COX-1 more than COX-2 and shunting arachidonate to LOX physiologically (176). Lipidomics analysis even allowed the elucidation of the role of liver X receptors (LXRs) in the biosynthesis of insulin-sensitizing ω -3 fatty acids (177).

As our understanding of biology has grown, our interests in applying those insights to questions with biomedical implications have grown in parallel. For example, inflammatory hyperalgesia and various forms of pain induce essential bioactive lipid production in the spinal cord and spinal fluid. Over the years with my UCSD colleague Tony Yaksh, we’ve characterized PLA₂ enzymes in spinal cord and spinal fluid (178, 179) as well as examining the role of resulting eicosanoids (180). This included the discovery of the novel role of spinal 12-lipoxygenase-derived hepoxilins A₃ and B₃, which activate the TRPV1 and TRPA1 receptors, in inflammatory hyperalgesia (181). It was shown that intrathecal administration of these hepoxilins induces hyperalgesia in rats. The specific 12-lipoxygenase enzyme eLOX3 responsible for the generation of hepoxilins has also recently been identified (182) along with the particular glial cell source in spinal cord. In another application, persistent effects after the resolution of inflammation in serum-transferred arthritis were analyzed (183). Even the consequences for cancer resulting from the up-regulation of the cyclooxygenase in certain cell types are better understood by the incorporation of lipidomics analysis of the oxygenated PUFAs (184, 185). These results demonstrate the utility of a comprehensive lipidomics approach for identifying potential contributors to disease pathology, and we hope that they will facilitate the development of more precisely targeted treatment strategies.

During these years, we have also been reminded about the importance of thinking beyond enzymatic reactions. Specifically, we investigated the oxidation of LDL lipids to explain the effect on macrophage activation and its implications for atherosclerosis. The first critical finding was that when human LDL is oxidized, significant amounts of the PUFA in the *sn*-2 position of 1-stearoyl, 2-arachidonoyl phosphatidylcholine become oxidized and yield 1-stearoyl, 2-oxovaleroyl phosphatidylcholine as a



FIGURE 12. Lipid Mediators in Health and Disease II: A Tribute to Edward Dennis and 7th International Conference on Phospholipase A₂ and Lipid Mediators held May 19–20, 2016 at Scripps Seaside Forum in La Jolla, California. Honorary Chair Bengt Samuelsson and Organizing Committee Chair Nicolas G. Bazan (LSU Health New Orleans) and his international committee of Jerold Chun (Scripps Research Institute), Jesper Z. Haeggström (Karolinska Institutet), Timothy Hla (Weill Cornell Medical College), Charles N. Serhan (Harvard Medical School), and Takao Shimizu (University of Tokyo) are pictured along with invited speakers, chairs, and participants.

major product. Furthermore, this product can easily undergo aldol condensations and also form Schiff bases with lysine side chains of apoproteins in the LDL (186). Such products are recognized by receptors and anti-phospholipid antibodies (187–189), and lipid complexes of phosphorylcholine were found to be pattern recognition ligands (190).

The LIPID MAPS lipidomics approaches enabled our discovery that oxidized PUFAs in LDL could also be found esterified to cholesterol (191), and lipidomics studies of zebrafish fed a high fat diet revealed oxidized phospholipids and cholesteryl esters (192). Subsequent lipidomics studies revealed that such oxidized phospholipids and cholesterol esters bind to apoproteins (193) and have multiple effects on macrophage signaling (194). Significantly, these oxidized species can be found in human coronary artery disease (195).

Finally, the power of lipidomics analysis has also been applied to fluxomics using unlabeled natural molecules to analyze the eicosanoid cascade. By taking advantage of the specificity of receptors to activate PLA₂, the release of free

arachidonic acid can be initiated (196). Fluxomics analysis allows the temporal course of metabolites in the eicosanoid cascade initiated by the activation of PLA₂ to be followed quantitatively. In recent experiments, we've compared primary macrophages including tissue-resident, thioglycolate-elicited, and bone marrow-derived, with cell lines (197), leading to a general model describing the fluxes of eicosanoids initiated by priming with a TLR4 agonist followed by activation with a purinergic P2X₇ agonist (198). Furthermore, this general approach has been integrated with transcriptional regulation and protein induction using a "directed proteomics" approach (199). The resultant competing eicosanoid fluxes of the multiple competing ω -3 and ω -6 fatty acid substrates under physiological conditions in the eicosanoid cascade can now be quantitatively compared (200).

Circling Back to Chirality

It is a delight to me that as we identify chiral eicosanoids, lipidomics studies have brought me back to my early fas-

cination with stereochemistry. Enzymatically produced eicosanoids always turn out to be chiral compounds as one might expect, but interestingly, certain drugs can alter the chirality of enzymatically produced eicosanoids. A recent example is the effect of aspirin described above, which acetylates the cyclooxygenase enzyme and changes the chirality of a COX side product lipoxin A₄ to epi-lipoxin A₄, implicated in the resolution of inflammation (81, 171). Another example is the identification of a particular oxidized lipid that corrects a defect in peroxisome proliferator-activated receptor- γ (PPAR- γ) signaling in *Cftr*-deficient mice (201). Chirality determination is also important because oxidized lipids can be produced non-enzymatically in response to stress and are associated with many diseases including non-alcoholic fatty liver disease (NAFLD) and non-alcoholic steatohepatitis (NASH), as characterized by LIPID MAPS (202). These products can be differentiated from the chirality that results from enzymatically produced products of cellular signaling pathways. The most recent work using quantitative UPLC/MS approaches resulted in the identity of a panel of eicosanoids in human plasma that can differentiate non-alcoholic fatty liver (NAFL) from NASH as a non-invasive biomarker important for NAFLD assessment and treatment (203).

When I was initially enticed by stereochemistry in college, I didn't expect it to occupy so much of my career. Indeed, my research trajectory has gone far beyond what I dreamed of then. I am humbled that the early connections that my colleagues and I described between PLA₂ action and lipid mediators have helped build this exciting field, and I am grateful to my scientific colleagues—including many former students and scientific collaborators—who organized a recent conference in my honor (Fig. 12). Today my laboratory is focused on enzyme- and drug-induced stereochemical changes of bioactive lipid molecules involved in inflammation and its resolution as well as on membrane-induced allosteric effects on enzymes that act on lipids, and these studies are now enhanced by lipidomic approaches to specificity. And although I can't claim to have predicted the future, I can say that I was prepared from the beginning to question established scientific thought. By the time I defended my Ph.D. thesis, William Doering, the professor who had so stimulated my fascination with the enormous stereo-complexity of simple molecules in three-dimensional space, had moved from Yale to Harvard and served on my doctoral thesis committee. During my defense, he asked why I had included my failed Cumulative Exam proposal on phosphorus isomers in my the-

sis as an Appendix. I explained jokingly that I was still challenging my grade!

Acknowledgments—I thank my early scientific mentors Harry H. Wasserman (Yale), Frank H. Westheimer (Harvard), Eugene P. Kennedy (Harvard Medical School), and as a new faculty member especially Nathan O. Kaplan and Morris Friedkin (UCSD) as well as George Carman (Rutgers) and numerous other colleagues and collaborators at UCSD and around the world. I'm especially grateful to those who participated in my LIPID MAPS initiative with whom I have worked very closely in recent years. Special thanks go to the many undergraduates, over 30 graduate students, over 50 postdoctoral scholars, several project scientists, many staff research associates, and numerous visiting scientists and sabbatical visitors (many of whom are cited in the references) who graced my laboratory. I especially appreciate the continuity and wise counsel provided by the long-time presence of Ray Deems (over three decades) and now Oswald Quehenberger (almost a decade) in my laboratory.

The NIH, NSF, and Lilly Research Laboratories and numerous other non-profit organizations and for-profit companies have provided the essential support for my laboratory. I want to explicitly express my appreciation to the National Institute of General Medical Sciences (NIGMS) for continuous individual predoctoral and postdoctoral fellowships, RO1 support for over 40 years (RO1 GM20501-40), and over 10 years of support for the LIPID MAPS "Glue" grant (US4 GM069338). I especially thank the *Journal of Biological Chemistry*, a journal in which over 70 publications on my work have appeared, for inviting me to write this Reflection.

Address correspondence to: edennis@ucsd.edu.

References

- Westheimer, F. H. (2003) Musings. *J. Biol. Chem.* **278**, 11729–11730
- Dennis, E. A., and Westheimer, F. H. (1966) The geometry of the transition state in the hydrolysis of phosphate esters. *J. Am. Chem. Soc.* **88**, 3432–3433
- Roberts, G. C., Dennis, E. A., Meadows, D. H., Cohen, J. S., and Jardetzky, O. (1969) The mechanism of action of ribonuclease. *Proc. Natl. Acad. Sci. U.S.A.* **62**, 1151–1158
- Dennis, E. A., and Kennedy, E. P. (1972) Intracellular sites of lipid synthesis and the biogenesis of mitochondria. *J. Lipid Res.* **13**, 263–267
- Kennedy, E. P. (2001) Hitler's gift and the era of biosynthesis. *J. Biol. Chem.* **276**, 42619–42631
- Singer, S. J., and Nicolson, G. L. (1972) The fluid mosaic model of the structure of cell membranes. *Science* **175**, 720–731
- Warner, T. G., and Dennis, E. A. (1973) New sensitive assay for phosphatidylserine decarboxylase based on the detection of CO₂ from nonradiolabeled phosphatidylserine. *J. Lipid Res.* **14**, 595–598
- Dennis, E. A. (1973) Kinetic dependence of phospholipase A₂ activity on the detergent Triton X-100. *J. Lipid Res.* **14**, 152–159
- Dennis, E. A., Cao, J., Hsu, Y. H., Magrioti, V., and Kokotos, G. (2011) Phospholipase A₂ enzymes: physical structure, biological function, disease implication, chemical inhibition, and therapeutic intervention. *Chem. Rev.* **111**, 6130–6185
- Dennis, E. A., and Owens, J. M. (1973) Studies on mixed micelles of Triton X-100 and phosphatidylcholine using nuclear magnetic resonance techniques. *J. Supramol. Struct.* **1**, 165–176
- Ribeiro, A. A., and Dennis, E. A. (1975) Proton magnetic resonance relaxation studies on the structure of mixed micelles of Triton X-100 and dimyristoyl-phosphatidylcholine. *Biochemistry* **14**, 3746–3755
- Robson, R. J., and Dennis, E. A. (1977) The size, shape, and hydration of nonionic surfactant micelles. Triton X-100. *J. Phys. Chem.* **81**, 1075–1078
- Roberts, M. F., and Dennis, E. A. (1977) Proton nuclear magnetic resonance demonstration of conformationally nonequivalent phospholipid fatty acid chains in mixed micelles. *J. Am. Chem. Soc.* **99**, 6142–6143
- Roberts, M. F., Bothner-By, A. A., and Dennis, E. A. (1978) Magnetic non-equivalence within fatty acyl chains of phospholipids in membrane models: ¹H nuclear magnetic resonance studies of the α -methylene groups. *Biochemistry* **17**, 935–942

15. De Bony, J., and Dennis, E. A. (1981) Magnetic nonequivalence of the two fatty acid chains in phospholipids of small unilamellar vesicles and mixed micelles. *Biochemistry* **20**, 5256–5260
16. Whiles, J. A., Deems, R., Vold, R. R., and Dennis, E. A. (2002) Bicelles in structure-function studies of membrane-associated proteins. *Bioorg. Chem.* **30**, 431–442
17. Robson, R. J., and Dennis, E. A. (1983) Micelles of nonionic detergents and mixed micelles with phospholipids. *Acc. Chem. Res.* **16**, 251–258
18. Lichtenberg, D., Robson, R. J., and Dennis, E. A. (1983) Solubilization of phospholipids by detergents: structural and kinetic aspects. *Biochim. Biophys. Acta* **737**, 285–304
19. Dennis, E. A. (1973) Phospholipase A₂ activity towards phosphatidylcholine in mixed micelles: surface dilution kinetics and the effect of thermotropic phase transitions. *Arch Biochem. Biophys.* **158**, 485–493
20. Deems, R. A., Eaton, B. R., and Dennis, E. A. (1975) Kinetic analysis of phospholipase A₂ activity toward mixed micelles and its implications for the study of lipolytic enzymes. *J. Biol. Chem.* **250**, 9013–9020
21. Carman, G. M., Deems, R. A., and Dennis, E. A. (1995) Lipid signaling enzymes and surface dilution kinetics. *J. Biol. Chem.* **270**, 18711–18714
22. Eaton, B. R., and Dennis, E. A. (1976) Analysis of phospholipase C (*Bacillus cereus*) action toward mixed micelles of phospholipid and surfactant. *Arch Biochem. Biophys.* **176**, 604–609
23. Warner, T. G., and Dennis, E. A. (1975) Action of the highly purified, membrane-bound enzyme phosphatidylserine decarboxylase *Escherichia coli* toward phosphatidylserine in mixed micelles and erythrocyte ghosts in the presence of surfactant. *J. Biol. Chem.* **250**, 8004–8009
24. Roberts, M. F., Deems, R. A., and Dennis, E. A. (1977) Dual role of interfacial phospholipid in phospholipase A₂ catalysis. *Proc. Natl. Acad. Sci. U.S.A.* **74**, 1950–1954
25. Changeux, J. P. (2013) The concept of allosteric interaction and its consequences for the chemistry of the brain. *J. Biol. Chem.* **288**, 26969–26986
26. Roberts, M. F., Adamich, M., Robson, R. J., and Dennis, E. A. (1979) Phospholipid activation of cobra venom phospholipase A₂: 1. Lipid-lipid or lipid-enzyme interaction. *Biochemistry* **18**, 3301–3308
27. Adamich, M., Roberts, M. F., and Dennis, E. A. (1979) Phospholipid activation of cobra venom phospholipase A₂: 2. Characterization of the phospholipid-enzyme interaction. *Biochemistry* **18**, 3308–3314
28. Boegeman, S. C., Deems, R. A., and Dennis, E. A. (2004) Phospholipid binding and the activation of Group IA secreted phospholipase A₂. *Biochemistry* **43**, 3907–3916
29. Fremont, D. H., Anderson, D. H., Wilson, I. A., Dennis, E. A., and Xuong, N. H. (1993) Crystal structure of phospholipase A₂ from Indian cobra reveals a trimeric association. *Proc. Natl. Acad. Sci. U.S.A.* **90**, 342–346
30. Segelke, B. W., Nguyen, D., Chee, R., Xuong, N. H., and Dennis, E. A. (1998) Structures of two novel crystal forms of *Naja naja naja* phospholipase A₂ lacking Ca²⁺ reveal trimeric packing. *J. Mol. Biol.* **279**, 223–232
31. Lefkowitz, L. J., Deems, R. A., and Dennis, E. A. (1999) Expression of Group IA phospholipase A₂ in *Pichia pastoris*: identification of a phosphatidylcholine activator site using site-directed mutagenesis. *Biochemistry* **38**, 14174–14184
32. Kelley, M. J., Crowl, R. M., and Dennis, E. A. (1992) Renaturation of cobra venom phospholipase A₂ expressed from a synthetic gene in *Escherichia coli*. *Biochim. Biophys. Acta* **1118**, 107–115
33. Roberts, M. F., Deems, R. A., and Dennis, E. A. (1977) Spectral perturbations of the histidine and tryptophan in cobra venom phospholipase A₂ upon metal ion and mixed micelle binding. *J. Biol. Chem.* **252**, 6011–6017
34. Plückthun, A., and Dennis, E. A. (1982) Role of monomeric activators in cobra venom phospholipase A₂ action. *Biochemistry* **21**, 1750–1756
35. Plückthun, A., and Dennis, E. A. (1985) Activation, aggregation, and product inhibition of cobra venom phospholipase A₂ and comparison with other phospholipases. *J. Biol. Chem.* **260**, 11099–11106
36. Plückthun, A., and Dennis, E. A. (1981) ³¹P nuclear magnetic resonance study on the incorporation of monomeric phospholipids into nonionic detergent micelles. *J. Phys. Chem.* **85**, 678–683
37. Plückthun, A., Rohlf, R., Davidson, F. F., and Dennis, E. A. (1985) Short-chain phosphatidylethanolamines: physical properties and susceptibility of the monomers to phospholipase A₂ action. *Biochemistry* **24**, 4201–4208
38. Lombardo, D., Fanni, T., Plückthun, A., and Dennis, E. A. (1986) Rate-determining step in phospholipase A₂ mechanism: ¹⁸O isotope exchange determined by ¹³C NMR. *J. Biol. Chem.* **261**, 11663–11666
39. Plückthun, A., and Dennis, E. A. (1982) Acyl and phosphoryl migration in lysophospholipids: importance in phospholipid synthesis and phospholipase specificity. *Biochemistry* **21**, 1743–1750
40. Hendrickson, H. S., and Dennis, E. A. (1984) Kinetic analysis of the dual phospholipid model for phospholipase A₂ action. *J. Biol. Chem.* **259**, 5734–5739
41. Hendrickson, H. S., and Dennis, E. A. (1984) Analysis of the kinetics of phospholipid activation of cobra venom phospholipase A₂. *J. Biol. Chem.* **259**, 5740–5744
42. Kensil, C. R., and Dennis, E. A. (1979) Action of cobra venom phospholipase A₂ on the gel and liquid crystalline states of dimyristoyl and dipalmitoyl phosphatidylcholine vesicles. *J. Biol. Chem.* **254**, 5843–5848
43. Adamich, M., and Dennis, E. A. (1978) Exploring the action and specificity of cobra venom phospholipase A₂ toward human erythrocytes, ghost membranes, and lipid mixtures. *J. Biol. Chem.* **253**, 5121–5125
44. Kensil, C. R., and Dennis, E. A. (1981) Alkaline hydrolysis of phospholipids in model membranes and the dependence on their state of aggregation. *Biochemistry* **20**, 6079–6085
45. Jarvis, A. A., Cain, C., and Dennis, E. A. (1984) Purification and characterization of a lysophospholipase from human amniotic membranes. *J. Biol. Chem.* **259**, 15188–15195
46. Zhang, Y. Y., and Dennis, E. A. (1988) Purification and characterization of a lysophospholipase from a macrophage-like cell line P388D₁. *J. Biol. Chem.* **263**, 9965–9972
47. Stafford, R. E., Fanni, T., and Dennis, E. A. (1989) Interfacial properties and critical micelle concentration of lysophospholipids. *Biochemistry* **28**, 5113–5120
48. Wang, A., Deems, R. A., and Dennis, E. A. (1997) Cloning, expression, and catalytic mechanism of murine lysophospholipase I. *J. Biol. Chem.* **272**, 12723–12729
49. Wang, A., Loo, R., Chen, Z., and Dennis, E. A. (1997) Regiospecificity and catalytic triad of lysophospholipase I. *J. Biol. Chem.* **272**, 22030–22036
50. Dennis, E. A. (ed) (1991) *Phospholipases, Methods in Enzymology*, Vol. 197, 640 pages, Academic Press, Orlando
51. Dennis, E. A. (2015) Introduction to Thematic Review Series: Phospholipases: central role in lipid signaling and disease. *J. Lipid Res.* **56**, 1245–1247
52. Roberts, M. F., Otnaess, A. B., Kensil, C. A., and Dennis, E. A. (1978) The specificity of phospholipase A₂ and phospholipase C in a mixed micellar system. *J. Biol. Chem.* **253**, 1252–1257
53. Balboa, M. A., Balsinde, J., Dennis, E. A., and Insel, P. A. (1995) A phospholipase D-mediated pathway for generating diacylglycerol in nuclei from Madin-Darby canine kidney cells. *J. Biol. Chem.* **270**, 11738–11740
54. Balboa, M. A., Balsinde, J., Dillon, D. A., Carman, G. M., and Dennis, E. A. (1999) Proinflammatory macrophage-activating properties of the novel phospholipid diacylglycerol pyrophosphate. *J. Biol. Chem.* **274**, 522–526
55. Johnson, C. A., Balboa, M. A., Balsinde, J., and Dennis, E. A. (1999) Regulation of cyclooxygenase-2 expression by phosphatidate phosphohydrolase in human amniotic WISH cells. *J. Biol. Chem.* **274**, 27689–27693
56. Grkovich, A., Johnson, C. A., Buczynski, M. W., and Dennis, E. A. (2006) Lipopolysaccharide-induced cyclooxygenase-2 expression in human U937 macrophages is phosphatidic acid phosphohydrolase-1-dependent. *J. Biol. Chem.* **281**, 32978–32987
57. Balsinde, J., and Dennis, E. A. (1996) Bromoenol lactone inhibits magnesium-dependent phosphatidate phosphohydrolase and blocks triacylglycerol biosynthesis in mouse P388D₁ macrophages. *J. Biol. Chem.* **271**, 31937–31941
58. Balboa, M. A., Balsinde, J., and Dennis, E. A. (1998) Involvement of phosphatidate phosphohydrolase in arachidonic acid mobilization in human amniotic WISH cells. *J. Biol. Chem.* **273**, 7684–7690
59. Davidson, F. F., and Dennis, E. A. (1990) Evolutionary relationships and implications for the regulation of phospholipase A₂ from snake venom to human secreted forms. *J. Mol. Evol.* **31**, 228–238
60. Davidson, F. F., Dennis, E. A., Powell, M., and Glenney, J. R., Jr. (1987) Inhibition of phospholipase A₂ by “lipocortins” and calpactins: an effect of binding to substrate phospholipids. *J. Biol. Chem.* **262**, 1698–1705
61. Davidson, F. F., and Dennis, E. A. (1989) Biological relevance of lipocortins and related proteins as inhibitors of phospholipase A₂. *Biochem. Pharmacol.* **38**, 3645–3651
62. Davidson, F. F., Lister, M. D., and Dennis, E. A. (1990) Binding and inhibition studies on lipocortins using phosphatidylcholine vesicles and phospholipase A₂ from snake venom, pancreas, and a macrophage-like cell line. *J. Biol. Chem.* **265**, 5602–5609
63. Yu, L., Deems, R. A., Hajdu, J., and Dennis, E. A. (1990) The interaction of phospholipase A₂ with phospholipid analogues and inhibitors. *J. Biol. Chem.* **265**, 2657–2664
64. Yu, L., and Dennis, E. A. (1991) Critical role of a hydrogen bond in the interaction of phospholipase A₂ with transition-state and substrate analogues. *Proc. Natl. Acad. Sci. U.S.A.* **88**, 9325–9329

65. Yu, L., and Dennis, E. A. (1992) Defining the dimensions of the catalytic site of phospholipase A_2 using amide substrate analogues. *J. Am. Chem. Soc.* **114**, 8757–8763
66. Plesniak, L. A., Boegeman, S. C., Segelke, B. W., and Dennis, E. A. (1993) Interaction of phospholipase A_2 with thioether amide containing phospholipid analogues. *Biochemistry* **32**, 5009–5016
67. Plesniak, L. A., Yu, L., and Dennis, E. A. (1995) Conformation of micellar phospholipid bound to the active site of phospholipase A_2 . *Biochemistry* **34**, 4943–4951
68. Lombardo, D., and Dennis, E. A. (1985) Cobra venom phospholipase A_2 inhibition by manoalide: a novel type of phospholipase inhibitor. *J. Biol. Chem.* **260**, 7234–7240
69. Reynolds, L. J., Morgan, B. P., Hite, G. A., Mihelich, E. D., and Dennis, E. A. (1988) Phospholipase A_2 inhibition and modification by manoalogue. *J. Am. Chem. Soc.* **110**, 5172–5177
70. Reynolds, L. J., Mihelich, E. D., and Dennis, E. A. (1991) Inhibition of venom phospholipases A_2 by manoalide and manoalogue: stoichiometry of incorporation. *J. Biol. Chem.* **266**, 16512–16517
71. Washburn, W. N., and Dennis, E. A. (1990) A novel general approach for the assay and inhibition of hydrolytic enzymes: utilizing suicide inhibitory bifunctionally linked substrates (SIBLINKS) exemplified by a phospholipase A_2 assay. *J. Am. Chem. Soc.* **112**, 2040–2041
72. Washburn, W. N., and Dennis, E. A. (1990) Suicide inhibitory bifunctionally linked substrates (SIBLINKS) as phospholipase A_2 inhibitors. *J. Am. Chem. Soc.* **112**, 2042–2043
73. Washburn, W. N., and Dennis, E. A. (1991) Suicide-inhibitory bifunctionally linked substrates (SIBLINKS) as phospholipase A_2 inhibitors: mechanistic implications. *J. Biol. Chem.* **266**, 5042–5048
74. Yu, L., and Dennis, E. A. (1993) Effect of polar head groups on the interactions of phospholipase A_2 with phosphonate transition-state analogues. *Biochemistry* **32**, 10185–10192
75. Barden, R. E., Darke, P. L., Deems, R. A., and Dennis, E. A. (1980) Interaction of phospholipase A_2 from cobra venom with Cibacron Blue F3GA. *Biochemistry* **19**, 1621–1625
76. Barlow, P. N., Lister, M. D., Sigler, P. B., and Dennis, E. A. (1988) Probing the role of substrate conformation in phospholipase A_2 action on aggregated phospholipids using constrained phosphatidylcholine analogues. *J. Biol. Chem.* **263**, 12954–12958
77. Hazlett, T. L., and Dennis, E. A. (1985) Aggregation studies on fluorescein-coupled cobra venom phospholipase A_2 . *Biochemistry* **24**, 6152–6158
78. Dennis, E. A. (1987) Regulation of eicosanoid production: role of phospholipases and inhibitors. *Nature Biotechnology* **5**, 1294–1300
79. Vane, J. R. (1971) Inhibition of prostaglandin synthesis as a mechanism of action for aspirin-like drugs. *Nat. New Biol.* **231**, 232–235
80. Samuelsson, B. (2012) Role of basic science in the development of new medicines: examples from the eicosanoid field. *J. Biol. Chem.* **287**, 10070–10080
81. Dennis, E. A., and Norris, P. C. (2015) Eicosanoid storm in infection and inflammation. *Nat. Rev. Immunol.* **15**, 511–523
82. Ackermann, E. J., Kempner, E. S., and Dennis, E. A. (1994) Ca^{2+} -independent cytosolic phospholipase A_2 from macrophage-like P388D₁ cells: isolation and characterization. *J. Biol. Chem.* **269**, 9227–9233
83. Dennis, E. A. (1994) Diversity of group types, regulation, and function of phospholipase A_2 . *J. Biol. Chem.* **269**, 13057–13060
84. Dennis, E. A. (1997) The growing phospholipase A_2 superfamily of signal transduction enzymes. *Trends Biochem. Sci.* **22**, 1–2
85. Six, D. A., and Dennis, E. A. (2000) The expanding superfamily of phospholipase A_2 enzymes: classification and characterization. *Biochim. Biophys. Acta* **1488**, 1–19
86. Schaloske, R. H., and Dennis, E. A. (2006) The phospholipase A_2 superfamily and its group numbering system. *Biochim. Biophys. Acta* **1761**, 1246–1259
87. Balboa, M. A., Balsinde, J., Winstead, M. V., Tischfield, J. A., and Dennis, E. A. (1996) Novel Group V phospholipase A_2 involved in arachidonic acid mobilization in murine P388D₁ macrophages. *J. Biol. Chem.* **271**, 32381–32384
88. Balboa, M. A., Balsinde, J., Jones, S. S., and Dennis, E. A. (1997) Identity between the Ca^{2+} -independent phospholipase A_2 enzymes from P388D₁ macrophages and Chinese hamster ovary cells. *J. Biol. Chem.* **272**, 8576–8580
89. Tang, J., Kriz, R. W., Wolfman, N., Shaffer, M., Sehra, J., and Jones, S. S. (1997) A novel cytosolic calcium-independent phospholipase A_2 contains eight ankyrin motifs. *J. Biol. Chem.* **272**, 8567–8575
90. Balsinde, J., Barbour, S. E., Bianco, I. D., and Dennis, E. A. (1994) Arachidonic acid mobilization in P388D₁ macrophages is controlled by two distinct Ca^{2+} -dependent phospholipase A_2 enzymes. *Proc. Natl. Acad. Sci. U.S.A.* **91**, 11060–11064
91. Balsinde, J., Bianco, I. D., Ackermann, E. J., Conde-Frieboes, K., and Dennis, E. A. (1995) Inhibition of calcium-independent phospholipase A_2 prevents arachidonic acid incorporation and phospholipid remodeling in P388D₁ macrophages. *Proc. Natl. Acad. Sci. U.S.A.* **92**, 8527–8531
92. Balsinde, J., and Dennis, E. A. (1997) Function and inhibition of intracellular calcium-independent phospholipase A_2 . *J. Biol. Chem.* **272**, 16069–16072
93. Balsinde, J., Balboa, M. A., and Dennis, E. A. (1997) Antisense inhibition of Group VI Ca^{2+} -independent phospholipase A_2 blocks phospholipid fatty acid remodeling in murine P388D₁ macrophages. *J. Biol. Chem.* **272**, 29317–29321
94. Yang, H. C., Mosior, M., Ni, B., and Dennis, E. A. (1999) Regional distribution, ontogeny, purification, and characterization of the Ca^{2+} -independent phospholipase A_2 from rat brain. *J. Neurochem.* **73**, 1278–1287
95. Ross, M. I., Deems, R. A., Jesaitis, A. J., Dennis, E. A., and Ulevitch, R. J. (1985) Phospholipase activities of the P388D₁ macrophage-like cell line. *Arch. Biochem. Biophys.* **238**, 247–258
96. Balsinde, J., and Dennis, E. A. (1996) Distinct roles in signal transduction for each of the phospholipase A_2 enzymes present in P388D₁ macrophages. *J. Biol. Chem.* **271**, 6758–6765
97. Lister, M. D., Deems, R. A., Watanabe, Y., Ulevitch, R. J., and Dennis, E. A. (1988) Kinetic analysis of the Ca^{2+} -dependent, membrane-bound, macrophage phospholipase A_2 and the effects of arachidonic acid. *J. Biol. Chem.* **263**, 7506–7513
98. Lister, M. D., Glaser, K. B., Ulevitch, R. J., and Dennis, E. A. (1989) Inhibition studies on the membrane-associated phospholipase A_2 *in vitro* and prostaglandin E_2 production *in vivo* of the macrophage-like P388D₁ cell: effects of manoalide, 7,7-dimethyl-5,8-eicosadienoic acid, and *p*-bromophenacyl bromide. *J. Biol. Chem.* **264**, 8520–8528
99. Glaser, K. B., Asmis, R., and Dennis, E. A. (1990) Bacterial lipopolysaccharide priming of P388D₁ macrophage-like cells for enhanced arachidonic acid metabolism: platelet-activating factor receptor activation and regulation of phospholipase A_2 . *J. Biol. Chem.* **265**, 8658–8664
100. Asmis, R., Randriamampita, C., Tsien, R. Y., and Dennis, E. A. (1994) Intracellular Ca^{2+} , inositol 1,4,5-trisphosphate and additional signalling in the stimulation by platelet-activating factor of prostaglandin E_2 formation in P388D₁ macrophage-like cells. *Biochem. J.* **298**, 543–551
101. Balboa, M. A., Balsinde, J., Johnson, C. A., and Dennis, E. A. (1999) Regulation of arachidonic acid mobilization in lipopolysaccharide-activated P388D₁ macrophages by adenosine triphosphate. *J. Biol. Chem.* **274**, 36764–36768
102. Balsinde, J., Balboa, M. A., and Dennis, E. A. (1998) Functional coupling between secretory phospholipase A_2 and cyclooxygenase-2 and its regulation by cytosolic Group IV phospholipase A_2 . *Proc. Natl. Acad. Sci. U.S.A.* **95**, 7951–7956
103. Balsinde, J., Balboa, M. A., and Dennis, E. A. (2000) Identification of a third pathway for arachidonic acid mobilization and prostaglandin production in activated P388D₁ macrophage-like cells. *J. Biol. Chem.* **275**, 22544–22549
104. Shinohara, H., Balboa, M. A., Johnson, C. A., Balsinde, J., and Dennis, E. A. (1999) Regulation of delayed prostaglandin production in activated P388D₁ macrophages by Group IV cytosolic and Group V secretory phospholipase A_2 s. *J. Biol. Chem.* **274**, 12263–12268
105. Balsinde, J., Shinohara, H., Lefkowitz, L. J., Johnson, C. A., Balboa, M. A., and Dennis, E. A. (1999) Group V phospholipase A_2 -dependent induction of cyclooxygenase-2 in macrophages. *J. Biol. Chem.* **274**, 25967–25970
106. Kessen, U. A., Schaloske, R. H., Stephens, D. L., Killermann Lucas, K., and Dennis, E. A. (2005) PGE₂ release is independent of upregulation of Group V phospholipase A_2 during long-term stimulation of P388D₁ cells with LPS. *J. Lipid Res.* **46**, 2488–2496
107. Shirai, Y., Balsinde, J., and Dennis, E. A. (2005) Localization and functional interrelationships among cytosolic Group IV, secreted Group V, and Ca^{2+} -independent Group VI phospholipase A_2 s in P388D₁ macrophages using GFP/RFP constructs. *Biochim. Biophys. Acta* **1735**, 119–129
108. Balboa, M. A., Shirai, Y., Gaietta, G., Ellisman, M. H., Balsinde, J., and Dennis, E. A. (2003) Localization of Group V phospholipase A_2 in caveolin-enriched granules in activated P388D₁ macrophage-like cells. *J. Biol. Chem.* **278**, 48059–48065
109. Liscovitch, M., and Cantley, L. C. (1994) Lipid second messengers. *Cell* **77**, 329–334
110. Bradshaw, R. A., and Dennis, E. A. (2003) Cell signaling: yesterday, today and tomorrow. in *HandBook of Cell Signaling* (Dennis, E. A., and Bradshaw, R. A. eds), Vol. 1, pp. 1–3, Academic Press, an imprint of Elsevier Science, Orlando, FL
111. Bradshaw, R. A., and Dennis, E. A. (2003) *Handbook of Cell Signaling*, Vol 1, 747 pages; Vol 2, 899 pages; and Vol 3, 709 pages, Academic Press, an imprint of Elsevier Science, Orlando, FL

112. Balsinde, J., Balboa, M. A., Insel, P. A., and Dennis, E. A. (1999) Regulation and inhibition of phospholipase A₂. *Annu. Rev. Pharmacol. Toxicol.* **39**, 175–189
113. Barbour, S. E., and Dennis, E. A. (1993) Antisense inhibition of Group II phospholipase A₂ expression blocks the production of prostaglandin E₂ by P388D₁ cells. *J. Biol. Chem.* **268**, 21875–21882
114. Ackermann, E. J., Conde-Frieboes, K., and Dennis, E. A. (1995) Inhibition of macrophage Ca²⁺-independent phospholipase A₂ by bromoenol lactone and trifluoromethyl ketones. *J. Biol. Chem.* **270**, 445–450
115. Conde-Frieboes, K., Reynolds, L. J., Lio, Y., Hale, M., Wasserman, H. H., and Dennis, E. A. (1996) Activated ketones as inhibitors of intracellular Ca²⁺-dependent and Ca²⁺-independent phospholipase A₂. *J. Am. Chem. Soc.* **118**, 5519–5525
116. Lio, Y. C., Reynolds, L. J., Balsinde, J., and Dennis, E. A. (1996) Irreversible inhibition of Ca²⁺-independent phospholipase A₂ by methyl arachidonoyl fluorophosphonate. *Biochim. Biophys. Acta* **1302**, 55–60
117. Kokotos, G., Kotsivolou, S., Six, D. A., Constantinou-Kokotou, V., Beltzner, C. C., and Dennis, E. A. (2002) Novel 2-oxoamide inhibitors of human Group IVA phospholipase A₂. *J. Med. Chem.* **45**, 2891–2893
118. Kokotos, G., Six, D. A., Loukas, V., Smith, T., Constantinou-Kokotou, V., Hadjipavlou-Litina, D., Kotsivolou, S., Chiou, A., Beltzner, C. C., and Dennis, E. A. (2004) Inhibition of Group IVA cytosolic phospholipase A₂ by novel 2-oxoamides *in vitro*, in cells, and *in vivo*. *J. Med. Chem.* **47**, 3615–3628
119. Stephens, D., Barbayianni, E., Constantinou-Kokotou, V., Peristeraki, A., Six, D. A., Cooper, J., Harkewicz, R., Deems, R. A., Dennis, E. A., and Kokotos, G. (2006) Differential inhibition of Group IVA and Group VIA phospholipases A₂ by 2-oxoamides. *J. Med. Chem.* **49**, 2821–2828
120. Six, D. A., Barbayianni, E., Loukas, V., Constantinou-Kokotou, V., Hadjipavlou-Litina, D., Stephens, D., Wong, A. C., Magrioti, V., Moutevelis-Minakakis, P., Baker, S. F., Dennis, E. A., and Kokotos, G. (2007) Structure-activity relationship of 2-oxoamide inhibition of Group IVA cytosolic phospholipase A₂ and Group V secreted phospholipase A₂. *J. Med. Chem.* **50**, 4222–4235
121. Baskakis, C., Magrioti, V., Cotton, N., Stephens, D., Constantinou-Kokotou, V., Dennis, E. A., and Kokotos, G. (2008) Synthesis of polyfluoro ketones for selective inhibition of human phospholipase A₂ enzymes. *J. Med. Chem.* **51**, 8027–8037
122. Kokotos, G., Hsu, Y. H., Burke, J. E., Baskakis, C., Kokotos, C. G., Magrioti, V., and Dennis, E. A. (2010) Potent and selective fluoroketone inhibitors of Group VIA calcium-independent phospholipase A₂. *J. Med. Chem.* **53**, 3602–3610
123. Kokotos, G., Feuerherm, A. J., Barbayianni, E., Shah, I., Sæther, M., Magrioti, V., Nguyen, T., Constantinou-Kokotou, V., Dennis, E. A., and Johansen, B. (2014) Inhibition of Group IVA cytosolic phospholipase A₂ by thiazolyl ketones *in vitro*, *ex vivo*, and *in vivo*. *J. Med. Chem.* **57**, 7523–7535
124. Yaksh, T. L., Kokotos, G., Svensson, C. I., Stephens, D., Kokotos, C. G., Fitzsimmons, B., Hadjipavlou-Litina, D., Hua, X. Y., and Dennis, E. A. (2006) Systemic and intrathecal effects of a novel series of phospholipase A₂ inhibitors on hyperalgesia and spinal prostaglandin E₂ release. *J. Pharmacol. Exp. Ther.* **316**, 466–475
125. López-Vales, R., Navarro, X., Shimizu, T., Baskakis, C., Kokotos, G., Constantinou-Kokotou, V., Stephens, D., Dennis, E. A., and David, S. (2008) Intracellular phospholipase A₂ Group IVA and Group VIA play important roles in Wallerian degeneration and axon regeneration after peripheral nerve injury. *Brain* **131**, 2620–2631
126. Kalyvas, A., Baskakis, C., Magrioti, V., Constantinou-Kokotou, V., Stephens, D., López-Vales, R., Lu, J. Q., Yong, V. W., Dennis, E. A., Kokotos, G., and David, S. (2009) Differing roles for members of the phospholipase A₂ superfamily in experimental autoimmune encephalomyelitis. *Brain* **132**, 1221–1235
127. López-Vales, R., Ghasemlou, N., Redensek, A., Kerr, B. J., Barbayianni, E., Antonopoulou, G., Baskakis, C., Rathore, K. I., Constantinou-Kokotou, V., Stephens, D., Shimizu, T., Dennis, E. A., Kokotos, G., and David, S. (2011) Phospholipase A₂ superfamily members play divergent roles after spinal cord injury. *FASEB J.* **25**, 4240–4252
128. Reynolds, L. J., Kempner, E. S., Hughes, L. L., and Dennis, E. A. (1995) Inactivation of secretory phospholipase A₂ by ionizing radiation. *Biophys. J.* **68**, 2108–2114
129. Lio, Y. C., and Dennis, E. A. (1998) Interfacial activation, lysophospholipase and transacylase activity of Group VI Ca²⁺-independent phospholipase A₂. *Biochim. Biophys. Acta* **1392**, 320–332
130. Pickard, R. T., Chiou, X. G., Striffler, B. A., DeFelippis, M. R., Hyslop, P. A., Tebbe, A. L., Yee, Y. K., Reynolds, L. J., Dennis, E. A., Kramer, R. M., and Sharp, J. D. (1996) Identification of essential residues for the catalytic function of 85-kDa cytosolic phospholipase A₂: probing the role of histidine, aspartic acid, cysteine, and arginine. *J. Biol. Chem.* **271**, 19225–19231
131. Loo, R. W., Conde-Frieboes, K., Reynolds, L. J., and Dennis, E. A. (1997) Activation, inhibition, and regioselectivity of the lysophospholipase activity of the 85-kDa Group IV cytosolic phospholipase A₂. *J. Biol. Chem.* **272**, 19214–19219
132. Mosior, M., Six, D. A., and Dennis, E. A. (1998) Group IV cytosolic phospholipase A₂ binds with high affinity and specificity to phosphatidylinositol 4,5-bisphosphate resulting in dramatic increases in activity. *J. Biol. Chem.* **273**, 2184–2191
133. Balsinde, J., Balboa, M. A., Li, W. H., Llopis, J., and Dennis, E. A. (2000) Cellular regulation of cytosolic Group IV phospholipase A₂ by phosphatidylinositol bisphosphate levels. *J. Immunol.* **164**, 5398–5402
134. Six, D. A., and Dennis, E. A. (2003) Essential Ca²⁺-independent role of the Group IVA cytosolic phospholipase A₂ C₂ domain for interfacial activity. *J. Biol. Chem.* **278**, 23842–23850
135. Cao, J., Burke, J. E., and Dennis, E. A. (2013) Using hydrogen/deuterium exchange mass spectrometry to define the specific interactions of the phospholipase A₂ superfamily with lipid substrates, inhibitors, and membranes. *J. Biol. Chem.* **288**, 1806–1813
136. Burke, J. E., and Dennis, E. A. (2009) Phospholipase A₂ structure/function, mechanism, and signaling. *J. Lipid Res.* **50**, (suppl.) S237–S242
137. Burke, J. E., and Dennis, E. A. (2009) Phospholipase A₂ biochemistry. *Cardiovasc. Drugs Ther.* **23**, 49–59
138. Harkewicz, R., and Dennis, E. A. (2011) Applications of mass spectrometry to lipids and membranes. *Annu. Rev. Biochem.* **80**, 301–325
139. Burke, J. E., Karbarz, M. J., Deems, R. A., Li, S., Woods, V. L., Jr., and Dennis, E. A. (2008) Interaction of Group IA phospholipase A₂ with metal ions and phospholipid vesicles probed with deuterium exchange mass spectrometry. *Biochemistry* **47**, 6451–6459
140. Hsu, Y. H., Burke, J. E., Stephens, D. L., Deems, R. A., Li, S., Asmus, K. M., Woods, V. L., Jr., and Dennis, E. A. (2008) Calcium binding rigidifies the C₂ domain and the intradomain interaction of GIVA phospholipase A₂ as revealed by hydrogen/deuterium exchange mass spectrometry. *J. Biol. Chem.* **283**, 9820–9827
141. Burke, J. E., Hsu, Y. H., Deems, R. A., Li, S., Woods, V. L., Jr., and Dennis, E. A. (2008) A phospholipid substrate molecule residing in the membrane surface mediates opening of the lid region in Group IVA cytosolic phospholipase A₂. *J. Biol. Chem.* **283**, 31227–31236
142. Hsu, Y. H., Burke, J. E., Li, S., Woods, V. L., Jr., and Dennis, E. A. (2009) Localizing the membrane binding region of Group VIA Ca²⁺-independent phospholipase A₂ using peptide amide hydrogen/deuterium exchange mass spectrometry. *J. Biol. Chem.* **284**, 23652–23661
143. Cao, J., Hsu, Y. H., Li, S., Woods, V. L., and Dennis, E. A. (2011) Lipoprotein-associated phospholipase A₂ interacts with phospholipid vesicles via a surface-disposed hydrophobic α -helix. *Biochemistry* **50**, 5314–5321
144. Cao, J., Hsu, Y. H., Li, S., Woods, V. L., Jr., and Dennis, E. A. (2013) Structural basis of specific interactions of Lp-PLA₂ with HDL revealed by hydrogen/deuterium exchange mass spectrometry. *J. Lipid Res.* **54**, 127–133
145. Pang, X. Y., Cao, J., Addington, L., Lovell, S., Battaile, K. P., Zhang, N., Rao, J. L., Dennis, E. A., and Moise, A. R. (2012) Structure/function relationships of adipose phospholipase A₂ containing a Cys-His-His catalytic triad. *J. Biol. Chem.* **287**, 35260–35274
146. Bucher, D., Hsu, Y. H., Mouchlis, V. D., Dennis, E. A., and McCammon, J. A. (2013) Insertion of the Ca²⁺-independent phospholipase A₂ into a phospholipid bilayer via coarse-grained and atomistic molecular dynamics simulations. *PLoS Comput. Biol.* **9**, e1003156
147. Mouchlis, V. D., Bucher, D., McCammon, J. A., and Dennis, E. A. (2015) Membranes serve as allosteric activators of phospholipase A₂, enabling it to extract, bind, and hydrolyze phospholipid substrates. *Proc. Natl. Acad. Sci. U.S.A.* **112**, E516–E525
148. Burke, J. E., Babakhani, A., Gorfie, A. A., Kokotos, G., Li, S., Woods, V. L., Jr., McCammon, J. A., and Dennis, E. A. (2009) Location of inhibitors bound to Group IVA phospholipase A₂ determined by molecular dynamics and deuterium exchange mass spectrometry. *J. Am. Chem. Soc.* **131**, 8083–8091
149. Hsu, Y. H., Bucher, D., Cao, J., Li, S., Yang, S. W., Kokotos, G., Woods, V. L., Jr., McCammon, J. A., and Dennis, E. A. (2013) Fluoroketone inhibition of Ca²⁺-independent phospholipase A₂ through binding pocket association defined by hydrogen/deuterium exchange and molecular dynamics. *J. Am. Chem. Soc.* **135**, 1330–1337
150. Mouchlis, V. D., Limnios, D., Kokotou, M. G., Barbayianni, E., Kokotos, G., McCammon, J. A., and Dennis, E. A. (2016) Development of potent and selective inhibitors for Group VIA calcium-independent phospholipase A₂

- guided by molecular dynamics and structure-activity relationships. *J. Med. Chem.* **59**, 4403–4414
151. Mouchlis, V. D., Morisseau, C., Hammock, B. D., Li, S., McCammon, J. A., and Dennis, E. A. (2016) Computer-aided drug design guided by hydrogen/deuterium exchange mass spectrometry: a powerful combination for the development of potent and selective inhibitors of Group VIA calcium-independent phospholipase A₂. *Bioorg. Med. Chem.* **24**, 4801–4811
 152. Dove, A. (2015) Greasing the wheels of lipidomics. *Science* **347**, 788–790
 153. Dennis, E. A. (2009) Lipidomics joins the omics evolution. *Proc. Natl. Acad. Sci. U.S.A.* **106**, 2089–2090
 154. Raetz, C. R., Garrett, T. A., Reynolds, C. M., Shaw, W. A., Moore, J. D., Smith, D. C., Jr., Ribeiro, A. A., Murphy, R. C., Ulevitch, R. J., Fearn, C., Reichart, D., Glass, C. K., Benner, C., Subramaniam, S., Harkewicz, R., Bowers-Gentry, R. C., Buczynski, M. W., Cooper, J. A., Deems, R. A., and Dennis, E. A. (2006) Kdo₂-Lipid A of *Escherichia coli*, a defined endotoxin that activates macrophages via TLR-4. *J. Lipid Res.* **47**, 1097–1111
 155. Dennis, E. A., Deems, R. A., Harkewicz, R., Quehenberger, O., Brown, H. A., Milne, S. B., Myers, D. S., Glass, C. K., Hardiman, G., Reichart, D., Merrill, A. H., Jr., Sullards, M. C., Wang, E., Murphy, R. C., Raetz, C. R., Garrett, T. A., Guan, Z., Ryan, A. C., Russell, D. W., McDonald, J. G., Thompson, B. M., Shaw, W. A., Sud, M., Zhao, Y., Gupta, S., Maurya, M. R., Fahy, E., and Subramaniam, S. (2010) A mouse macrophage lipidome. *J. Biol. Chem.* **285**, 39976–39985
 156. Andreyev, A. Y., Shen, Z., Guan, Z., Ryan, A., Fahy, E., Subramaniam, S., Raetz, C. R., Briggs, S., and Dennis, E. A. (2010) Application of proteomic marker ensembles to subcellular organelle identification. *Mol. Cell. Proteomics* **9**, 388–402
 157. Andreyev, A. Y., Fahy, E., Guan, Z., Kelly, S., Li, X., McDonald, J. G., Milne, S., Myers, D., Park, H., Ryan, A., Thompson, B. M., Wang, E., Zhao, Y., Brown, H. A., Merrill, A. H., Raetz, C. R., Russell, D. W., Subramaniam, S., and Dennis, E. A. (2010) Subcellular organelle lipidomics in TLR-4-activated macrophages. *J. Lipid Res.* **51**, 2785–2797
 158. Spann, N. J., Garmire, L. X., McDonald, J. G., Myers, D. S., Milne, S. B., Shibata, N., Reichart, D., Fox, J. N., Shaked, I., Heudobler, D., Raetz, C. R., Wang, E. W., Kelly, S. L., Sullards, M. C., Murphy, R. C., Merrill, A. H., Jr., Brown, H. A., Dennis, E. A., Li, A. C., Ley, K., Tsimikas, S., Fahy, E., Subramaniam, S., Quehenberger, O., Russell, D. W., and Glass, C. K. (2012) Regulated accumulation of desmosterol integrates macrophage lipid metabolism and inflammatory responses. *Cell* **151**, 138–152
 159. Shibata, N., Carlin, A. F., Spann, N. J., Saijo, K., Morello, C. S., McDonald, J. G., Romanoski, C. E., Maurya, M. R., Kaikkonen, M. U., Lam, M. T., Crotti, A., Reichart, D., Fox, J. N., Quehenberger, O., Raetz, C. R., Sullards, M. C., Murphy, R. C., Merrill, A. H., Jr., Brown, H. A., Dennis, E. A., Fahy, E., Subramaniam, S., Cavener, D. R., Spector, D. H., Russell, D. W., and Glass, C. K. (2013) 25-Hydroxycholesterol activates the integrated stress response to reprogram transcription and translation in macrophages. *J. Biol. Chem.* **288**, 35812–35823
 160. Quehenberger, O., Armando, A. M., Brown, A. H., Milne, S. B., Myers, D. S., Merrill, A. H., Bandyopadhyay, S., Jones, K. N., Kelly, S., Shaner, R. L., Sullards, C. M., Wang, E., Murphy, R. C., Barkley, R. M., Leiker, T. J., Raetz, C. R., Guan, Z., Laird, G. M., Six, D. A., Russell, D. W., McDonald, J. G., Subramaniam, S., Fahy, E., and Dennis, E. A. (2010) Lipidomics reveals a remarkable diversity of lipids in human plasma. *J. Lipid Res.* **51**, 3299–3305
 161. Quehenberger, O., and Dennis, E. A. (2011) The human plasma lipidome. *N. Engl. J. Med.* **365**, 1812–1823
 162. Fahy, E., Subramaniam, S., Brown, H. A., Glass, C. K., Merrill, A. H., Jr., Murphy, R. C., Raetz, C. R., Russell, D. W., Seyama, Y., Shaw, W., Shimizu, T., Spener, F., van Meer, G., VanNieuwenhze, M. S., White, S. H., Witztum, J. L., and Dennis, E. A. (2005) A comprehensive classification system for lipids. *J. Lipid Res.* **46**, 839–861
 163. Fahy, E., Subramaniam, S., Murphy, R. C., Nishijima, M., Raetz, C. R., Shimizu, T., Spener, F., van Meer, G., Wakelam, M. J., and Dennis, E. A. (2009) Update of the LIPID MAPS comprehensive classification system for lipids. *J. Lipid Res.* **50**, (suppl.) S9–S14
 164. Buczynski, M. W., Dumlao, D. S., and Dennis, E. A. (2009) Thematic Review Series: Proteomics: an integrated omics analysis of eicosanoid biology. *J. Lipid Res.* **50**, 1015–1038
 165. Harkewicz, R., Fahy, E., Andreyev, A., and Dennis, E. A. (2007) Arachidonate-derived dihomoprostaglandin production observed in endotoxin-stimulated macrophage-like cells. *J. Biol. Chem.* **282**, 2899–2910
 166. Harkewicz, R., Du, H., Tong, Z., Alkuraya, H., Bedell, M., Sun, W., Wang, X., Hsu, Y. H., Esteve-Rudd, J., Hughes, G., Su, Z., Zhang, M., Lopes, V. S., Molday, R. S., Williams, D. S., Dennis, E. A., and Zhang, K. (2012) Essential role of ELOVL4 protein in very long chain fatty acid synthesis and retinal function. *J. Biol. Chem.* **287**, 11469–11480
 167. Wang, Y., Armando, A. M., Quehenberger, O., Yan, C., and Dennis, E. A. (2014) Comprehensive ultra-performance liquid chromatographic separation and mass spectrometric analysis of eicosanoid metabolites in human samples. *J. Chromatogr. A* **1359**, 60–69
 168. Dumlao, D. S., Buczynski, M. W., Norris, P. C., Harkewicz, R., and Dennis, E. A. (2011) High-throughput lipidomic analysis of fatty acid derived eicosanoids and *N*-acylethanolamines. *Biochim. Biophys. Acta* **1811**, 724–736
 169. Quehenberger, O., Armando, A. M., and Dennis, E. A. (2011) High sensitivity quantitative lipidomics analysis of fatty acids in biological samples by gas chromatography-mass spectrometry. *Biochim. Biophys. Acta* **1811**, 648–656
 170. Baker, P. R., Armando, A. M., Campbell, J. L., Quehenberger, O., and Dennis, E. A. (2014) Three-dimensional enhanced lipidomics analysis combining UPLC, differential ion mobility spectrometry, and mass spectrometric separation strategies. *J. Lipid Res.* **55**, 2432–2442
 171. Norris, P. C., Gosselin, D., Reichart, D., Glass, C. K., and Dennis, E. A. (2014) Phospholipase A₂ regulates eicosanoid class switching during inflammasome activation. *Proc. Natl. Acad. Sci. U.S.A.* **111**, 12746–12751
 172. Buczynski, M. W., Stephens, D. L., Bowers-Gentry, R. C., Grkovich, A., Deems, R. A., and Dennis, E. A. (2007) TLR-4 and sustained calcium agonists synergistically produce eicosanoids independent of protein synthesis in RAW264.7 cells. *J. Biol. Chem.* **282**, 22834–22847
 173. Blaho, V. A., Buczynski, M. W., Brown, C. R., and Dennis, E. A. (2009) Lipidomic analysis of dynamic eicosanoid responses during the induction and resolution of Lyme arthritis. *J. Biol. Chem.* **284**, 21599–21612
 174. Dumlao, D. S., Cunningham, A. M., Wax, L. E., Norris, P. C., Hanks, J. H., Halpin, R., Lett, K. M., Blaho, V. A., Mitchell, W. J., Fritsche, K. L., Dennis, E. A., and Brown, C. R. (2012) Dietary fish oil substitution alters the eicosanoid profile in ankle joints of mice during Lyme infection. *J. Nutr.* **142**, 1582–1589
 175. Tam, V. C., Quehenberger, O., Oshansky, C. M., Suen, R., Armando, A. M., Treuting, P. M., Thomas, P. G., Dennis, E. A., and Aderem, A. (2013) Lipidomic profiling of influenza infection identifies mediators that induce and resolve inflammation. *Cell* **154**, 213–227
 176. Norris, P. C., and Dennis, E. A. (2012) Omega-3 fatty acids cause dramatic changes in TLR4 and purinergic eicosanoid signaling. *Proc. Natl. Acad. Sci. U.S.A.* **109**, 8517–8522
 177. Li, P., Spann, N. J., Kaikkonen, M. U., Lu, M., Oh, D. Y., Fox, J. N., Bandyopadhyay, G., Talukdar, S., Xu, J., Lagakos, W. S., Patsouris, D., Armando, A., Quehenberger, O., Dennis, E. A., Watkins, S. M., Auwerx, J., Glass, C. K., and Olefsky, J. M. (2013) NCoR repression of LXR α restricts macrophage biosynthesis of insulin-sensitizing omega 3 fatty acids. *Cell* **155**, 200–214
 178. Svensson, C. I., Lucas, K. K., Hua, X. Y., Powell, H. C., Dennis, E. A., and Yaksh, T. L. (2005) Spinal phospholipase A₂ in inflammatory hyperalgesia: role of the small, secretory phospholipase A₂. *Neuroscience* **133**, 543–553
 179. Lucas, K. K., Svensson, C. I., Hua, X. Y., Yaksh, T. L., and Dennis, E. A. (2005) Spinal phospholipase A₂ in inflammatory hyperalgesia: role of Group IVA cPLA₂. *Br. J. Pharmacol.* **144**, 940–952
 180. Buczynski, M. W., Svensson, C. I., Dumlao, D. S., Fitzsimmons, B. L., Shim, J. H., Scherbart, T. J., Jacobsen, F. E., Hua, X. Y., Yaksh, T. L., and Dennis, E. A. (2010) Inflammatory hyperalgesia induces essential bioactive lipid production in the spinal cord. *J. Neurochem.* **114**, 981–993
 181. Gregus, A. M., Doolen, S., Dumlao, D. S., Buczynski, M. W., Takasusuki, T., Fitzsimmons, B. L., Hua, X. Y., Taylor, B. K., Dennis, E. A., and Yaksh, T. L. (2012) Spinal 12-lipoxygenase-derived hepxilin A₃ contributes to inflammatory hyperalgesia via activation of TRPV1 and TRPA1 receptors. *Proc. Natl. Acad. Sci. U.S.A.* **109**, 6721–6726
 182. Gregus, A. M., Dumlao, D. S., Wei, S. C., Norris, P. C., Catella, L. C., Meyerstein, F. G., Buczynski, M. W., Steinauer, J. J., Fitzsimmons, B. L., Yaksh, T. L., and Dennis, E. A. (2013) Systematic analysis of rat 12/15-lipoxygenase enzymes reveals critical role for spinal eLOX3 hepxilin synthase activity in inflammatory hyperalgesia. *FASEB J.* **27**, 1939–1949
 183. Christianson, C. A., Dumlao, D. S., Stokes, J. A., Dennis, E. A., Svensson, C. I., Corr, M., and Yaksh, T. L. (2011) Spinal TLR4 mediates the transition to a persistent mechanical hypersensitivity after the resolution of inflammation in serum-transferred arthritis. *Pain* **152**, 2881–2891
 184. Jiao, J., Mikulec, C., Ishikawa, T. O., Magyar, C., Dumlao, D. S., Dennis, E. A., Fischer, S. M., and Herschman, H. (2014) Cell-type-specific roles for COX-2 in UVB-induced skin cancer. *Carcinogenesis* **35**, 1310–1319
 185. Jiao, J., Ishikawa, T. O., Dumlao, D. S., Norris, P. C., Magyar, C. E., Mikulec, C., Catapang, A., Dennis, E. A., Fischer, S. M., and Herschman, H. R. (2014) Targeted deletion and lipidomic analysis identify epithelial cell COX-2 as a major driver of chemically induced skin cancer. *Mol. Cancer Res.* **12**, 1677–1688

186. Friedman, P., Horkko, S., Steinberg, D., Witztum, J. L., and Dennis, E. A. (2002) Correlation of antiphospholipid antibody recognition with the structure of synthetic oxidized phospholipids. Importance of Schiff base formation and aldol condensation. *J. Biol. Chem.* **277**, 7010–7020
187. Hörkkö, S., Bird, D. A., Miller, E., Itabe, H., Leitinger, N., Subbanagounder, G., Berliner, J. A., Friedman, P., Dennis, E. A., Curtiss, L. K., Palinski, W., and Witztum, J. L. (1999) Monoclonal autoantibodies specific for oxidized phospholipids or oxidized phospholipid-protein adducts inhibit macrophage uptake of oxidized low-density lipoproteins. *J. Clin. Invest.* **103**, 117–128
188. Bird, D. A., Gillotte, K. L., Hörkkö, S., Friedman, P., Dennis, E. A., Witztum, J. L., and Steinberg, D. (1999) Receptors for oxidized low-density lipoprotein on elicited mouse peritoneal macrophages can recognize both the modified lipid moieties and the modified protein moieties: implications with respect to macrophage recognition of apoptotic cells. *Proc. Natl. Acad. Sci. U.S.A.* **96**, 6347–6352
189. Chang, M. K., Bergmark, C., Laurila, A., Hörkkö, S., Han, K. H., Friedman, P., Dennis, E. A., and Witztum, J. L. (1999) Monoclonal antibodies against oxidized low-density lipoprotein bind to apoptotic cells and inhibit their phagocytosis by elicited macrophages: evidence that oxidation-specific epitopes mediate macrophage recognition. *Proc. Natl. Acad. Sci. U.S.A.* **96**, 6353–6358
190. Boullier, A., Friedman, P., Harkewicz, R., Hartvigsen, K., Green, S. R., Almazan, F., Dennis, E. A., Steinberg, D., Witztum, J. L., and Quehenberger, O. (2005) Phosphocholine as a pattern recognition ligand for CD36. *J. Lipid Res.* **46**, 969–976
191. Harkewicz, R., Hartvigsen, K., Almazan, F., Dennis, E. A., Witztum, J. L., and Miller, Y. I. (2008) Cholesteryl ester hydroperoxides are biologically active components of minimally oxidized low density lipoprotein. *J. Biol. Chem.* **283**, 10241–10251
192. Fang, L., Harkewicz, R., Hartvigsen, K., Wiesner, P., Choi, S. H., Almazan, F., Pattison, J., Deer, E., Sayaphupha, T., Dennis, E. A., Witztum, J. L., Tsimikas, S., and Miller, Y. I. (2010) Oxidized cholesteryl esters and phospholipids in zebrafish larvae fed a high cholesterol diet: macrophage binding and activation. *J. Biol. Chem.* **285**, 32343–32351
193. Leibundgut, G., Scipione, C., Yin, H., Schneider, M., Boffa, M. B., Green, S., Yang, X., Dennis, E., Witztum, J. L., Koschinsky, M. L., and Tsimikas, S. (2013) Determinants of binding of oxidized phospholipids on apolipoprotein (a) and lipoprotein (a). *J. Lipid Res.* **54**, 2815–2830
194. Choi, S. H., Yin, H., Ravandi, A., Armando, A., Dumlaio, D., Kim, J., Almazan, F., Taylor, A. M., McNamara, C. A., Tsimikas, S., Dennis, E. A., Witztum, J. L., and Miller, Y. I. (2013) Polyoxxygenated cholesterol ester hydroperoxide activates TLR4 and SYK dependent signaling in macrophages. *PLoS ONE* **8**, e83145
195. Ravandi, A., Leibundgut, G., Hung, M. Y., Patel, M., Hutchins, P. M., Murphy, R. C., Prasad, A., Mahmud, E., Miller, Y. I., Dennis, E. A., Witztum, J. L., and Tsimikas, S. (2014) Release and capture of bioactive oxidized phospholipids and oxidized cholesteryl esters during percutaneous coronary and peripheral arterial interventions in humans. *J. Am. Coll. Cardiol.* **63**, 1961–1971
196. Gupta, S., Maurya, M. R., Stephens, D. L., Dennis, E. A., and Subramaniam, S. (2009) An integrated model of eicosanoid metabolism and signaling based on lipidomics flux analysis. *Biophys. J.* **96**, 4542–4551
197. Norris, P. C., Reichart, D., Dumlaio, D. S., Glass, C. K., and Dennis, E. A. (2011) Specificity of eicosanoid production depends on the TLR-4-stimulated macrophage phenotype. *J. Leukoc. Biol.* **90**, 563–574
198. Kihara, Y., Gupta, S., Maurya, M. R., Armando, A., Shah, I., Quehenberger, O., Glass, C. K., Dennis, E. A., and Subramaniam, S. (2014) Modeling of eicosanoid fluxes reveals functional coupling between cyclooxygenases and terminal synthases. *Biophys. J.* **106**, 966–975
199. Sabidó, E., Quehenberger, O., Shen, Q., Chang, C. Y., Shah, I., Armando, A. M., Andreyev, A., Vitek, O., Dennis, E. A., and Aebersold, R. (2012) Targeted proteomics of the eicosanoid biosynthetic pathway completes an integrated genomics-proteomics-metabolomics picture of cellular metabolism. *Mol. Cell. Proteomics* **11**, M111.014746
200. Gupta, S., Kihara, Y., Maurya, M. R., Norris, P. C., Dennis, E. A., and Subramaniam, S. (2016) Computational modeling of competitive metabolism between ω -3- and ω -6-polyunsaturated fatty acids in inflammatory macrophages. *J. Phys. Chem. B* **120**, 8346–8353
201. Harmon, G. S., Dumlaio, D. S., Ng, D. T., Barrett, K. E., Dennis, E. A., Dong, H., and Glass, C. K. (2010) Pharmacological correction of a defect in PPAR- γ signaling ameliorates disease severity in *Cftr*-deficient mice. *Nat. Med.* **16**, 313–318
202. Gorden, D. L., Myers, D. S., Ivanova, P. T., Fahy, E., Maurya, M. R., Gupta, S., Min, J., Spann, N. J., McDonald, J. G., Kelly, S. L., Duan, J., Sullards, M. C., Leiker, T. J., Barkley, R. M., Quehenberger, O., Armando, A. M., Milne, S. B., Mathews, T. P., Armstrong, M. D., Li, C., Melvin, W. V., Clements, R. H., Washington, M. K., Mendonsa, A. M., Witztum, J. L., Guan, Z., Glass, C. K., Murphy, R. C., Dennis, E. A., Merrill, A. H., Jr., Russell, D. W., Subramaniam, S., and Brown, H. A. (2015) Biomarkers of NAFLD progression: a lipidomics approach to an epidemic. *J. Lipid Res.* **56**, 722–736
203. Loomba, R., Quehenberger, O., Armando, A., and Dennis, E. A. (2015) Polyunsaturated fatty acid metabolites as novel lipidomic biomarkers for noninvasive diagnosis of nonalcoholic steatohepatitis. *J. Lipid Res.* **56**, 185–192