

## UC Santa Barbara

### UC Santa Barbara Previously Published Works

**Title**

Selenocysteine Substitution in a Class I Ribonucleotide Reductase

**Permalink**

<https://escholarship.org/uc/item/9g07w4zh>

**Journal**

Biochemistry, 58(50)

**ISSN**

0006-2960

**Authors**

Greene, Brandon L  
Stubbe, JoAnne  
Nocera, Daniel G

**Publication Date**

2019-12-17

**DOI**

10.1021/acs.biochem.9b00973


Peer reviewed

## Selenocysteine Substitution in a Class I Ribonucleotide Reductase

Brandon L. Greene,<sup>\*,†</sup> JoAnne Stubbe,<sup>‡,§</sup> and Daniel G. Nocera<sup>||</sup><sup>†</sup>Department of Chemistry and Biochemistry, University of California, Santa Barbara, Santa Barbara, California 93106, United States<sup>‡</sup>Department of Chemistry and <sup>§</sup>Department of Biology, Massachusetts Institute of Technology, Cambridge, Massachusetts 20139, United States<sup>||</sup>Department of Chemistry and Chemical Biology, Harvard University, Cambridge, Massachusetts 02138, United States

## Supporting Information

**ABSTRACT:** Ribonucleotide reductases (RNRs) employ a complex radical-based mechanism during nucleotide reduction involving multiple active site cysteines that both activate the substrate and reduce it. Using an engineered allo-tRNA, we substituted two active site cysteines with distinct function in the class Ia RNR of *Escherichia coli* for selenocysteine (U) via amber codon suppression, with efficiency and selectivity enabling biochemical and biophysical studies. Examination of the interactions of the C<sub>439</sub>U  $\alpha_2$  mutant protein with nucleotide substrates and the cognate  $\beta_2$  subunit demonstrates that the endogenous Y<sub>122</sub><sup>•</sup> of  $\beta_2$  is reduced under turnover conditions, presumably through radical transfer to form a transient U<sub>439</sub><sup>•</sup> species. This putative U<sub>439</sub><sup>•</sup> species is formed in a kinetically competent fashion but is incapable of initiating nucleotide reduction via 3'-H abstraction. An analogous C<sub>225</sub>U  $\alpha_2$  protein is also capable of radical transfer from Y<sub>122</sub><sup>•</sup>, but the radical-based substrate chemistry partitions between turnover and stalled reduction akin to the reactivity of mechanism-based inhibitors of RNR. The results collectively demonstrate the essential role of cysteine redox chemistry in the class I RNRs and establish a new tool for investigating thiol radical reactivity in biology.



Redox active amino acids endow enzymes with intrinsic cofactors and play critical roles in a plethora of enzymatic reactions.<sup>1</sup> Among the redox active amino acids, cysteine (C) is unique in that the thiol side chain can supply an electron (and a proton) individually, forming a thiyl radical, or in tandem with a second sterically accessible cysteine, forming a disulfide bond. Both thiol–thiyl radical and thiol–disulfide redox reactions involve proton-coupled electron transfer (PCET) and are exploited extensively in enzyme catalysis. Contrary to the thiol/disulfide couple, the role of thiyl radicals in enzymology remains poorly understood, yet thiyl radicals continue to be invoked in a broad range of enzymatic transformations. Identifying and defining the role of thiyl radicals is challenging for several distinct reasons. First, the one-electron reduction potential ( $E^0$ ) of the cysteine thiyl radical is the highest of the physiologically relevant redox active amino acid radicals, which increase in the following order: selenocysteine (U)<sup>2</sup> < tyrosine (Y) < tryptophan (W) < glycine (G) < cysteine (C) (pH 7).<sup>1</sup> Second, thiyl radicals are quite reactive toward elements of all proteins, including C=O and C–H bonds.<sup>3</sup> Third, thiyl radicals are challenging to detect by conventional biophysical spectroscopic techniques, including ultraviolet–visible (UV–vis) absorption due to low extinction coefficients,<sup>4</sup> and paramagnetic techniques due to broadening.<sup>5</sup> Lastly, the controlled generation of thiyl radicals requires site-specific delivery of a potent oxidant, often through endothermic radical transfer (RT) from another protein, cofactor, or substrate-based radical. These properties of the


protein-based thiyl radical present significant barriers to the study of their function in biology.

Both thiol–thiyl radical and thiol–disulfide redox reactions figure prominently in the function of ribonucleotide reductase (RNR), which catalyzes the reduction of nucleotides (di- or triphosphates) to deoxynucleotides, a committed step in *de novo* DNA biosynthesis and repair (Figure 1A).<sup>6,7</sup> Early identification of conserved and essential cysteines of the enzyme,<sup>8,9</sup> structural homology of the active site (Figure 1B),<sup>10</sup> and reactivity studies<sup>11–13</sup> suggest that a radical-based mechanism is employed by all RNRs involving a conserved cysteine on the “top face” that forms a thiyl radical and activates the substrate toward reduction by abstracting the 3'-H of the nucleotide. Two additional cysteines, or a cysteine, methionine, and formate, serve as radical substrate reductants located in the “bottom face”. The mechanism of cysteine oxidation on the top face has formed the basis of class differentiation. Class I RNRs utilize a second subunit harboring a redox active (metallo)cofactor for thiyl radical generation; class II RNRs utilize adenosylcob(II)alamin (AdoCbl), and class III RNRs utilize a radical-SAM activating enzyme that produces a glycy radical.

Received: October 30, 2019

Revised: November 25, 2019

Published: November 27, 2019



**Figure 1.** RNR mechanism of nucleotide reduction and active site structural homology. (A) General reaction catalyzed by RNR in all organisms. N is a nucleoside base. (B) Structural alignment of class Ia [*E. coli*, green; Protein Data Bank (PDB) entry 1RLR], class II (*L. leichmannii*, orange; PDB entry 1L1L), and class III (Enterobacteria phage T4, light blue; PDB entry 1B8B) RNRs aligned based on the “top” and “bottom-face” cysteines. Top-face cysteines are shown on the top of the image, and bottom-face cysteines (disulfide for both class I and II and cysteine and methionine for class III) on the bottom. Some of the class III RNRs contain a methionine in place of one of the cysteines, and formate serves as the ultimate electron donor. Cysteine/methionine sulfurs are shown as yellow spheres. (C) Putative mechanism of nucleotide reduction following the generation of the thiyl radical in step i. The cysteine numbering corresponds to *E. coli* class Ia.

Thiyl radical-based catalysis on the top face has been most clearly demonstrated in the class II RNRs. The AdoCbl-dependent nature of the class II enzymes, and fortuitous reaction kinetics, proved to be essential in trapping a thiyl radical during turnover by rapid freeze-quench (RFQ) EPR spectroscopy.<sup>14</sup> Analysis of the early reaction products resulting from mixing *Lactobacillus leichmannii* class II ribonucleotide triphosphate reductase (RTPR), substrate ATP, and AdoCbl yielded an exchange-coupled cob(II)-alamin-thiyl radical (C<sub>408</sub>-S•, *L. leichmannii* numbering), generated in a kinetically competent fashion, and shown to be chemically competent for nucleotide reduction.<sup>14,15</sup> All available data thus far indicate that the thiyl radical abstracts the substrate 3'-H (Figure 1C, step i) in all RNRs, but this has yet to be explicitly demonstrated outside this specific class II enzyme.

The role of the bottom-face cysteines in substrate reduction has also been investigated, requiring perturbations to the enzyme active site or substrate to observe intermediates. Using several mechanism-based inhibitors (MBIs), it was shown that removal of the 2'-OH is assisted by proton transfer (PT) from one of the bottom-face cysteines in both class I and class II RNRs (Figure 1C, step ii).<sup>11</sup> The resulting thiolate then reduces the 3'-keto-2'-deoxynucleotide radical (3) to form a disulfide radical anion (step iii), which then reduces the 3'-ketone regenerating the 3'-hydroxyl radical (5, step iv). Site-directed mutagenesis (SDM) affecting the proton inventory in the active site stabilizes the disulfide radical anion, which has been characterized in the *Escherichia coli* class Ia RNR (E<sub>441</sub>Q α<sub>2</sub>) by high-field pulsed EPR.<sup>16</sup> Unfortunately, the disulfide radical anion, a key intermediate in the proposed mechanism, has been observed only by altering protein or substrate.

We sought to develop methods for examining the PCET chemistry of cysteines in the active site with minimal protein or substrate perturbations. We now demonstrate the use of U as a cysteine surrogate with distinct electron and proton affinities, thus enabling the rational perturbation of the PCET chemistry at both the top and bottom face of the RNR active site. U is advantageous as it has a depressed U• reduction potential relative to both C• and Y•.<sup>2,17</sup> We targeted the α<sub>2</sub> subunit cysteines of the paradigmatic class Ia RNR from *E. coli*, C<sub>439</sub> (C<sub>439</sub>U, top face) and C<sub>225</sub> (C<sub>225</sub>U, bottom face), for mutagenesis via a recently developed amber suppression protocol<sup>18</sup> in good yield and purity. Mixing of the U-labeled protein with the diferric-tyrosyl radical harboring β<sub>2</sub> subunit, substrate, and effector results in RT from the tyrosyl radical in β (Y<sub>122</sub>•) to the active site through a pathway of redox active amino acids (Y<sub>122</sub>•[β] ↔ W<sub>48</sub>•[β] ↔ Y<sub>356</sub>[β] ↔ Y<sub>731</sub>[α] ↔ Y<sub>730</sub>[α] ↔ C<sub>439</sub>(U)[α])<sup>19</sup> and provides kinetic and chemical evidence for the involvement of thiyl radical chemistry on the top and bottom face. The results are consistent with reaction energetics arising from redox and pK<sub>a</sub> contributions, and the implication of this new method reported herein is discussed with regard to mechanistic consensus among all RNRs, and the broader implications of the method as a new tool to examine the multitude of thiyl radical enzymes.

## ■ MATERIALS AND METHODS

**Materials.** Adenosine, adenosine 5'-triphosphate (ATP), agarose, ampicillin, L-arabinose, 1 mg/L 99.99% arsenic in a 2% HNO<sub>3</sub> ICP-MS standard, bovine pancreas chymotrypsin (>40 units/mg), cytidine, cytosine, cytosine 5'-diphosphate (CDP), deoxycytidine, ethylenediaminetetraacetic acid (EDTA), glycerol (99%), guanidinium hydrochloride (GdHCl), 4-(2-

hydroxyethyl)-1-piperazineethanesulfonic acid (HEPES acid), kanamycin, LB medium, magnesium sulfate heptahydrate ( $\text{MgSO}_4$ ), methanol (HPLC grade), reduced nicotinamide adenine dinucleotide phosphate (NADPH), ruthenium(II) hexamine dichloride, 1 mg/L 99.99% selenium in 2%  $\text{HNO}_3$  ICP-MS standard, L-selenocystine, sodium selenite, and bovine pancreas trypsin (>6000 units/mg) were purchased from Sigma-Aldrich at reagent grade purity or higher. [ $5\text{-}^3\text{H}$ ]CDP was purchased from ViTrax and diluted with “cold” CDP to a final concentration of 32100 cpm/nmol. Calf intestine alkaline phosphatase (AP, 1 unit/ $\mu\text{L}$ ) and 10 $\times$  AP dephosphorylation buffer were purchased from Roche. Isopropyl  $\beta$ -D-1-thiogalactopyranoside (IPTG, 98%) and dithiothreitol (DTT, 98%) were purchased from Promega. NdeI, XhoI, NcoI, Gibson Master Mix, GoTaq Green, and Phusion polymerase were purchased from New England Biolabs. Primers were purchased from Integrative DNA Technologies (IDT). Template pET-28a plasmids for wild type ( $\text{His}$ ) $_6$ -WT  $\alpha_2$  were available from previous studies,<sup>20</sup> and the pSec-EVOL2 plasmid was kindly provided by the Söll lab from previous studies.<sup>18</sup> The pET-14b plasmid was purchased from EMD-Millipore. *E. coli* thioredoxin (Tr, 40  $\mu\text{mol min}^{-1} \text{mg}^{-1}$ ), thioredoxin reductase (TrR, 1800  $\mu\text{mol min}^{-1} \text{mg}^{-1}$ ), wt  $\beta_2$  (1.1  $\text{Y}^*/\beta_2$ , 6000  $\text{nmol min}^{-1} \text{mg}^{-1}$ ), photo $\beta_2$ , and 2,3,5-F $_3$ Y $_{356}$  photo $\beta_2$  and 2,3,6-F $_3$ Y $_{356}$  photo $\beta_2$  were available from previous studies.<sup>20–22</sup> Milli-Q water (18 M $\Omega$ ) was used for the preparation of all solutions.

**Construction of Plasmids.** The previously developed pET-28a *nrdA* plasmid was subcloned into pET-14b by Gibson assembly.<sup>23</sup> The *nrdA* fragment of the pET-28a construct was amplified by PCR with the following primers: forward, 5'-ctttgtagcagccgatcctcgcagCCGTAGGCCGGATAAG-3'; reverse, 3'-caatcgtcggcctaggagctcGGCATCCGGCCTATTCCGC-5'. Lowercase bases indicate the pET-14b complementary multiple-cloning site sequence, and uppercase bases indicate *nrdA*. The amplified *nrdA* fragment was extracted and purified by a Qiagen DNA gel purification kit. A pET-14b plasmid was doubly digested with XhoI and NcoI restriction enzymes as per the manufacturer's instructions and purified similarly. Gibson assembly of the two DNA fragments was performed as per the manufacturer's instructions. The assembly product was directly transformed into XL10 competent cells and streaked on fresh ampicillin plates. Colonies were screened for cloned *nrdA* pET-14b by PCR using the T7 promoter/terminator primers and GoTaq Green polymerase, performed as directed by the manufacturer, and 1% agarose gel electrophoresis of the colony PCR products identified several successfully cloned plasmids. Colonies containing the *nrdA* pET-14b were selected for further growth and DNA isolation by a Qiagen mini-prep kit. Sanger sequencing through Quintara Bio confirmed the *nrdA* pET-14b authenticity.

The wild type (wt) *nrdA* pET-14b was subjected to three individual SDM protocols to construct C $_{439}$ S, C $_{439}$ U, and C $_{225}$ U *nrdA* pET-14b with the following primers: C $_{225}$ U, 5'-AGCTCCTAGGTACTGATC-3' (forward) and 3'-AAGTCGAGGATCCATGAC-5' (reverse); C $_{439}$ U, 5'-AACCTGTAGCTGGAGATA-3' (forward) and 3'-AGATTGGACATCGACCTC-5' (reverse); C $_{439}$ S, 5'-AACCTGTCCCTGGAGATA-3' (forward) and 3'-AGATTGGACAGGGACCTC-5' (reverse). Bold bases represent sites of mutation. SDM by PCR was performed using Phusion polymerase as per the manufacturer's instructions, and the success of the reaction estimated by 1% agarose gel electrophoresis. Plasmids were amplified in XL10 cells; DNA

was isolated by the Qiagen mini-prep kit, and their sequence was confirmed by Sanger sequencing by Quintara Bio.

**Protein Expression.** The C $_{439}$ U RNR was produced by co-transformation of ME6 cells<sup>18</sup> (containing a scar-less quadruple deletion  $\Delta\text{selABC } \Delta\text{fdhF}$  from *E. coli* strain BW25113) with the pSec-EVOL2 and C $_{439}$ U pET-14b *nrdA* plasmids via heat shock by immersion of the samples in a 42  $^\circ\text{C}$  water bath for 30 s and then placed on ice for 2 min. Following transformation, the cell suspension was diluted 1:1 (v:v) with SOC medium and allowed to recover for 20 min at 37  $^\circ\text{C}$  prior to being plated onto LB plates supplemented with 50  $\mu\text{g/mL}$  kanamycin and 100  $\mu\text{g/mL}$  ampicillin. Single colonies were selected for further growth. To test expression conditions, small scale growth (10 mL) was employed, where a single colony was inoculated into LB medium containing 50  $\mu\text{g/mL}$  kanamycin, 100  $\mu\text{g/mL}$  ampicillin, and various levels of  $\text{Na}_2\text{SeO}_3$  (0–100  $\mu\text{M}$ ). Temperature, shaking, induction levels, and induction timing were optimized on this scale and evaluated by SDS-PAGE analysis of total protein expression and amber codon suppression. Shaking rates of 200 rpm and inoculation temperatures of 37  $^\circ\text{C}$  were found to be appropriate. Protein expression in 10 mL cultures did not scale to larger volumes, all other things being equal. We found that it was essential to inoculate single colonies directly into the larger cultures (>1 L) necessary for protein isolation and purification for biophysical methods. For large scale growth (4  $\times$  2 L), single colonies were inoculated into 2 L of the medium described above in a 6 L Erlenmeyer flask at 37  $^\circ\text{C}$  shaking at 200 rpm. At an OD $_{600}$  of 0.1 (~6 h), 30  $\mu\text{M}$   $\text{Na}_2\text{SeO}_3$  was added. At an OD $_{600}$  of 0.4, 0.1% L-arabinose was added to induce pSec-EVOL2, and the temperature was decreased to 30  $^\circ\text{C}$ . Then the pET-14b *nrdA* plasmid was induced by addition of 0.25 mM IPTG at an OD $_{600}$  of 0.8, and the culture was maintained for an additional 20 h. Cells were harvested by centrifugation, collected, flash-frozen in liquid N $_2$ , and stored until needed. An identical protocol was used for expression of all *nrdA* constructs for the sake of consistency.

Protein purification was performed identically as previously reported,<sup>21</sup> with the exception of an increased level of reductant in the final buffer exchange, yielding protein stored in 2 mM DTT, 50 mM HEPES, and 5% glycerol (pH 7.6). The truncated proteins of both C $_{439}$ U and C $_{225}$ U expression were insoluble and precipitated in the cell pellet or in the DNA pellet during purification. All proteins were flash-frozen in liquid N $_2$  following purification and stored until they were used.

**Selenium Quantitation.** The incorporation of selenium into  $\alpha$  was quantitated by inductively coupled plasma mass spectrometry (ICP-MS) on an Agilent 7900 instrument. The  $\text{Na}_2\text{SeO}_3$  used as the selenium source during protein expression was used as a selenium standard for calibration, and  $^{75}\text{As}$  was used as an internal standard in all samples. The instrument response for all isotopes except  $^{80}\text{Se}$  yielded a linear response ( $R = 0.99$ ) between from 0.5 and  $\gg 100$  ppb for all isotopes examined ( $^{74}\text{Se}$ ,  $^{76}\text{Se}$ ,  $^{77}\text{Se}$ ,  $^{78}\text{Se}$ , and  $^{82}\text{Se}$ ). Due to the overlap of the most abundant isotope ( $^{80}\text{Se}$ ) with an  $\text{Ar}_2^+$  product of the plasma,  $^{78}\text{Se}$  was routinely used (23.7% relative abundance) for quantitation. Protein samples analyzed by ICP-MS were digested by trypsin and chymotrypsin to improve solubility in the 2%  $\text{HNO}_3$  preparation solution. Protein aliquots corresponding to a theoretical final concentration of 100 ppb  $^{78}\text{Se}$  in 1.5 mL (typically 2–10  $\mu\text{L}$ ) were digested in 150  $\mu\text{L}$  of 100 mM Tris, 10 mM  $\text{CaCl}_2$ , and 2 M  $\text{GdHCl}$  (pH

8.0) with 1 unit of both trypsin and chymotrypsin for 24 h. The resulting digested peptide solution was diluted with 2% HNO<sub>3</sub> to 1.5 mL supplemented with 50 ppb arsenic in 2% HNO<sub>3</sub> and directly analyzed by ICP-MS. The ion integration time was 0.3 s, and ion counts, observed at <sup>78</sup>Se unless otherwise stated, were converted to parts per billion via a standard curve.

**Steady State Assays.** Assays were performed as previously described in a coupled assay followed spectrophotometrically.<sup>21</sup> Briefly, 0.2 μM α<sub>2</sub>, 2.5 μM β<sub>2</sub>, 1 mM CDP, 3 mM ATP, 60 μM Tr, 1 μM TrR, and 0.2 mM NADPH were mixed in assay buffer [50 mM HEPES, 15 mM MgSO<sub>4</sub>, 1 mM EDTA, and 5% glycerol (pH 7.6)], and NADPH consumption was monitored spectrophotometrically by a Cary 50 UV–vis spectrometer at 340 nm over 1 min. One unit of activity is defined as 1 nmol of CDP reduced per minute per milligram of protein. Assays were performed in triplicate, and data are reported as the mean with the error reported as one standard deviation.

**Single-Turnover Assays.** All α<sub>2</sub> samples were buffer exchanged from storage buffer into assay buffer under an inert atmosphere in a glovebox maintained at 4 °C to remove excess DTT unless otherwise noted. The buffer-exchanged α<sub>2</sub> was then added to a solution containing 10 μM β<sub>2</sub>, 0.2 mM [5-<sup>3</sup>H]CDP (32100 cpm/nmol), and 1 mM ATP to a final concentration of 10 μM α<sub>2</sub> and a final volume of 60 μL. The reaction was quenched by addition of 10 μL of 2% HClO<sub>4</sub> for 5–10 min, and then the mixture neutralized by addition of 5 μL of 0.5 M KOH. The precipitated protein was removed by centrifugation, and 50 μL of the supernatant was dephosphorylated by either (A) dilution to 450 μL in 1× AP dephosphorylation buffer supplemented with 14 units of AP or (B) dilution to 60 μL in 10× AP dephosphorylation buffer and supplemented with 14 units of AP. Solutions from both protocols were then reacted for 2 h at 37 °C.

Protocol A was employed for product isolation by the method of Steeper and Steward.<sup>24</sup> Radioactivity was used to quantitate CDP reduction products and measured by scintillation counting. Results are reported as the mean of three independent measurements, and the reported error reflects a single standard deviation.

Protocol B was employed for product isolation and characterization by HPLC on a Waters 717/515/2487 HPLC instrument equipped with an Absorbosphere Nucleotide-Nucleoside column (250 mm × 4.6 mm); 45 μL of a dephosphorylated reaction mixture (*vide supra*) was loaded onto the column equilibrated with 5 mM KP<sub>i</sub> (pH 6.8). The flow rate was maintained at 2 mL/min, and a stepwise gradient was applied as follows: isocratic 0% MeOH from 0 to 3 min, linear gradient from 0% to 5% methanol from 3 to 15 min, and linear gradient from 5% to 20% MeOH from 15 to 20 min. The eluent was monitored by a photodiode array from 220 to 400 nm, and HPLC traces are reported from absorption at 270 nm.

**X-Band EPR Spectroscopy.** Samples prepared for EPR spectroscopy were prepared by protocols C and D. In protocol C, wt β<sub>2</sub> was mixed with CDP and ATP, and reaction was initiated with addition of C<sub>439</sub>(U/S) α<sub>2</sub> to a final concentration of 50 μM α<sub>2</sub>β<sub>2</sub>, 1 mM CDP, and 3 mM ATP in assay buffer in a total volume of 220 μL and the mixture then transferred to a Wilmad 4 mm precision quartz EPR tube, all under an inert atmosphere in a glovebox at 4 °C. The reaction was quenched by hand following transfer (~15 s) into liquid N<sub>2</sub>, and the

mixture stored cryogenically. In protocol D, the EPR tube was initially loaded with wt β<sub>2</sub> (19 μL), and a premade solution of C<sub>439</sub>(U/S) α<sub>2</sub>, CDP, and ATP in assay buffer supplemented with 20% glycerol (201 μL) was swiftly added in a single step in the EPR tube and rapidly frozen in liquid N<sub>2</sub>-cooled isopentane. This protocol decreased the quenching time to ~3 s. Samples were measured directly after being frozen on a Bruker EMX X-band EPR spectrometer equipped with a recirculating helium cryostat-cooled ER4199HS cavity. The following general EPR parameters were used: microwave frequency, 9.3–9.8 GHz; modulation amplitude, 1.00 G; modulation frequency, 100 kHz; time constant, 40.96 ms. The temperature and microwave power were varied over a wide range, and individual spectra are reported with corresponding values.

**Stopped-Flow (SF) UV–Vis Spectroscopy.** SF measurements were performed on an Applied Photophysics DX 17MV instrument, and data were recorded on a personal computer. The reaction mixture was thermostated at 25 °C by a recirculating water bath. The contents of two syringes (A and B) were mixed in a 1:1 (v:v) ratio to a total volume of 60 μL. Syringe A contained 10–20 μM α<sub>2</sub> mutant of interest and 6 mM ATP in assay buffer, and syringe B contained 10–20 μM wt β<sub>2</sub> and 2 mM CDP in assay buffer. Transients were monitored at 340 nm ( $\epsilon_{\text{Se}^\bullet} \sim 1100 \text{ M}^{-1} \text{ cm}^{-1}$ , estimated by comparing to absorption at 460 nm), 460 nm ( $\epsilon_{\text{Se}^\bullet} = 560 \text{ M}^{-1} \text{ cm}^{-1}$ ),<sup>2</sup> and 412 nm ( $\epsilon_{\text{Y}^\bullet} = 3600 \text{ M}^{-1} \text{ cm}^{-1}$ ) at 4 nm resolution. The data represent the average of at least eight transients, and individual fitting parameters for mono-, bi-, and triexponential fits to experimental data starting at 3 ms are reported.

**Transient Absorption Spectroscopy.** Transient absorption (TA) lifetime and spectral measurements were performed on a home-built nanosecond time-resolved instrument described previously.<sup>25</sup> Samples were prepared with 30 μM photoβ<sub>2</sub>, 50 μM α<sub>2</sub>, 1 mM CDP, 3 mM ATP, and 10 mM Ru(NH<sub>3</sub>)<sub>6</sub>Cl<sub>3</sub> in assay buffer. The sample volume (650 μL) was recirculated and filtered by an inline 0.22 μm PFE membrane filter (Millipore Sigma). The probe light was generated by a Xe-arc lamp. For TA lifetime measurements, the observation wavelength was selected by the spectrophotometer ( $\lambda_{\text{obs}} = 430 \text{ nm}$  for 2,3,6-F<sub>3</sub>Y<sub>356</sub> photoβ<sub>2</sub> at 0.45 nm resolution). Transient signals were collected at 2 ns digital resolution and linearly binned to 100 ns resolution to decrease noise. One thousand shots were averaged in 100-shot collection intervals to ensure samples did not degrade during the course of the measurement. TA spectra were collected on a thermoelectrically cooled CCD camera with 0.45 nm resolution (as determined by slits), and pixels were binned to a final resolution of 2 nm (4 pixels/nm) from 1 to 1.05 μs after excitation. Again, 1000 shots were averaged, and spectra were collected in a  $-\log[(\text{pump on:probe on})/(\text{pump off:probe off}) - (\text{pump off:probe on})/(\text{pump off:probe off})]$  controlled by shutters and delay generators to eliminate signatures induced by the probe light.

## RESULTS

**Protein Preparation and Characterization.** Two of the three catalytic cysteines within the α<sub>2</sub> subunit of the *E. coli* class Ia RNR, C<sub>439</sub> and C<sub>225</sub>, were chosen for substitution with U, representing redox active residues on both top and bottom faces, respectively, of the active site of RNR. To accomplish U substitution for C, a newly developed strategy for heterologous

selenoprotein expression was employed, wherein U is site-specifically installed by amber codon suppression. We thus cloned the *nrdA* ( $\alpha_2$ ) RNR gene into a pET-14b plasmid and performed site-directed mutagenesis to generate the (His)<sub>6</sub>-C<sub>439</sub>Z or (His)<sub>6</sub>-C<sub>225</sub>Z  $\alpha_2$ , where Z indicates the amber codon mutation. A second plasmid was employed for selenocysteine biosynthesis and site-specific incorporation into the amber codon-labeled  $\alpha_2$  variant. This second plasmid controls expression of an engineered tRNA (allo-tRNA<sup>UTu</sup>)<sup>18</sup> and proteins SelA (*Aeromonas salmonicida* subsp. *pectinolytica* 34mel), SelD (*A. salmonicida*), and Tr (*Treponema denticola*). The allo-tRNA<sup>UTu</sup> is crucial in that it has been engineered to interact with *E. coli*'s native elongation factor Tu (EF-Tu), circumventing the necessity for a selenocysteine insertion sequence (SECIS) structural recognition element in the gene of interest mRNA, thus eliminating the need for the bacterial SelB enzyme. Allo-tRNA<sup>UTu</sup> was also chosen for its inherent capability to be charged with serine by *E. coli* serine tRNA synthetase (serRS). Once charged with serine, S-allo-tRNA<sup>UTu</sup> is converted to U-allo-tRNA<sup>UTu</sup> by SelA, with the selenium source being selenophosphate derived from SelD and selenide. Tr serves to maintain U in the reduced state.

Evaluation of C<sub>439</sub>U expression and proper U incorporation was optimized on the 10 mL scale under a variety of conditions and analyzed by SDS-PAGE (Figure S1). From the gel analysis expression of both C<sub>439</sub>U  $\alpha_2$  and selenocysteine/tRNA encoding plasmids are required for full length C<sub>439</sub>U  $\alpha_2$  expression, and that expression of both plasmids results in an ~1:1 ratio of truncated to full length protein. Encouraged by this result, we attempted to scale up expression by sequential culture saturation (5 mL → 100 mL → 2 L) for biophysical characterization. Interestingly, the level of expression was significantly lower at larger scales (>1 L) by this method, resulting in very low yields (<0.5 mg/L) and a decrease in both total protein expression and amber codon suppression. We observed expressed protein yield could be recovered to some extent by inoculating a 2 L culture with a single colony from a freshly prepared plate (transformed, plated, and grown for ~10 h) with total expression levels and amber codon suppression efficiency similar to small scale expression by SDS-PAGE. The truncated protein precipitated during cell lysis, and the resultant full length protein, following Ni affinity chromatography, was estimated to be >95% pure by SDS-PAGE (Figure S2).


Due to the nature of the selenocysteine biosynthesis machinery and incorporation method, it is possible to incorporate S rather than U at the stop codon site. This can occur if SelA/SelD activity cannot compete with S-allo-tRNA<sup>UTu</sup> charging by serRS and amber codon suppression. Fortunately, S is redox inert under the conditions employed here, rendering C<sub>439</sub>S  $\alpha_2$  a mere loss in overall yield, minimizing complications in the subsequent analysis. The efficiency of U versus S incorporation was evaluated by inductively coupled plasma mass spectrometry (ICP-MS) analysis of the purified protein to quantitate [Se] relative to [ $\alpha_2$ ] (Table S1). Following digestion of the purified  $\alpha_2$  by trypsin and chymotrypsin, ICP-MS analysis revealed that selenocysteine incorporation was dependent on  $\alpha_2$  expression levels; induction with 1 mM IPTG exhibited a Se/ $\alpha$  ratio of 0.54 (0.06), whereas with 0.25 mM IPTG, Se/ $\alpha$  ratios of >0.9 were achieved. The lower IPTG concentration came at the cost of total protein isolated (1–2 mg/g of cell paste) but improved the amber suppression efficiency. The pET-14b *nrdA* vector of

the C<sub>439</sub>U  $\alpha_2$  gene was also found to be somewhat “leaky”, expressing  $\alpha_2$  in the absence of IPTG at an effective IPTG level of 100  $\mu$ M. Under similar conditions, C<sub>225</sub>U was poorly expressed, but pure protein with a Se/ $\alpha$  ratio of 0.82 (0.07) was produced at 0.5 mg/g of cell paste. The origin of this difference is still under investigation. It is also possible that U can be overincorporated. While unlikely, we examined this possibility by expressing both wt and C<sub>439</sub>S  $\alpha_2$  under identical conditions of U incorporation. Both proteins exhibited a Se/ $\alpha$  ratio of <0.05, verifying that U is site-specifically incorporated at the amber codon. These proteins were employed as controls in further experiments. Care was taken to keep all selenoproteins microaerobic with excess (2 mM) DTT to ensure selenium remained in the selenol/selenide oxidation state.


**Top-Face Reactivity.** To assess the effect of U incorporation at the top-face C<sub>439</sub> on radical-based substrate activation, we performed multiturnover activity assays aerobically on C<sub>439</sub>U  $\alpha_2$ . No activity was found above background [20 (30) units] in these experiments, identical to the activity observed for C<sub>439</sub>S (Table S2) and consistent with contaminating wt protein. By contrast, the activity measured for wild type  $\alpha_2$  produced under identical conditions is 1920 (30) units, nearly identical to the known activity of this subunit isolated under standard conditions (2200 units).<sup>26</sup> These data imply that the C<sub>439</sub>U mutation results in either an  $\alpha_2\beta_2$  complex that is incapable of RT, a RT event that is arrested at C<sub>439</sub>U (C<sub>439</sub>U<sup>\*</sup>), or noncanonical radical chemistry with the substrate/enzyme that does not couple to bottom-face oxidation. It is also possible that, despite the presence of excess reductant, the selenocysteine has been oxidized by molecular oxygen to the corresponding selenic or seleninic acid.

Radical-based reactions in enzyme active sites can often access off-pathway outcomes, either quenching the radical or generating new products that can inactivate the enzyme; indeed, this is the case with RNR. To address this possibility in C<sub>439</sub>U  $\alpha_2$ , we performed single-turnover assays in which the external reductant system (Tr, TrR, NADPH) is omitted from the reaction, and only the endogenous reducing equivalents, C<sub>225</sub>/C<sub>462</sub> in the bottom face and a second cysteine pair in the C-terminus, can drive nucleotide reduction. In these experiments, one  $\alpha_2$  can therefore theoretically produce four dNDPs. In practice, values between 1.5 and 3.6 dCDPs/ $\alpha_2$  are obtained. Consistent with this product yield, wild type  $\alpha_2$  produced under U incorporation conditions yielded 1.6 dCDPs/ $\alpha_2$ . The C<sub>439</sub>U  $\alpha_2$  gave no observable product when submitted to single-turnover assay conditions in an inert atmosphere, either by the method of Steeper and Stuart (Table S2)<sup>24</sup> or by HPLC analysis of the reaction products (Figure 2 and Figure S3). Addition of DTT did not affect single-turnover reactivity. These data suggest that either RT is impeded altogether by the C<sub>439</sub>U mutation or RT is arrested at this site.

To evaluate the potential for C<sub>439</sub>U to act as a radical trap, we turned to X-band EPR spectroscopy. Under conditions similar to those employed in single-turnover experiments, C<sub>439</sub>U and C<sub>439</sub>S  $\alpha_2$  were mixed separately with  $\beta_2$ , CDP, and ATP and quenched by hand in liquid nitrogen after 10–15 s in an X-band EPR tube and submitted to X-band EPR analysis (Figure 3). Thiyl and selenyl radicals, while extremely rare in the literature, are generally observed to have strong g anisotropy and relatively fast spin relaxation rates.<sup>4,27–30</sup> Our EPR analysis reveals that the reaction of C<sub>439</sub>U  $\alpha_2$  with wt  $\beta_2$  (1.1 Y<sup>\*</sup>/ $\beta_2$ ), CDP, and ATP yields a net loss of ~60% of the



**Figure 2.** HPLC chromatograms of single-turnover reaction products for 10  $\mu\text{M}$  wt (blue),  $\text{C}_{225}\text{U}$  (green), and  $\text{C}_{439}\text{U}$  (red)  $\alpha_2$  with 10  $\mu\text{M}$   $\beta_2$  and an authentic standard (black). The standard contained 20  $\mu\text{M}$  cytosine ( $t_{\text{R}} = 2.6$  min), cytidine ( $t_{\text{R}} = 4.6$  min), and deoxycytidine ( $t_{\text{R}} = 7.3$  min). Peak splitting was observed at all flow rates and all concentrations suggesting column degradation [channeling (see Figure S3)]. The asterisk indicates the residual protein absorption profile that elutes in the void volume.




**Figure 3.** X-Band EPR spectra of the  $\text{Y}_{122}^{\bullet}$  of wt  $\beta_2$  in complex with  $\text{C}_{439}\text{S}$  (black) and  $\text{C}_{439}\text{U}$  (red)  $\alpha_2$ . Samples were prepared by protocol C, resulting in 50  $\mu\text{M}$   $\text{C}_{439}\text{X}$  ( $\text{X} = \text{S}$  or  $\text{U}$ )  $\alpha_2$ , 50  $\mu\text{M}$  wt  $\beta_2$ , 1 mM CDP, and 3 mM ATP in assay buffer. Spectra were recorded at 80 K with a modulation frequency of 100 kHz, a modulation amplitude of 1 G, and a microwave power of 20  $\mu\text{W}$ .

$\text{Y}_{122}^{\bullet}$  signature as determined by double integration of the derivative spectrum, and no new resonance consistent with a selenyl radical was observed. Further experiments with 2-deoxy-2-azido cytidine diphosphate ( $\text{N}_3\text{CDP}$ ), a mechanism-based inhibitor that forms a distinct N-centered substrate radical following activation by 3'-H abstraction during the reaction with wt  $\alpha_2$  and  $\beta_2$ ,<sup>31</sup> yielded slightly lower levels of  $\text{Y}_{122}^{\bullet}$  loss [ $\sim 40\%$  (Figure S4)], but no new resonance was observed. Radical trapping experiments were performed under a variety of conditions: (i) with and without reductant and (ii) in a completely inert atmosphere. In either case, no new species were observed in the EPR spectrum. We also varied the quenching method to access faster quenching times of  $\sim 3$  s to no avail.

The kinetics of RT in the *E. coli* class Ia RNR have been characterized at each Y within the RT pathway by stopped-flow (SF) UV-vis; these experiments reveal that RT is gated by conformational changes that enable long-range PCET at  $\sim 8$   $\text{s}^{-1}$ .<sup>21,26,32</sup> Here we applied SF UV-vis to define the kinetics of RT from  $\text{Y}_{122}^{\bullet}[\beta]$  to  $\text{C}_{439}\text{U}[\alpha]$  by monitoring both the sharp absorption feature of  $\text{Y}_{122}^{\bullet}$  ( $\lambda_{\text{max}} = 412$  nm, and  $\epsilon_{412} \sim 3600$   $\text{M}^{-1} \text{cm}^{-1}$ )<sup>33</sup> and the broad selenyl radical absorption (300–


500 nm;  $\epsilon < 1000$   $\text{M}^{-1} \text{cm}^{-1}$ ).<sup>2</sup> Mixing of  $\text{C}_{439}\text{U}$   $\alpha_2$  and ATP with  $\beta_2$  and CDP to a final concentration of 10  $\mu\text{M}$   $\alpha_2\beta_2$ , 1 mM CDP, and 3 mM ATP resulted in a decay of the 412 nm feature with bimolecular kinetics characterized by a  $k_{\text{obs1}}$  of 7.1 (0.1)  $\text{s}^{-1}$  and a  $k_{\text{obs2}}$  of 0.40 (0.01)  $\text{s}^{-1}$  (Figure 4, red). A



**Figure 4.** Stopped-flow UV-vis kinetics for wt  $\beta_2$  and effector mixed with  $\text{C}_{439}\text{U}$   $\alpha_2$  and substrate. Transients at 412 nm (left axis) are colored red, and those at 340 nm (right axis) blue, with associated biexponential fits in black. Proteins at a final concentration of 10  $\mu\text{M}$ .

temporally concerted induced absorption signal was observed at both 340 nm [Figure 4, blue;  $k_{\text{obs1}} = 6.5$  (0.3)  $\text{s}^{-1}$ , and  $k_{\text{obs2}} = 0.42$  (0.03)  $\text{s}^{-1}$ ] and 460 nm (Figure S5), with a markedly lower intensity. This induced absorption feature at 460 nm appeared to be stable for 10 s and decayed with a rate of 0.02  $\text{s}^{-1}$  (Figure S6). The induced absorbances at 340 and 460 nm, while consistent with a weak absorption profile of  $\text{C}_{439}\text{U}^{\bullet}$ , also coincide with the broad absorption profile of the di-iron cluster. Control experiments with wt  $\alpha_2$  probed at 340 and 460 nm yielded kinetically similar yet markedly weaker signals, thus complicating interpretation (Figure S7). Fitting amplitudes and rates are listed in Table S3.

The rate-determining conformational gating of RT and active site chemistry during the reaction of  $\alpha_2$  with  $\beta_2$  precludes insight into the role of the  $\text{C}_{439}\text{U}$  substitution in long-range PCET.<sup>26</sup> We have previously employed a photosensitized  $\beta_2$ , termed photo $\beta_2$ , to rapidly generate kinetically and chemically competent radicals on the RT pathway at the subunit interface in an intact  $\alpha_2\beta_2$  complex.<sup>34</sup> This photo $\beta_2$  is constructed by the covalent attachment of a rhenium(I) 1,10-phenanthroline tricarbonyl bromomethylpyridine chromophore ( $[\text{Re}^{\text{I}}]$ ) adjacent to the pathway residue  $\text{Y}_{356}$  through a  $\text{S}_{355}\text{C}$  substitution. Upon excitation of  $[\text{Re}^{\text{I}}]$ , the excited state can directly oxidize  $\text{Y}_{356}$  followed by charge recombination, or the  $[\text{Re}^{\text{I}}]^*$  state can be oxidatively quenched, resulting in a  $[\text{Re}^{\text{II}}]$  that can subsequently oxidize  $\text{Y}_{356}$  to form a long-lived  $\text{Y}_{356}^{\bullet}$  for spectroscopic and kinetic characterization.<sup>20,21</sup> To address the influence of the  $\text{C}_{439}\text{U}$  mutation in  $\alpha_2$  on RT kinetics in the absence of conformational gating, we employed a 2,3,6- $\text{F}_3\text{Y}_{356}$ -photo $\beta_2$ . The 2,3,6- $\text{F}_3\text{Y}_{356}$  substitution enables specific spectroscopic observation of 2,3,6- $\text{F}_3\text{Y}_{356}^{\bullet}$  among the potential  $\text{Y}^{\bullet}$  background due to the distinct absorption profile<sup>21</sup> and also increases the one-electron reduction potential of the (fluoro)phenoxy radical relative to tyrosine by 135 mV,<sup>35</sup> biasing forward RT into  $\alpha_2$  and the remainder of the RT pathway. By monitoring the kinetics of 2,3,6- $\text{F}_3\text{Y}_{356}^{\bullet}$  in the presence of the radical trap  $\text{C}_{439}\text{U}$   $\alpha_2$  and comparing this rate to that obtained in the presence of the RT pathway block,  $\text{Y}_{731}\text{F}$ , we can deduce the rate of PCET across the subunit interface.<sup>21</sup> Figure 5 shows the kinetics of 2,3,6- $\text{F}_3\text{Y}_{356}^{\bullet}$  decay



**Figure 5.** Transient absorption kinetics of 2,3,6-F<sub>3</sub>Y<sub>356</sub><sup>•</sup> decay (red) in the presence of C<sub>439</sub>U α<sub>2</sub> substrate, and effector in spectroscopy buffer and the associated monoexponential fit. Data represent the average of three independent trials, and the fit is displayed for illustrative purposes.


in the presence of C<sub>439</sub>U α<sub>2</sub> at pH 8.2. The rate of decay was determined to be  $9.9 (0.1) \times 10^4 \text{ s}^{-1}$  ( $k_{\text{obs}}$ ) as compared to the rate in the presence of Y<sub>731</sub>F previously determined to be  $6.8 (0.3) \times 10^4 \text{ s}^{-1}$  ( $k_0$ ), yielding a radical injection rate  $k_{\text{PCET}}$  of  $3.1 (0.5) \times 10^4 \text{ s}^{-1}$ . This observation is consistent with the data presented above, indicating that C<sub>439</sub>U is redox active and modulates RT analogous to 3-aminotyrosine (NH<sub>2</sub>Y).<sup>21</sup>

**Bottom-Face Reactivity.** The redox properties of C<sub>225</sub>U α<sub>2</sub> are distinct from the C<sub>439</sub>U, C<sub>225</sub>S, or wt counterparts. Single-turnover and multiturnover assays performed either aerobically or anaerobically rendered no NDP reduction activity, or very little in the case of C<sub>225</sub>U. Single-turnover activity was observed under anaerobic conditions if the C<sub>225</sub>U α<sub>2</sub> protein was directly used from DTT-containing storage buffer, resulting in a final DTT concentration of 200 μM in the assay mixture. The residual DTT can serve as an external reductant, allowing the oxidized cysteines of the bottom face to be re-reduced for subsequent turnovers, yielding a maximal

theoretical turnover of 24 dCDPs/α<sub>2</sub>. Under these conditions, reanalysis of the single-turnover activity of C<sub>225</sub>U revealed 7.9 (0.9) dCDPs/α<sub>2</sub>, similar to the activity observed for wt α<sub>2</sub> (11 dCDPs/α<sub>2</sub>) as determined by the method of Steeper and Stuart (Table S2).<sup>24</sup> This method of turnover quantitation does not discriminate between dCDP and cytosine production, a byproduct of substrate degradation due to incomplete reduction of the 3'-keto intermediate (Figure 1C, intermediate 4), resulting in the release of pyrophosphate, cytosine (base), and furanone (Figure S8).<sup>36–38</sup> The formation of cytosine and dCDP by C<sub>225</sub>U α<sub>2</sub> was evidenced by HPLC analysis of the reaction products under similar conditions (Figure 2, green). This reactivity is analogous to that of 2'-deoxy-2'-X cytidine diphosphate (X = F or Cl), mechanism-based inhibitors of RNR. Efforts to characterize intermediates during substrate activation and selenyl radical chemistry in the C<sub>225</sub>U α<sub>2</sub> protein were unsuccessful as determined by X-band EPR following the mixing of C<sub>225</sub>U α<sub>2</sub> with wt β<sub>2</sub>, CDP, and ATP (Figure S9) on the 3–15 s time scale.

## DISCUSSION

The thiol radical form of cysteine is the active cofactor implicated in RNR and in glycol radical enzymes (GREs), a growing class of enzymes that are important in anaerobic metabolism.<sup>39,40</sup> In particular, GREs are pivotal in host–microbiome interactions in the anaerobic environment of the gut, where they endow microorganisms with the ability to exploit host-derived metabolites<sup>41</sup> and xenobiotics<sup>42</sup> for energy and biosynthesis. The reactivity of thiol radicals represents a natural target for small molecule mechanism-based inhibitor design and is a target of current chemotherapeutic drugs.<sup>43,44</sup> The rational design of inhibitors and the better understanding of thiol radicals in biology require a detailed understanding of their reactivity that has proven to be experimentally challenging. In this work, we substitute the redox active



**Figure 6.** Proposed reaction sequences for C<sub>439</sub>U (top) and C<sub>225</sub>U (bottom) α<sub>2</sub>.



cysteines of the paradigmatic thiol radical enzyme RNR of *E. coli* for selenocysteine in an attempt to attenuate radical reactivity such that intermediates may be observed and reactivity trends revealed. Selenocysteine has been demonstrated to be quite distinct in reactivity relative to cysteine, particularly in its ability to scavenge reactive oxygen species reversibly and have diminished activity toward deleterious radical-based protein degradation.<sup>45</sup> We have shown that the method of site-specific U incorporation developed by Söll et al.<sup>18</sup> is robust in producing relatively large quantities of quantitatively substituted selenoproteins on the scale necessary for biochemical and biophysical characterization. While a strategy was ultimately developed for large scale expression, significant challenges were encountered en route. The observation of poor expression following sequential culture saturations as opposed to direct inoculation of the large volume medium preparations suggests that either replication events or growth stagnation affects protein expression. This observation must be reconciled with the additional observation of no observable effect on growth rates in the presence of antibiotics that select for each plasmid. We do not observe any issues with growth or expression using just pET-based vectors for *nrdA* expression, indicating that the likely culprit is the pSec-EVOL2 vector instability. The pSec-EVOL2 plasmid contains a high-copy number origin of replication that may place some burden on the co-transformed cells. We hypothesize that horizontal gene transfer of the kanamycin resistance cassette during stationary phase conditions and the subsequent elimination of the pSec-EVOL2 plasmid during replication may rationalize the observed phenomena.

An interesting and unexpected consequence of the C<sub>439</sub>U mutation is that multiple methods of assessing RT provide strong kinetic evidence that radicals on pathway appear to funnel toward U<sub>439</sub>; however, no U<sub>439</sub>• is directly observed, and no product deoxynucleotide is formed. SF UV–vis and transient absorption spectroscopies demonstrate that Y<sub>122</sub>• and Y<sub>356</sub>•, respectively, are highly susceptible to reduction by RT and behave analogously to prior observations of radical trap NH<sub>2</sub>Y-labeled α<sub>2</sub> proteins.<sup>21,26</sup> We have exhaustively sought to observe the putative C<sub>439</sub>U• intermediate by SF UV–vis, EPR, and transient absorption (TA) using a photoβ<sub>2</sub> construct. SF UV–vis data indicate that a new optical feature is formed concomitant to Y<sub>122</sub>• reduction that resembles an authentic U•, but no evidence of this species was obtained by either EPR or TA. We hypothesize that either the radical exhibits an extreme level of broadening of the EPR signal rendering it unresolved in the current experiments or the transiently formed U• reacts with another element of the protein structure on a rapid time scale and is reduced (Figure 6). Thus, the reactivity of the corresponding putative U<sub>439</sub>• appears to be quite high but distinct from C•, allowing insight to be gained regarding top-face thiol radical catalysis. This is not unexpected as the C–H bond dissociation energy (BDE) of the 3′-H is estimated to be ~390 kJ/mol for aliphatic alcohols,<sup>46</sup> whereas the BDE of Se–H of U is ~310 kJ/mol and the Se•/Se<sup>–</sup> couple ~500 mV lower than that of the corresponding S•/SH.<sup>2</sup> With regard to the latter, the reactivity of S• toward elements of protein structures is well-documented and they can attack C–H bonds of the protein backbone, with known reactivity toward the relatively weak amino acid α-H (BDE ~ 355 kJ/mol).<sup>47,48</sup> The reactivity is kinetically accelerated due to polar resonance effects of the SH/S• exchange.<sup>49,50</sup> Furthermore, C• acts as a nucleophile in the GRE pyruvate formate lyase, where it is

proposed to attack the carbonyl carbon resulting in a transient substrate-based oxyl radical.<sup>51,52</sup> Analysis of the active site of the *E. coli* α<sub>2</sub> near C<sub>439</sub> does not immediately reveal a likely site of the backbone or side chain attack. Furthermore, selenyl radical attack of the protein backbone should produce a new radical, likely a carbon-centered radical, which is not observed under any of the conditions applied herein. It is also possible that the protonation state of U<sub>439</sub>, anticipated to be a selenate, affects PCET during radical transfer, as C<sub>439</sub> is both the proton and electron donor during radical transfer from Y<sub>730</sub>• to C<sub>439</sub>. In this scenario, the Y<sub>730</sub>• → C<sub>439</sub> would require protonation of Y<sub>730</sub>–O•. The H-bonding environment of Y<sub>730</sub>–O• has been examined by electron nuclear double resonance (ENDOR)<sup>53,54</sup> and hyperfine sublevel correlation spectroscopy (HYSCORE) and contains H-bonds from both Y<sub>731</sub> and C<sub>439</sub> as well as a water molecule observed in several X-ray structures.<sup>55,56</sup> Current evidence points to water as a proton donor during reduction of Y<sub>356</sub>,<sup>54</sup> and thus, this is a plausible alternative proton donor to Y<sub>730</sub> during radical transfer to U<sub>439</sub>.

When cysteine-based thiols are involved in enzymatic catalysis, they often serve as electron donors or nucleophiles. In such instances, the substitution for selenocysteine may impact the reaction outcome, significantly affecting some reactions through the “seleno effect”.<sup>57</sup> This effect, measured as  $k_s/k_{se}$  for the reaction of interest, can be significantly greater than or less than 1, depending on the reaction mechanism and role of the particular chalcogen. For the catalytic C/U at position 439 of α<sub>2</sub>, the seleno effect appears to be quite large ( $k_s/k_{se} \gg 1$ ), rendering the reaction mute in the seleno form. This is anticipated to be a conserved effect for all thiol radical enzymes that involve C–H activation, including RNR and most GREs. Conversely, during substrate radical reduction (Figure 1C, steps ii–iv), the bottom-face cysteines serve to reduce the substrate. Given the higher nucleophilicity of U relative to C, and the lower reduction potential of the mixed selenosulfide relative to the canonical disulfide, it may be expected here that the seleno effect of the overall reaction for C<sub>225</sub>U would be low ( $k_s/k_{se} < 1$ ). Kinetic analysis of the seleno effect in this instance is impossible due to the conformational gating of the overall reaction,<sup>58</sup> and the rapid active site chemistry,<sup>59</sup> but we do observe substrate reduction at the earliest time points examined, suggesting that  $k_s/k_{se}$  is not significantly increased.

The proton inventory during substrate radical reduction is essential to the proper orchestration of radical chemistry at the active site. This has been examined in several ways, including active site mutations that inhibit side chain ionization (C<sub>225</sub>S, C<sub>462</sub>S, and E<sub>441</sub>Q)<sup>9,60,61</sup> and mechanism-based inhibitor substrates whose leaving groups are not protonated upon release (2′-deoxy-2′-X, where X = Cl or F). In the latter case, the substrate radical can lose the 2′-X as X<sup>–</sup>, whereas the native substrate loses the 2′-OH as H<sub>2</sub>O, taking a proton of the active site from C<sub>225</sub>. On the basis of the pK<sub>a</sub> of selenocysteine (5.2–5.4), it is anticipated that the U<sub>225</sub> side chain is deprotonated, absent significant pK<sub>a</sub> perturbations at this site due to the protein environment. This would render U<sub>225</sub> unable to protonate the 2′-OH and alter the proton inventory. Thus, with mechanism-based inhibitors, the proton inventory is too large, whereas in the C<sub>225</sub>U protein, it is too small. The selenate is a better reductant than the corresponding thiolate and may be expected to reduce the substrate radical, but by ET rather than by PCET. Product analysis of the reaction of C<sub>225</sub>U with CDP, ATP, and wt β<sub>2</sub> shows that dCDP is indeed

produced but that the vast majority of substrate decomposes from the radical/3'-keto state, producing cytosine. The small fraction of dCDP formed in the single-turnover reaction of C<sub>225</sub>U could be due to a small fraction of selenol (SeH) that can donate an H atom during reduction of the 2' radical, or it could be due to a small quantity of contaminating wt protein; at this point, we cannot disentangle these two possibilities. SDS-PAGE analysis of the resultant protein indicated no cleavage of the C<sub>225</sub>U protein following the reaction (data not shown), which is distinct behavior relative to the C<sub>225</sub>S protein, which is cleaved near the 225 site during the reaction with wt  $\beta_2$ , substrate, and effector.<sup>15</sup> Furthermore, EPR analysis of the reaction mixture following single-turnover preparations of C<sub>225</sub>U  $\alpha_2$  showed little radical loss over the course of the reaction. Collectively, these data suggest that the substrate radical generated by H atom abstraction by the C<sub>439</sub> thiyl radical is not reduced by the U<sub>225</sub> in the absence of a selenol but rather by the top-face C<sub>439</sub>, resulting in a furanone that decomposes to release cytosine, 2-methylene-3(2H)-furanone, and pyrophosphate, leaving the protein-based radical intact (Figure 6). Inhibition on the longer time scale is caused by alkylation of the active site or C-terminal cysteines. The analogous reaction products produced by wt  $\alpha_2$  with mechanism-based inhibitors (excess H<sup>+</sup>) and the C<sub>225</sub>U  $\alpha_2$  with native substrates (deficient H<sup>+</sup>) illustrate the essential role of proton coupling in PCET during radical-based active site catalysis.

## CONCLUSIONS

The results presented herein have established a facile method for site-specifically substituting redox active cysteines for selenocysteine. They also reveal for the first time the seleno effect on thiyl radical chemistry in a thiyl radical enzyme and render parallels between observations made on the thiyl radical reactivity of the class II RNRs with those of the class I enzymes, relevant to human health and common bacterial infections. The successful generation and analysis of these selenoproteins suggest that this technique may be broadly useful for interpreting reactivity trends in the diverse GREs, which can cleave C-C, C-N, C-O, C-S, and C-P bonds through diverse mechanisms that are challenging to examine by the conventional techniques of biochemistry and biophysics.

## ASSOCIATED CONTENT

### Supporting Information

The Supporting Information is available free of charge at <https://pubs.acs.org/doi/10.1021/acs.biochem.9b00973>.

SDS-PAGE analysis of selenoprotein expression and purification, selenium quantitation, activity measurements, HPLC standards, EPR spectrum of C<sub>439</sub>S/C<sub>439</sub>U  $\alpha_2$  and wt  $\beta_2$  with N<sub>3</sub>CDP, SF UV-vis kinetics of C<sub>439</sub>U  $\alpha_2$ , SF UV-vis kinetics of wt  $\alpha_2$ , mechanism of RNR inhibition by 2'-F/Cl-substituted nucleotides, EPR spectrum of C<sub>225</sub>U  $\alpha_2$  and wt  $\beta_2$  with CDP, and mechanism of RNR inhibition by 2'-N<sub>3</sub>-substituted nucleotides (PDF)

### Accession Codes

The UniProt accession numbers for *E. coli* class Ia RNR  $\alpha_2$  and  $\beta_2$  are P00452 and P69924, respectively.

## AUTHOR INFORMATION

### Corresponding Author

\*E-mail: [greene@chem.ucsb.edu](mailto:greene@chem.ucsb.edu).

### ORCID

Brandon L. Greene: 0000-0001-6981-5900

JoAnne Stubbe: 0000-0001-8076-4489

Daniel G. Nocera: 0000-0001-5055-320X

### Funding

This work was supported by National Institutes of Health Grants GM047274 (D.G.N.) and GM029595 (J.S.).

### Notes

The authors declare no competing financial interest.

## ACKNOWLEDGMENTS

The authors gratefully acknowledge Dr. Ana Crnković and Prof. Dieter Söll for providing the pSec-EVOL2 plasmid, ME6 cells, and generous insight and advice.

## ABBREVIATIONS

RNR, ribonucleotide reductase; U, selenocysteine; C, cysteine; PCET, proton-coupled electron transfer; Y, tyrosine; W, tryptophan; G, glycine; RT, radical transfer; AdoCbl, adenosylcob(II)alamin; RFQ, rapid freeze quench; EPR, electron paramagnetic resonance; RTPR, ribonucleotide triphosphate reductase; ATP, adenosine triphosphate; MBI, mechanism-based inhibitor; PT, proton transfer; SDM, site-directed mutagenesis; CDP, cytosine 5'-diphosphate; EDTA, ethylenediaminetetraacetic acid; ENDOR, electron nuclear double resonance; GdHCl, guanidinium hydrochloride; HEPES, 4-(2-hydroxyethyl)-1-piperazineethanesulfonic acid; HYSCORE, hyperfine sublevel correlation; NADPH, reduced nicotinamide adenine dinucleotide phosphate; AP, calf intestine alkaline phosphatase; IPTG, isopropyl  $\beta$ -D-1-thiogalactopyranoside; DTT, dithiothreitol; Tr, *E. coli* thioredoxin; TrR, *E. coli* thioredoxin reductase; PCR, polymerase chain reaction; SDS-PAGE, sodium dodecyl sulfate-polyacrylamide gel electrophoresis; ICP-MS, inductively coupled plasma mass spectrometry; HPLC, high-performance liquid chromatography; SF, stopped-flow; TA, transient absorption; EF-Tu, elongation factor Tu; SECIS, selenocysteine insertion sequence; serRS, serine tRNA synthetase; [Re<sup>I</sup>], rhenium(I) 1,10-phenanthroline tricarbonyl bromomethylpyridine chromophore; NH<sub>2</sub>Y, 3-aminotyrosine; GREs, glycy radical enzymes; BDE, bond dissociation energy.

## REFERENCES

- (1) Stubbe, J., and van der Donk, W. A. (1998) Protein radicals in enzyme catalysis. *Chem. Rev.* 98, 705–762.
- (2) Nauser, T., Dockheer, S., Kissner, R., and Koppenol, W. H. (2006) Catalysis of Electron Transfer by Selenocysteine. *Biochemistry* 45, 6038–6043.
- (3) Schöneich, C. (2011) Cysteine residues as catalysts for covalent peptide and protein modification: a role for thiyl radicals? *Biochem. Soc. Trans.* 39, 1254–1259.
- (4) Everett, S. A., Schöneich, C., Stewart, J. H., and Asmus, K. D. (1992) Perthiyl Radicals, Trisulfide Radical Ions, and Sulfate Formation. A Combined Photolysis and Radiolysis Study on Redox Processes with Organic Di- and Trisulfides. *J. Phys. Chem.* 96, 306–314.
- (5) van Gastel, M., Lubitz, W., Lassmann, G., and Neese, F. (2004) Electronic Structure of the Cysteine Thiyl Radical: A DFT and Correlated *ab Initio* Study. *J. Am. Chem. Soc.* 126, 2237–2246.

- (6) Reichard, P. (1993) From RNA to DNA, why so many ribonucleotide reductases? *Science* 260, 1773–1777.
- (7) Stubbe, J. A., and van der Donk, W. A. (1995) Ribonucleotide reductases: radical enzymes with suicidal tendencies. *Chem. Biol.* 2, 793–801.
- (8) Aberg, A., Hahne, S., Karlsson, M., Larsson, A., Ormö, M., Ahgren, A., and Sjöberg, B. M. (1989) Evidence for two different classes of redox-active cysteines in ribonucleotide reductase of *Escherichia coli*. *J. Biol. Chem.* 264, 12249–12252.
- (9) Mao, S. S., Holler, T. P., Yu, G. X., Bollinger, J. M., Jr., Booker, S., Johnston, M. I., and Stubbe, J. (1992) A Model for the Role of Multiple Cysteine Residues Involved in Ribonucleotide Reduction: Amazing and Still Confusing. *Biochemistry* 31, 9733–9743.
- (10) Sintchak, M. D., Arjara, G., Kellogg, B. A., Stubbe, J., and Drennan, C. L. (2002) The crystal structure of class II ribonucleotide reductase reveals how an allosterically regulated monomer mimics a dimer. *Nat. Struct. Biol.* 9, 293–300.
- (11) Harris, G., Ator, M., and Stubbe, J. (1984) Mechanism of Inactivation of *Escherichia coli* and *Lactobacillus leichmannii* Ribonucleotide Reductases by 2'-Chloro-2'-deoxynucleotides: Evidence for Generation of 2-Methylene-3(2H)-furanone. *Biochemistry* 23, 5214–5225.
- (12) Thelander, L., and Larsson, B. (1976) Active site of ribonucleoside diphosphate reductase from *Escherichia coli*. Inactivation of the enzyme by 2'-substituted ribonucleoside diphosphates. *J. Biol. Chem.* 251, 1398–1405.
- (13) Mao, S. S., Holler, T. P., Bollinger, M. J., Jr., Yu, G. X., Johnston, M. I., and Stubbe, J. (1992) Interaction of C225SR1 Mutant Subunit of Ribonucleotide Reductase with R2 and Nucleoside Diphosphates: Tales of a Suicidal Enzyme. *Biochemistry* 31, 9744–9751.
- (14) Licht, S., Gerfen, G. J., and Stubbe, J. A. (1996) Thiyl Radicals in Ribonucleotide Reductases. *Science* 271, 477–481.
- (15) Gerfen, G. J., Licht, S., Willems, J. P., Hoffman, B. M., and Stubbe, J. (1996) Electron Paramagnetic Resonance Investigations of a Kinetically Competent Intermediate Formed in Ribonucleotide Reduction: Evidence for a Thiyl Radical-Cob(II)alamin Interaction. *J. Am. Chem. Soc.* 118, 8192–8197.
- (16) Lawrence, C. C., Bennati, M., Obias, H. V., Bar, G., Griffin, R. G., and Stubbe, J. (1999) High-field EPR detection of a disulfide radical anion in the reduction of cytidine 5'-diphosphate by the E441Q R1 mutant of *Escherichia coli* ribonucleotide reductase. *Proc. Natl. Acad. Sci. U. S. A.* 96, 8979–8984.
- (17) Steinmann, D., Nauser, T., and Koppenol, W. H. (2006) Kinetics of tyrosyl radical repair by selenocysteine. *Free Radical Res.* 40, S136–S136.
- (18) Mukai, T., Sevostyanova, A., Suzuki, T., Fu, X., and Söll, D. (2018) A facile method for producing selenocysteine-containing proteins. *Angew. Chem., Int. Ed.* 57, 7215–7219.
- (19) Minnihan, E. C., Nocera, D. G., and Stubbe, J. (2013) Reversible, Long-Range Radical Transfer in *E. coli* Class Ia Ribonucleotide Reductase. *Acc. Chem. Res.* 46, 2524–2535.
- (20) Greene, B. L., Stubbe, J., and Nocera, D. G. (2018) Photochemical Rescue of a Conformationally Inactivated Ribonucleotide Reductase. *J. Am. Chem. Soc.* 140, 15744–15752.
- (21) Greene, B. L., Taguchi, A. T., Stubbe, J., and Nocera, D. G. (2017) Conformationally Dynamic Radical Transfer within Ribonucleotide Reductase. *J. Am. Chem. Soc.* 139, 16657–16665.
- (22) Olshansky, L., Stubbe, J., and Nocera, D. G. (2016) Charge-Transfer Dynamics at the  $\alpha/\beta$  Subunit Interface of a Photochemical Ribonucleotide Reductase. *J. Am. Chem. Soc.* 138, 1196–1205.
- (23) Gibson, D. G., Young, L., Chuang, R. Y., Venter, J. C., Hutchison, C. A., and Smith, H. O. (2009) Enzymatic assembly of DNA molecules up to several hundred kilobases. *Nat. Methods* 6, 343–345.
- (24) Steeper, J. R., and Steuart, C. D. (1970) A rapid assay for CDP reductase activity in mammalian cell extracts. *Anal. Biochem.* 34, 123–130.
- (25) Pizano, A. A., Olshansky, L., Holder, P. G., Stubbe, J., and Nocera, D. G. (2013) Modulation of Y356 Photooxidation in *E. coli* Class Ia Ribonucleotide Reductase by Y731 Across the  $\alpha2:\beta2$  Interface. *J. Am. Chem. Soc.* 135, 13250–13253.
- (26) Minnihan, E. C., Seyedsayamdost, M. R., Uhlin, U., and Stubbe, J. (2011) Kinetics of Radical Intermediate Formation and Deoxynucleotide Production in 3-Aminotyrosine-Substituted *Escherichia coli* Ribonucleotide Reductases. *J. Am. Chem. Soc.* 133, 9430–9440.
- (27) Nelson, D. J., Petersen, R. L., and Symons, M. C. R. (1977) Unstable Intermediates. Part 178. The Structure of Intermediates Formed in the Radiolysis of Thiols. *J. Chem. Soc., Perkin Trans. 2* 2, 2005–2015.
- (28) Budzinski, E. E., and Box, H. C. (1971) Oxidation and Reduction of Organic Compounds by Ionizing Radiation: L-Penicillamine Hydrochloride. *J. Phys. Chem.* 75, 2564–2570.
- (29) Kennedy, M. C., Antholine, W. E., and Beinert, H. (1997) An EPR Investigation of the Products of the Reaction of Cytosolic and Mitochondrial Aconitases with Nitric Oxide. *J. Biol. Chem.* 272, 20340–20347.
- (30) Kolberg, M., Bleifuss, G., Gräslund, A., Sjöberg, B. M., Lubitz, W., Lendzian, F., and Lassmann, G. (2002) Protein thiyl radicals directly observed by EPR spectroscopy. *Arch. Biochem. Biophys.* 403, 141–144.
- (31) van der Donk, W. A., Stubbe, J., Gerfen, G. J., Bellew, B. F., and Griffin, R. G. (1995) EPR Investigations of the Inactivation of *Escherichia coli* Ribonucleotide Reductase with 2'-Azido-2'-Deoxyuridine 5'-Diphosphate: Evidence for the Involvement of the Thiyl Radical of C225-R1. *J. Am. Chem. Soc.* 117, 8908–8916.
- (32) Kasanmascheff, M., Lee, W., Nick, T. U., Stubbe, J., and Bennati, M. (2016) Radical transfer in *E. coli* ribonucleotide reductase: a NH2Y731/R411A- $\alpha$  mutant unmasks a new conformation of the pathway residue 731. *Chem. Sci.* 7, 2170–2178.
- (33) Bollinger, J. M., Jr., Hangtong, W., Ravi, N., Huynh, B. H., Edmondson, D., and Stubbe, J. (1995) Use of rapid kinetics methods to study the assembly of the diferric-tyrosyl radical cofactor of *E. coli* ribonucleotide reductase. *Methods Enzymol.* 258, 278–303.
- (34) Pizano, A. A., Lutterman, D. A., Holder, P. G., Teets, T. S., Stubbe, J., and Nocera, D. G. (2012) Photo-ribonucleotide reductase  $\beta2$  by selective cysteine labeling with a radical phototrigger. *Proc. Natl. Acad. Sci. U. S. A.* 109, 39–43.
- (35) Ravichandran, K. R., Zong, A. B., Taguchi, A. T., Nocera, D. G., Stubbe, J., and Tommos, C. (2017) Formal Reduction Potentials of Difluorotyrosine and Trifluorotyrosine Protein Residues: Defining the Thermodynamics of Multistep Radical Transfer. *J. Am. Chem. Soc.* 139, 2994–3004.
- (36) Thelander, L., Larsson, B., Hobbs, J., and Eckstein, F. (1976) Active Site of Ribonucleoside Diphosphate Reductase from *Escherichia coli*. *J. Biol. Chem.* 251, 1398–1405.
- (37) Stubbe, J., and Kozarich, J. W. (1980) Inorganic Pyrophosphate Is Released from 2'-Chloro-2'-deoxyuridine 5'-Diphosphate by Ribonucleoside Diphosphate Reductase. *J. Am. Chem. Soc.* 102, 2505–2507.
- (38) Stubbe, J., and Kozarich, J. W. (1980) Fluoride, Pyrophosphate, and Base Release from  $\dot{Z}$ -Deoxy- $\dot{Z}$ -fluoronucleoside 5'-Diphosphates by Ribonucleoside-Diphosphate Reductase. *J. Biol. Chem.* 255, 5511–5513.
- (39) Backman, L. R. F., Funk, M. A., Dawson, C. D., and Drennan, C. L. (2017) New tricks for the glycy radical enzyme family. *Crit. Rev. Biochem. Mol. Biol.* 52, 674–695.
- (40) Selmer, T., Pierik, A. J., and Heider, J. (2005) New glycy radical enzymes catalysing key metabolic steps in anaerobic bacteria. *Biol. Chem.* 386, 981–988.
- (41) Levin, B. J., Huang, Y. Y., Peck, S. C., Wei, Y., Martinez-del Campo, A., Marks, J. A., Franzosa, E. A., Huttenhower, C., and Balskus, E. P. (2017) A prominent glycy radical enzyme in human gut microbiomes metabolizes trans-4-hydroxy-L-proline. *Science* 355, eaai8386.

- (42) Koppel, N., Maini Rekdal, V., and Balskus, E. P. (2017) Chemical transformation of xenobiotics by the human gut microbiota. *Science* 356, eaag2770.
- (43) Hertel, L. W., Boder, G. B., Kroin, J. S., Rinzel, S. M., Poore, G. A., Todd, G. C., and Grindey, G. B. (1990) Evaluation of the antitumor activity of gemcitabine (2', 2'-difluoro-2'-deoxycytidine). *Cancer Res.* 50, 4417–4422.
- (44) Wang, J., Lohman, G. J. S., and Stubbe, J. (2009) Mechanism of Inactivation of Human Ribonucleotide Reductase with p53R2 by Gemcitabine 5'-Diphosphate. *Biochemistry* 48, 11612–11621.
- (45) Reich, H. J., and Hondal, R. J. (2016) Why Nature Chose Selenium. *ACS Chem. Biol.* 11, 821–841.
- (46) McMillen, D. F., and Golden, D. M. (1982) Hydrocarbon bond dissociation energies. *Annu. Rev. Phys. Chem.* 33, 493–532.
- (47) Mozziconacci, O., Williams, T. D., Kerwin, B. A., and Schoneich, C. (2008) Reversible intramolecular hydrogen transfer between protein cysteine thiol radicals and  $\alpha$ C–H bonds in insulin: control of selectivity by secondary structure. *J. Phys. Chem. B* 112, 15921–15932.
- (48) Rauk, A., Yu, D., Taylor, J., Shustov, G. V., Block, D. A., and Armstrong, D. A. (1999) Effects of structure on  $\alpha$ C–H bond enthalpies of amino acid residues: relevance to H transfers in enzyme mechanisms and protein oxidation. *Biochemistry* 38, 9089–9096.
- (49) Dang, H.-S., Roberts, B. P., and Tocher, D. A. (2001) Selective radical-chain epimerization at electron-rich chiral tertiary C–H centres using thiols as protic polarity reversal catalysts. *J. Chem. Soc., Perkin Trans. 1*, 2452–2461.
- (50) Reid, D. L., Shustov, G. V., Armstrong, D. A., Rauk, A., Schuchmann, M. N., Akhlaq, S., and von Sonntag, C. (2002) H-atom abstraction from thiols by C-centered radicals: a theoretical and experimental study of reaction rates. *Phys. Chem. Chem. Phys.* 4, 2965–2974.
- (51) Becker, A., Fritz-Wolf, K., Kabsch, W., Knappe, J., Schultz, S., and Volker Wagner, A. F. (1999) Structure and mechanism of the glycyl radical enzyme pyruvate formate-lyase. *Nat. Struct. Biol.* 6, 969–975.
- (52) Becker, A., and Kabsch, W. (2002) X-ray structure of pyruvate formate-lyase in complex with pyruvate and CoA. How the enzyme uses the Cys-418 thiol radical for pyruvate cleavage. *J. Biol. Chem.* 277, 40036–40042.
- (53) Argirević, T., Riplinger, C., Stubbe, J., Neese, F., and Bennati, M. (2012) ENDOR Spectroscopy and DFT Calculations: Evidence for the Hydrogen-Bond Network Within  $\alpha 2$  in the PCET of *E. coli* Ribonucleotide Reductase. *J. Am. Chem. Soc.* 134, 17661–17670.
- (54) Nick, T. U., Lee, W., Koßmann, S., Neese, F., Stubbe, J., and Bennati, M. (2015) Hydrogen Bond Network between Amino Acid Radical Intermediates on the Proton-Coupled Electron Transfer Pathway of *E. coli*  $\alpha 2$  Ribonucleotide Reductase. *J. Am. Chem. Soc.* 137, 289–298.
- (55) Uhlin, U., and Eklund, H. (1994) Structure of ribonucleotide reductase protein R1. *Nature* 370, 533–539.
- (56) Yokoyama, K., Uhlin, U., and Stubbe, J. (2010) Site-Specific Incorporation of 3-Nitrotyrosine as a Probe of pK<sub>a</sub> Perturbation of Redox-Active Tyrosines in Ribonucleotide Reductase. *J. Am. Chem. Soc.* 132, 8385–8397.
- (57) O'Keefe, J. P., Dustin, C. M., Barber, D., Snider, G. W., and Hondal, R. J. (2018) A “Seleno Effect” Differentiates the Roles of Redox Active Cysteine Residues in *Plasmodium falciparum* Thio-redoxin Reductase. *Biochemistry* 57, 1767–1778.
- (58) Ge, J., Yu, G., Ator, M. A., and Stubbe, J. (2003) Pre-Steady-State and Steady-State Kinetic Analysis of *E. coli* Class I Ribonucleotide Reductase. *Biochemistry* 42, 10071–10083.
- (59) Licht, S., and Stubbe, J. (1999) Mechanistic Investigations of Ribonucleotide Reductase. *Comp. Nat. Prod. Chem.* 5, 163–203.
- (60) van der Donk, W., Yu, G., Silva, D. J., Stubbe, J., McCarthy, J. R., Jarvi, E. T., Matthews, D. P., Resvick, R. J., and Wagner, E. (1996) Inactivation of Ribonucleotide Reductase by (E)-2'-Fluoromethylene-2'-deoxycytidine 5'-Diphosphate: A Paradigm for Nucleotide Mechanism-Based Inhibitors. *Biochemistry* 35, 8381–8391.
- (61) Persson, A. L., Eriksson, M., Katterle, B., Pötsch, S., Sahlin, M., and Sjöberg, B. M. (1997) A New Mechanism-based Radical Intermediate in a Mutant R1 Protein Affecting the Catalytically Essential Glu441 in *Escherichia coli* Ribonucleotide Reductase. *J. Biol. Chem.* 272, 31533–31541.