

## **UC Irvine**

### **Western Journal of Emergency Medicine: Integrating Emergency Care with Population Health**

#### **Title**

Achieving the Triple Aim Through Informed Consent for Computed Tomography

#### **Permalink**

<https://escholarship.org/uc/item/9g14874z>

#### **Journal**

Western Journal of Emergency Medicine: Integrating Emergency Care with Population Health, 16(7)

#### **ISSN**

1936-900X

#### **Authors**

Carney, Dylan  
Rodriguez, Robert M.

#### **Publication Date**

2015

#### **DOI**

10.5811/westjem.2015.12.29466

#### **Copyright Information**

Copyright 2015 by the author(s). This work is made available under the terms of a Creative Commons Attribution License, available at <https://creativecommons.org/licenses/by/4.0/>

Peer reviewed

# Achieving the Triple Aim Through Informed Consent for Computed Tomography

Dylan Carney, MD  
Robert M. Rodriguez, MD

University of California, San Francisco, Department of Emergency Medicine, San Francisco, California

*Section Editor:* Mark I. Langdorf, MD, MHPE

Submission history: Submitted December 4, 2015; Revision received December 8, 2015; Accepted December 9, 2015

Electronically published December 16, 2015

Full text available through open access at [http://escholarship.org/uc/uciem\\_westjem](http://escholarship.org/uc/uciem_westjem)

DOI: 10.5811/westjem.2015.12.29466

[West J Emerg Med. 2015;16(7):1030-1032.]

At the end of a particularly busy shift, you meet Mary, a 24 year-old female with no past medical history, who presents with six hours of crampy, intermittent, periumbilical abdominal pain but no associated fever, nausea, vomiting, diarrhea or anorexia. Her vital signs are normal and her abdominal and gynecological exams are notable only for mild, diffuse abdominal tenderness without rebound or guarding. Her lab results and urinalysis are unremarkable, and her pain improves somewhat with intravenous pain medications. You explain to the patient that you have a low suspicion for an intraabdominal emergency, but cannot be certain without a computed tomography (CT) scan. "I'll do whatever you recommend," she replies. The patient ultimately gets a CT, which is normal, and she is discharged 30 minutes later with a diagnosis of nonspecific abdominal pain.

Emergency department (ED) clinicians see patients like Mary every day- patients for whom our experience and clinical suspicion of a truly emergent condition is low, albeit not negligible, and for whom there are no consensus, evidence-based guidelines or algorithms to guide the use of advanced imaging. The decision to image is, at times, dictated by systems-factors, such as difficulties arranging for adequate follow-up. Fear of litigation has been cited as a common driver of excessive diagnostic testing.<sup>1</sup> Ultimately, diagnostic CT use in the ED is on a steep rise, which, combined with unchanging prevalence of disease, results in greater exposure to CT risks and costs with lower corresponding diagnostic yield.<sup>2-5</sup>

The risks (and costs) of CT are undeniably real and should not be ignored or minimized. Extrapolating from data from atomic bomb survivors, Smith-Bindman, Brenner and others have calculated the risks of cancer development associated with CT and estimated that up to 2% of all cancers in the US are attributable to CT scans.<sup>6,7</sup> Beyond these theoretical extrapolations, Mathews and others demonstrated a dose-response risk of cancer development associated with CT scans in a large cohort of patients in Australia.<sup>8</sup>

Concurring with these concerns about CT, a number of professional medical societies and various governing bodies (BEIR- The National Research Council's Biological Effects of Ionizing Radiation VII Report, UNSCEAR- The United Nations Scientific Committee on the Effects of Atomic Radiation, and IRCP- The International Commission on Radiological Protection) have targeted cutbacks in unnecessary CT imaging and reduction of radiation doses with CT as specialty-wide goals.<sup>9-12</sup> The American College of Surgeons and the American College of Emergency Physicians have both included reductions in CT as part of their Choosing Wisely campaigns.<sup>13,14</sup>

Merck and colleagues in this issue of *Western Journal of Emergency Medicine*, note the risks of CT, comparing them to the risks associated with blood transfusion (for which clinicians universally obtain informed consent).<sup>15</sup> This is a thought-provoking and rational link that, combined with the fact that patients (and physicians) are unaware of CT risks, leads to the logical next-step of their novel trial of informed consent for abdominal/pelvic CT. In this multiphase, observational cohort study, the authors collected data on abdominal pain patients and built a multivariate logistic regression model to assess probability of CT utilization as a function of history, exam findings, diagnostic testing and disposition. Patients who had CT scans were included in a second multivariable model that estimated the likelihood of having a positive scan. In the next phase of the study, emergency providers used a one-page, standardized, written informed consent tool, which included potential biological risks and diagnostic benefits of CT, to engage patients in shared decision making. The authors report that the tool took less than one minute to use and was minimally disruptive to provider workflow. Patients in this implementation phase were stratified as "low" or "high" risk based on clinical factors (focal or rebound tenderness and the presence of a rigid abdomen). The investigators then built a logistic regression model to assess CT utilization among the low

and high risk groups after controlling for confounders with propensity score matching.<sup>15</sup>

Their results were striking. While CT utilization was unaffected in the high-risk group, the investigators noted a 24% reduction in CT utilization among low-risk patients after the implementation of their written informed consent protocol. Notably, they found no difference in adverse events or patient return visits within 30 days among the nearly 4,000 patients included in the study, indicating that their protocol was both safe and effective.<sup>15</sup>

Overall, these findings introduce a novel way to improve patient-centered care. Their informed-consent intervention was simple, safe, fast, and effective, achieving several goals of the Institute for Health Improvement's Triple Aims: (1) improving population health, (2) reducing costs and (3) enhancing the patient experience.<sup>16</sup> Decreasing the rate of low-yield CT scans may decrease costs and improve outcomes by minimizing unnecessary ionizing radiation and decreasing potentially unnecessary workups of incidental findings. At least as important are the potential positive effects of shared decision-making. Prior investigators have shown that patients want to be informed of the risks (and costs) of CT whenever possible and that many patients would prefer to avoid imaging when their risk of life-threatening injury is low.<sup>17</sup> Informed consent for CT may therefore improve patient satisfaction and patient care experience.

Should informed consent for CT be routinely obtained? Should it even become a standard of care? Although there are certainly some cases in which consent may not be feasible, most scenarios for CT use in the ED are truly not emergent enough to preclude informed consent/shared decision-making. We await further examinations of this important topic, most notably a large, multicenter study of shared-decision making in pediatric head trauma.<sup>18</sup> In the meantime, the work of Merck et al is compelling enough that we would advocate providing consent for CT in low risk abdominal pain cases.

---

*Address for Correspondence:* Dylan Carney, MD, University of California, San Francisco, Department of Emergency Medicine, San Francisco, California. Email:

*Conflicts of Interest:* By the WestJEM article submission agreement, all authors are required to disclose all affiliations, funding sources and financial or management relationships that could be perceived as potential sources of bias. The authors disclosed none.

*Copyright:* © 2015 Carney et al. This is an open access article distributed in accordance with the terms of the Creative Commons Attribution (CC BY 4.0) License. See: <http://creativecommons.org/licenses/by/4.0/>

---

## REFERENCES

- Bishop TF, Federman AD, Keyhani S. Physicians' views on defensive medicine: a national survey. *Arch Intern Med.* 2010;170(12):1081-1083.
- Korley FK, Pham JC, Kirsch TD. Use of advanced radiology during visits to US emergency departments for injury-related conditions, 1998-2007. *JAMA.* 2010;304(13):1465-1471.
- Broder J, Warshauer DM. Increasing utilization of computed tomography in the adult emergency department, 2000-2005. *Emerg Radiol.* 2006;13(1):25-30.
- Larson DBB, Johnson LWW, Schnell BMM, et al. National Trends in CT Use in the Emergency Department: 1995-2007. *Radiology.* 2011;258(1527-1315 (Electronic)):164-173.
- Hussein W, Mullins PM, Alghamdi K, et al. Trends in Advanced Computed Tomography Use for Injured Patients in United States Emergency Departments: 2007-2010. *Acad Emerg Med.* 2015;22(6):663-669.
- Smith-Bindman R, Lipson J, Marcus R, et al. Radiation Dose Associated With Common Computed Tomography Examinations and the Associated Lifetime Attributable Risk of Cancer. *Arch Intern Med.* 2009;169(22):2078-2086.
- Brenner DJ, Hall EJ. Computed tomography--an increasing source of radiation exposure. *N Engl J Med.* 2007;357(22):2277-2284.
- Mathews JD, Forsythe AV, Brady Z, et al. Cancer risk in 680 000 people exposed to computed tomography scans in childhood or adolescence: data linkage study of 11 million Australians. *BMJ.* 2013;346.
- Lam DL, Larson DB, Eisenberg JD, et al. Communicating Potential Radiation-Induced Cancer Risks From Medical Imaging Directly to Patients. *Am J Roentgenol.* 2015;1-9.
- Report of the United Nations Scientific Committee on the Effects of Atomic Radiation to the General Assembly*;2013. [http://www.unscear.org/docs/reports/2013/13-85418\\_Report\\_2013\\_GA\\_Report.pdf](http://www.unscear.org/docs/reports/2013/13-85418_Report_2013_GA_Report.pdf).
- IRCP. *Proceedings of the Second International Symposium on the System of Radiological Protection.* Vol 43.;2014.
- Committee to Assess Health Risks from Exposure to Low Levels of Ionizing Radiation. *Health Risks from Exposure to Low Levels of Ionizing Radiation - BEIR VII Phase 2.*; 2006.
- ACEP. American College of Emergency Physicians; Five Things Physicians and Patients Should Question. American Board of Internal Medicine, Choosing Wisely Campaign. Available at: <http://www.choosingwisely.org/wp-content/uploads/2015/02/ACEP-Choosing-Wisely-List.pdf>. Published 2013. Accessed Dec 1, 2015.
- American College of Surgeons. American College of Surgeons; Five Things Physicians and Patients Should Question. American Board of Internal Medicine, Choosing Wisely Campaign. Available at: <https://www.facs.org/~media/files/education/acclist.ashx>. Published 2013. Accessed Dec 1, 2015.
- Merck LH, Ward LA, Applegate KE. Written Informed Consent for Computed Tomography of the Abdomen / Pelvis is Associated with Decreased CT Utilization in Low-Risk Emergency Department Patients. 2006.
- Institute for Healthcare Improvement. The Triple Aim. *Healthc Exec.* 2009;64-66. [www.ache.org](http://www.ache.org).
- Rodriguez RM, Henderson TM, Ritchie AM, et al. Patient preferences

- and acceptable risk for computed tomography in trauma. *Injury*. 2014;45(9):1345-1349.
18. Hess EP, Wyatt KD, Kharbanda AB, et al. Effectiveness of the head CT choice decision aid in parents of children with minor head trauma: study protocol for a multicenter randomized trial. *Trials*. 2014;15(1):253.