

Lawrence Berkeley National Laboratory

Recent Work

Title

GROUND-STATE ELECTRONIC STRUCTURE OF ERBIUM AND NUCLEAR SPINS OF Er169, Er171 AND GD159

Permalink

<https://escholarship.org/uc/item/9hs9039p>

Authors

Cabezas, Amado
Lindgren, Ingvar
Marrus, Richard.

Publication Date

1960-04-04

DISCLAIMER

This document was prepared as an account of work sponsored by the United States Government. While this document is believed to contain correct information, neither the United States Government nor any agency thereof, nor the Regents of the University of California, nor any of their employees, makes any warranty, express or implied, or assumes any legal responsibility for the accuracy, completeness, or usefulness of any information, apparatus, product, or process disclosed, or represents that its use would not infringe privately owned rights. Reference herein to any specific commercial product, process, or service by its trade name, trademark, manufacturer, or otherwise, does not necessarily constitute or imply its endorsement, recommendation, or favoring by the United States Government or any agency thereof, or the Regents of the University of California. The views and opinions of authors expressed herein do not necessarily state or reflect those of the United States Government or any agency thereof or the Regents of the University of California.

Ground-State Electronic Structure of Erbium and Nuclear Spins of Er^{169} ,
 Er^{171} , and Gd^{159} . * AMADO Y. CABEZAS, INGVAR P. K. LINDGREN,**

AND RICHARD MARRUS, Lawrence Radiation Laboratory and Department

of Physics, Berkeley, California. -- Atomic-beam magnetic resonance

studies of neutron-activated erbium and gadolinium have yielded three

nuclear spins. These are Er^{169} ($I=1/2$), Er^{171} ($I=5/2$) and Gd^{159} ($I=3/2$).

These measurements are in agreement with the assignments made from
beta-ray spectroscopy and with the level diagram proposed by Nilsson.¹

Measurements of the erbium spins were made by observing the Zeeman

transitions in a state characterized by $J = 6$ and $g_J = 1.164(5)$. It is inferred

that this state almost certainly arises from almost pure L-S coupling among

the electrons of the configuration $(4f)^{12} (6s)^2$ to the Hund's Rule ground state

3H_6 . Observations of gadolinium resonances were made in the five J states

arising from the ground state 9D . The observed resonances are in agree-
ment with g_J values previously measured.²

* This research was sponsored by the Atomic Energy Commission and the
Office of Naval Research.

** On leave from the Institute of Physics, University of Uppsala, Uppsala, Sweden.

¹ S. G. Nilsson, Kgl. Danske Videnskab. Selskab Mat.-fys. Medd. 29, No. 16
(1955).

² K. F. Smith and I. J. Spaulding (Cambridge University, Cambridge, England),
private communication.

A. P. S. Meeting-Montreal, Canada, June 1960.