

Lawrence Berkeley National Laboratory

Recent Work

Title

RYDBERG SPECTROSCOPY OF H₂ VIA STEPWISE RESONANT TWO-PHOTON ION-PAIR (H⁺+H⁻) PRODUCTION

Permalink

<https://escholarship.org/uc/item/9j15x936>

Author

Kung, A.H.

Publication Date

1985-10-01



Lawrence Berkeley Laboratory

UNIVERSITY OF CALIFORNIA

Materials & Molecular Research Division

RECEIVED
LAWRENCE
BERKELEY LABORATORY
DEC 12 1985
PHYSICS AND
ASTRONOMY SECTION

Submitted to Physical Review Letters

RYDBERG SPECTROSCOPY OF H₂ VIA STEPWISE RESONANT
TWO-PHOTON ION-PAIR (H⁺+H⁻) PRODUCTION

A.H. Kung, R.H. Page, R.J. Larkin, Y.R. Shen,
and Y.T. Lee

October 1985

For Reference
Not to be taken from this room



DISCLAIMER

This document was prepared as an account of work sponsored by the United States Government. While this document is believed to contain correct information, neither the United States Government nor any agency thereof, nor the Regents of the University of California, nor any of their employees, makes any warranty, express or implied, or assumes any legal responsibility for the accuracy, completeness, or usefulness of any information, apparatus, product, or process disclosed, or represents that its use would not infringe privately owned rights. Reference herein to any specific commercial product, process, or service by its trade name, trademark, manufacturer, or otherwise, does not necessarily constitute or imply its endorsement, recommendation, or favoring by the United States Government or any agency thereof, or the Regents of the University of California. The views and opinions of authors expressed herein do not necessarily state or reflect those of the United States Government or any agency thereof or the Regents of the University of California.

RYDBERG SPECTROSCOPY OF H_2 VIA STEPWISE RESONANT
TWO-PHOTON ION-PAIR ($H^+ + H^-$) PRODUCTION

A.H. Kung

San Francisco Laser Center, Department of Chemistry
University of California, Berkeley, California 94720

Ralph H. Page^a, R.J. Larkin^b, Y.R. Shen^a, and Yuan T. Lee^c
Materials and Molecular Research Division
Lawrence Berkeley Laboratory, University of California
Berkeley, California 94720


ABSTRACT

High resolution spectra of new Rydberg states of H_2 in the XUV region were obtained by two-step doubly-resonant excitation followed by H^- ion detection. Resolved Fano line profiles showing a sign reversal of the profile index along the Rydberg series were observed for the first time in H_2 . Dynamical information such as lifetimes and relative transition strengths was also derived.

Recently, extensive spectroscopic work on H_2 , especially in the range of photon energy below 17eV, has led to a good understanding of the autoionization process through Rydberg transitions that converge to $H_2^+(X^2\Sigma_g^+)$ and up to $v^+=7.1-5$. Extension of the same level of understanding to series that converge to higher vibrational states ($v^+>7$) and to other important processes such as ion-pair formation and dissociative ionization is, however, inhibited by the limited amount of data available above 17eV. Spectroscopic study of these highly excited states is made difficult by weak transition probability and the lack of intense light sources at these frequencies. With the recent advances in the development of tunable coherent XUV sources,^{6,7} it is now possible to apply nonlinear optical spectroscopy to the investigation of these highly excited states. By using resonant two-photon excitation,⁸ the weak transition probabilities due to small Franck-Condon factors for single-photon excitation can be greatly improved. New excited states whose transitions cannot be reached from the ground state can be studied. The spectrum can also be simplified by selective labelling of the rotational states. In this paper, we report the first application of two-step doubly-resonant excitation followed by negative (H^-) ion detection to study the Rydberg states of H_2 above 17eV. High resolution spectra of new Rydberg states with high convergence limits (to $H_2^+ X^2\Sigma_g^+, v^+=9, N^+=1,3$) were obtained and analyzed. The Fano line profiles were resolved and reversal of the profile index along the series was observed, the

latter for the first time in H_2 . Dynamical information such as lifetimes and relative transition strengths from low electronic states to Rydberg states and the ion-pair dissociation continuum was deduced. Use of H^- detection was essential in this study because the alternative method of probing these Rydberg states via autoionization and detecting H_2^+ often yields a very poor spectrum dominated by an intense ionization continuum. The H^- continuum arising from direct ion-pair formation, on the other hand, is relatively weak, rendering the observation of the Rydberg structure comparatively easy. Ion-pair formation via single-photon excitation was previously used by Chupka, et al. to study Rydberg states of H_2 .⁹ No lineshape or dynamical information was reported because of weaker signal and lower resolution than in the experiment reported here.

Described below are the two-photon excitation scheme and the subsequent ionization or dissociation of H_2 applicable to this experiment (see Figure 1):



The dominant channel in the region of interest is the formation of H_2^+ by direct ionization. The threshold for this process is 15.43eV. In the range of a few eV above this threshold, the spectrum obtained by detecting H_2^+ consists of auto-

ionizing lines superimposed on a continuum; the latter generally dominates as the excitation energy increases.² Ion-pair formation has a threshold at 17.34eV. The intensity of the ion-pair channel immediately above its threshold is low. The interference of Rydberg states with this ion-pair continuum allows us to probe the Rydberg line spectrum in this region with high accuracy by measuring the excitation frequency dependence of the H^- ion signal.⁹

In the experiment, the frequency-tripled output of a frequency-doubled Nd:YAG laser-pumped dye laser (laser 1) was tuned to selectively excite either the $v'=2, J'=1$ level of the $C^1\Pi_u$ state of a pulsed beam of H_2 at $103509.24 \text{ cm}^{-1}$ or the $v'=12, J'=0$ level of the $B^1\Sigma_u^+$ state at $103700.91 \text{ cm}^{-1}$.¹⁰ From these levels, the molecules were further excited by a second laser (laser 2). This second laser was a frequency-doubled Nd:YAG laser-pumped dye laser, which was tuned to have the sum of the frequencies of lasers 1 and 2 span the region from below the ion-pair formation threshold to well above that threshold. The H^- ions produced were detected by a time-of-flight mass spectrometer. The ion signals were averaged using a boxcar-integrator and digitized for analysis. The signal level was as big as 100 ion counts per laser pulse. Wavelength calibration of laser 2 was performed using the optogalvanic effect with an argon discharge lamp.

Figure 2(a) presents a portion of the spectrum obtained by tuning laser 2 with $C^1\Pi_u$ ($v'=2, J'=1$) as the intermediate state. It exhibits many sharp dips superimposed on a structured continuum. Analysis shows that the dips can be assigned to transitions to the

high n states of two distinct Rydberg series converging to the ion core levels specified by the rotational quantum numbers $N^+=1$ and $N^+=3$ and the vibrational quantum number $v^+=9$. The effective principal quantum number n^* of the observed states were determined and are listed in Table 1. Here, n^* is defined by the Rydberg series formula $\nu = S - R/(n^*)^2$, where ν is the measured frequency in inverse centimeters, S is the theoretical series limit given by Wind¹¹ and $R=109707.42\text{cm}^{-1}$ is the Rydberg constant for H_2 . Our present signal-to-noise ratio limits the determination of n^* to ~ 36 . Figure 2(b) gives the spectrum in the same region with $B^1\Sigma_u^+$ ($v'=12, J'=0$) as the intermediate state. The lines in this spectrum can be identified as arising from transition to the same Rydberg series displayed in Fig. 2(a). Their position relative to the ionization level of the series ($v^+=9, N^+=1,3$) agree with those deduced from Figure 2(a) to within 1 cm^{-1} , which is well within our experimental resolution of 1.5 cm^{-1} .

From Table 1, the quantum defect for the $N^+=1$ series is estimated to be 0.1 ± 0.1 , whereas that for the $N^+=3$ series is 0.7 ± 0.15 . The perturbations among adjacent Rydberg states in H_2 , the limited signal-to-noise ratio for the higher n states, and the limited frequency resolution have prevented us from determining the orbital quantum number (s or d) of these series and from obtaining finer limits for the quantum defects. However, these measurements are an important first step in providing data for spectroscopists to use to improve our understanding of these high-lying states of H_2 . Some additional lines which do not

belong to these series also appear in the spectra. These, along with the broad structures in the spectra, are not yet assigned. The broad structures are likely to arise from transitions to low n states of Rydberg series with v^+ higher than 9.

While the Rydberg line positions in the two cases with different intermediate states agree, the lineshapes observed are very different. Those in Fig. 2(a) are sharp dips but those in Fig. 2(b) have asymmetric Fano profiles with changing profile indices. We have used Fano's lineshape formula^{12,13} which takes into account the interaction of discrete autoionizing states with a continuum, to fit individually the observed lines with $n \geq 25$ in Figure 2 (those lying above the $v^+=9$, $N^+=1$ series limit of $139746.58 \text{ cm}^{-1}$). The results for Fig. 2(b) are shown in Fig. 3. Values of the profile index q and linewidths deduced are given in Table 2. Here $q = A * |\langle i/z/n \rangle| / |\langle i/z/a \rangle|$, where $\langle i/z/n \rangle$ is the dipole matrix element from the intermediate state $\langle i \rangle$ to the discrete autoionizing state $\langle n \rangle$, $\langle i/z/a \rangle$ the matrix element from $\langle i \rangle$ to the continuum state $\langle a \rangle$, and A is independent of state $\langle i \rangle$ and is inversely proportional to the strength of the interaction between states $\langle n \rangle$ and $\langle a \rangle$. The laser linewidth has been deconvolved to obtain the transition widths. We did not try to fit the lines with $n < 25$ because their profiles are distorted by strong perturbation between the $N^+=1$ and $N^+=3$ series. The lifetimes obtained from the inverse linewidths are also listed in Table 2.

An interesting result from the lineshape analysis is the revelation of a change of the sign of the profile index q (q

reversal) at $n \geq 31$ (see Figure 3). This sign reversal arises from a relative phase shift in the two transition matrix elements that determine q . It therefore must be the result of the interaction of the $N^+=3$ series with a broad state of finite bandwidth¹⁴ which appears to be centered around the $n=30$ transition. Unfortunately, the origin of this broad state is not yet clear. Connerade has previously reported evidence of similar q -reversal in the Rydberg spectra of Tl¹⁴ and of CH₃Br.¹⁵ This is, however, the first time that q reversal has been observed for H₂ or any diatomic molecule.

The observed lineshapes also reveal important information about the relative transition probabilities between high-lying electronic states of H₂. It is known that $C^1\Pi_u$ is a "Rydberg-like" state ($n=2$) whereas $B^1\Sigma_u^+$ is an "ion-pair-like" state.¹⁶ This suggests transitions from $B^1\Sigma_u^+$ to the ion-pair continuum should be stronger than those from $C^1\Pi_u$ to the ion-pair continuum.

This was indeed observed in our experiment: the H⁻ continuum signal was ten times stronger when exciting via the B level than via the C level. On the other hand, one would also expect transitions from $C^1\Pi_u$ to the high-lying Rydberg states to be stronger than those from $B^1\Sigma_u^+$ to the Rydberg states. Experimentally, however, we found that the reverse was true, as implied by the values of the Fano parameters q listed in Table II. The result $q \approx 0$ for the $C^1\Pi_u$ case indicates that the transitions from $C^1\Pi_u$ to the Rydberg states are even weaker by about an order of magnitude than the transitions from $B^1\Sigma_u^+$ to the ion-pair continuum. This can

be easily understood if we realize that in our case, transitions from $C^1\Pi_u$ to the Rydberg states should have small Franck-Condon factors because of the larger change in the vibrational quantum number ($v=2 \leftrightarrow v=9$). If it were not for the Fano coupling between the Rydberg states and the ion-pair continuum, we would not have observed these transitions. For the transitions from B to the Rydberg states, Δv is only 3 in our case. This is consistent with the observation of Pratt, et al., who, using multiphoton photoelectron spectroscopy, have shown a near unity Franck-Condon factor for transitions from $C^1\Pi_u$ to high-lying Rydberg states with $\Delta v=0$.¹⁷

We have reported here the first successful attempt in extending high-resolution spectroscopy to the XUV region above 17eV using the technique of doubly resonant excitation coupled with negative ion detection. Interesting new information about high-lying Rydberg states of H_2 and their coupling with the ion-pair continuum was obtained. The technique could be extended to shorter wavelength regions for H_2 and is generally applicable to studies of highly-excited states of other molecules.

Acknowledgement

This work was supported by the Director, Office of Energy Research, Office of Basic Energy Sciences of the U.S. Department of Energy under contract DE-ACO3-76SF-00098. A. H. K. was also supported by the National Science Foundation under grant CHE 83-03208.

^aAlso affiliated with the Department of Physics, University of California, Berkeley.

^bPresent address: United Technologies Research Center, Optics and Applied Technology Laboratory, West Palm Beach, FL 33402.

^cAlso affiliated with the Department of Chemistry, University of California, Berkeley.

REFERENCES

1. G. Herzberg and Ch. Jungen, *J. Mol Spectrosc.* 41, 425 (1972).
2. P.M. Dehmer and W.A. Chupka, *J. Chem. Phys.* 65, 2243, (1976).
3. Ch. Jungen and D. Dill, *J. Chem. Phys.* 73, 3338 (1980).
4. M. Raoult and Ch. Jungen, *J. Chem Phys.* 74, 3388 (1981).
5. H. Rottke and K. H. Welge, *J. Physique, Colloque* C1, 127 (1985).
6. E.E. Marinero, C.T. Rettner, R.N. Zare, and A.H. Kung, *Chem. Phys. Lett.* 95, 486 (1983), and C.T. Rettner, E.E. Marinero, R.N. Zare, and A.H. Kung, *J. Phys. Chem.* 88, 4459 (1984).
7. R. Hilbig, A. Lago, and R. Wallenstein, *Optics Comm.* 49, 297 (1984).
8. D.J. Kligler and C.K. Rhodes, *Phys. Rev. Lett.* 40, 309 (1978).
9. W.A. Chupka, P.M. Dehmer, and W.T. Jivery, *J. Chem. Phys.* 63, 3929 (1975).
10. I. Dabrowski and G. Herzberg, *Can. J. Phys.* 52, 1110 (1974).
11. H. Wind, *J. Chem. Phys.* 43, 2956 (1956).
12. U. Fano and J.W. Cooper, *Rev. Mod. Phys.* 40, 441 (1968).
13. M.B. Robin, "Higher Excited States of Polyatomic Molecules," p. 32, Academic Press, New York (1974).
14. J.P. Connerade, *Proc. R. Soc. A* 362, 361 (1978).
15. J.P. Connerade, M.A. Baig, and S.P. McGlynn, *J. Phys. B* 14, L 67 (1981).
16. G. Herzberg, "Spectra of Diatomic Molecules," Van Nostrand Reinhold, 2nd edition, New York, 1950.
17. S.T. Pratt, P.M. Dehmer, and J.L. Dehmer, *Chem. Phys. Lett.*, 105, 28 (1984).
18. T.E. Sharp, *Atomic Data* 2, 119 (1971).

Table 1: Rydberg State Identification in H₂

Line Position cm ⁻¹	n*
Series converging to H ₂ ⁺ (v ⁺ =9, N ⁺ =1) at 139746.58 cm ⁻¹	
139581.8	25.80
139598.0	27.17
139608.1	28.15
139617.2	29.12
139625.6	30.11
139633.8	31.19
139640.2	32.11
139645.8	32.99
139651.5	33.97
Series converging to H ₂ ⁺ (v ⁺ =9, N ⁺ =3) at 139910.77 cm ⁻¹	
139604.5	18.93
139633.9	19.91
139661.9	21.00
139681.1	21.86
139700.4	22.84
139716.8	23.78
139732.5	24.81
139746.5	25.84
139757.6	26.76
139768.2	27.74
139777.5	28.69
139786.0	29.65
139794.0	30.65
139801.8	31.73
139808.6	32.77
139814.1	33.69
139819.0	34.58
139824.2	35.60

Table 2: Fano Profile Parameters for Rydberg Series
 Converging to $H_2^+(v^+=9, N^+=3)$

n	linecenter(± 0.5) cm^{-1}	linewidth cm^{-1}	lifetime psec	$q(+0.05)$ ($C^1\Pi_u$) ^a	$q(\pm 0.05)$ ($B^1\Sigma_u^+$) ^a
25	139746.5	3.0 \pm 0.5	1.7	b	0.40
26	139757.6	1.5 \pm 0.3	3.5	0.00	0.2
27	139768.2	1.2 \pm 0.2	4.3	0.00	0.60
28	139777.5	0.8 \pm 0.1	6.5	-0.10	0.55
29	139786.0	0.6 \pm 0.1	8.7	0.10	0.10
30	139794.0	2.0 \pm 0.3	2.6	0.00	0.00
31	139801.8	1.4 \pm 0.2	3.7	0.10	-0.30
32	139808.6	3.0 \pm 0.5	1.7	-0.10	-0.30
33	139814.1	0.4 \pm 0.1	13.1	0.00	-0.80
34	139819.0	0.3 \pm 0.1	17.5	b	-0.80
35	139824.2	1.1 \pm 0.2	4.7	b	-0.15

^aintermediate state

^bnot measured

FIGURE CAPTIONS

- Fig. 1: Potential energy diagram for H_2 showing the levels of interest, from Sharp.¹⁸
- Fig. 2: Resonant two-photon ion-pair spectra of H_2 from the onset of H^- production threshold, indicating transitions assigned to series that converge to $H_2^+(X^2\Sigma_g^+, v^+=9, n^+=1,3)$: (a) intermediate state: $C^1\Pi_u (v'=2, J'=1)$. (b) intermediate state: $B^1\Sigma_u^+ (v'=12, J'=0)$. Spectrum (a) is obtained with 20 times higher gain than (b).
- Fig. 3: Expanded ion pair production spectrum of H_2 between the $N^+=1$ and $N^+=3$ series limits showing the fitted Fano line profiles: ————calculated profile, ● experimental points. Arrows indicate fitted linecenter position.




Figure 1

XBL 857-6414A




Figure 2a





Figure 2b

139549 622 694 767 139840
TOTAL PHOTON ENERGY (cm^{-1})

Figure 3



This report was done with support from the Department of Energy. Any conclusions or opinions expressed in this report represent solely those of the author(s) and not necessarily those of The Regents of the University of California, the Lawrence Berkeley Laboratory or the Department of Energy.

Reference to a company or product name does not imply approval or recommendation of the product by the University of California or the U.S. Department of Energy to the exclusion of others that may be suitable.

*LAWRENCE BERKELEY LABORATORY
TECHNICAL INFORMATION DEPARTMENT
UNIVERSITY OF CALIFORNIA
BERKELEY, CALIFORNIA 94720*