

# UCLA

## UCLA Previously Published Works

### Title

GEMINI PLANET IMAGER OBSERVATIONS OF THE AU MICROSCOPII DEBRIS DISK:  
ASYMMETRIES WITHIN ONE ARCSECOND

### Permalink

<https://escholarship.org/uc/item/9k21k2z5>

### Journal

The Astrophysical Journal Letters, 811(2)

### ISSN

2041-8205

### Authors

Wang, Jason J  
Graham, James R  
Pueyo, Laurent  
[et al.](#)

### Publication Date

2015-10-01

### DOI

10.1088/2041-8205/811/2/119

Peer reviewed

## GEMINI PLANET IMAGER OBSERVATIONS OF THE AU MICROSCOPII DEBRIS DISK: ASYMMETRIES WITHIN ONE ARCSECOND

JASON J. WANG<sup>1</sup>, JAMES R. GRAHAM<sup>1</sup>, LAURENT PUEYO<sup>2</sup>, ERIC L. NIELSEN<sup>3,4</sup>, MAX MILLAR-BLANCHAER<sup>5</sup>, ROBERT J. DE ROSA<sup>1</sup>,  
PAUL KALAS<sup>1</sup>, S. MARK AMMONS<sup>6</sup>, JOANNA BULGER<sup>7</sup>, ANDREW CARDWELL<sup>8</sup>, CHRISTINE CHEN<sup>2</sup>, EUGENE CHIANG<sup>1</sup>,  
JEFFREY K. CHILCOTE<sup>9</sup>, RENÉ DOYON<sup>10</sup>, ZACHARY H. DRAPER<sup>11,12</sup>, GASPARD DUCHÊNE<sup>1,13</sup>, THOMAS M. ESPOSITO<sup>14</sup>,  
MICHAEL P. FITZGERALD<sup>14</sup>, STEPHEN J. GOODSSELL<sup>8,15</sup>, ALEXANDRA Z. GREENBAUM<sup>2,16</sup>, MARKUS HARTUNG<sup>8</sup>, PASCALE HIBON<sup>8</sup>,  
SASHA HINKLEY<sup>17</sup>, LI-WEI HUNG<sup>14</sup>, PATRICK INGRAHAM<sup>18</sup>, JAMES E. LARKIN<sup>14</sup>, BRUCE MACINTOSH<sup>4,6</sup>, JEROME MAIRE<sup>9</sup>,  
FRANCK MARCHIS<sup>3</sup>, CHRISTIAN MAROIS<sup>11,12</sup>, BRENDA C. MATTHEWS<sup>11,12</sup>, KATIE M. MORZINSKI<sup>19</sup>, REBECCA OPPENHEIMER<sup>20</sup>,  
JENNY PATIENCE<sup>7</sup>, MARSHALL D. PERRIN<sup>2</sup>, ABHIJITH RAJAN<sup>7</sup>, FREDRIK T. RANTAKYRÖ<sup>8</sup>, NARU SADAKUNI<sup>8</sup>, ANDREW SERIO<sup>8</sup>,  
ANAND SIVARAMAKRISHNAN<sup>2</sup>, RÉMI SOUMMER<sup>2</sup>, SANDRINE THOMAS<sup>18</sup>, KIMBERLY WARD-DUONG<sup>7</sup>,  
SLOANE J. WIKTOROWICZ<sup>21</sup>, AND SCHUYLER G. WOLFF<sup>2,16</sup>

<sup>1</sup> Astronomy Department, University of California, Berkeley, Berkeley, CA 94720, USA

<sup>2</sup> Space Telescope Science Institute, 3700 San Martin Drive, Baltimore, MD 21218, USA

<sup>3</sup> SETI Institute, Carl Sagan Center, 189 Bernardo Avenue, Mountain View, CA 94043, USA

<sup>4</sup> Kavli Institute for Particle Astrophysics and Cosmology, Stanford University, Stanford, CA 94305, USA

<sup>5</sup> Department of Astronomy & Astrophysics, University of Toronto, Toronto, ON M5S 3H4, Canada

<sup>6</sup> Lawrence Livermore National Laboratory, 7000 East Avenue, Livermore, CA 94040, USA

<sup>7</sup> School of Earth and Space Exploration, Arizona State University, P.O. Box 871404, Tempe, AZ 85287, USA

<sup>8</sup> Gemini Observatory, Casilla 603, La Serena, Chile

<sup>9</sup> Dunlap Institute for Astronomy & Astrophysics, University of Toronto, 50 St. George Street, Toronto, ON M5S 3H4, Canada

<sup>10</sup> Institut de Recherche sur les Exoplanètes, Département de Physique, Université de Montréal, Montréal, QC H3C 3J7, Canada

<sup>11</sup> University of Victoria, 3800 Finnerty Road, Victoria, BC V8P 5C2, Canada

<sup>12</sup> National Research Council of Canada Herzberg, 5071 West Saanich Road, Victoria, BC V9E 2E7, Canada

<sup>13</sup> Université Grenoble Alpes/CNRS, Institut de Planétologie et d'Astrophysique de Grenoble, F-38000 Grenoble, France

<sup>14</sup> Department of Physics and Astronomy, University of California, Los Angeles, Los Angeles, CA 90095, USA

<sup>15</sup> Center for Advanced Instrumentation, Durham University, Durham DH1 3LE, UK

<sup>16</sup> Physics and Astronomy Department, Johns Hopkins University, Baltimore, MD 21218, USA

<sup>17</sup> University of Exeter, Astrophysics Group, Physics Building, Stocker Road, Exeter EX4 4QL, UK

<sup>18</sup> Large Synoptic Survey Telescope, 950 N Cherry Avenue, Tucson, AZ 85719, USA

<sup>19</sup> Steward Observatory, 933 N. Cherry Avenue, University of Arizona, Tucson, AZ 85721, USA

<sup>20</sup> American Museum of Natural History, New York, NY 10024, USA

<sup>21</sup> Department of Astronomy, University of California, Santa Cruz, 1156 High Street, Santa Cruz, CA 95064, USA

*Received 2015 June 1; accepted 2015 August 18; published 2015 September 23*

### ABSTRACT

We present Gemini Planet Imager (GPI) observations of AU Microscopii, a young M dwarf with an edge-on, dusty debris disk. Integral field spectroscopy and broadband imaging polarimetry were obtained during the commissioning of GPI. In our broadband imaging polarimetry observations, we detect the disk only in total intensity and find asymmetries in the morphology of the disk between the southeast (SE) and northwest (NW) sides. The SE side of the disk exhibits a bump at  $1''$  (10 AU projected separation) that is three times more vertically extended and three times fainter in peak surface brightness than the NW side at similar separations. This part of the disk is also vertically offset by  $69 \pm 30$  mas to the northeast at  $1''$  when compared to the established disk midplane and is consistent with prior Atacama Large Millimeter/submillimeter Array and *Hubble Space Telescope*/Space Telescope Imaging Spectrograph observations. We see hints that the SE bump might be a result of detecting a horizontal sliver feature above the main disk that could be the disk backside. Alternatively, when including the morphology of the NW side, where the disk midplane is offset in the opposite direction  $\sim 50$  mas between  $0''.4$  and  $1''.2$ , the asymmetries suggest a warp-like feature. Using our integral field spectroscopy data to search for planets, we are 50% complete for  $\sim 4 M_{\text{Jup}}$  planets at 4 AU. We detect a source, resolved only along the disk plane, that could either be a candidate planetary mass companion or a compact clump in the disk.

*Key words:* circumstellar matter – instrumentation: adaptive optics – methods: data analysis – planet–disk interactions – stars: individual (AU Mic) – techniques: high angular resolution

### 1. INTRODUCTION

Debris disks are comprised of rocky and icy bodies, left over from planet formation. The planetesimals in these disks collide, fragment, and produce micron-sized grains that scatter optical and near-infrared radiation from the star, resulting in a detectable nebulosity. Grains are influenced by the gravitational pull of larger bodies in the system as well as radiative forces and corpuscular drag due to the host star. These effects can

induce morphology in debris disks such as midplane warps, as in  $\beta$  Pictoris (Mouillet et al. 1997); bow structures, as seen in HD 61005 (Hines et al. 2007); and narrow, offset rings, such as in Fomalhaut (Kalas et al. 2005).

The debris disk around AU Microscopii (AU Mic) is interesting for multiple reasons. At a distance of 9.9 pc, we are able to probe closer to the star than most systems. As a member of the  $\beta$  Pictoris moving group (Navascués et al. 1999), its

young age of  $23 \pm 3$  Myr (Mamajek & Bell 2014) makes it a prime candidate for directly detecting young planets still hot from their recent formation. It is also one of the very few M dwarfs with a debris disk imaged in scattered light. Since its ground-based discovery by Kalas et al. (2004), this edge-on disk has been studied in scattered light using both adaptive optics (AO) and the *Hubble Space Telescope* (Liu 2004; Krist et al. 2005; Metchev et al. 2005; Fitzgerald et al. 2007; Graham et al. 2007; Schneider et al. 2014), as well as in millimeter thermal emission with interferometric arrays (Wilner et al. 2012; MacGregor et al. 2013). From modeling these observations, it is believed AU Mic harbors a “birth ring” of planetesimals at  $\sim 40$  AU, inward of which is empty and outward of which is comprised of micron-sized dust particles placed on barely bound orbits (Augereau & Beust 2006; Strubbe & Chiang 2006). Within the disk, there are also local enhancements of surface brightness and deviations in the disk’s vertical structure (e.g., Fitzgerald et al. 2007; Schneider et al. 2014). These features are likely non-axisymmetric structures located at or exterior to the birth ring and seen in projection (Graham et al. 2007).

The Gemini Planet Imager (GPI; Macintosh et al. 2014) is a high-contrast imaging instrument on Gemini South with a high-order AO system, a coronagraph to suppress starlight, and a science instrument that features both integral field spectroscopy (spectral mode; Larkin et al. 2014) and broadband imaging polarimetry (polarimetry mode; Perrin et al. 2015). GPI gives us the opportunity to probe closer in and deeper than previous optical or near-infrared observations of AU Mic. We use GPI in both spectral and polarimetry modes to study the morphology of the debris disk and search for planets close in to the star.

## 2. OBSERVATIONS AND DATA REDUCTION

AU Mic was observed as part of commissioning the GPI instrument. On 2014 May 12, twenty-seven 60 s spectral mode frames were taken in the *K1* spectral band ( $1.90\text{--}2.19 \mu\text{m}$ ) at an airmass  $< 1.01$ . Images of the sky and instrument thermal background were not taken with the observations, but thermal background data were taken earlier that night and used in the data reduction. Forty-four 60 s polarimetry mode observations were taken on 2014 May 15 in the *H*-band at an airmass  $\leq 1.015$ . The GPI waveplate was rotated by  $22.5^\circ$  between frames. AU Mic transits almost directly overhead at Cerro Pachon, so a total of  $154^\circ$  of field rotation was obtained in each data set for angular differential imaging (ADI; Marois et al. 2006). For the *H*-band data, four frames were discarded due to anomalies associated with commissioning tests.

The raw data consist of individual spectra in spectral mode and pairs of spots for the two orthogonal polarizations dispersed by the Wollaston prism in polarimetry mode. Each spectrum or spot pair corresponds to a spatial resolution element, or spaxel, in a data cube created by the GPI Data Reduction Pipeline (DRP; Perrin et al. 2014). The spectral mode data were dark subtracted, corrected for shifts in the position of individual spectra on the detector due to instrument flexure (Wolff et al. 2014) with an *H*-band argon arc lamp exposure taken immediately before the observing sequence, wavelength calibrated with *K1*-band argon arc lamp data, fixed for bad pixels in the 2D data, assembled into a spectral data cube, thermal background subtracted, fixed for bad pixels in the 3D data cube, and corrected for distortion.


The reduction of the polarimetry mode data follows the steps described in Perrin et al. (2015). To create polarimetry data cubes where the third dimension contains two orthogonal polarizations, the data were dark subtracted, flexure corrected using a cross-correlation routine (Draper et al. 2014), fixed for bad pixels in the 2D data, assembled into a data cube using a model of the point-spread function (PSF) of the spot pair, fixed for bad pixels in the 3D data cube, and corrected for distortion. The polarimetry data cubes were then collapsed in the polarization dimension to obtain 2D total intensity images. For our total intensity reductions, we do not construct the full Stokes data cube, but one is made with the GPI DRP to search for a polarized signal.

To combine and photometrically calibrate our data, we used the GPI DRP to measure the four fiducial diffraction or “satellite” spots, which are centered on the occulted star and imprinted with its attenuated spectrum (Wang et al. 2014). In spectral mode, the satellite spot fluxes were used to calibrate the data photometrically relative to the occulted star. For each data cube, the location of the occulted star at each wavelength was found using a least squares fit to all of the satellite spots’ positions and the magnitude of the atmospheric differential refraction. In polarimetry mode, the satellite spot flux calibration is an ongoing task, so no flux-calibrated polarimetry mode data are presented here. The central star in the polarimetry data was located using a radon-transform-based algorithm (Wang et al. 2014; Pueyo et al. 2015). The precision on the star center is 0.7 and 0.9 mas for spectral and polarimetry mode, respectively (Wang et al. 2014).

To remove the stellar PSF, we used `pyKLIP` (Wang et al. 2015), a Python implementation of the Karhunen–Loève Image Projection (KLIP) algorithm (Soummer et al. 2012; Pueyo et al. 2015). To distinguish astrophysical sources from the stellar PSF, we used ADI for the polarimetry mode data and both ADI and spectral differential imaging (SDI; Marois et al. 2000) for the spectral mode data. Multiple PSF-subtracted images were made depending on the purpose of the reduction. For all the subtractions, we divided the PSF into annuli, divided each annulus into azimuthal sectors, and ran KLIP on each sector.

The main parameters we adjusted were the number of modes used from the Karhunen–Loève (KL) transform and an exclusion criterion for reference PSFs, similar to  $N_0$  in Lafrenière et al. (2007), defined by the number of pixels a hypothetical astrophysical source would move azimuthally, radially, or both due to ADI/SDI. For the *H*-band total intensity data, we used 3 KL modes and a 10-pixel exclusion for the diffuse left side of the disk presented in Figure 1(a) and 4 KL modes and a 6-pixel exclusion for the more compact right side presented in Figure 1(a). We also searched for fainter features using 10 KL modes and a 3-pixel exclusion as shown in Figure 1(b). For the *K1*-band data, we used 10 KL modes and a 3-pixel exclusion criterion to characterize a compact candidate source shown in Figure 1(c) and 20 KL modes and a 1.5 pixel exclusion criteria to filter out extended disk features and quantify our sensitivity to point sources in Section 4.

The frames were rotated to orient the disk horizontally using a disk position angle (PA) of  $128.41^\circ$  (MacGregor et al. 2013) derived from observations with the Atacama Large Millimeter/submillimeter Array (ALMA) before being mean combined to produce the images in Figure 1.



**Figure 1.** Reduced images of the AU Mic disk. The images are rotated so that the disk is horizontal. Images (a) and (b) are from the  $H$ -band total intensity broadband observations. Image (a) is optimized for the bright main belt of the disk. The azimuthal structure outside of the horizontal disk plane is due to unsubtracted speckles. Image (b) is used to look for fainter features. The curly brace indicates the position of the sliver feature we detect. Image (c) is from the  $K1$ -band spectral mode observations. The arrow indicates the candidate compact source that we detect.

We did not detect any polarized signal (5% upper limit on the polarization fraction at  $0''.5$ ), consistent with a low expected polarization for forward scattered light at small projected angular separations (Graham et al. 2007) and reduced polarization sensitivity due to image persistence in the HAWAII 2-RG detector (Smith et al. 2008), for which calibration work is ongoing.

### 3. ANALYSIS OF THE DEBRIS DISK

All three reductions in Figure 1 show stark asymmetries between the southeast (SE) and northwest (NW) sides of the disk. To quantify these asymmetries, we used the conservative  $H$ -band reduction in Figure 1(a) and fit Gaussian vertical profiles to the disk at various separations. We fit between  $0''.4$  and  $1''.25$ . The inner radius is determined by the increasing noise at small angular separations, and the outer radius is set by GPI's finite field of view. Due to the relatively low signal-to-noise and ADI self-subtraction in the vertical wings of the disk, a Gaussian provides good fits with well-behaved residuals and performs as well as alternate profiles such as the Lorentzian used by Graham et al. (2007). The primary aim of the Gaussian fit was to measure the disk's midplane vertical offset and FWHM. We also obtained the peak brightness of the disk as part of the fit and calculated the vertically integrated flux by summing over the flux within one FWHM centered about the measured disk midplane.

Our reductions are subject to varying levels of disk self-subtraction, which bias our measurements. To correct for this effect, we repeated the analysis with model disks injected separately at 1 of 5 non-overlapping positions, resulting in 10 sides for analysis. The disk model is a symmetric, edge-on, optically thin ring parameterized by a power-law grain size distribution, a Henyey–Greenstein scattering function, and a Gaussian vertical profile. We adopted values from the porous water-ice model from Graham et al. (2007) and varied both the scale height and the surface brightness to accommodate the range of measured values. As GPI only sees the inner arcsecond of the disk, the main contribution to the flux in the models is the front edge of the disk with scattering angles


$<15^\circ$ . The mean FWHM, peak brightness, and integrated vertical flux we measured from the model disks were used to correct biases in our measured values as a function of separation. The scatter of all four quantities in the model disks were used as the uncertainties in our measurements.

We present measurements of the midplane offset and FWHM as a function of separation with  $1\sigma$  error bars in Figure 2. The vertical midplane offset peaks around  $1''$  on the SE side, with an offset from the average ALMA midplane of  $69 \pm 30$  mas to the northeast at  $1''$  separation. The midplane offset is roughly constant between  $0''.4$  and  $1''.2$  on the NW side, with an offset of  $47 \pm 8$  mas to the southwest at  $1''$ . The FWHM is roughly constant on the NW side and consistent with measurements at larger separations but triples on the SE side past  $0''.8$  to a peak FWHM of  $570 \pm 56$  mas. Due to unsubtracted residual PSF features, the disk brightness as a function of separation that we measure is not reliable. However, the overall brightness asymmetry between the two sides as seen in Figure 1 is real: on average, the peak surface brightness is three times larger on the NW side, but the vertically integrated brightnesses are consistent with being equal.

From the aggressive reduction in Figure 1(b), we see a dark gap and a faint horizontal sliver on the SE side above the main disk. This sliver is only at  $3.1\sigma$  significance when comparing the total flux of this feature to randomly summing up noise in the same manner. However, it emerges in successive reductions that progressively remove speckle noise and does not point radially toward the star, like residual speckles. We cannot rule out that the apparent morphology is influenced by KLIP self-subtraction. We also do not observe any indication of a corresponding feature on the NW side, which could be hidden underneath the oversubtraction of the brighter main belt of the disk.

### 4. SENSITIVITY TO PLANETS

The achieved sensitivity for our  $K1$  data was calculated following the procedure outlined by Mawet et al. (2014) and the  $5\sigma$  equivalent false positive threshold is plotted in Figure 3.



**Figure 2.** Measurements of the morphology of the disk in the  $H$ -band with the negative and positive separations for the SE and NW sides, respectively. At each separation, we plot the disk midplane vertical offset and FWHM with  $1\sigma$  error bars as blue squares. The faded circles represent the FWHM before correcting for biases. Adjacent points are correlated so that the plot is well sampled.

Flux attenuation due to PSF subtraction was quantified by injecting and recovering the brightnesses of simulated planets. To translate our contrast curve to limits on possible planets in the system, we ran a Monte Carlo analysis as described by Nielsen & Close (2010) to determine the completeness of our data. Since GPI is not sensitive to planets at large orbital distances due to its restricted field of view, we also included the limits from the Near-Infrared Coronagraphic Imager (NICI; Wahhaj et al. 2013) in Figure 3. With the improved inner working angle, the GPI data are 50% complete for  $4M_{\text{Jup}}$  planets with semimajor axes of 4 AU assuming COND models (Baraffe et al. 2003).


We did find one compact  $K1$  candidate, marked in Figure 1(c), at a separation of  $544 \pm 4$  mas, a PA of  $130^\circ 2 \pm 0^\circ 9$ , and a  $K1$  contrast of  $1.1 \pm 0.3 \times 10^{-5}$ . Although this source is detected with  $5\sigma$  confidence in  $K1$ , there is no corresponding source at  $H$ . Because polarimetry mode flux calibration is still ongoing, we cannot currently quantify the constraints placed by our  $H$ -band data. According to COND models (Baraffe et al. 2003), this source would correspond to a  $2\text{--}3M_{\text{Jup}}$  planet. However, inspection of Figure 1(c) reveals that the clump is extended along the disk. The FWHM perpendicular to the disk is  $\approx 70$  mas and consistent with being unresolved when compared to the satellite spot PSF; parallel to the disk, the measured FWHM is  $\approx 160$  mas giving an intrinsic size of  $\approx 140$  mas (1.4 AU), which suggests a local enhancement in the disk similar to those reported at larger projected separations (e.g., Liu 2004; Fitzgerald et al. 2007; Schneider et al. 2014). We are confident in the measured elongation because it is significantly larger than a residual speckle, which,

given the  $K1$  bandwidth, would subtend about 100 mas at this angular separation.


## 5. DISCUSSION

The AU Mic debris disk has significant asymmetries in the morphology and surface brightness between the SE and NW sides of the disk. In Figure 4, we compare our observations with 1.3 mm continuum data from ALMA (MacGregor et al. 2013) and optical imaging using the Space Telescope Imaging Spectrograph (STIS) on the *Hubble Space Telescope* (Schneider et al. 2014). The ALMA data probe larger planetesimals insensitive to the variable radiation pressure and stellar wind drag of AU Mic, making it easier to disentangle dynamical and radiative forces on the debris disk. STIS, like GPI, traces scattered light from small grains, but with a larger field of view.

The disk midplane offsets, FWHM enhancement, and brightness asymmetries we find agree well with the published STIS results from Schneider et al. (2014). The most prominent common feature is the faint, extended, and vertically offset bump on the SE side at  $1''$ . This feature is also continuous in the STIS data where it is seen to taper back down near  $2''$  from the star. Since the SE bump is also consistent with the most prominent undulation in the ALMA contours, it is plausible that the feature is associated with gravitational perturbations of the parent bodies. The statistical significance of this millimeter feature is questioned by MacGregor et al. (2013); however, the correspondence between ALMA and shorter-wavelength observations lends credence to its reality. Assuming the millimeter grains are not displaced from the birth ring, the correlation between scattered light and millimeter-emission



**Figure 3.** (Top) Sensitivity to point sources as a function of projected separation using the K1-band spectral mode data. (Bottom) Completeness for planets as a function of semimajor axis and mass in the AU Mic system based on contrast curves from NICI (Wahhaj et al. 2013) and GPI (this paper). Completeness is calculated using a Monte Carlo method described by Nielsen & Close (2010) where planets with random orbits are generated, the contrast curves determine whether they are detected, and COND models (Baraffe et al. 2003) are used to convert from luminosity to mass.



**Figure 4.** Comparison of GPI data with previous observations. GPI H-band data from Figure 1(a) is shown in the center with contours from 1.3 mm ALMA observations (MacGregor et al. 2013) overlaid on top using a linear scaling and an image from STIS (Schneider et al. 2014) in the background. The southeast (left) bump between  $-1''.5$  and  $-1''$  in the disk is apparent in all three observations. The midplane offset is also continuous between the STIS and GPI data.

suggests that the SE bump is located in the  $\sim 40$  AU birth ring and is just seen projected at  $\sim 10$  AU separation. As both the STIS and ALMA data were taken within four years of the GPI observations, we cannot yet distinguish between orbital motions at 10 and 40 AU.

The compact K1 candidate in our data could be another clump of disk material or a planet with a circumplanetary disk. The candidate must be a transient feature unless the object is self-gravitating. If the projected size of this source is representative of its radial extent ( $\delta a$ ), Keplerian shear at an orbital separation ( $a$ ) of 40 AU will completely erase this feature in only  $(2/3)(a/\delta a) = 19(a/40 \text{ AU})(1.4 \text{ AU}/\delta a)$  orbits. However, a 1.4 AU Hill sphere at 40 AU is established by a  $0.08(M_*/0.6 M_\odot) M_{\text{Jup}}$  object, which is well below our detection limits. Follow-up observations are required though for verification and to determine common proper motion.

Ignoring the possible detection of the sliver in Figure 1(b), the vertical midplane offset variations on the two sides of the disk in opposite directions hint at a vertical warp in the disk. Applying the analytical model from Dawson et al. (2011) to the AU Mic system, any inclined planet between an Earth and Neptune mass could induce the warp depending on the orbital separation and dynamical timescale. The candidate companion we see at a projected separation of 5.5 AU can be slightly lower than an Earth mass and still be responsible for the warp. The FWHM and flux asymmetries in the disk could be due to dynamical heating from planets or parent bodies in the ring, which can puff up the disk vertically but keep the total amount of starlight scattered constant.

If the sliver feature is real, it may be the backside of the disk (the bright main belt would be the front side). In this case, the variations in the disk could be explained by an inclined ring with azimuthally varying brightness due to clumps in the disk, like those seen at further separations (Fitzgerald et al. 2007). Essentially, on the SE side, the backside is visible due to a local brightness enhancement that is absent on the NW side. However, without a corresponding detection on the NW, the backside interpretation is uncertain.

Our GPI observations show that there remain many unanswered questions regarding this extensively studied debris disk. The stark asymmetries between the two sides of the disk and correlation with millimeter observations hint at gravitational interactions due to perturbers. With deeper imaging by new near-infrared high-contrast imaging instruments and higher-resolution data in the millimeter by ALMA, we may soon have a much better understanding of the physics in the system.

We thank M. MacGregor and G. Schneider for making available to us the reduced ALMA and STIS data, respectively. This research was supported in part by NASA NNX15AD95G, NASA NNX11AD21G, NSF AST-0909188, NSF AST-1413718, and the University of California LFRP-118057. The GPI project has been supported by Gemini Observatory, which is operated by AURA, Inc., under a cooperative agreement with the NSF on behalf of the Gemini partnership: the NSF (USA), the National Research Council (Canada), CONICYT (Chile), the Australian Research Council (Australia), MCTI (Brazil), and MINCYT (Argentina). This research has made use of the SIMBAD database, operated at CDS, Strasbourg, France.

*Facility:* Gemini:South (GPI).

## REFERENCES

- Augereau, J.-C., & Beust, H. 2006, *A&A*, 455, 987  
 Baraffe, I., Chabrier, G., Barman, T. S., Allard, F., & Hauschildt, P. H. 2003, *A&A*, 402, 701

- Barrado y Navascués, D., Stauffer, J. R., Song, I., & Caillault, J.-P. 1999, *ApJ*, **520**, 123
- Dawson, R. I., Murray-Clay, R. A., & Fabrycky, D. C. 2011, *ApJL*, **743**, L17
- Draper, Z. H., Marois, C., Wolff, S., et al. 2014, *Proc. SPIE*, **9147**, 91474Z
- Fitzgerald, M. P., Kalas, P. G., Duchêne, G., Pinte, C., & Graham, J. R. 2007, *ApJ*, **670**, 536
- Graham, J. R., Kalas, P. G., & Matthews, B. C. 2007, *ApJ*, **654**, 595
- Hines, D. C., Schneider, G., Hollenbach, D., et al. 2007, *ApJL*, **671**, L165
- Kalas, P., Graham, J. R., & Clampin, M. 2005, *Natur*, **435**, 1067
- Kalas, P., Liu, M. C., & Matthews, B. C. 2004, *Sci*, **303**, 1990
- Krist, J. E., Ardila, D. R., Golimowski, D. A., et al. 2005, *ApJ*, **129**, 1008
- Lafrenière, D., Marois, C., Doyon, R., Nadeau, D., & Artigau, É 2007, *ApJ*, **660**, 770
- Larkin, J. E., Chilcote, J. K., & Aliado, T. 2014, *Proc. SPIE*, **9147**, 91471K
- Liu, M. C. 2004, *Sci*, **305**, 1442
- MacGregor, M. A., Wilner, D. J., Rosenfeld, K. A., et al. 2013, *ApJL*, **762**, L21
- Macintosh, B. A., Graham, J. R., Ingraham, P., et al. 2014, *PNAS*, **111**, 12661
- Mamajek, E. E., & Bell, C. P. M. 2014, *MNRAS*, **445**, 2169
- Marois, C., Doyon, R., Racine, R., & Nadeau, D. 2000, *PASP*, **112**, 91
- Marois, C., Lafreniere, D., Doyon, R., Macintosh, B., & Nadeau, D. 2006, *ApJ*, **641**, 556
- Mawet, D., Milli, J., Wahhaj, Z., et al. 2014, *ApJ*, **792**, 97
- Metchev, S. A., Eisner, J. A., Hillenbrand, L. A., & Wolf, S. 2005, *ApJ*, **622**, 451
- Mouillet, D., Larwood, J. D., Papaloizou, J. C. B., & Lagrange, A. M. 1997, *MNRAS*, **292**, 896
- Nielsen, E. L., & Close, L. M. 2010, *ApJ*, **717**, 878
- Perrin, M. D., Duchêne, G., Millar-Blanchaer, M., et al. 2015, *ApJ*, **799**, 182
- Perrin, M. D., Maire, J., Ingraham, P., et al. 2014, *Proc. SPIE*, **9147**, 91473J
- Pueyo, L., Soummer, R., Hoffman, J., et al. 2015, *ApJ*, **803**, 31
- Schneider, G., Grady, C. A., Hines, D. C., et al. 2014, *ApJ*, **148**, 59
- Smith, R. M., Zavodny, M., Rahmer, G., & Bonati, M. 2008, *Proc. SPIE*, **7021**, 70210J
- Soummer, R., Pueyo, L., & Larkin, J. 2012, *ApJL*, **755**, L28
- Strubbe, L. E., & Chiang, E. I. 2006, *ApJ*, **648**, 652
- Wahhaj, Z., Liu, M. C., Nielsen, E. L., et al. 2013, *ApJ*, **773**, 179
- Wang, J. J., Rajan, A., Graham, J. R., et al. 2014, *Proc. SPIE*, **9147**, 914755
- Wang, J. J., Ruffio, J.-B., de Rosa, R. J., et al. 2015, Astrophysics Source Code Library, record ascl: 1506.001
- Wilner, D. J., Andrews, S. M., MacGregor, M. A., & Hughes, A. M. 2012, *ApJL*, **749**, L27
- Wolff, S. G., Perrin, M. D., Maire, J., et al. 2014, *Proc. SPIE*, **9147**, 91477H