Lawrence Berkeley National Laboratory

Recent Work

Title

TRLFS, EXAFS and IR studies on the structural determination of the extracted complexes of Ln(III) and An(III) with Cyanex301 and Cyanex302

Permalink

https://escholarship.org/uc/item/9kp2m9n2

Author

Rao, Linfeng

Publication Date

2003-04-25

TRLFS, EXAFS and IR Studies on the Structural Determination of the Extracted Complexes of Ln(III) and An(III) with Cyanex301 and Cyanex302

Guoxin Tian¹, Takaumi Kimura,² Zenko Yoshida², Yongjun Zhu³, Linfeng Rao¹

¹Glenn T. Seaborg Center, Lawrence Berkeley National Laboratory, 1 Cyclotron Road, Berkeley, CA 94720

²Advanced Science Research Center, Japan Atomic Energy Research Institute, Tokai-mura, Ibaraki 319-11, Japan

³Institute of Nuclear Energy Technology, Tsinghua University, Beijing, 100084, China

The structure of the extracted complexes of lanthanides, Ln(III) (Ln = Sm, Eu, Tb, Dy), and Cm(III) with Cyanex301 and Cyanex302 was investigated using Time-Resolved Laser-induced Fluorescence Spectroscopy (TRLFS), EXAFS and FT-IR spectroscopy. The results indicate that the composition of the extracted Ln(III)/Cyanex301 complexes is different from that of the An(III)/Cyanex301 complexes. Data suggest that the formula of the Ln(III) complexes is LnL₃•2H₂O or LnL₄•H₂O, where L stands for Cyanex301. However, the formula for the An(III)/Cyanex301 complexes is AnL₄, with no H₂O but only 8 sulfur atoms from the 4 Cyanex301 ligands coordinated to An(III), suggested by the TRLFS data for Cm(III) and previous EXAFS and IR data for Am(III). In the case of Cyanex302 complexes, data indicate that there are 3 oxygen from 3 Cyanex302 and 3 to 5 H₂O coordinated to Ln(III) or An(III), suggesting that Cyanex302 is monodentate in the complexes and the molecular formula is ML₃•nH₂O (n=3-5).