

# UCSF

## UC San Francisco Previously Published Works

### Title

A kinetic study of the murine mixed lymphocyte reaction by 5,6-carboxyfluorescein diacetate succinimidyl ester labeling

### Permalink

<https://escholarship.org/uc/item/9n76w40n>

### Journal

Journal of Immunological Methods, 279(1-2)

### ISSN

0022-1759

### Authors

Chen, Jeng Chang  
Chang, M L  
Muench, Marcus O.

### Publication Date

2003-08-01

Peer reviewed



# A kinetic study of the murine mixed lymphocyte reaction by 5,6-carboxyfluorescein diacetate succinimidyl ester labeling

Jeng-Chang Chen<sup>a,b,1</sup>, Ming-Ling Chang<sup>c,d,1</sup>, Marcus O. Muench<sup>e,\*</sup>

<sup>a</sup>Fetal Treatment Center, Department of Surgery, University of California, San Francisco, CA, USA

<sup>b</sup>Department of Surgery, Chang Gung Children's Hospital, Taoyuan, Taiwan

<sup>c</sup>Liver Center, Department of Medicine, University of California, San Francisco, CA, USA

<sup>d</sup>Department of Hepatology, Chang Gung Memorial Hospital, Taoyuan, Taiwan

<sup>e</sup>Department of Laboratory Medicine, University of California, 533 Parnassus Avenue, Room U-440, San Francisco, CA 94143-0793, USA

Received 6 December 2002; received in revised form 16 May 2003; accepted 19 May 2003

## Abstract

Alternatives to the use of radioisotopes to measure cell proliferation in mixed lymphocyte reactions (MLR) are desirable to avoid the hazards and costs associated with radioisotope use. The versatile fluorochrome 5,6-carboxyfluorescein diacetate succinimidyl ester (CFSE) has been used to measure MLR and provides the opportunity to measure several different growth parameters. This study was aimed at determining which growth parameter is most practical and suitable for measuring murine MLR. The parameters measured were: the relative number of daughter T-cells, the relative number and frequency of reactive T-cell precursors and the relative number of mitotic events. Responder cells were CFSE-labeled unfractionated splenocytes from C57BL/6 mice. Stimulator cells included irradiated splenocytes from C57BL/6 (control), B6D2F<sub>1</sub> (haplo-allogeneic) or FVB/N (allogeneic) mice. Cultures were harvested daily for 1 week. Stimulator T-cells rapidly declined to less than 0.2–0.3% of the mixed population by day 2 of culture. Experimental groups had a significantly higher number of daughter T-cells and mitotic events after 2 days of culture with the number of daughter T-cells climbing exponentially after 5 days of culture. The number and frequency of reactive T-cell precursors were significantly higher in experimental groups on days 2–3, but this difference became insignificant by day 4. Among all the parameters, the relative number of daughter T-cells was the most practical for measuring MLR, after 5 days of culture, based upon the growth kinetics of responder T-cells and the survival of the stimulator cells.

© 2003 Published by Elsevier Science B.V.

**Keywords:** 5,6-carboxyfluorescein diacetate succinimidyl ester; Mixed lymphocyte reaction; Mouse

**Abbreviations:** MLR, mixed lymphocyte reactions; CFSE, 5,6-Carboxyfluorescein diacetate succinimidyl ester; <sup>3</sup>H-TdR, <sup>3</sup>H-thymidine; PBS, phosphate buffer saline; BSA, bovine serum albumin; FBS, fetal bovine serum; PI, propidium iodide.

\* Corresponding author. Tel.: +1-415-476-8420; fax: +1-415-476-2956.

E-mail address: muench@itsa.ucsf.edu (M.O. Muench).

<sup>1</sup> These authors were visiting scholars at the University of California, San Francisco, CA from Taiwan during the period of October 16, 2000 to October 15, 2002.

## 1. Introduction

The mixed lymphocyte reaction (MLR) measures the proliferative response of responder T cells against antigens present on allogeneic stimulator cells. It is presumed that the MLR is an in vitro analog of in vivo alloreactivity, and is widely used in transplantation

37 immunology to measure recipient T cell responses  
38 against donor tissues due to the mismatch of MHC  
39 antigens. Cell proliferation is an essential part of an  
40 immune response which can be measured in an  
41 allogeneic mixed lymphocyte culture as an indicator  
42 of immune reaction. There are numerous methods to  
43 measure changes in cell number. The most widely  
44 used method to measure proliferation is  $^3\text{H}$ -thymidine  
45 ( $^3\text{H}$ -TdR) incorporation. However, due to the hazards  
46 and cost associated with the use and disposal of  
47 tritium, nonradioactive alternatives have been pur-  
48 sued.

49 Nonradioactive methods include the use of a range  
50 of fluorochromes, which have been used for measur-  
51 ing the proliferation of lymphocytes as well as for  
52 tracking migration and positioning in vivo (Parish,  
53 1999). Among them, 5,6-carboxyfluorescein diacetate  
54 succinimidyl ester (CFSE) stands out as being the  
55 most versatile in terms of long-term cell tracking and  
56 quantifying proliferation either in vivo or in vitro  
57 (Lyons, 1999; Parish, 1999). CFSE can spontaneously  
58 and irreversibly couple to intracellular proteins and is  
59 equally distributed between two daughter cells when  
60 cells divide. Thus, proliferating cells can be tracked  
61 by flow cytometry based upon the sequential loss of  
62 fluorescence intensity (Lyons, 2000; Lyons and Par-  
63 ish, 1994). Furthermore, multiparameter flow cytom-  
64 etry allows for the examination of cell subsets within  
65 the dividing cell population as well as monitoring  
66 phenotypic changes associated with activation and  
67 cell division.

68 CFSE staining has been used as a replacement for  
69  $^3\text{H}$ -TdR incorporation to measure the MLR of cells of  
70 human and nonhuman-primate origins (Matthews et  
71 al., 2000; Popma et al., 2000; Nitta et al., 2001). In  
72 most cases, the outcome of the MLR has been  
73 determined based on the total numbers of daughter  
74 T-cells generated. However, the ability of CFSE to  
75 track cell division allows for other parameters of cell  
76 proliferation to be measured. These include estimates  
77 of the number and frequency of T-cell precursors that  
78 have responded in the MLR as well as the number of  
79 total mitotic events that have occurred. The utility of  
80 measuring these parameters in analyzing murine MLR  
81 has not been closely examined. Therefore, we studied  
82 the kinetics of murine T-cell responses in allogeneic  
83 MLR using CFSE labeling and multiparameter flow  
84 cytometry.

## 2. Materials and methods 85

### 2.1. Responder and stimulator 86

87  
88 Two inbred strains of mice, C57BL/6 (H-2<sup>b</sup>) and  
89 FVB/N (H-2<sup>q</sup>), and one hybrid strain of mice B6D2F<sub>1</sub>  
90 (C57BL/6 × DBA/2, H-2<sup>b/d</sup>) were purchased from  
91 Charles River (Wilmington, MA) or Simonsen (Gilroy,  
92 CA). They were 8–30 weeks of age when sacrificed  
93 and their spleens were harvested. Responder T-cells  
94 were from the C57BL/6 strain, whereas stimulators  
95 included C57BL/6 (syngeneic), B6D2F<sub>1</sub> (haplo-allo-  
96 geneic) and FVB/N (allogeneic) strains. Splenocytes  
97 were harvested under sterile conditions and passaged  
98 through 70 μm cell strainers (Becton Dickinson, Frank-  
99 lin Lakes, NJ). Red cells were depleted by chemical  
100 lysis using ACK buffer, pH 7.2–7.4, consisting of 0.15  
101 M NH<sub>4</sub>Cl, 1.0 mM KHCO<sub>3</sub> and 0.1 mM Na<sub>2</sub>EDTA  
102 (Sigma, St. Louis, MO). The cell suspension was then  
103 washed twice with phosphate buffer saline (PBS)  
104 containing 0.3% bovine serum albumin (BSA) (PBS/  
105 BSA) (Roche Molecular Biochemicals, Indianapolis,  
106 IN). Responder cells were suspended at  $2 \times 10^7$  cells/  
107 ml in PBS/BSA for CFSE labeling, whereas stimulator  
108 cells received 3000 cGy irradiation and were then  
109 resuspended at  $6 \times 10^6$  cells/ml in culture medium.  
110

### 2.2. CFSE labeling of responder cells 111

112 CFSE labeling of responder cells was undertaken as  
113 previously described (Lyons, 2000; Lyons and Parish,  
114 1994). Immediately before labeling, 5 mM CFSE stock  
115 (Molecular Probes, Eugene, OR) in DMSO (Fisher  
116 Scientific, Fair Lawn, NJ) was thawed and diluted to  
117 10 μM in a volume of PBS/BSA equal to that in which  
118 the responder cells were suspended. The two equal vo-  
119 lumes were mixed to initiate labeling and periodically  
120 agitated at room temperature for 10 min. The labeling  
121 process was quenched by adding an equal volume of  
122 heat-inactivated fetal bovine serum (FBS, Hyclone,  
123 Logan, UT) to the sample. After 1 min, the CFSE-la-  
124 beled cells were washed twice, recounted and adjusted  
125 to a concentration of  $2 \times 10^6$  cells/ml in culture media.  
126

### 2.3. MLR 127

128 The MLR was performed in 96-well U-bottom  
129 microtiter plates (Costar, Cambridge, MA). Culture

130 medium consisted of RPMI 1640 supplemented with  
131 10% heat-inactivated FBS, sodium pyruvate (110 mg/  
132 l), nonessential amino acids, L-glutamate (1 mM), 2-  
133 mercaptoethanol ( $5 \times 10^{-5}$  M), penicillin (50 U/ml)  
134 and streptomycin (50  $\mu$ g/ml) and N-acetyl cysteine  
135 (Sigma; 10 mmol/l, pH adjusted to 7.2). CFSE-labeled  
136 C57BL/6 responder cells were plated at  $1 \times 10^6$  cells/  
137 ml in a volume of 250  $\mu$ l per well and cocultured at a  
138 ratio of 1:3 with 3000 cGy irradiated C57BL/6,  
139 B6D2F<sub>1</sub> or FVB/N stimulator cells. The plates were  
140 then placed in a humidified 37 °C, 5% CO<sub>2</sub> incubator.  
141 The control experiment for accessing background fluo-  
142 rescence was set up by coculture of unlabeled C57BL/  
143 6 responder cells and irradiated stimulator cells under  
144 the same conditions. In order to assess the kinetic  
145 division of the responder cells and the viability of stim-  
146 ulator cells, cells were harvested from quadruplet  
147 wells on a daily basis.

#### 148 2.4. Flow cytometric analysis

150 Cells harvested from each well were suspended in  
151 culture supernatant from the clone 2.4G2 hybridoma  
152 cell line (ATCC, Manassas, VA) that produces mAb  
153 against Fc $\gamma$ II/Fc $\gamma$ III receptors to block nonantigen-  
154 specific binding of immunoglobulins. CFSE-labeled  
155 cells were incubated on ice for 30 min with PE  
156 conjugated anti-CD3 (Caltag Laboratories, Burlin-  
157 game, CA). Cells were then washed three times in  
158 PBS/BSA with 0.01% NaN<sub>3</sub> (Sigma). Cells from each  
159 well were suspended in 150  $\mu$ l PBS/BSA with 0.01%  
160 NaN<sub>3</sub> and 2  $\mu$ g/ml propidium iodide (PI, Molecular  
161 Probes). Three-color flow cytometry was performed  
162 on a FACScan (BD Biosciences, San Jose, CA).  
163 Events for each well were acquired at a fixed speed  
164 (high setting) for 1 min to measure an equal volume  
165 from each sample, thereby providing a basis for the  
166 relative comparison of data collected for each sample.  
167 Dead cells were excluded from analysis based on  
168 staining with PI.

#### 169 2.5. Data and statistical analyses

171 Data were analyzed using CellQuest acquisition  
172 and analysis software (BD Biosciences). The total  
173 numbers of events (cells) were determined by analyz-  
174 ing the data using dot plots and rectangular regions to  
175 define the cell populations. Histograms were used to

176 track the divisions of CFSE-labeled cells where each  
177 peak of CFSE-fluorescence defined a single round of  
178 cell division. All data presented are from analyses of  
179 live (PI<sup>-</sup>) cells only.

180 The relative numbers of T-cell precursors required  
181 for generating these daughter cells under each division  
182 peak was calculated by dividing the number of daugh-  
183 ter-cell events by 2 raised to the power of the given  
184 round of division ( $2^n$ ). The sum of all the calculated  
185 numbers of precursors from each division peak was  
186 used to represent the number of reactive T-cell pre-  
187 cursors. Furthermore, the frequency of reactive T-cell  
188 precursors was estimated by dividing the number of  
189 reactive T-cell precursors by the number of total T-  
190 cells, which equals the number of undivided T-cells  
191 plus the number of reactive T-cell precursors. Alterna-  
192 tively, the number of total live T-cells measured on the  
193 first day of culture was also used to estimate the  
194 frequency of reactive T-cell precursors. The frequency  
195 values were multiplied by 100 to be represented as a  
196 percentage of all T cells. In addition, the mitotic events  
197 under each peak can also be determined by subtracting  
198 the number of reactive T-cell precursors from the  
199 number of daughter T-cells under each peak (Wells  
200 et al., 1997). Thus, the relative number of total mitotic  
201 events can be calculated as the sum of individual  
202 mitotic events under each peak, which equals the  
203 number of daughter T-cells minus the number of  
204 reactive T-cell precursors.

205 Independent sample *t*-test was used for comparing  
206 means of events and ratios between control and ex-  
207 perimental groups. Paired-samples *t*-test was used to  
208 make a comparison between the two precursor fre-  
209 quencies generated on the basis of different denomi-  
210 nators, one derived from the total T-cell precursors on  
211 the indicated day and the other from the mean undi-  
212 vided T-cells on day 1. A *P*-value < 0.05 was consid-  
213 ered statistically significant.

### 214 3. Results

#### 215 3.1. Proliferation of CD3<sup>+</sup> responder cells

216 Mixed lymphocyte cultures were harvested after 1–  
217 7 days and stained with CD3-PE to identify the T-cell  
218 component of the cultures. CD3 staining and CFSE  
219 fluorescence were detected by flow cytometry (Fig. 1).  
220




Fig. 1. Flow cytometric analysis of mixed lymphocyte cultures. Progressive loss of CFSE fluorescence by C57BL/6 responder  $CD3^+$  T-cells, indicated by the regions R3 and R4, is shown for days 1, 3 and 6 of culture. Viable FVB/N stimulator cells are indicated by regions R5 and R6. The stimulator cells were mostly located at the area of low fluorescent intensity ( $<10^1$ ).

221 Daughter T-cells, derived from the responder spleno-  
 222 cytes, could be differentiated from undivided T-cells  
 223 by the intensity of CFSE staining. The relative numb-  
 224 ers of daughter T-cells could be easily determined by  
 225 electronic gating, as shown in Fig. 1. In the case of  
 226 alloreactive MLR, the numbers of daughter T-cells  
 227 showed a slow increase until day 4 and then an expo-  
 228 nential expansion after day 5 (Fig. 2A). The syngeneic  
 229 control group (C57BL/6) generated significantly lower  
 230 numbers of daughter T-cells than the haploidentical

B6D2F<sub>1</sub> and completely MHC-mismatched FVB/N  
 groups beginning on the second day of culture on-  
 wards. There were no significant differences in the  
 numbers of daughter T-cells between the B6D2F<sub>1</sub> and  
 FVB/N groups at any time point. Thus, a T-cell pro-  
 liferative response to MHC antigens could be readily  
 detected by this method.

Survival of the undivided CD3<sup>+</sup> cells in the mixed  
 cultures was also analyzed (Fig. 2B). A linear decrease  
 in the number of unresponsive T-cells was observed in  
 the C57BL/6 group over the 7 days of culture ana-  
 lyzed. However, the B6D2F<sub>1</sub> and FVB/N groups had  
 comparatively steady numbers of undivided CD3<sup>+</sup>  
 cells until day 3. A slight drop followed in these  
 groups on day 4, and then a linear decrease in the  
 number of unresponsive T-cells was seen as in the  
 C57BL/6 control group.

The kinetics of CD3<sup>+</sup> cell proliferation was ana-  
 lyzed in greater detail using histograms to view the  
 levels of CFSE fluorescence (Fig. 3). On day 1 (24 h of  
 culture), CD3<sup>+</sup> responder cells were uniformly stained  
 with CFSE, indicating that these cells had not yet  
 begun to divide. By the second day of culture, CD3<sup>+</sup>  
 responders, stimulated by B6D2F<sub>1</sub> or FVB/N spleno-  
 cytes, had begun to divide for up to three rounds of  
 mitosis (Fig. 3 and data not shown). In contrast,  
 responder T-cells stimulated with syngeneic C57BL/  
 6 cells did not show any evidence of cell division for  
 the first 2 days (Fig. 3). On days 3 and 4, four to six  
 rounds of divisions could be detected in all the three  
 groups. On day 5, CD3<sup>+</sup> cells of all three groups had  
 divided at least for seven to eight times, resulting in a  
 loss of fluorescent intensity on some responder cells to  
 background levels. Thus, at this time point further  
 resolution of subsequent cell division could not be  
 accurately measured. On days 6–7, almost all the  
 daughter T-cells, stimulated by B6D2F<sub>1</sub> and FVB/N  
 splenocytes, had reached the area of background  
 fluorescence due to many rounds of divisions.

Measured estimates of the relative number and  
 frequency of reactive T-cell precursors are listed for  
 a representative experiment in Table 1. These calcu-  
 lated measures are based on measurements of viable  
 (PI<sup>-</sup>) CD3<sup>+</sup> cells. A significantly higher number and  
 frequency of reactive T-cell precursors were observed  
 in the B6D2F<sub>1</sub> and FVB/N groups compared to the  
 C57BL/6 group on days 2–3. However, the frequency  
 of reactive T-cell precursors did not differ between





Fig. 2. Kinetics of expansion and survival of responder T-cells in mixed lymphocyte cultures. Accumulations of daughter T-cells from the responder population are shown in (A) and survival of undivided T-cells are shown in (B). Data are shown as the mean  $\pm$  2.0 standard errors of the mean (SE) of measurements made on replicate cultures.



t1.1 Table 1

t1.2 Relative numbers and frequencies of reactive T-cell precursors as well as the number of mitotic events from MLR analyzed at 1–4 days of culture

t1.3	Day	Stimulator	Relative Number of T-cell Precursors	Estimated T-cell Precursor Frequency (%) <sup>a</sup>	Relative Number of Mitotic Events
t1.4	1	C57BL/6	4.13 ± 0.80	0.05 ± 0.01/0.05 ± 0.01	4.13 ± 0.80
t1.5		B6D2F1	2.38 ± 0.32	0.03 ± 0.01/0.03 ± 0.00	2.38 ± 0.31
t1.6		FVB/N	3.38 ± 0.55	0.05 ± 0.01/0.04 ± 0.01	3.38 ± 0.55
t1.7	2	C57BL/6	5.00 ± 0.79	0.07 ± 0.01/0.06 ± 0.01 <sup>b</sup>	5.00 ± 0.79
t1.8		B6D2F1	54.50 ± 1.85 <sup>c</sup>	0.68 ± 0.02 <sup>c</sup> /0.65 ± 0.02 <sup>c</sup>	80.75 ± 2.39 <sup>c</sup>
t1.9		FVB/N	47.25 ± 2.46 <sup>c</sup>	0.64 ± 0.02 <sup>c</sup> /0.60 ± 0.03 <sup>c</sup>	67.25 ± 2.25 <sup>c</sup>
t1.10	3	C57BL/6	40.50 ± 2.06	0.63 ± 0.04/0.47 ± 0.02 <sup>b</sup>	140.00 ± 10.79
t1.11		B6D2F1	152.75 ± 11.18 <sup>c</sup>	1.87 ± 0.10 <sup>c</sup> /1.82 ± 0.13 <sup>c</sup>	759.75 ± 75.93 <sup>c</sup>
t1.12		FVB/N	129.00 ± 3.49 <sup>c</sup>	1.69 ± 0.04 <sup>c</sup> /1.65 ± 0.04 <sup>c</sup>	573.50 ± 32.17 <sup>c</sup>
t1.13	4	C57BL/6	141.00 ± 15.15	2.37 ± 0.18/1.63 ± 0.18 <sup>b</sup>	591.00 ± 76.73
t1.14		B6D2F1	157.75 ± 3.59	2.08 ± 0.04/1.88 ± 0.04 <sup>b</sup>	1411.75 ± 59.31 <sup>c</sup>
t1.15		FVB/N	160.25 ± 6.20	2.24 ± 0.10/2.05 ± 0.08 <sup>b</sup>	1720.75 ± 101.25 <sup>c</sup>

t1.16 Data are shown as the mean ± standard error of the mean for quadruplicate analyses performed on live cells.

t1.17 <sup>a</sup> T-cell precursor frequencies were calculated by two methods. The first number represents the frequency calculated using the sum of reactive T-cell precursors and non-responsive T-cells measured on the day when the culture was harvested as the denominator. The second value was calculated using the mean number of undivided T-cells on day 1, when responder T-cells had not yet divided, as the denominator.t1.18 <sup>b</sup>  $P < 0.05$  comparison between first and second numbers, paired-samples t-test.t1.19 <sup>c</sup>  $P < 0.05$  compared with the C57BL/6 control group, independent-samples t-test.

279 control and experimental groups by the fourth day of  
 280 culture. The frequencies of T-cell precursors were  
 281 estimated by two methods, as described in Materials  
 282 and methods. The linear decrease in undivided CD3<sup>+</sup>  
 283 cells in the control group resulted in an overestimation  
 284 of T-cell precursor frequency on days 2–4; whereas  
 285 the relatively slow decrease of undivided CD3<sup>+</sup> cells in  
 286 the experimental groups did not notably influence the  
 287 frequency calculations until day 4.

288 The relative numbers of mitotic events that occurred  
 289 in the cultures could also be calculated (Table 1). The  
 290 B6D2F<sub>1</sub> and FVB/N groups exhibited higher division  
 291 of T-cells on days 2–4 compared with the control  
 292 group. This pattern was consistent with the measures  
 293 of the relative number of daughter T-cells (Fig. 2A),  
 294 which measures the outcome of these early mitotic  
 295 events as a latter accumulation of daughter T-cells. By  
 296 the 5th day of culture, it was difficult to accurately  
 297 estimate the number and frequency of reactive T-cell

precursors as well as the number of mitotic events. This  
 is because T cells under peak 6 might contain daughter  
 cells that had undergone more than six rounds of  
 division and can no longer be discriminated from  
 background fluorescence as indicated using an un-  
 stained control (Fig. 3).

### 3.2. Proliferation of responder CD3<sup>-</sup> cells

Responder CD3<sup>-</sup> cells also started to proliferate in  
 the mixed culture beginning on day 2, reached a peak  
 around days 3–4, and subsequently declined to a  
 plateau (Fig. 4A). This proliferation pattern did not  
 parallel that of responder CD3<sup>+</sup> cells. In B6D2F<sub>1</sub> and  
 FVB/N groups, the ratio of CD3<sup>+</sup> to CD3<sup>-</sup> responder  
 cells was usually less than 1 over the first 3 days,  
 gradually increased to five on day 5, and abruptly  
 amplified to over 10 on day 7. In contrast, the ratios  
 from C57BL/6 control group sustained below 1 until

Fig. 3. Histogram plots gated on CD3<sup>+</sup> cells show the kinetics of cell division associated with an MLR. Histograms of CFSE fluorescence for T-cells stimulated by haploallogeneic B6D2F<sub>1</sub> splenocytes are shown in the left column and histograms for the syngeneic control group are shown for comparison in the right column. A peak of low fluorescence (arrow) represents viable stimulator T-cells observed during the first 2 days of culture. On days 5 and 7, cells with ≥ 6 rounds of division had reached a fluorescent intensity similar to the background autofluorescence of unstained responder T-cells (thin line). This overlap limited the resolution of further cycles of cell division and marked the maximal cycle that could be distinguished.






Fig. 4. Proliferation and contribution of CD3<sup>-</sup> responder cells to mixed lymphocyte cultures. Accumulations of daughter CD3<sup>-</sup> cells (non-T-cells) from the responder population are shown in (A). The ratio of daughter CD3<sup>+</sup> to CD3<sup>-</sup> cells is shown in (B). Data are shown as the mean  $\pm$  2.0 SE.

316 day 4, slightly increased to the peak of 3 on day 6, and  
 317 then went down on day 7 (Fig. 4B).

318

### 319 3.3. Viability of stimulator cells

320 Viable stimulator cells in the mixed cultures were  
 321 observed in the early days of culture and could be identified  
 322 by their lack of CFSE staining. They remained  
 323 after 1 day of culture, comprising both CD3<sup>-</sup> cells and  
 324 CD3<sup>+</sup> cells (Fig. 1). CD3<sup>-</sup> stimulator cells exhibited a

325 much slower decline and represented 2–3% of the total  
 326 cell population on day 3 of culture (Fig. 5A). However,  
 327 a rapid decline of viable CD3<sup>+</sup> cells to 0.2–0.3% of all  
 328 viable cells was observed by the second day of culture,  
 329 and less than 0.1% stimulator CD3<sup>+</sup> cells were present  
 330 on days 3–4 (Fig. 5B). An MLR, in which the  
 331 responders were not stained with CFSE, was also  
 332 undertaken to trace the viability of stimulator cells by  
 333 flow cytometry using differences in MHC antigens (H-  
 334 2K<sup>b, d, and q</sup>). Total viable stimulator cells in the mixed  
 335 culture accounted for 11.5–23.6% on day 1, 2.9–7.6%  
 336 on day 2, 0.5–0.7% on day 3, 0.5–1% on day 4, 0.2–  
 337 0.5% on day 5 and 0.1–0.3% on days 6–7 (data not  
 338 shown).




Fig. 5. Survival of stimulator cells in mixed lymphocyte cultures. The percentages of viable CD3<sup>-</sup> stimulator cells are shown in (A) and the percentages of viable stimulator CD3<sup>+</sup> T-cells are shown in (B). Data are shown as the mean  $\pm$  2.0 SE.

339 **4. Discussion**

340 Cell proliferation encompasses DNA synthesis,  
341 mitosis and an increase in cell number.  $^3\text{H-TdR}$  incor-  
342 poration measures cell proliferation by the incorpora-  
343 tion of  $^3\text{H-TdR}$  into DNA while cells are dividing, and  
344 has been a standard method for measuring cell prolif-  
345 eration in mixed lymphocyte cultures for decades. In  
346 view of the increasing costs of waste disposal and  
347 protective precautions associated with radioisotope  
348 use, developing an alternative for quantify cell prolif-  
349 eration, which can be as effective as, but less hazard-  
350 ous than  $^3\text{H-TdR}$ , is welcomed. CFSE labeling stands  
351 out in many aspects as a versatile alternative to  
352 conventional  $^3\text{H-TdR}$  incorporation. Most importantly,  
353 this method was shown to correlate well with  $^3\text{H-TdR}$   
354 incorporation (Fulcher and Wong, 1999; Popma et al.,  
355 2000). To extend the adoption of CFSE for use in  
356 measuring murine MLR, we closely examined a num-  
357 ber of parameters associated with cell division to  
358 determine the optimal timing and methods to measure  
359 murine MLR.

360  $^3\text{H-TdR}$  incorporation only measures cell division  
361 during a brief period of culture and provides no  
362 information on the types of cells growing or their  
363 degree of response. In contrast, CFSE labeling can  
364 provide more information such as the relative number  
365 of reactive T-cell precursors, the estimated frequency  
366 of reactive T-cell precursors, the relative number of  
367 mitotic events and relative number of daughter T-cells  
368 associated with an MLR. The proliferation of CFSE-  
369 labeled responder cells could be traced for at least 1  
370 week, providing a detailed picture of the different  
371 stages of an MLR. Moreover, immunophenotypic  
372 analyses of the dividing cells insured that the mea-  
373 surement was specific to  $\text{CD3}^+$  T-cells. This might be  
374 crucial in some cases because unfractionated tissues  
375 such as spleen, lymph node or fetal liver (Harris et al.,  
376 1994) have been used as the source of responder T  
377 cells and substantial non-T-cell proliferation might  
378 occur from such tissues. Under these circumstances,  
379 data obtained by  $^3\text{H-TdR}$  incorporation might vari-  
380 ously reflect proliferation of both T and non-T cells.  
381 According to our results, non-T-cells displayed a  
382 different pattern of proliferation from T-cells. Clearly,  
383 non-T-cells are most apt to contribute to overall  
384 proliferation in the early days of a splenic MLR.  
385 Furthermore, cell proliferation in the control group

tended to be more the result of non-T-cells than in the  
experimental groups, which could be readily distin-  
guished and disregarded by the flow cytometric  
method.

The simplest method of quantifying an MLR using  
CFSE was found to be measurement of the accumula-  
tion of daughter  $\text{CD3}^+$  cells. The magnitude of the T-  
cell proliferative response elicited by various stimula-  
tor conditions mainly depended upon the rounds of  
division the T-cell precursors had undergone rather  
than upon the frequency of reactive T-cell precursors.  
This argument was supported by the presence of two  
distinct peaks on days 6 and 7 in the histogram plots  
shown in Fig. 3. One peak represented quiescent T-  
cells and the other was daughter T-cells that had  
undergone at least seven to eight rounds of division.  
There was a relative paucity of daughter cells in  
between. T-cell proliferation was asynchronous, to a  
degree, likely due to variation in the time of entry into  
the first division cycle (Hasbold et al., 1999). None-  
theless, new T-cell precursors did not appear to be  
recruited after day 5 of culture. Accordingly, the  
alloresponse to MHC-mismatched stimulators led to  
an exponential expansion of daughter T-cells by eliciting  
more rounds of division than the nonspecific  
response in syngeneic group. Modest T-cell prolifera-  
tion in the control group appeared later than in the  
experimental groups, becoming evident on day 3.

The number and frequency of reactive T cell pre-  
cursors in an MLR could also be estimated using flow  
cytometry. Our estimated frequencies of reactive T-cell  
precursors was about 2%, which was within some  
published estimates of the percentage of allorespon-  
sive precursors (Ford and Atkins, 1973; Ford et al.,  
1975; Matzinger and Bevan, 1977; Sherman and  
Chattopadhyay, 1993). These measurements were best  
performed on the third day of culture rather than at  
later time points that are better suited for measuring the  
accumulation of daughter T cells. This is because  
differences between the control and experimental  
groups were evident on days 2 and 3, but by day 4,  
these differences disappeared. The weak background  
proliferation observed in the controls by day 4 of  
culture might be due to nonspecific activation of the  
T cells by cytokines produced in the cultures or FBS  
used to supplement cultures. Another reason for early  
analysis of the cultures is that the death of some of the  
daughter cells and nonresponsive T-cells affects the

estimates of the size and frequency of the precursor pool. Dead (PI<sup>+</sup>) daughter and nonresponder cells were observed, but it was necessary to disregard these cells for analyses as the PI staining and high nonspecific binding of antibody by dead cells interferes with the analysis of CD3 expression. Since the estimation of the frequency of T-cell precursors relies on an accurate measurement of the total number of live T cells used to initiate the cultures, we tested two methods by which to calculate precursor frequencies. One method relied on a single flow cytometric analysis, which was used to estimate the total pool of T cells based on the calculated number of reactive T-cell precursors plus the number of undivided T-cells. A second method used an additional analysis after 1 day of culture, when the T-cell precursor pool had not yet begun to divide, to measure the total number of live T-cells. We chose to make this analysis after 1 day rather than on day 0 of culture as the CFSE staining procedure may result in the death of some T cells that is not immediately evident. Both methods of calculating the size of the total T-cell pool resulted in similar estimates of reactive T-cell precursor frequencies for the experimental groups, but the control group had a faster loss of nonresponsive T cells leading to higher estimates of precursor frequencies beginning on the second day of culture. Thus, choosing which method to use to calculate the size of the total T-cell pool depends on the types of MLR being compared.

Being able to estimate the number or frequency of reactive T-cell precursors may have specific application in studies aimed at measuring these values. However, this study clearly indicates that measuring the relative number of daughter T-cells is the best parameter for analyzing the magnitude of an MLR (Matthews et al., 2000; Popma et al., 2000). It is simple as well as straightforward without the requirement of extensive calculations. The number of mitotic events can also be used for comparing in vitro MLR because its pattern generally paralleled that of the accumulation of daughter T-cells at least during the first 4 days of cultures. However, it offered no clear advantages over the measuring the number of daughter T-cells and required labor-intensive calculations. Using the number of daughter T-cells, the differences between the control and experimental groups could be distinguished beginning on the second day of culture onwards, and were more pronounced after 5 days of

culture. Furthermore, irradiated stimulators were nearly absent from culture after 3 days. Thus, quantifying MLR by the number of daughter T-cells after 5 days of culture could be easily performed with minimal interference from stimulator cells. These findings support the use of a flow cytometry as a reliable method to analyze murine MLR based on CFSE staining.

### Acknowledgements

We thank Drs. Alicia Bárcena and Lewis Lanier for advice on this study. We also thank Dr. Y.W. Kan for his advice and support. Dr. Paul Dazin also provided valuable assistance with the operation of the flow cytometry equipment and software.

This work was supported by grant CMRP1052 (JCC) from Chang Gung University, Taoyuan, Taiwan and by NIH grant DK59301 (MOM).

### References

- Ford, W.L., Atkins, R.C., 1973. The proportion of lymphocytes capable of recognizing strong transplantation antigens in vivo. *Adv. Exp. Med. Biol.* 29, 255.
- Ford, W.L., Simmonds, S.J., Atkins, R.C., 1975. Early cellular events in a systemic graft-vs.-host reaction: II. Autoradiographic estimates of the frequency of donor lymphocytes which respond to each Ag-B-determined antigenic complex. *J. Exp. Med.* 141, 681.
- Fulcher, D., Wong, S., 1999. Carboxyfluorescein succinimidyl ester-based proliferative assays for assessment of T cell function in the diagnostic laboratory. *Immunol. Cell Biol.* 77, 559.
- Harris, D., LoCascio, J., Besencon, F., 1994. Analysis of the alloreactive capacity of human umbilical cord blood: implications for graft-versus-host disease. *Bone Marrow Transplant.* 14, 545.
- Hasbold, J., Gett, A.V., Rush, J.S., Deenick, E., Avery, D., Jun, J., Hodgkin, P.D., 1999. Quantitative analysis of lymphocyte differentiation and proliferation in vitro using carboxyfluorescein diacetate succinimidyl ester. *Immunol. Cell Biol.* 77, 516.
- Lyons, A.B., 1999. Divided we stand: tracking cell proliferation by flow cytometry. *Immunol. Cell Biol.* 77, 509.
- Lyons, A.B., 2000. Analysing cell division in vivo and in vitro using flow cytometric measurement of CFSE dye dilution. *J. Immunol. Methods* 243, 147.
- Lyons, A.B., Parish, C.R., 1994. Determination of lymphocyte division by flow cytometry. *J. Immunol. Methods* 171, 131.
- Matthews, N.C., Wadhwa, M., Bird, C., Borrás, F.E., Navarrete, C.V., 2000. Sustained expression of CD154 (CD40L) and proinflammatory cytokine production by alloantigen-stimulated umbilical cord blood T cells. *J. Immunol.* 164, 6206.

- 529 Matzinger, P., Bevan, M.J., 1977. Hypothesis: why do so many lymphocytes respond to major histocompatibility antigens? *Cell Immunol.* 29, 1.
- 530
- 531
- 532 Nitta, Y., Nelson, K., Andrews, R.G., Thomas, R., Gaur, L.K., Allen, M.D., 2001. CFSE dye dilution mixed lymphocyte reactions quantify donor-specific alloreactive precursors in non-human primate cardiac graft rejection. *Transplant. Proc.* 33, 326.
- 533
- 534
- 535
- 536 Parish, C.R., 1999. Fluorescent dyes for lymphocyte migration and proliferation studies. *Immunol. Cell Biol.* 77, 499.
- 537
- 538
- 539 D., Moore, J.S., Rosengard, B.R., 2000. Immune monitoring in xenotransplantation: the multiparameter flow cytometric mixed lymphocyte culture assay. *Cytometry* 42, 277.
- 540
- 541
- 542 Sherman, L.A., Chattopadhyay, S., 1993. The molecular basis of allorecognition. *Annu. Rev. Immunol.* 11, 385.
- 543
- 544 Wells, A.D., Gudmundsdottir, H., Turka, L.A., 1997. Following the fate of individual T cells throughout activation and clonal expansion. Signals from T cell receptor and CD28 differentially regulate the induction and duration of a proliferative response. *J. Clin. Invest.* 100, 3173.
- 545
- 546
- 547
- 548

UNCORRECTED PROOF