

# UC San Diego

## UC San Diego Previously Published Works

### Title

Enhanced Upper-Airway Muscle Responsiveness Is a Distinct Feature of Overweight/Obese Individuals without Sleep Apnea

### Permalink

<https://escholarship.org/uc/item/9pg8m00p>

### Journal

American Journal of Respiratory and Critical Care Medicine, 190(8)

### ISSN

1073-449X

### Authors

Sands, Scott A  
Eckert, Danny J  
Jordan, Amy S  
et al.

### Publication Date

2014-10-15

### DOI

10.1164/rccm.201404-0783oc

Peer reviewed



# Enhanced Upper-Airway Muscle Responsiveness Is a Distinct Feature of Overweight/Obese Individuals without Sleep Apnea

Scott A. Sands<sup>1,2</sup>, Danny J. Eckert<sup>1,3</sup>, Amy S. Jordan<sup>1,4</sup>, Bradley A. Edwards<sup>1</sup>, Robert L. Owens<sup>1</sup>, James P. Butler<sup>1</sup>, Richard J. Schwab<sup>5</sup>, Stephen H. Loring<sup>6</sup>, Atul Malhotra<sup>1,7</sup>, David P. White<sup>1</sup>, and Andrew Wellman<sup>1</sup>

<sup>1</sup>Division of Sleep Medicine, Brigham and Women's Hospital and Harvard Medical School, Boston, Massachusetts; <sup>2</sup>Department of Allergy Immunology and Respiratory Medicine and Central Clinical School, The Alfred and Monash University, Melbourne, Australia; <sup>3</sup>Neuroscience Research Australia and the School of Medical Sciences, University of New South Wales, Sydney, Australia; <sup>4</sup>School of Psychological Science, University of Melbourne, Melbourne, Australia; <sup>5</sup>Division of Sleep Medicine, Department of Medicine, University of Pennsylvania, Philadelphia, Pennsylvania; <sup>6</sup>Department of Anesthesia and Critical Care, Beth Israel Deaconess Medical Center and Harvard Medical School, Boston, Massachusetts; and <sup>7</sup>Division of Pulmonary and Critical Care, University of Southern California San Diego, La Jolla, California

## Abstract

**Rationale:** Body habitus is a major determinant of obstructive sleep apnea (OSA). However, many individuals do not have OSA despite being overweight/obese (body mass index > 25 kg/m<sup>2</sup>) for reasons that are not fully elucidated.

**Objectives:** To determine the key physiologic traits (upper-airway anatomy/collapsibility, upper-airway muscle responsiveness, chemoreflex control of ventilation, arousability from sleep) responsible for the absence of OSA in overweight/obese individuals.

**Methods:** We compared key physiologic traits in 18 overweight/obese subjects without apnea (apnea-hypopnea index < 15 events per hour) with 25 overweight/obese matched patients with OSA (apnea-hypopnea index ≥ 15 events per hour) and 11 normal-weight nonapneic control subjects. Traits were measured by repeatedly lowering continuous positive airway pressure to subtherapeutic levels for 3 minutes during non-REM sleep.

**Measurements and Main Results:** Overweight/obese subjects without apnea exhibited a less collapsible airway than overweight/obese patients with apnea (critical closing pressure:  $-3.7 \pm 1.9$  vs.  $0.6 \pm 1.2$  cm H<sub>2</sub>O;  $P = 0.003$ ; mean  $\pm$  95% confidence interval),

but a more collapsible airway relative to normal-weight control subjects ( $-8.8 \pm 3.1$  cm H<sub>2</sub>O;  $P < 0.001$ ). Notably, overweight/obese subjects without apnea exhibited a threefold greater upper-airway muscle responsiveness than both overweight/obese patients with apnea ( $\Delta$ genioglossus EMG/ $\Delta$ epiglottic pressure:  $-0.49$  [ $-0.22$  to  $-0.79$ ] vs.  $-0.15$  [ $-0.09$  to  $-0.22$ ] %max/cm H<sub>2</sub>O;  $P = 0.008$ ; mean [95% confidence interval]) and normal-weight control subjects ( $-0.16$  [ $-0.04$  to  $-0.30$ ] %max/cm H<sub>2</sub>O;  $P = 0.02$ ). Loop gain was elevated (more negative) in both overweight/obese groups and normal-weight control subjects ( $P = 0.02$ ). Model-based analysis demonstrated that overweight/obese individuals without apnea rely on both more favorable anatomy and collapsibility and enhanced upper-airway dilator muscle responses to avoid OSA.

**Conclusions:** Overweight/obese individuals without apnea have a moderately compromised upper-airway structure that is mitigated by highly responsive upper-airway dilator muscles to avoid OSA. Elucidating the mechanisms underlying enhanced muscle responses in this population may provide clues for novel OSA interventions.

**Keywords:** apnea phenotypes; upper airway muscles; obesity; control of breathing; mathematical model

(Received in original form April 29, 2014; accepted in final form September 1, 2014)

Supported by the National Institutes of Health (5R01HL048531-16, R01HL102321, 1R01HL090897-01A2, 1K24HL093218-01A1, and 1P01HL095491). S.A.S. was supported by an American Heart Association Postdoctoral Fellowship 11POST7360012, and is currently supported by a National Health and Medical Research Council of Australia (NHMRC) Early Career Fellowship (1053201) and R. G. Menzies award. D.J.E. is supported by an NHMRC R. D. Wright Fellowship (1049814). A.S.J. is supported by an Australian Research Council Future Fellowship (100100203). B.A.E. is supported by the NHMRC C. J. Martin Overseas Biomedical Fellowship (1035115).

Author Contributions: Conception of the current substudy, S.A.S., R.J.S., D.P.W., and A.W. Conception and experimental design of parent study, D.J.E., A.S.J., A.M., D.P.W., and A.W. Data collection and analysis, S.A.S., D.J.E., A.S.J., B.A.E., and A.W. Statistical analysis, mathematical-model analysis, and initial drafting of the manuscript, S.A.S., D.P.W., and A.W. All authors contributed substantially to the interpretation of findings and to the editing of the manuscript for important intellectual content.

Correspondence and requests for reprints should be addressed to Scott A. Sands, Ph.D., Division of Sleep Medicine, Brigham and Women's Hospital and Harvard Medical School, 221 Longwood Avenue, Boston, MA 02115. E-mail: sasands@partners.org

This article has an online supplement, which is accessible from this issue's table of contents at [www.atsjournals.org](http://www.atsjournals.org)

Am J Respir Crit Care Med Vol 190, Iss 8, pp 930–937, Oct 15, 2014

Copyright © 2014 by the American Thoracic Society

Originally Published in Press as DOI: 10.1164/rccm.201404-0783OC on September 5, 2014

Internet address: [www.atsjournals.org](http://www.atsjournals.org)

## At a Glance Commentary

### Scientific Knowledge on the

**Subject:** Obesity is a major risk factor for obstructive sleep apnea (OSA). However, many individuals who are overweight/obese do not develop OSA for reasons that are not fully understood.

### What This Study Adds to the

**Field:** Overweight/obese individuals without apnea exhibit markedly enhanced upper-airway muscle responsiveness compared with overweight/obese patients with OSA, and also compared with normal-weight nonapneic control subjects. These characteristics tend to mitigate OSA by counteracting a moderate anatomical compromise. Quantitatively, the more favorable anatomy and enhanced muscle responsiveness in overweight/obese individuals without apnea are both essential for protecting against the development of OSA.

Obstructive sleep apnea (OSA) is a prevalent disease with major cardiovascular and neurocognitive consequences. Obesity is a major risk factor for OSA, and most individuals manifest OSA with sufficient additional body weight (1). However, many overweight/obese individuals do not have OSA for reasons that remain unclear.

Mechanistically, OSA is caused primarily by a narrow, collapsible upper airway (2, 3). Because of increased fat

deposition, overweight/obese individuals generally have a more collapsible airway than normal-weight individuals (4–8), yet the modest relationship between body mass index (BMI) and upper-airway collapsibility (3, 9) indicates that some individuals gain weight without major upper-airway compromise.

Whether a collapsible airway ultimately yields OSA also depends on additional nonanatomic physiologic traits (2, 4, 10–17). In individuals with a highly collapsible airway, enhanced upper-airway dilator muscle responsiveness during sleep is theoretically necessary to provide a level of ventilation that is compatible with stable breathing and sleep. Indeed, evidence indicates that overweight/obese individuals without apnea have a greater capacity to activate upper-airway muscles and increase airflow during sleep compared with BMI-matched patients with OSA (4, 14). However, other nonanatomic traits, such as a higher (less-sensitive) arousal threshold or a reduced ventilatory drive response to asphyxia (lower loop gain), can also protect against OSA (11, 18, 19).

Accordingly, we sought to determine which key physiologic traits protect overweight/obese individuals from developing OSA. We tested the primary hypotheses that overweight/obese individuals without apnea not only have a less collapsible upper airway (lower critical closing pressure [P<sub>crit</sub>]), but also have more responsive pharyngeal muscles (greater genioglossus EMG [EMG<sub>gg</sub>] response to negative pressure) than patients with OSA. We studied 11 new participants

and combined their data with our larger phenotype study (2). In total, OSA traits were examined in 18 overweight/obese (BMI > 25 kg/m<sup>2</sup>) nonapneic control subjects (apnea-hypopnea index [AHI] < 15 events per hour) and compared with 25 overweight/obese patients with OSA (AHI > 15) and 11 normal-weight (BMI < 25 kg/m<sup>2</sup>) control subjects.

## Methods

### Subjects

All overweight/obese nonapneic control subjects aged 25–70 years from our larger study (2) were included. Patients with OSA were selected to match this group for BMI, age, and sex. All normal-weight nonapneic control subjects were included for comparison. The 11 new participants were studied to better match the groups for sex and enhance statistical power. For details, see the online supplement.

### Polysomnography and Physiologic Traits

Details have been described previously (2, 16). Briefly, participants presented for three overnight sleep studies. First, routine polysomnography determined the presence or absence of OSA. Hypopneic events (>10 s) were scored as those that caused a 3% oxygen desaturation or arousal from sleep (20). Subsequently, over 2 nights we measured key physiologic traits by repeatedly lowering continuous positive airway pressure (CPAP) below therapeutic levels to challenge the pharyngeal airway; drops were performed for 3-minute

**Table 1.** Patient Characteristics

| Characteristics                       | Overweight/Obese Subjects without Apnea (n = 18) | Overweight/Obese Patients with OSA (n = 25) | Normal-Weight Control Subjects without Apnea (n = 11) |
|---------------------------------------|--|---|---|
| Male:female*                          | 9:9  | 16:9  | 6:5   |
| Age, yr                               | 44 ± 5   | 48 ± 4                                      | 41 ± 8  |
| Body mass index, kg · m <sup>-2</sup> | 31 ± 2   | 32 ± 1                                      | 22 ± 1 <sup>†</sup>                                   |
| AHI total, h <sup>-1</sup>            | 8.0 ± 2.1  | 41.5 ± 8.8 <sup>‡</sup>                     | 4.6 ± 3.0   |
| AHI supine non-REM, h <sup>-1</sup>   | 6.3 ± 2.4  | 42.0 ± 9.4 <sup>‡</sup>                     | 3.7 ± 3.0   |
| AHI supine REM, h <sup>-1</sup>       | 20.0 ± 6.5                                       | 43.9 ± 8.5 <sup>‡</sup>                     | 10.5 ± 5.9  |

*Definition of abbreviations:* AHI = apnea-hypopnea index; OSA = obstructive sleep apnea.

Values are mean ± 95% confidence interval.

Body mass index is the weight divided by the squared height.

AHI during REM was available in 14 of 18, 21 of 25, and 8 of 11 participants, respectively.

\*Fisher exact test, not significant ( $P = 0.6$ ). One-way analysis of variance with Student-Newman-Keuls *post hoc* analysis was used for continuous variables.

<sup>†</sup> $P < 0.001$  normal-weight versus both overweight/obese groups.

<sup>‡</sup> $P < 0.001$  overweight/obese patients with OSA versus both nonapneic groups.

intervals during supine non-REM sleep. We measured ventilation and CPAP level by a nasal mask, epiglottic pressure (Pepi), and EMGgg using fine-wire electrodes. Traits were measured as follows (see Figure E1 in the online supplement) (2, 16, 17):

1. Airway collapsibility: the critical CPAP level  $P_{crit}$  yielding zero airflow through a passive airway (x-intercept of a linear regression between peak-inspiratory flow vs. mask pressure on the third to fifth breaths following the CPAP drop, if breaths were flow limited).
2. Upper-airway muscle responsiveness: the increase in peak EMGgg (%maximum) per change in nadir Pepi across each CPAP drop.
3. Loop gain: the increase in ventilatory drive (measured as the ventilatory overshoot following a switch to optimal CPAP) in response to a steady-state reduction in ventilation.
4. Arousal threshold: the nadir Pepi on the breath preceding arousal from sleep.

### Model-based Analysis


To examine how the traits manifest the absence of OSA in overweight/obese individuals without apnea, we combined the traits using a mathematical model that graphically illustrates their contribution to OSA (16, 17) (see Figure E2). For this purpose we remeasured the traits in units of ventilation as follows (16):

1. Anatomy/collapsibility: the ventilation " $V_{passive}$ " through a passive airway (CPAP = 0 cm  $H_2O$ ) at eupneic ventilatory drive (y-intercept of a plot of ventilation vs. CPAP).
2. Upper-airway muscle effectiveness ("upper-airway gain"): the compensatory increase in ventilation across the drop (the activated level at the end of the drop minus the "passive" ventilation at the start) per increase in ventilatory drive (measured as the ventilatory overshoot following a switch to optimal CPAP).
3. Arousal threshold: the ventilatory drive preceding arousal.

The traits were subsequently combined to calculate two intermediate physiologic parameters: "Vactive" is the ventilation that can be achieved through a maximally active airway without arousal, and "Varousal" is the minimum ventilation that can be tolerated without arousal. The difference between Varousal and Vactive, called the physiologic "gap," predicts whether stable breathing is possible or patients will exhibit OSA (17). A positive "gap" quantifies the degree of ventilatory insufficiency and predicts the presence of OSA (the ventilation needed to avoid arousal cannot be achieved through the activated airway;  $V_{active} < V_{arousal}$ ). A negative "gap" quantifies the degree of ventilatory reserve and predicts that stable breathing is possible without arousal ( $V_{active} > V_{arousal}$ ).

### Statistical Analysis

One-way analysis of variance (ANOVA) assessed trait differences between groups. As necessary, transforms were applied before



**Figure 1.** Physiologic differences between overweight/obese nonapneic individuals (nOSA BMI > 25), overweight/obese patients with OSA (OSA BMI > 25), and normal-weight control subjects (nOSA control). (A) The upper airway is less collapsible ( $P_{crit}$  is lower) in overweight/obese nOSA versus overweight/obese OSA, but is more collapsible than normal-weight control subjects. (B) Upper-airway dilator muscles are markedly more responsive in overweight/obese nOSA compared with both overweight/obese OSA and normal-weight control subjects. Responsiveness is defined as the genioglossus electromyogram (EMGgg) response to negative epiglottic pressure,  $\Delta EMGgg/\Delta Pepi$  (EMGgg is reported relative to maximum achievable activity). (C) Loop gain is elevated (more negative) in both overweight/obese groups versus nOSA control subjects. (D) Overweight/obese nOSA individuals exhibit a similar arousal threshold to nOSA control subjects. The arousal threshold is elevated (more negative) in OSA versus nOSA control subjects. Men and women are denoted by circles and diamonds, respectively. Mean data are illustrated by horizontal bars. Data in B and D were square-root transformed before statistical analysis to achieve normally distributed data; these data are plotted on a square-root scale. Data in C were log-transformed before statistical analysis; these data are plotted on a logarithmic scale. One-way analysis of variance with Student-Newman-Keuls *post hoc* analysis were used to compare groups. Measures of upper-airway responsiveness could be made in 17 of 18 (nOSA BMI > 25), 23 of 25 (OSA BMI > 25), and 10 of 11 (nOSA control) individuals. Measures of loop gain could be made in 14 of 18 (nOSA BMI > 25), 23 of 25 (OSA BMI > 25), and 8 of 11 (nOSA control) individuals. BMI = body mass index; OSA = obstructive sleep apnea;  $P_{crit}$  = critical closing pressure.

statistical analysis to provide normally distributed data (square root transform for upper-airway muscle responsiveness [EMG<sub>gg</sub>/Pepi] and arousal threshold [Pepi], and log<sub>10</sub> transform for loop gain); reported means and 95% confidence intervals reflect back-transformed values. Two-way ANOVA assessed trait differences between groups adjusting for sex. Multiple logistic regression examined whether enhanced muscle responsiveness and effectiveness contributed to the absence of OSA in addition to upper-airway collapsibility (Pcrit). *P* less than 0.05 was considered statistically significant.

## Results

Subject characteristics are detailed in Table 1. Despite similar BMI, overweight/obese subjects without apnea have a less-collapsible airway (lower Pcrit) compared with overweight/obese patients with apnea (Figure 1A). However, the upper-airway anatomy/collapsibility of overweight/obese subjects without apnea remained moderately compromised (higher Pcrit) versus normal-weight control subjects. Overweight/obese subjects without apnea exhibited approximately threefold greater upper-airway muscle responsiveness versus patients with OSA and normal-weight control subjects (Figure 1B). Loop gain was similar between overweight/obese subjects without apnea and patients with OSA; both groups exhibited elevated loop gain versus normal-weight control subjects (Figure 1C). Arousal thresholds were elevated in patients with OSA versus normal-weight control subjects (Figure 1D). After adjusting for potential sex effects (two-way ANOVA), the aforementioned group differences remained significant for all traits; there were no significant effects of sex or significant interactions between sex and group.

Measurement of the traits for model-based analysis revealed similar results. The upper-airway muscle effectiveness (upper-airway gain) was threefold greater in overweight/obese subjects without apnea compared with patients with apnea (Figure 2). *V*<sub>passive</sub> was significantly reduced in overweight/obese OSA versus normal-weight control subjects ( $34 \pm 17\%$  vs.  $78 \pm 9\%$ ; *P* < 0.01; mean  $\pm$  95% confidence interval); *V*<sub>passive</sub> in overweight/obese subjects without apnea


( $56 \pm 18\%$ ) was midway between overweight/obese OSA and normal-weight control subjects. The ventilatory drive at arousal was similar between groups.

Model-based analyses are illustrated in Figures 3A–3C. Because of the enhanced muscle responses and moderately compromised anatomy, the overweight/obese subjects without apnea exhibited a small negative “gap” (–11% of eupneic ventilation; calculated from group summary data) consistent with the absence of OSA. By contrast, the overweight/obese patients with apnea exhibited a large positive “gap” (+38% of eupneic ventilation) consistent with OSA. As expected, the normal-weight nonapneic control subjects had a large negative gap (–21% of eupneic ventilation) consistent with a large reserve against OSA (see Figure 3D for individual data). We calculated that if the overweight/obese subjects without apnea had the same upper-airway responsiveness and effectiveness or the same upper-airway collapsibility (*V*<sub>passive</sub>) as the group with apnea, they would have a positive gap (+16% or +11% of eupneic ventilation) and would exhibit OSA (Figure 4).

Further analysis (multiple logistic regression) illustrates that upper-airway muscle responsiveness (Figure 5A) and effectiveness (Figure 5B) help to explain the presence of OSA beyond upper-airway collapsibility alone. Individuals with greater upper-airway collapsibility require a greater upper-airway muscle responsiveness and effectiveness to avoid OSA.

## Discussion

The major finding of the current study is that enhanced muscle responsiveness is a distinct feature of overweight/obese individuals without moderate-severe OSA. Overweight/obese individuals without apnea exhibit approximately three times greater pharyngeal muscle responsiveness during sleep compared with their OSA counterparts and normal-weight control subjects. Our study also confirmed that overweight/obese individuals without apnea have an upper-airway structural advantage (lower Pcrit) relative to overweight/obese patients with OSA (4–8); we extend these findings to demonstrate that the collapsibility of overweight/obese individuals without apnea is considerably compromised (higher Pcrit) when considered alongside normal-weight




**Figure 2.** Upper-airway muscle effectiveness (upper-airway gain) is threefold greater in overweight/obese individuals without apnea (nOSA BMI > 25) versus overweight/obese patients with OSA (OSA BMI > 25). Data from normal-weight control subjects (nOSA control) are shown for comparison. Median data are illustrated by horizontal bars. Men and women are denoted by circles and diamonds, respectively. Measures of upper-airway muscle effectiveness could be made in only 14 of 18 (nOSA BMI > 25), 23 of 25 (OSA BMI > 25), and 8 of 11 (nOSA control) individuals because of difficulties measuring loop gain. Groups were compared using one-way analysis of variance on ranks with Dunn *post hoc* analysis. BMI = body mass index; OSA = obstructive sleep apnea.

control subjects. Indeed, the mean Pcrit of –3.7 cm H<sub>2</sub>O in overweight/obese subjects without apnea was within the vulnerable range for moderate-severe OSA (–5 cm H<sub>2</sub>O or above) (2, 9). Overall, our data show that overweight/obese individuals without apnea escape OSA via highly effective upper-airway muscle responses that compensate for a moderately compromised anatomy.

### Novel Physiologic Insights

Examination of our model-based analysis demonstrates that overweight/obese individuals without apnea are reliant on their enhanced muscle responsiveness to avoid OSA. We note that overweight/obese individuals without apnea have a minimal



**Figure 3.** Combining the four physiologic traits graphically using a mathematical model illustrates how they interact to manifest the absence or presence of obstructive sleep apnea (OSA). (A–C) Group data are shown for each trait measured in units of ventilation. Veupnea is the ventilation on optimal continuous positive airway pressure. Vpassive is the functional anatomy/collapsibility represented as the ventilation off continuous positive airway pressure at normal ventilatory drive. The *upper-airway gain line* describes the functional compensatory increase in ventilation with increased ventilatory drive (which activates upper-airway muscles); this line indicates the achievable ventilation. The *loop gain line* illustrates the level of ventilatory drive that will eventually develop for any reduction in ventilation (e.g., because of rising carbon dioxide). If these diagonal lines intersect at a level of ventilatory drive below the arousal threshold (vertical dashed line) then stable breathing is possible during sleep; otherwise, OSA should occur. (A) On average, the physiologic traits of overweight/obese subjects without apnea (nOSA) are consistent with stable breathing: Note that the upper airway is so effective (steep slope of *upper-airway gain line*) that stable breathing occurs (intersection of *diagonal lines* before the arousal threshold), despite a moderately compromised anatomy. The maximum ventilation achievable without arousal (Vactive) marginally exceeds the minimum level of ventilation that can be tolerated without arousal (Varousal). (B) The traits of overweight/obese patients with apnea are consistent with OSA. Stable breathing is not possible because the achievable ventilation (Vactive) is below the level needed to prevent arousal (Varousal). (C) The traits of healthy normal-weight control subjects are consistent with stable breathing. The favorable anatomy provides for stable breathing regardless of their modest upper-airway responsiveness (the slope is no steeper than in OSA). (D) Ultimately, the “gap” (Varousal – Vactive) predicts whether OSA will occur (mean data are illustrated by *horizontal bars*). In OSA, the “gap” is positive and significantly greater than in overweight/obese subjects without apnea (analysis of variance), in whom the gap is negative. Men and women are denoted by *circles* and *diamonds*, respectively. Of note, measures of the “gap” could be made in 14 of 18 (nOSA BMI > 25), 23 of 25 (OSA BMI > 25), and 8 of 11 (nOSA control) individuals because of difficulties measuring loop gain.  $V_{active} = V_{passive} + (\text{upper-airway gain}) \times (\text{arousal threshold} - 100)$ .  $Varousal = 100 - (\text{loop gain})^{-1} \times (\text{arousal threshold} - 100)$ . BMI = body mass index.

reserve (the “gap” is approximately 10% of eupneic ventilation) to protect against the development of OSA, such that any added deficiency would cause OSA. Using the model, we showed that if overweight/obese individuals without apnea are given the same upper-airway responsiveness or collapsibility as the group with apnea, they would exhibit OSA (Figure 4). Consistent with this view, in REM sleep, when muscle activity becomes more severely compromised (15, 21), the otherwise nonapneic group exhibited OSA of moderate severity (Table 1). By contrast, the groups with minimal muscle responsiveness did not exhibit an increased AHI in REM. These findings taken together provide conclusive, quantitative evidence of the essential role of upper-airway muscle responsiveness for the prevention of OSA in those with a vulnerable anatomy.


Based on the near-identical muscle responsiveness of overweight/obese patients with OSA and normal-weight control subjects (Figure 1B), our study demonstrates that impaired genioglossus

muscle responsiveness is probably not a primary cause of OSA in overweight/obese individuals *per se*. Rather, the reduced upper-airway compensatory responses in this population only seem to be reduced in previous work (4), because responsiveness has been compared with weight-matched subjects without apnea, who this study shows have a “supranormal” responsiveness. Thus, we interpret that OSA occurs with obesity primarily because of its effect on upper-airway structure/collapsibility (and elevated loop gain, noting the difference in Figure 1C). However, enhanced upper-airway muscle responsiveness, when present, acts as a key effect modifier that is capable of averting OSA in many individuals.

Our data confirm previous work illustrating that individuals without apnea can achieve greater upper-airway dilator muscle activation during sleep (increased EMGgg and ventilation above passive conditions) versus BMI-matched patients

with OSA (4, 14). We emphasize, however, that the increase in EMGgg and ventilation during sleep reflects two key traits: the upper-airway responsiveness and the arousal threshold (i.e.,  $V_{active} - V_{passive} = [\text{upper-airway gain}] \times [\text{arousal threshold} - 100]$ ). A novel finding of our study is that, among the overweight/obese population, individuals without apnea exhibit an enhanced upper-airway responsiveness *per se* (change in EMGgg per unit change in Pepi), whereas the arousal threshold is similar between groups.

Our data also suggest that loop gain is increased in overweight/obese individuals (vs. normal-weight control subjects) regardless of whether or not such individuals have OSA. An increase in loop gain may be an important additional mechanism by which individuals who gain weight become predisposed to OSA. A possible mechanism is the increased circulating leptin in overweight individuals, which is known to raise chemosensitivity (22).



**Figure 4.** Essential roles of the greater upper-airway muscle responsiveness and reduced upper-airway collapsibility in overweight/obese subjects without apnea versus patients with apnea. The physiologic traits of subjects without apnea are combined mathematically (Figure 3A) to illustrate why stable breathing is possible without arousal from sleep in overweight/obese individuals without obstructive sleep apnea (OSA); note that the intersection of the *loop gain* line and *upper-airway gain* line (stable breathing point) lies to the left of the arousal threshold (dashed vertical line). (A) If overweight/obese subjects without apnea had the upper-airway effectiveness (or “gain”) of the OSA group (\*upper-airway gain; solid line) then the maximum achievable ventilation during non-REM sleep ( $V_{active}$ ) would instead be below the minimum tolerable ventilation ( $V_{arousal}$ ), providing a physiologic “gap” (+16% of  $V_{eupnea}$ ) that cannot be overcome without arousal. Hence, the effective upper-airway muscle response in subjects without apnea can be considered essential for the observed absence of OSA. (B) A similar result is observed for the anatomy/collapsibility. If  $V_{passive}$  is reduced to the level of the OSA group (by 22% of eupneic ventilation; difference in mean values), the gap would become positive (+11% of  $V_{eupnea}$ ) such that OSA would occur. Thus the less-vulnerable anatomy (higher  $V_{passive}$ ) in overweight/obese subjects without apnea is also essential for their avoidance of OSA.

### Mechanisms of Advantageous Physiology in Overweight and Obese Individuals without Apnea

The mechanisms of less-pronounced upper-airway collapsibility in overweight/obese subjects without apnea versus weight-matched patients with apnea are currently being elucidated. Compared with patients with OSA, weight-matched individuals without apnea tend to have reduced fat distribution to the neck (23), and at a local pharyngeal level exhibit reduced tongue fat (8) and parapharyngeal fat pad volumes (5–8, 24, 25). Overweight/obese individuals without apnea also have craniofacial features that provide a greater anatomic reserve for fat deposition by increased intramandibular volume (7, 26, 27). These advantageous structural differences are likely under genetic control given the familial nature of upper-airway structure and OSA (28–31).

The origin of enhanced muscle responsiveness in overweight/obese individuals without apnea is poorly

understood. Although a more-effective translation from neuromuscular activity (EMG<sub>gg</sub>) to increased airflow (e.g., reduced mechanical load, increased muscle strength and coordination, reduced fatigability) may mitigate OSA in many individuals (32–34), our group data indicate that much of the enhanced functional responsiveness (upper-airway gain) in overweight/obese individuals without apnea is neurophysiologic in origin (the EMG<sub>gg</sub> response to negative pressure). We base this interpretation on the finding that the EMG<sub>gg</sub> responsiveness is enhanced by a similar magnitude (threefold) as the functional effectiveness (Figure 2 vs. Figure 1B). Enhanced responsiveness may occur at the level of the negative pressure sensory afferents or the medullary negative pressure and CO<sub>2</sub> inputs to the hypoglossal motor nucleus (e.g., via the nucleus of the solitary tract and peribex region [35]). It is also unclear whether the mechanism responsible is inherent (e.g., genetic) or adaptive (e.g., a neuroplastic


recalibration of reflexes consequent to weight gain). The absence of large responsiveness values in normal-weight control subjects suggests an adaptive mechanism may be at play that is largely absent in patients with OSA. We also considered the possibility that OSA itself may reduce muscle responsiveness, perhaps by sleep deprivation (36, 37) or neurologic damage (38). However, the current study examined patients with OSA who were treated with CPAP to minimize such effects. Moreover, available evidence suggests that CPAP treatment of OSA has little impact on upper-airway neuromuscular behavior (threshold ventilatory drive required for airway opening) (39). Elucidating the precise mechanism involved may provide for novel therapeutic strategies to enhance muscle responses and treat OSA.

### Clinical Implications

Our data illustrate the magnitude of the interventions to the traits that are sufficient for effective OSA resolution in overweight/obese patients with apnea. That is, a threefold increase in dilator muscle responsiveness and an approximately 4 cm H<sub>2</sub>O decrease in  $P_{crit}$  (or increased passive ventilation of ~20% of eupneic levels) should be sufficient to provide patients with apnea with the physiology of individuals without apnea and resolve OSA. We envisage that, in some individuals, OSA treatment could be achieved in this manner by improving anatomy (e.g., with positional therapy or mandibular advancement) in combination with improved muscle responses via novel methods to activate, sensitize, or even train the upper-airway muscles (40–42).

### Conclusions

The current study demonstrates that overweight/obese individuals without apnea counteract a moderate anatomic deficit with powerfully enhanced upper-airway dilator muscle responses during sleep to avoid OSA; by contrast there was no evidence of lowered loop gain or elevated arousal threshold compared with patients with OSA. The mechanisms responsible for enhanced responsiveness may provide clues for future treatment. ■



**Figure 5.** Upper airway collapsibility ( $P_{crit}$ ) and upper airway dilator muscle behavior determine the presence or absence of obstructive sleep apnea (OSA). (A) Upper airway responsiveness and  $P_{crit}$ . (B) Upper airway effectiveness (upper airway gain) and  $P_{crit}$ . Lines based on logistic regression illustrate the boundary between overweight/obese patients with OSA and overweight/obese individuals without OSA. Individuals with more collapsible airways require the most responsive and effective upper airway muscles to avoid OSA. Three outliers are labeled in A: (i) one individual had minimal responsiveness but avoided OSA likely because of a moderate upper airway effectiveness (+0.6) and very low loop gain (-2), two individuals with good responsiveness developed OSA likely caused by (ii) a poor upper airway muscle effectiveness (-0.2) and elevated loop gain (-6), and (iii) a low arousal threshold (-8 cm H<sub>2</sub>O). Upper airway responsiveness data were square-root transformed to achieve normally distributed data; these data are shown on a square-root scale. Regression models: (A)  $Y = 2.36 + (0.38 \times P_{crit}) + (3.19 \times \text{Responsiveness}^{0.5})$  [ $P_{crit}$ :  $P = 0.02$ ; Responsiveness:  $P = 0.04$ ], (B)  $Y = 3.31 + (0.63 \times P_{crit}) - (3.56 \times \text{Effectiveness})$  [ $P_{crit}$ :  $P = 0.02$ ; Effectiveness:  $P = 0.03$ ]. BMI = body mass index; EMGgg = genioglossus EMG;  $P_{crit}$  = critical closing pressure.

**Author disclosures** are available with the text of this article at [www.atsjournals.org](http://www.atsjournals.org).

**Acknowledgment:** The authors are grateful for the technical support provided by Lauren Hess, Diandra Grinage, Erik

Smales, Geoffrey Kehlman, Karen Stevenson, Louise Dover, Salonee Parikh, and Scott Smith.

## References

- Peppard PE, Young T, Barnet JH, Palta M, Hagen EW, Hla KM. Increased prevalence of sleep-disordered breathing in adults. *Am J Epidemiol* 2013;177:1006–1014.
- Eckert DJ, White DP, Jordan AS, Malhotra A, Wellman A. Defining phenotypic causes of obstructive sleep apnea. Identification of novel therapeutic targets. *Am J Respir Crit Care Med* 2013;188:996–1004.
- Jordan AS, Wellman A, Edwards JK, Schory K, Dover L, MacDonald M, Patel SR, Fogel RB, Malhotra A, White DP. Respiratory control stability and upper airway collapsibility in men and women with obstructive sleep apnea. *J Appl Physiol* (1985) 2005;99:2020–2027.
- McGinley BM, Schwartz AR, Schneider H, Kirkness JP, Smith PL, Patil SP. Upper airway neuromuscular compensation during sleep is defective in obstructive sleep apnea. *J Appl Physiol* (1985) 2008;105:197–205.
- Pahkala R, Seppa J, Ikonen A, Smirnov G, Tuomilehto H. The impact of pharyngeal fat tissue on the pathogenesis of obstructive sleep apnea. *Sleep Breath* 2014;18:275–282.
- Li Y, Lin N, Ye J, Chang Q, Han D, Sperry A. Upper airway fat tissue distribution in subjects with obstructive sleep apnea and its effect on retropalatal mechanical loads. *Respir Care* 2012;57:1098–1105.
- Kim C, Jackson N, Marcus CL, Pien GW, Staley B, Sun Y, Wieland W, Schwab RJ. Relationship between upper airway collapsibility ( $p_{crit}$ ) and pharyngeal structures [abstract]. *Am J Respir Crit Care Med* 2012;185:A2430.



8. Kim C, Keenan BT, Jackson N, Chan EL, Staley B, Poptani H, Torigian DA, Pack AI, Schwab RJ. Tongue fat and its relationship to obstructive sleep apnea. *Sleep* (In press)
9. Kirkness JP, Schwartz AR, Schneider H, Punjabi NM, Maly JJ, Laffan AM, McGinley BM, Magnuson T, Schweitzer M, Smith PL, *et al.* Contribution of male sex, age, and obesity to mechanical instability of the upper airway during sleep. *J Appl Physiol* (1985) 2008; 104:1618–1624.
10. Wellman A, Jordan AS, Malhotra A, Fogel RB, Katz ES, Schory K, Edwards JK, White DP. Ventilatory control and airway anatomy in obstructive sleep apnea. *Am J Respir Crit Care Med* 2004; 170:1225–1232.
11. Eckert DJ, Owens RL, Kehlmann GB, Wellman A, Rahangdale S, Yim-Yeh S, White DP, Malhotra A. Eszopiclone increases the respiratory arousal threshold and lowers the apnoea/hypopnoea index in obstructive sleep apnoea patients with a low arousal threshold. *Clin Sci (Lond)* 2011;120:505–514.
12. Edwards BA, Sands SA, Eckert DJ, White DP, Butler JP, Owens RL, Malhotra A, Wellman A. Acetazolamide improves loop gain but not the other physiological traits causing obstructive sleep apnoea. *J Physiol* 2012;590:1199–1211.
13. Younes M, Loewen AH, Ostrowski M, Laprairie J, Maturino F, Hanly PJ. Genioglossus activity available via non-arousal mechanisms vs. that required for opening the airway in obstructive apnea patients. *J Appl Physiol* (1985) 2012;112:249–258.
14. Patil SP, Schneider H, Marx JJ, Gladmon E, Schwartz AR, Smith PL. Neuromechanical control of upper airway patency during sleep. *J Appl Physiol* (1985) 2007;102:547–556.
15. Eckert DJ, McEvoy RD, George KE, Thomson KJ, Catcheside PG. Genioglossus reflex inhibition to upper-airway negative-pressure stimuli during wakefulness and sleep in healthy males. *J Physiol* 2007;581:1193–1205.
16. Wellman A, Eckert DJ, Jordan AS, Edwards BA, Passaglia CL, Jackson AC, Gautam S, Owens RL, Malhotra A, White DP. A method for measuring and modeling the physiological traits causing obstructive sleep apnea. *J Appl Physiol* (1985) 2011; 110:1627–1637.
17. Wellman A, Edwards BA, Sands SA, Owens RL, Nemati S, Butler J, Passaglia CL, Jackson AC, Malhotra A, White DP. A simplified method for determining phenotypic traits in patients with obstructive sleep apnea. *J Appl Physiol* (1985) 2013;114: 911–922.
18. Edwards BA, Connolly JG, Campana LM, Sands SA, Trinder JA, White DP, Wellman A, Malhotra A. Acetazolamide attenuates the ventilatory response to arousal in patients with obstructive sleep apnea. *Sleep* 2013;36:281–285.
19. Wellman A, Malhotra A, Jordan AS, Stevenson KE, Gautam S, White DP. Effect of oxygen in obstructive sleep apnea: role of loop gain. *Respir Physiol Neurobiol* 2008;162:144–151.
20. AASM. Sleep-related breathing disorders in adults: recommendations for syndrome definition and measurement techniques in clinical research. The Report of an American Academy of Sleep Medicine Task Force. *Sleep* 1999;22:667–689.
21. Wiegand L, Zwillich CW, Wiegand D, White DP. Changes in upper airway muscle activation and ventilation during phasic REM sleep in normal men. *J Appl Physiol* (1985) 1991;71:488–497.
22. O'donnell CP, Schaub CD, Haines AS, Berkowitz DE, Tankersley CG, Schwartz AR, Smith PL. Leptin prevents respiratory depression in obesity. *Am J Respir Crit Care Med* 1999;159: 1477–1484.
23. Hoffstein V, Mateika S. Differences in abdominal and neck circumferences in patients with and without obstructive sleep apnoea. *Eur Respir J* 1992;5:377–381.
24. Horner RL, Mohiaddin RH, Lowell DG, Shea SA, Burman ED, Longmore DB, Guz A. Sites and sizes of fat deposits around the pharynx in obese patients with obstructive sleep apnoea and weight matched controls. *Eur Respir J* 1989;2:613–622.
25. Schwab RJ, Pasirstein M, Pierson R, Mackley A, Hachadoorian R, Arens R, Maislin G, Pack AI. Identification of upper airway anatomic risk factors for obstructive sleep apnea with volumetric magnetic resonance imaging. *Am J Respir Crit Care Med* 2003;168:522–530.
26. Watanabe T, Isono S, Tanaka A, Tanzawa H, Nishino T. Contribution of body habitus and craniofacial characteristics to segmental closing pressures of the passive pharynx in patients with sleep-disordered breathing. *Am J Respir Crit Care Med* 2002;165:260–265.
27. Lee RW, Petocz P, Prvan T, Chan AS, Grunstein RR, Cistulli PA. Prediction of obstructive sleep apnea with craniofacial photographic analysis. *Sleep* 2009;32:46–52.
28. Redline S, Tishler PV, Tosteson TD, Williamson J, Kump K, Browner I, Ferrette V, Krejci P. The familial aggregation of obstructive sleep apnea. *Am J Respir Crit Care Med* 1995;151:682–687.
29. Schwab RJ, Pasirstein M, Kaplan L, Pierson R, Mackley A, Hachadoorian R, Arens R, Maislin G, Pack AI. Family aggregation of upper airway soft tissue structures in normal subjects and patients with sleep apnea. *Am J Respir Crit Care Med* 2006;173:453–463.
30. Strohl KP, Saunders NA, Feldman NT, Hallett M. Obstructive sleep apnea in family members. *N Engl J Med* 1978;299:969–973.
31. Chi L, Comyn FL, Keenan BT, Cater J, Maislin G, Pack AI, Schwab RJ. Heritability of craniofacial structures in normal subjects and patients with sleep apnea. *Sleep* (In press)
32. Brown EC, Cheng S, McKenzie DK, Butler JE, Gandevia SC, Bilston LE. Respiratory movement of upper airway tissue in obstructive sleep apnea. *Sleep* 2013;36:1069–1076.
33. Bilston LE, Gandevia SC. Biomechanical properties of the human upper airway and their effect on its behavior during breathing and in obstructive sleep apnea. *J Appl Physiol* (1985) 2014;116:314–324.
34. Carrera M, Barbé F, Sauleda J, Tomás M, Gómez C, Santos C, Agustí AG. Effects of obesity upon genioglossus structure and function in obstructive sleep apnoea. *Eur Respir J* 2004;23:425–429.
35. Chamberlin NL, Eikermann M, Fassbender P, White DP, Malhotra A. Genioglossus pre-motoneurons and the negative pressure reflex in rats. *J Physiol* 2007;579:515–526.
36. Leiter JC, Knuth SL, Bartlett D Jr. The effect of sleep deprivation on activity of the genioglossus muscle. *Am Rev Respir Dis* 1985;132: 1242–1245.
37. Eckert DJ, Parikh S, White DP, Jordan AS, Merchia P, Malhotra A. Sleep deprivation impairs genioglossus muscle responsiveness [abstract]. *Am J Respir Crit Care Med* 2011;183:A6163.
38. Saboisky JP, Stashuk DW, Hamilton-Wright A, Carusona AL, Campana LM, Trinder J, Eckert DJ, Jordan AS, McSharry DG, White DP, *et al.* Neurogenic changes in the upper airway of patients with obstructive sleep apnea. *Am J Respir Crit Care Med* 2012;185:322–329.
39. Loewen A, Ostrowski M, Laprairie J, Atkar R, Gnitecki J, Hanly P, Younes M. Determinants of ventilatory instability in obstructive sleep apnea: inherent or acquired? *Sleep* 2009;32:1355–1365.
40. Grace KP, Hughes SW, Horner RL. Identification of a pharmacological target for genioglossus reactivation throughout sleep. *Sleep* 2014; 37:41–50.
41. Strollo PJ Jr, Soose RJ, Maurer JT, de Vries N, Cornelius J, Froyovich O, Hanson RD, Padhya TA, Steward DL, Gillespie MB, *et al.*; STAR Trial Group. Upper-airway stimulation for obstructive sleep apnea. *N Engl J Med* 2014;370:139–149.
42. Puhan MA, Suarez A, Lo Cascio C, Zahn A, Heitz M, Braendli O. Didgeridoo playing as alternative treatment for obstructive sleep apnoea syndrome: randomised controlled trial. *BMJ* 2006;332: 266–270.