

# UC San Diego

## UC San Diego Previously Published Works

### Title

Associations Between HIV Stigma and Mental Health Among Older Hispanics and Whites Living With HIV

### Permalink

<https://escholarship.org/uc/item/9qc7z84h>

### Journal

Innovation in Aging, 5(Supplement\_1)

### ISSN

2399-5300

### Authors

Karcher, Kayle  
Kamalyan, Lily  
Gonzalez, Veronica  
et al.

### Publication Date

2021-12-17

### DOI

10.1093/geroni/igab046.3463

Peer reviewed

mortality. Models were repeated, stratifying by median daily total time spent sedentary and active.

**Results:** After 9.4±3.7 years of follow-up, 1,487 (51.0%) men died. Men averaged 16.9±5.1 and 8.2±4.2 sedentary and active bouts/day, respectively. After full covariate adjustment, each quartile reflecting a higher sedentary (Q4 vs Q1 HR: 0.68, 95%CI: 0.58-0.81, p-trend<0.001) and active bout (Q4 vs Q1 HR: 0.57, 95%CI: 0.48-0.68, p-trend<0.001) frequency was associated with lower mortality risk. There was no evidence that effects differed by total sedentary time (p-interaction for sedentary bout frequency and total sedentary time>0.05).

**Conclusions:** More frequent, prolonged sedentary and active bouts are associated with a lower mortality risk in older men and is not moderated by total sedentary time.

#### ASSOCIATIONS BETWEEN HIV STIGMA AND MENTAL HEALTH AMONG OLDER HISPANICS AND WHITES LIVING WITH HIV

Kayle Karcher,<sup>1</sup> Lily Kamalyan,<sup>2</sup> Veronica Gonzalez,<sup>3</sup> Lilla Brody,<sup>4</sup> Robert Heaton,<sup>5</sup> Raeanne Moore,<sup>6</sup> Dilip Jeste,<sup>7</sup> and Maria Marquine,<sup>7</sup> 1. *University of California-San Diego, San Bernardino, California, United States*, 2. *San Diego State University/University of California San Diego Joint Doctoral Program in Clinical Psychology, San Diego, California, United States*, 3. *UCSD, San Diego, California, United States*, 4. *Weill Cornell Medicine, New York, New York, United States*, 5. *University of California San Diego, San Diego, California, United States*, 6. *UC San Diego, San Diego, California, United States*, 7. *University of California San Diego, La Jolla, California, United States*

Hispanics/Latinos/as/x (henceforth Hispanics) have higher rates of HIV infection than non-Hispanic (NH) Whites, particularly in older age. People living with HIV (PWH) are at increased risk of stigma and poor mental health, but these associations have not been thoroughly examined in older PWH. We investigated ethnic differences in HIV stigma and its association with mental health in older Hispanic and NH White PWH. Participants included 116 PWH ages 50-75 (58 Hispanic and 58 NH White) from southern California (for the overall cohort: 82.7% male; 57.7% AIDS, 93.9% on antiretroviral therapy). Participants completed self-report measures of HIV-stigma, depression (Beck Depression Inventory-II; BDI-II), and cumulative alcohol use (i.e., lifetime total quantity/total days). Covariates examined included sociodemographic and HIV-disease characteristics. An independent sample t-test showed no significant ethnic differences in HIV stigma ( $p=.82$ ). Separate multivariable linear regression models on mental health outcomes (adjusting for significant covariates) showed no significant interaction between HIV stigma and ethnicity on BDI-II scores ( $p=.83$ ) or cumulative alcohol use ( $p=.51$ ). Follow up models removing the interaction term, showed that increased HIV stigma was associated with higher BDI-II scores ( $B=0.34$ , 95% CI=0.21-0.48;  $p<.001$ ), but not with cumulative alcohol use ( $p=.49$ ) in the overall sample. Findings indicate a significant link between HIV stigma and depression symptoms in older PWH, with comparable associations among Hispanics and NH Whites. Future studies examining factors that may moderate the link between HIV stigma and depression in diverse older

PWH would help guide the development of interventions aimed at improving mental health in this population.

#### BIOPSYCHOSOCIAL CORRELATES OF COGNITIVE FUNCTION AMONG KOREAN OLDER ADULTS: HISTORY OF HYPERTENSION AND DIABETES

Kyuyoung Cho,<sup>1</sup> and Hye Won Chai,<sup>2</sup> 1. *Dong-A University, Busan, Pusan-jikhalsi, Republic of Korea*, 2. *The University of Texas at Austin, Austin, Texas, United States*

Based on biopsychosocial perspectives on health, this study examined risk and protective factors of cognitive function among Korean older adults. Specifically, we focused on comparing the role of these factors based on the respondents' history of having hypertension or diabetes. This study used 2009 Korean National Health Insurance Service data that included a sample of older adults who maintained qualification for health insurance and medical aid in 2002 ( $n=26,242$ ). Cognitive function was measured using KDSQ-C and biopsychosocial factors included metabolic syndrome, drinking, smoking, and walking. The sample was divided into two groups based on their medical history, and thus four sets of linear regression models were analyzed to explore the associations between biopsychosocial factors and cognitive functioning. Among individuals with a history of hypertension, metabolic syndrome, drinking, and walking were associated with cognitive functioning. For those without a history of hypertension, only drinking and walking were associated with cognitive functioning. For diabetes, smoking and walking were associated with cognitive functioning among older adults with a history of diabetes. For those without a history of diabetes, drinking and walking were associated with cognitive functioning. In sum, metabolic syndrome was a particularly significant correlate of cognitive function among Korean older adults with a history of hypertension. Walking was a consistently significant factor regardless of medical history. These results highlight the importance of considering medical history of chronic conditions such as hypertension and diabetes in identifying factors associated with older adults' cognitive function and further developing tailored prevention programs for cognitive decline.

#### CAN A DATA-DRIVEN MEASURE OF NEUROANATOMIC DEMENTIA RISK BE CONSIDERED A MEASURE OF BRAIN AGING?

Ramon Casanova,<sup>1</sup> Andrea Anderson,<sup>2</sup> Jamie Justice,<sup>2</sup> Gwen Windham,<sup>3</sup> Rebecca Gottesman,<sup>4</sup> Thomas Mosley,<sup>5</sup> Lynne Wagenknecht,<sup>6</sup> and Stephen Kritchevsky,<sup>2</sup> 1. *Wake Forest University, Winston-Salem, North Carolina, United States*, 2. *Wake Forest School of Medicine, Wake Forest School of Medicine, North Carolina, United States*, 3. *University of Mississippi Medical Center, Jackson, Mississippi, United States*, 4. *NIH, Bethesda, Maryland, United States*, 5. *The University of Mississippi Medical Center, Jackson, Mississippi, United States*, 6. *Wake Forest School of Medicine, Winston-Salem, North Carolina, United States*

There is an increasing interest in identifying aging-related factors which may be permissive of Alzheimer's Disease (AD) emergence. We previously used machine learning to derive an index of neuroanatomic risk of dementia called