

# Lawrence Berkeley National Laboratory

## Recent Work

### Title

CANTORI FOR SYMPLECTIC MAPS

### Permalink

<https://escholarship.org/uc/item/9r66m544>

### Authors

Chen, Q.  
MacKay, R.B.  
Meiss, J.D.

### Publication Date

1989-05-01



# Lawrence Berkeley Laboratory

UNIVERSITY OF CALIFORNIA

## Accelerator & Fusion Research Division

**For Reference**

Not to be taken from this room


Submitted to Nonlinearity

**Cantori for Symplectic Maps**

Q. Chen, R.S. MacKay, and J.D. Meiss

May 1989

RECEIVED  
FEB 2 1989  
BOOKS & PERIODICALS



LBL-27299  
91

## **DISCLAIMER**

This document was prepared as an account of work sponsored by the United States Government. While this document is believed to contain correct information, neither the United States Government nor any agency thereof, nor the Regents of the University of California, nor any of their employees, makes any warranty, express or implied, or assumes any legal responsibility for the accuracy, completeness, or usefulness of any information, apparatus, product, or process disclosed, or represents that its use would not infringe privately owned rights. Reference herein to any specific commercial product, process, or service by its trade name, trademark, manufacturer, or otherwise, does not necessarily constitute or imply its endorsement, recommendation, or favoring by the United States Government or any agency thereof, or the Regents of the University of California. The views and opinions of authors expressed herein do not necessarily state or reflect those of the United States Government or any agency thereof or the Regents of the University of California.

## Cantori for symplectic maps\*

Qi Chen<sup>1</sup>, R.S. MacKay<sup>2</sup>, and J.D. Meiss<sup>3</sup>

May 3, 1989

\* This work was supported by the U.S. Department of Energy under Contract DE-AC03-76SF00098 and grant DE-FG05-80ET-53088, the UK Science and Engineering Research Council, and a NATO grant, RG 85/0461, for international collaboration.

1 Institute for Fusion Studies, University of Texas, Austin, TX 78712

2 Mathematics Institute, University of Warwick, Coventry CV4 7AL, ENGLAND

3 Lawrence Berkeley Laboratory, University of California, Berkeley, CA 94720. Permanent Address: Institute for Fusion Studies, University of Texas, Austin, Tx 78712

# Cantori for symplectic maps\*

Qi Chen<sup>1</sup>, R.S. MacKay<sup>2</sup>, and J.D. Meiss<sup>3</sup>

May 3, 1989

**Abstract:** We construct invariant sets for the 2d-dimensional generalization of the sawtooth map which are semi-conjugate to any incommensurate rotation vector. When  $d \leq 2$  we show that these are Cantor sets. These invariant sets are hyperbolic, and we give a structural stability argument to show the existence of cantori for a non-trivial class of smooth symplectic maps.

---

\* This work was supported by the US Department of Energy under contract DE-AC03-76SF00098 and grant DE-FG05-80ET-53088, the UK Science and Engineering Research Council, and a NATO grant, RG 85/0461, for international collaboration.

<sup>1</sup> Institute for Fusion Studies, University of Texas, Austin, TX 78712

<sup>2</sup> Mathematics Institute, University of Warwick, Coventry CV4 7AL, ENGLAND

<sup>3</sup> Lawrence Berkeley Laboratory, University of California, Berkeley, CA 94720. Permanent Address: Institute for Fusion Studies, University of Texas, Austin, TX 78712

## §1. Introduction

Aubry and Mather[1, 2] show that orbits with any rotation number exist for area-preserving twist maps. Extension of this theory to symplectic maps with more than one degree of freedom appears to be a difficult task. Bangert[3] and Mather[4] have results about the set of rotation vectors for orbits of minimal action; however, an example of Hedlund[5] shows that in general one can not hope to obtain minimizing orbits with all rotation vectors. Bernstein and Katok[6] show that for maps close enough to integrable, the minimizing periodic orbits satisfy some regularity properties which are sufficient to allow the existence of a limiting orbit as the rotation vector approaches any limit. However, they can not prove anything about the rotation vectors of the limiting orbits.

Here we consider the opposite situation, when the "potential" dominates the "kinetic" energy. In this limit, the minimizing orbits of an area preserving map approach the minimizing orbits of the "sawtooth mapping" which is a piecewise linear, discontinuous mapping. As was shown independently by Aubry and Percival [7-10], an explicit formula can be obtained for orbits of the sawtooth mapping with irrational rotation number; these are dense on Cantor sets, and are called cantori. We consider the generalization of the sawtooth mapping to  $d$  degrees of freedom in §2, and obtain a formula for a set of orbits with incommensurate rotation vectors. When  $d \leq 2$ , these are shown to cover Cantor sets in §3.

We use the structural stability of hyperbolic sets to show in §4 that the cantori persist under perturbations of the mapping which can be arbitrary in a region around the discontinuity, but which are  $C^1$  small on the cantorus.

## §2. Many-dimensional sawtooth map

Let  $x = (x_1, x_2, \dots, x_d)$  represent a configuration point in  $\mathcal{R}^d$ . Consider a mapping with the generating function

$$h(x, x') = \frac{1}{2}|x - x'|^2 + \frac{\lambda}{2}\{x\}' Q \{x\} \quad (1)$$

where  $x, x' \in \mathcal{X}^d$ ,  $Q$  is a positive definite matrix,  $\lambda > 0$ , superscript  $t$  represents transpose, and  $\{x\}$  represents a fractional part of  $x$  with respect to the group of translations by integer vectors,  $\mathbb{Z}^d$ . For definiteness, choose some  $\sigma = (\sigma_1, \dots, \sigma_d)$  with  $\sigma_j \in \{+, -\}$  and define the fractional part

$$\{x\}^\sigma_i \equiv x_i - \left[ x_i + \frac{1}{2} \right]^{\sigma_i} \quad (2)$$

where  $[x]^+$  ( $[x]^-$ ) is the right (left) continuous integer part. A function  $f(x)$  for which  $f(x + \sigma \epsilon) \rightarrow f(x)$  as  $\epsilon \rightarrow 0$  for all  $x$  and  $\epsilon > 0$ , will be called  $\sigma$ -continuous. The fractional part  $\{x\}^\sigma$  is  $\sigma$ -continuous.

A trajectory generated by (1) consists of a configuration  $(x^s, x^{s+1}, \dots, x^t)$ ,  $s < t$  which is a stationary point of the action

$$W[x^s, x^{s+1}, \dots, x^t] = \sum_{j=s}^{t-1} h(x^j, x^{j+1})$$

Differentiation yields the second difference equation

$$x^{t+1} - 2x^t + x^{t-1} = \lambda Q\{x^t\} \quad (3)$$

provided none of the configuration points falls on the discontinuities. This defines a map  $S: (x^{t-1}, x^t) \rightarrow (x^t, x^{t+1})$  which is the 2d-dimensional generalization of the sawtooth mapping[10]. If  $Q$  is diagonal the components of (3) decouple and it can be treated as a pair area-preserving sawtooth mappings; however in general, (3) can not be decoupled.

Consider an incommensurate rotation vector  $\omega \in \mathcal{X}^d$  (i.e.  $k \cdot \omega \notin \mathbb{Z} \forall k \in \mathbb{Z}^d \setminus 0$ ). We look for invariant sets with this rotation vector given by a function  $x: \mathcal{X}^d \rightarrow \mathcal{X}^d$  where  $x(\theta + k) = x(\theta) + k$ ,  $\forall k \in \mathbb{Z}^d$ , and such that the orbit of  $x(\theta_0)$  is  $x(\theta_0 + \omega t)$ ,  $\forall \theta_0 \in \mathcal{X}^d$  and  $t \in \mathbb{Z}$ . By (3),  $x(\theta)$  must satisfy

$$x(\theta + \omega) - 2x(\theta) + x(\theta - \omega) = \lambda Q\{x(\theta)\} \quad (4)$$

Such sets exist if the parameter  $\lambda$  is large enough as is shown by the following

**Proposition:** There exists an  $L > 0$  such that for  $\lambda > L$  and any incommensurate  $\omega \in \mathcal{X}^d$ , there is an invariant set  $\tilde{M}_\omega$  of the sawtooth map of the form

$$\tilde{M}_\omega = \{(x^\sigma(\theta), x^\sigma(\theta+\omega)) \mid \theta \in \mathbb{R}^d, \sigma \in \{+, -\}^d\} \quad (5)$$

such that  $x^\sigma(\theta+k) = x^\sigma(\theta) + k \forall k \in \mathbb{Z}^d$  and that  $x^\sigma(\theta_0 + \omega t)$  is an orbit for all  $\theta_0 \in \mathbb{R}^d$  and  $\sigma \in \{+, -\}^d$ . The solution is given by

$$x^\sigma(\theta) = \theta - \sum_n B^{-1} b(n) B \{\theta+n\omega\}^\sigma \quad (6)$$

where  $B$  is an orthogonal matrix which diagonalizes  $Q$ , and  $b(n)$ ,  $n \in \mathbb{Z}$ , are diagonal matrices of the form

$$b(n) = \text{diag}(\alpha_i \rho_i^{-|n|}) \quad (7)$$

with  $\alpha_i$  and  $\rho_i$  positive (see (8)).

**Proof:** We first show that a solution to (4) can be found of the form (6) under the assumptions that the solution satisfies  $[x^\sigma(\theta)] = [\theta]$  and that  $x(\theta)$  has no points on the discontinuity set of  $\{ \}$ . Let  $x^\sigma(\theta) = \theta + \psi(\theta)$ ; by assumption  $\{x^\sigma(\theta)\}$  is independent of the choice of fractional part, so choose the  $\sigma$ -continuous one:

$$\begin{aligned} \{x^\sigma(\theta)\} &= x^\sigma(\theta) - [x^\sigma(\theta)]^\sigma \\ &= \theta + \psi(\theta) - [\theta]^\sigma \\ &= \{\theta\}^\sigma + \psi(\theta) \end{aligned}$$

Therefore by (4)  $\psi$  must satisfy

$$\psi(\theta+\omega) - 2\psi(\theta) + \psi(\theta-\omega) = \lambda Q[\{\theta\}^\sigma + \psi(\theta)]$$

A solution to this can be obtained from the ansatz

$$\psi(\theta) = \sum_{n=-\infty}^{\infty} a(n) \{\theta+n\omega\}^\sigma$$

providing

$$\begin{aligned} a(n-1) - 2a(n) + a(n+1) &= \lambda Q a(n) & n \neq 0 \\ a(-1) - 2a(0) + a(1) &= \lambda Q (I + a(0)) \end{aligned}$$

Since  $Q$  is symmetric, it can be diagonalized by an orthogonal coordinate change. Let  $Q = B^{-1}DB$ , where  $B^t = B^{-1}$  and  $D = \text{diag}(q_i)$  is a diagonal matrix; since  $Q$  is



positive definite, the entries  $q_i$  are positive reals. Defining  $b(n) = -Ba(n)B^{-1}$ , then the solution for  $b(n)$  which decays as  $n \rightarrow \pm \infty$  is given by (7) providing

$$\begin{aligned} \rho_i &= 1 + \frac{\lambda q_i}{2} + \left( \lambda q_i + \frac{(\lambda q_i)^2}{4} \right)^{1/2} \\ \alpha_i &= \left( 1 + \frac{4}{\lambda q_i} \right)^{-1/2} \end{aligned} \quad (8)$$

So we obtain  $x(\theta)$  of the form (6) providing the assumptions  $[x^\sigma(\theta)] = [\theta]$ , and  $x^\sigma(\theta)$  has no points on the discontinuity set are satisfied. To verify this we note the following properties of (6):

(a)  $x^\sigma(\theta)$  is  $\sigma$ -continuous. This follows because for  $\varepsilon > 0$ ,  $x^\sigma(\theta + \sigma\varepsilon) \rightarrow x^\sigma(\theta)$  as  $\varepsilon \rightarrow 0$ , since  $\{\theta\}^\sigma$  is  $\sigma$ -continuous.

(b) The derivative of  $x^\sigma(\theta)$  vanishes at all points of continuity. Since the series for  $x(\theta)$  is uniformly convergent, its derivative can be computed term by term. Evaluating the derivative of (6) at a point where  $\{\theta + n\omega\}^\sigma$  is continuous  $\forall n$  gives

$$\frac{\partial x}{\partial \theta} = I - B^{-1} \sum_n b(n)B = 0$$

which follows from the sum

$$\sum_n \rho_i^{-|n|} = \frac{\rho_i + 1}{\rho_i - 1} = \frac{1}{\lambda q_i} \left( \rho_i - \frac{1}{\rho_i} \right) = \frac{1}{\alpha_i} \quad (9)$$

Thus all changes in  $x^\sigma(\theta)$  occur at the discontinuities of  $\{\theta + n\omega\}^\sigma$ , that is the points

$$\theta_j + n\omega_j = m - 1/2$$

for any integer  $m$ . Such points are dense.

(c)  $x^{\sigma_j}(\theta)$  is a monotonic function of  $\theta_j$ . The jump in  $x^\sigma$  across a point  $\theta_j = m - n\omega_j - 1/2$ , holding all the other  $\theta_i$ ,  $i \neq j$ , fixed, is

$$\Delta x(n;j) = B^{-1}b(n)Be_j, \quad \forall \theta_i \neq j \text{ and } \sigma. \quad (10)$$

where  $e_j$  is the unit vector in the  $j$  direction. Thus  $\langle e_j, \Delta x(n;j) \rangle > 0$ , so  $x^\sigma_j$  strictly increases across such a discontinuity. Since the discontinuities are dense,  $x^\sigma_j$  increases between almost every pair of values of  $\theta_j$ .

(d)  $x(\theta)$  can be written as a sum of monotone vectors, each depending on a single  $\theta_j$ :

$$\begin{aligned} x^\sigma(\theta) &= \sum_{j=1}^d x^j(\theta_j) \\ x^j(\theta_j) &= \theta_j - B^{-1} \sum_n b(n) \{\theta_j + n\omega_j\}^{\sigma_j} B e_j \end{aligned} \quad (11)$$

We call a vector function  $x^j(\theta_j)$  monotone if  $\langle e_j, x^j(\theta_j + \delta) - x^j(\theta_j) \rangle \geq 0$ ,  $\forall \delta \geq 0$ . This is true for (11) as follows from (10).

A bound on the norm of  $x^j(\theta_j)$  can be obtained by noting that  $x^j(0) = 0$  and at any other point its value is the sum of the jumps  $\Delta x(n;j)$  for those  $n$  corresponding to discontinuities in  $\{\varphi + n\omega_j\}$  for  $\varphi \in [0, \theta_j]$ . Providing  $|\theta_j| < 1/2$ , the discontinuity corresponding to  $n = 0$  does not contribute. Furthermore  $x^j(\theta_j)$  is odd almost everywhere, indeed  $x^j(\theta_j + \varepsilon) \rightarrow -x^j(-\theta_j - \varepsilon)$  as  $\varepsilon \rightarrow 0^+$ , so only half of the remaining discontinuities contribute, thus

$$|x^j(\theta_j)| \leq \frac{1}{2} \sum_{n \neq 0} |\Delta x(n;j)| \leq \frac{1}{2} \sum_{n \neq 0} \sum_{i=1}^d \alpha_i \rho_i^{-|n|} = \frac{1}{2} \sum_{i=1}^d (1 - \alpha_i)$$

Thus from (11) the norm of  $x(\theta)$  itself is bounded by

$$|x(\theta)| \leq \frac{d}{2} \sum_{i=1}^d (1 - \alpha_i), \quad \text{for } |\theta_j| < 1/2 \quad (12)$$

This implies  $x(\theta)$  is certainly in the fundamental domain  $(-1/2, 1/2)^d$  when  $\theta$  is, providing  $\lambda$  satisfies

$$\lambda > L = \frac{4}{\min(q_i)} \left( \frac{d^4}{2d^2 - 1} - 1 \right) \quad (13)$$

which completes the proof ■

An example for  $d=2$  is shown in Fig. 1. Here we chose  $\lambda q = (0.02, 0.03)$ , and let  $B$  be a rotation by an angle  $0.5$ . The rotation vector is  $\omega = ((3-\sqrt{5})/2, \sqrt{2}-1)$ . Values for  $\theta$  were taken on a rectangular grid.

For  $d = 1$ , (13) implies that the solution (6) is valid for all  $\lambda > 0$ . For  $d > 1$ , this is not necessarily the case, and numerical calculation for  $d=2$  shows that if the  $q_i$  are too small then  $x^\sigma(\theta)$  can cross the discontinuity and therefore will not represent a valid solution. In practice the bound (13) is extremely weak; for  $d=2$  the set appears to remain in the fundamental domain until  $\lambda \min(q_i) \sim O(10^{-3})$ . Even in this case, however, it is possible to modify the fundamental domain  $(-1/2, 1/2)^d$  used in the definition of  $\{x\}$  to obtain a mapping for which (6) is valid for any  $\lambda$ .†

Note that (6) also gives invariant sets with commensurate rotation vector, whose topological form is different; in this paper we concentrate on the incommensurate case.

### §3 Cantori

In this section we consider the topological form of the set  $\tilde{M}_\omega$ . The sawtooth map commutes with the translations  $(x, x') \rightarrow (x+m, x'+m)$ , for  $m \in \mathbb{Z}^d$ , so it can be considered as a map of  $\mathbb{R}^d \times \mathbb{R}^d / \mathbb{Z}^d$  to itself.

**Proposition:** Let  $\omega$  be incommensurate and  $M_\omega = \tilde{M}_\omega / \mathbb{Z}^d$ , with  $\tilde{M}_\omega$  given by (5). Then for  $d \leq 2$ ,  $M_\omega$  is a Cantor set.

**Proof:** A Cantor set is a topological space which is non-empty, compact, totally disconnected and has no isolated points. Since  $\omega$  is incommensurate, it follows from property (a) of §2 that for all  $d$ ,  $M_\omega$  is non-empty, closed and has no isolated points. Furthermore, since the  $b(n)$  are summable,  $M_\omega$  is bounded. So it remains to prove that  $M$  is totally disconnected.

For  $d=1$ , this follows from the fact that the jumps (10) are positive and that  $\omega$  is incommensurate.

For  $d=2$ , consider a point  $\theta_j = n_j \omega_j + 1/2$ ,  $j = 1, 2$ . The vectors  $\Delta x(n_j; j)$  given by (10) span an area of positive orientation. This area is the same as that spanned by  $B \Delta x(n_j; j)$  since  $B \in SO(2)$ . Letting  $\varphi$  be the rotation angle, this area is given by the cross product

$$B \Delta x(n_1; 1) \times B \Delta x(n_2; 2) = \alpha_1 \alpha_2 [ \rho_1^{-|n_1|} \rho_2^{-|n_2|} \cos^2 \varphi + \rho_2^{-|n_1|} \rho_1^{-|n_2|} \sin^2 \varphi ] > 0 \quad (14)$$

---

† This follows from the Proposition of the next section: choose the boundaries of the fundamental domain to run in the gaps between the  $\gamma^1(0, \pm)$  curves and the  $\gamma^2(0, \pm)$  curves.

Define two curves  $\gamma^1(j, \sigma_1)$  by joining the gaps in the points of  $x^\sigma(j\omega_1 + 1/2, \theta_2)$ , for  $\theta_2 \in \mathcal{R}$  and  $\sigma_2 \in \{+, -\}$  with straight lines. By (c) these curves are graphs over the  $x_2$  axis. Similarly define the graphs  $\gamma^2(j, \sigma_2)$  over  $x_1$ . The positivity of (14) for all  $n_2$  implies that  $\gamma^1(n_1, +)$  and  $\gamma^1(n_1, -)$  do not intersect, and similarly for  $\gamma^2$ . Then (c) implies that the four curves  $\gamma^i(n_i, \pm)$ , as shown in Fig. 2, separate the points of  $M_\omega$  with  $\theta_i > n_i\omega_i$  and  $\theta_i < n_i\omega_i$  where  $i = 1, 2$ .

Since  $\omega_1$  and  $\omega_2$  are irrational, this implies that  $M_\omega$  is totally disconnected, thus it is a Cantor set ■

We believe that this result is also true for  $d > 2$ , but did not succeed in proving it.

## §5 Structural Stability

In this section we show that many smooth symplectic maps have invariant sets topologically equivalent to those found for the sawtooth map. Although the sawtooth map is discontinuous, the invariant sets  $M_\omega$  avoid a neighborhood of the discontinuity set so the map can be modified there in any way leaving  $M_\omega$  unchanged. In particular, there are  $C^\infty$  symplectic maps with the same  $M_\omega$ . Since the invariant sets of the sawtooth map are hyperbolic, with hyperbolicity constant  $\min(\rho_i)$ , they are structurally stable, that is  $M_\omega$  persists for all perturbations of the map which are  $C^1$  small enough in a neighborhood of  $M_\omega$ , see e.g. [11]. This allows us to deduce existence of topologically equivalent invariant sets for an open set of smooth symplectic maps of  $\mathcal{R}^d \times \mathcal{R}^d / \mathcal{Z}^d$ . This set includes non-trivial examples, as in the following

**Theorem:** Suppose  $f$  is a symplectic map with generating function  $h(x, x') = \frac{1}{2}|x-x'|^2 + \lambda V(x)$ ,  $x \in \mathcal{R}^d$ , such that  $V(x+k) = V(x) \forall k \in \mathcal{Z}^d$ , and  $V$  has a non-degenerate local minimum with  $D^3V(x) = 0$  at this minimum (e.g. because of symmetry). Then there exist invariant sets topologically equivalent to  $M_\omega$  for all  $\omega$ , provided  $\lambda$  is large enough.

**Proof:** With out loss of generality let  $x=0$  be the minimum of  $V$ ; and write the quadratic part of  $V$  there as  $\frac{1}{2} x^T Q x$ . For  $\theta$  in the fundamental domain, (12) implies that the invariant set  $M_\omega$  of the sawtooth map is contained within in the region  $R: \{|x| <$

$d^2/\lambda \min(q_i) \mid x \in \mathcal{X}^d$ . In this region  $\lambda V(x)$  is  $C^2$  close to its quadratic part, and so  $f$  is  $C^1$  close to the sawtooth mapping. Hence  $M_\omega$  persists for large enough  $\lambda$  ■

The above example can be modified to  $h(x,x') = T(x-x') + \lambda[V(x)+W(x')]$ , where  $T$  represents any positive definite quadratic form, since subtracting  $\lambda[W(x')-W(x)]$  from  $h$  does not change the orbits and  $T$  can be diagonalized as  $1/2|x-x'|^2$ . We believe that the restriction  $D^3V = 0$  is not necessary, since  $f$  would still be  $C^0$  close to the sawtooth map,  $S$ , on  $\mathbb{R}$  and the relative change in the derivative  $(Df DS^{-1} - I)$  would be small.

## §5 Questions

There are several remaining questions:

1) Is  $M_\omega$  a Cantor set when  $d > 2$ ? We suspect that this is the case. In particular, since  $M_\omega$  is hyperbolic and its orbits are semi-conjugate to a rotation, it has zero Hausdorff dimension[12].

2) Can the theorem be generalized to any  $V(x)$  with a non-degenerate minimum for which  $D^3V \neq 0$ ? Can one generalize to examples with a more general dependence of  $h$  on  $(x,x')$ ?

3) Do these orbits globally minimize the action? It is clear that they are local minima; however, we have not shown that variations which move a configuration point across the discontinuity necessarily do not decrease the action. Even if  $M_\omega$  consists of orbits of minimum action for the sawtooth do the perturbed sets for nearby maps consist of minimizing orbits? For  $d=1$ , Mather has given examples where the continuation of a hyperbolic Aubry-Mather set by structural stability no longer consists of minimizing orbits[13].

4) Can these Cantor sets be regarded as the ghosts which remain when invariant tori break up? Does every invariant torus have such a ghost, or something similar? In Figure 3 we display a periodic orbit of the 4 dimensional Froeschlé mapping[14] with primitive period 78635. When the parameter  $\lambda$  is small this orbit appear to nearly uniformly cover the fundamental domain; it is a close approximation to an invariant torus with incommensurate frequency. As  $\lambda$  increases the density of points becomes non-uniform, and low density regions begin to form. It appears that when  $\lambda$  is larger these low density regions will be empty, and the orbit will cover a Cantor set. Unfortunately in this situation the orbit becomes highly unstable, and

impossible to follow numerically.

## Acknowledgements

Support for this research was obtained from the US Department of Energy under contract DE-AC03-76SF00098 and grant DE-FG05-80ET-53088, the UK Science and Engineering Research Council, and a NATO grant, RG 85/0461, for international collaboration.

## References


- [1] Mather J N 1982 Existence of Quasi-Periodic Orbits for Twist Homeomorphisms of the Annulus, *Topology* **21** 457-467
- [2] Aubry S and Le Daeron P Y 1983 The Discrete Frenkel-Kontorova Model and Its Extensions, *Physica D* **8** 381-422
- [3] Bangert V 1988 Minimal Geodesics, Universität Bern *preprint*
- [4] Mather J N 1987 Minimal Measures, Princeton University *preprint*
- [5] Hedlund G A 1932 Geodesics on a Two-Dimensional Riemannian Manifold with Periodic Coefficients, *Ann. Math.* **33** 719-739
- [6] Bernstein D and Katok A 1987 Birkhoff Periodic Orbits for Small Perturbations of Completely Integrable Hamiltonian Systems with Convex Hamiltonians, *Invent. Math.* **88** 225
- [7] Aubry S 1978 On the Dynamics of Structural Phase Transitions, Lattice Locking, and Ergodic Theory, in *Solitons and Condensed Matter* ed A. R. Bishop and T. Schneider (New York:Springer-Verlag) 264-278
- [8] Aubry S 1982 The Devil's Staircase Transformation in Incommensurate Lattices, *Lect. Notes in Math.* **925** 221-245
- [9] Aubry S 1983 Exact Models with a Complete Devil's Staircase, *J. Phys. C* **16** 2497-2508
- [10] Percival I C 1979 Variational Principles for Invariant Tori and Cantori, in *Nonlinear Dynamics and the Beam-Beam Interaction* ed M. Month and J. C. Herrera (New York:Amer. Inst. Phys.) 302-310
- [11] Shub M 1986 *Global Stability of Dynamical Systems* (New York:Springer-Verlag)

- [12] Fathi A 1988 Expansiveness, Hyperbolicity and Hausdorff Dimension, University of Florida *preprint*
- [13] Mather J N 1988 *personal communication*
- [14] Kook H T and Meiss J D 1988 Periodic Orbits for Reversible, Symplectic Mappings, Physica D **in press**

### Figure Captions


- 1) Configuration space projection of a cantorus of the four dimensional sawtooth map.
- 2) Disconnectedness of the set  $M_\omega$ .
- 3,) Configuration space projection of a periodic orbit with rotation vector  $(17556/78635, 51016/78635)$  for the four dimensional Froeschlé map with parameters  $(a,b,c) = (0.5,0.35,0.02)$ , corresponding to  $\lambda q \approx (0.483,0.327)$ .






XBL 895-1733

Fig. 1



XBL 895-1734

Fig. 2



XBL 895-1735

Fig. 3

fig 3 Area study - 1900

LAWRENCE BERKELEY LABORATORY  
TECHNICAL INFORMATION DEPARTMENT  
1 CYCLOTRON ROAD  
BERKELEY, CALIFORNIA 94720