

UC Irvine

UC Irvine Previously Published Works

Title

What's the flux? Unraveling how CO2 fluxes from trees reflect underlying physiological processes

Permalink

<https://escholarship.org/uc/item/9sb6c4bw>

Journal

New Phytologist, 197(2)

ISSN

0028-646X

Authors

Trumbore, Susan E
Angert, Alon
Kunert, Norbert
[et al.](#)

Publication Date

2013

DOI

10.1111/nph.12065

Copyright Information

This work is made available under the terms of a Creative Commons Attribution License, available at <https://creativecommons.org/licenses/by/4.0/>

Peer reviewed

Commentary

What's the flux? Unraveling how CO₂ fluxes from trees reflect underlying physiological processes

Tree stems and branches emit carbon dioxide (CO₂) at rates that per unit area can rival emissions from leaves or the soil surface and summed over a forest stand can comprise 14–30% of the total CO₂ efflux (Chambers *et al.*, 2004; Ryan *et al.*, 2009). Stem CO₂ fluxes have predictable patterns of variation with growth rate, stand age, and elevation (Chambers *et al.*, 2004; Ryan *et al.*, 2009; Robertson *et al.*, 2010). Over the past decade observations of diel covariation of CO₂ efflux with sapflux rates measured in tree stems have led to the conclusion that internal transport of CO₂ within the stem strongly influences the measured CO₂ efflux at the surface (Teskey *et al.*, 2008). In this issue of *New Phytologist*, Bloemen *et al.* (pp. 555–565) report on a tracer experiment that demonstrates not only upward transport of ¹³CO₂ added to the transpiration stream, and emission of this label along the stem, but also fixation of a significant fraction of the added CO₂ in canopy branches, petioles and, to a minor extent, leaves. The study of Bloemen *et al.* adds to the growing literature that demonstrates the utility of isotope labeling studies to understand allocation and carbon (C) cycling in trees (Powers & Marshall, 2011; Epron *et al.*, 2012).

'Dynamic approaches for measuring continuous diurnal CO₂ fluxes and transport in the transpiration stream need to be more widely applied.'

Processes influencing stem CO₂ efflux

A number of factors can influence the efflux of CO₂ measured by a flux chamber covering a segment of tree stem (Fig. 1). The cambium is the site of formation of new tissue, that is, of growth, while maintenance respiration produces CO₂ in all living tissues. The C being respired may derive from recent photosynthetic products transported in the phloem (e.g. Powers & Marshall, 2011) and from storage reserves. The pathways for respiration may vary with time or tree species: recently ¹⁸O/¹⁶O measurements in oxygen (O₂) provided the first evidence for the alternative oxidase pathway contributing to respiration in some tree stems (Angert *et al.*, 2012a). CO₂ may also be locally fixed by photosynthetic tissues found under the bark before it is lost to the atmosphere.

Low rates of diffusion, especially across the cambium, can cause high CO₂ concentrations in stems, and internal O₂ concentrations can drop to very low levels (Spicer & Holbrook, 2005; Teskey *et al.*, 2008). CO₂ is highly soluble, and will dissolve in (or exsolve from) stem water, depending on local saturation conditions, which in turn are controlled by factors such as temperature and pH. Uptake of CO₂ directly from the soil atmosphere, once thought potentially important, has largely been shown to be minor (see summary in Bloemen *et al.*). Hence the source of CO₂ emitted to the atmosphere from the bark surface can reflect a combination of local growth and maintenance respiration, other local processes producing CO₂ (including potentially decomposition in heartwood) or CO₂ from respiration in other tissues (e.g. roots) that has been transported into the volume beneath a chamber in solution. However, there can also be net export in the xylem water stream, as indicated by the fate of the tracer added by Bloemen *et al.* The measured chamber flux at any given time is thus the complex result of transport in, transport out and respiration minus photosynthesis in local tissues. Use of a dark chamber will exclude local photosynthesis.

Observations of a relationship between sapflux and CO₂ efflux provide a clue as to whether CO₂ is net imported or exported from the volume of stem under a chamber attached to the stem surface (see Fig. 1, modified from Teskey *et al.*, 2008). Other evidence for net CO₂ transport away from the region of efflux measurement comes from lower-than-expected efflux rates compared with what is expected given the construction costs of wood (Ryan *et al.*, 2009), and potentially from higher efflux rates in canopy branches (Teskey *et al.*, 2008). Changes in local temperature and/or pH can change respiration rates and also cause changes in CO₂ solubility (Kunert & Mercado Cárdenas, 2012).

Stem anatomy, including bark thickness and tree hydraulics, likely influences the importance of the mechanisms and can help explain observations such as changes in CO₂ efflux with stand age or tree size, or differences between similar trees growing in different environments (Ryan *et al.*, 2009). Bloemen *et al.* report results from labeling *Populus deltoides*, the eastern cottonwood tree, which has very high transpiration rates and generally is found in riparian zones. As noted by Ubierna *et al.* (2009) most studies that have reported relationships between sapflux and CO₂ efflux have been made in tree species with high sapflux rates and small conducting area. By contrast, the large conifer trees investigated by Ubierna *et al.* (2009), with lower overall sapflux, did not demonstrate such relationships, and even crown removal did not change the rates of CO₂ efflux from stems they studied.

What do these results mean for interpretation of other ecosystem CO₂ efflux measurements?

A major conclusion of Bloemen *et al.* is that the transport of the tracer from the tree base to the canopy indicates that root respiration




Fig. 1 Sources, sinks and transport processes in tree stems result in the net CO_2 flux observed when a chamber encloses part of the stem surface (modified after Teskey *et al.*, 2008). The degree to which the stem CO_2 efflux records the underlying process of respiration (minus any fixation) depends on the saturation state and rate of transport of the xylem water. This balance can change from day to night and with altered temperature as sun warms the outer stem. Inset: Relationship between CO_2 efflux (E_{CO_2}) and O_2 influx (I_{O_2}) in the cases of net soluble C transport into or out of the volume covered by a stem respiration chamber. At times of low transport (e.g. at night), the two fluxes should be approximately equal. Note that, if respiration is the main process, a local imbalance in $E_{\text{CO}_2} : I_{\text{O}_2}$ means that there must be a compensating imbalance elsewhere – e.g. if CO_2 is transported to the canopy and emitted without consumption of O_2 .

can be a source of at least some of the CO_2 emitted in the canopy. While the high CO_2 concentrations at the base of trees do argue for a belowground source, Bloemen *et al.* did not successfully introduce enough label via roots to demonstrate definitively the transfer of root CO_2 up the stem. Aubrey & Teskey (2009) have argued that up to 50% of root respired CO_2 may be transported upward and diffuse out higher in the tree stem or in branches. Grossiord *et al.* (2012), using isotopic differences to distinguish plant and decomposition derived soil respiration, detected a daytime reduction in autotrophic respiration from soil, albeit the 'missing' root respiration they infer is transported up the tree stem amounted to only a 17% underestimation of the autotrophic CO_2 efflux on a daily basis.

Tracer studies by Powers & Marshall (2011) as well as Bloemen *et al.* show that ^{13}C -labeled CO_2 added to the xylem stream indeed is transported upward, emitted and a fraction refixed in the canopy. In the Bloemen *et al.* study, an estimated 6–17% of the added tracer was fixed in photosynthetic tissues in branches and petioles. Hence recycling of CO_2 within the plant is potentially quite important – perhaps especially so when CO_2 concentrations in the atmosphere were lower than those of today (Teskey *et al.*, 2008).

Hibberd & Quick (2002) provide an additional mechanism for internal C transport, based on the capture of CO_2 by PEP-carboxylase, after which it can be removed from the site of respiration as malate. Their labeling experiments indicate that malate transported in the xylem enters the bundle-sheath cells, and can be used for photosynthesis, in this 'C₄ like' mechanism. Bloemen *et al.* found that most of their labeled CO_2 was fixed in

branches and in leaf petioles, which agrees well with transported C being fixed in bundle-sheath cells.

How can we derive an estimate of the 'real' stem respiration flux?

The various effects of temperature and transpiration velocity can affect CO_2 efflux rates over a day–night cycle. One way to estimate fluxes might be to choose to sample at night, when transpiration flux is near zero (Teskey & McGuire, 2002). However, this is also the coolest time of day, so this might underestimate daytime respiration in tissues (such as the cambium) that may warm significantly over the daytime period (Kunert & Mercado Cárdenas, 2012). A second method is to measure CO_2 evolution or O_2 uptake (Teskey & McGuire, 2002; Spicer & Holbrook, 2005) on excised wood. Apart from damaging the tree (or the tissues with heat generated on sampling), such methods must be used with care as the degassing of high CO_2 in wood pores can initially yield too-high CO_2 fluxes (Teskey & McGuire, 2002).

Another possibility is to use *in situ* O_2 uptake as a measure of respiration (Angert & Sherer, 2011). Because O_2 is much less soluble in water than CO_2 , the molar flux of O_2 into stems should roughly equal that of CO_2 out if transport is minimal, given the stoichiometry of the respiration substrate in most woody tissues. In cases where CO_2 respired elsewhere is transported and emitted in the stem and canopy, we would expect CO_2 release to exceed O_2 uptake (i.e. $E_{\text{CO}_2}/I_{\text{O}_2} > 1$). In cases where locally derived CO_2 is exported, the ratio of CO_2 release to O_2 uptake will be ≤ 1 (Fig. 1 inset).

In Amazonian tropical forest trees, Angert *et al.* (2012b) found the CO₂ efflux from the stems was on average only 0.66 (± 0.18) of the O₂ influx, in other words about one third of the CO₂ respired in the stem section beneath the chamber is not locally emitted, but transported away. Some of this 'removed CO₂' can be carried by the xylem stream, as shown by Bloemen *et al.* However, the flux of CO₂ that can be removed in this way is constrained by the chemistry of the carbonate system, and is governed by the xylem pH which is seldom > 7 (Teskey *et al.*, 2008). Angert *et al.* (2012b) concluded, based on estimates of stem [CO₂] and xylem pH, that the rate of dissolved inorganic C export might not remove CO₂ at the rate required, and suggest that the C might be exported in organic (e.g. malate), rather than inorganic form.

Progress in understanding the sources and magnitudes of CO₂ fluxes in tree stems is being made rapidly, and linking actively respiring tissues local to, and remote from, the point of measurement. Future studies taking advantage of pulse labeling in trees with different water-use strategies would be useful for resolving conditions where stem xylem water transport in stems significantly impact soil CO₂ efflux. Radiocarbon measurements of stem CO₂ can help resolve questions about whether the C being respired (and potentially translocated) derives from storage reserves vs fresh photosynthetic products. Dynamic approaches for measuring continuous diurnal CO₂ fluxes and transport in the transpiration stream need to be more widely applied. New sensor methods for O₂ measurement can add information that allows separation of transport from local physiological processes. Future studies should also focus on quantitative measurements of the photosynthetic fraction supported by both inorganic, and organic C, transported internally in the xylem.

Meanwhile, we need to be careful about invoking the process 'respiration' when really we are measuring CO₂ flux. Ultimately the CO₂ emitted from a stem is produced by physiological processes, but the challenge remains identifying what portion is produced by local tissues, which will facilitate much-needed mechanistic understanding of factors controlling autotrophic respiration.

Susan E. Trumbore^{1*}, Alon Angert², Norbert Kunert¹,
Jan Muhr¹ and Jeffrey Q. Chambers³

¹Max-Planck Institute for Biogeochemistry, Jena, 07745, Germany;

²The Institute of Earth Sciences, The Hebrew University of Jerusalem, Jerusalem, Israel;

³Climate Sciences Department, Lawrence Berkeley National Laboratory, Berkeley, CA, 94720, USA

(*Author for correspondence: tel +49 3641 576164; email trumbore@bgc-jena.mpg.de)

References

- Angert A, Muhr J, Negron Juarez R, Alegria Munoz W, Kraemer G, Ramirez Santillan J, Chambers JQ, Trumbore SE. 2012a. The contribution of respiration in tree stems to the Dole Effect. *Biogeosciences* 9: 4037–4044.
- Angert A, Muhr J, Negron Juarez R, Alegria Munoz W, Kraemer G, Ramirez Santillan J, Chambers JQ, Trumbore SE. 2012b. Internal respiration of Amazon tree stems greatly exceeds external CO₂ efflux. *Biogeosciences Discussions* 9: 4979–4991.
- Angert A, Sherer Y. 2011. Determining the relationship between tree-stem respiration and CO₂ efflux by dO₂/Ar measurements. *Rapid Communications in Mass Spectrometry* 25: 1752–1756.
- Aubrey DP, Teskey RO. 2009. Root-derived CO₂ efflux via xylem stream rivals soil CO₂ efflux. *New Phytologist* 184: 35–40.
- Bloemen J, McGuire MA, Aubrey DP, Teskey RO, Steppe K. 2012. Transport of root-respired CO₂ via the transpiration stream affects above-ground C assimilation and CO₂ efflux in trees. *New Phytologist* 197: 555–565.
- Chambers JQ, Tribuzy ES, Toledo LC, Crispim BF, Higuchi N, dos Santos J, Araujo AC, Kruijt B, Nobre AD, Trumbore SE. 2004. Respiration from a tropical forest ecosystem: partitioning of sources and low carbon use efficiency. *Ecological Applications* 14(4 Suppl): S72–S88.
- Epron D, Bahn M, Derrien D, Lattanzi FA, Pompanen J, Gessler A, Höglberg P, Maillard P, Dannoura M, Gerant D *et al.* 2012. Pulse-labelling trees to study carbon allocation dynamics: a review of methods, current knowledge and future prospects. *Tree Physiology* 32: 776–798.
- Grossiord C, Mareschal L, Epron D. 2012. Transpiration alters the contribution of autotrophic and heterotrophic components of soil CO₂ efflux. *New Phytologist* 194: 647–653.
- Hibberd JM, Quick WP. 2002. Characteristics of C₄ photosynthesis in stems and petioles of C₃ flowering plants. *Nature* 415: 451–454.
- Kunert N, Mercado Cárdenas A. 2012. Effects of xylem water transport on CO₂ efflux of woody tissue in a tropical tree, Amazonas State, Brazil. *Hoehnea* 39: 139–144.
- Powers EM, Marshall JD. 2011. Pulse labeling of dissolved ¹³C-carbonate into tree xylem: developing a new method to determine the fate of recently fixed photosynthate. *Rapid Communications in Mass Spectrometry* 25: 33–40.
- Robertson AL, Malhi Y, Farfan F, Aragão LEOC, Espejo JES, Robertson MA. 2010. Stem respiration in tropical forests along an elevation gradient in the Andes. *Global Change Biology* 16: 3193–3204.
- Ryan MG, Cavaleri MA, Almeida AC, Penchel R, Senock RS, Stape JL. 2009. Wood CO₂ efflux and foliar respiration for Eucalyptus in Hawaii and Brazil. *Tree Physiology* 29: 1213–1222.
- Spicer R, Holbrook NM. 2005. Within-stem oxygen concentration and sap flow in four temperate tree species: does long-lived xylem parenchyma experience hypoxia? *Plant, Cell & Environment* 28: 192–201.
- Teskey RO, McGuire MA. 2002. Carbon dioxide transport in xylem causes errors in estimation of rates of respiration in stems and branches of trees. *Plant, Cell & Environment* 25: 1571–1577.
- Teskey RO, Saveyn A, Steppe K, McGuire MA. 2008. Origin, fate and significance of CO₂ in tree stems. *New Phytologist* 177: 17–32.
- Ubierna N, Kumar AS, Cernusak LA, Pangle RE, Gag PJ, Marshall JD. 2009. Storage and transpiration have negligible effects on d¹³C of stem CO₂ efflux in large conifer trees. *Tree Physiology* 29: 1563–1574.

Key words: isotope labeling, respiration, sapflux, soil respiration, stem CO₂ efflux.