

## **UC Irvine**

### **Western Journal of Emergency Medicine: Integrating Emergency Care with Population Health**

#### **Title**

Risk Perception of US-Mexico Border Crossers

#### **Permalink**

<https://escholarship.org/uc/item/9vj6927k>

#### **Journal**

Western Journal of Emergency Medicine: Integrating Emergency Care with Population Health, 9(1)

#### **ISSN**

1936-900X

#### **Authors**

DeLuca, Lawrence  
Bitar, Jamil  
Leeson, Kimberly  
et al.

#### **Publication Date**

2008

#### **Copyright Information**

Copyright 2008 by the author(s). All rights reserved unless otherwise indicated. Contact the author(s) for any necessary permissions. Learn more at <https://escholarship.org/terms>

Peer reviewed

**Background:** Violent behavior by patients is one of many occupational hazards faced by health care workers. Emergency department (ED) personnel are at high risk for patients carrying weapons, or exhibiting disruptive behavior or psychotic disorders. When systematic approaches to violent persons do not work, public safety officers (PSO) require additional means of elevated force to control dangerous behavior. The use of the electrical stun gun (TASER) offers an option that is more effective than baton but less lethal than a firearm. Its use has recently been criticized because of the association with deaths in custody.

**Methods:** We describe an approach to control workplace violence in a health care environment that includes staff education for early identification of potentially violent persons and initial approaches but allows for the use of TASER in select situations. We report the incidents of use of force in a Level 1 trauma center university hospital with 40,000 ED census.

**Results:** There were 107 PRE (12 month) and 149 POST (24 month) uses of force. During the POST, 92% were in clinical, 5 % in general public and 3% in exterior areas. Most involved patients (93%). In clinical areas, 56% were in the ED, 25 % inpatient and 11% outpatient areas. There were 30 displays and seven additional uses of the TASER, including two touches and five firings of probes, 77% for male subjects and 70% for psychiatric or ED patients. All displays or uses were reviewed in detail by multidisciplinary group and determined to be appropriate. There were no serious injuries in either safety personnel or patients that resulted from the use of the TASER. PSOs determined that the display of the TASER was able to de-escalate violent situations without the use of more elevated force.

**Conclusions:** A comprehensive approach to workplace violence that allows for the selected use of the TASER and requires mandatory reviews of all uses can be effectively implemented to help to control dangerous situations in health care environments.

---

## 18 Risk Perception of US-Mexico Border Crossers

Lawrence DeLuca, MD, EdD; Jamil Bitar, MD;  
Kimberly Leeson, MD; Samuel M. Keim, MD.  
*The University of Arizona*

---

**Background:** This study focused on risk perception of US-Mexico border crossers and builds on current research programs at The University of Arizona. No published studies have addressed specific risk processes (defined as perceived risk, intra-border crosser risk communications, Mexican government originated risk communications, and risk control actions) in US-Mexico border crossers.

**Objectives:** This project seeks to describe, analyze, and interpret border-crosser risk processes; and develop a

multidimensional model to describe border-crosser perceived risk and risk communications. Additionally, the main motivation for crossing will be investigated.

**Methods:** The project used rigorously coded qualitative but anonymous interview data obtained from up to 10 recent border-crossers to elicit information about domains of perceived risk and risk communications that can be incorporated into a proposed model and used for future research and refinement of border-crosser behavior models. Because of the qualitative design, thematic saturation occurred before 10 subjects were entered. Interview data were translated from Spanish to English and data extracted in an attempt to reach thematic saturation.

**Results:** A model of risk processes was created and suggestions for future behavioral interventions to reduce border crosser heat and injury related morbidity and mortality are presented.

**Conclusions:** Risk perception of US-Mexico border-crossers can be modeled using a qualitative methodology. Themes derived that were most important included desires of border-crossers to be re-united with family members living in the US regardless of risk and the state of limbo of recently deported border crossers.

---

## 19 Behaviors that Influence Crash Injury Risk in Latino Adolescent Males: Analysis of the 2005 National Youth Risk Behavior Survey (YRBS)

Federico Vaca, MD, MPH; Craig Anderson, PhD.  
*University of California, Irvine*

---

**Objective:** Motor vehicle crashes remain the leading cause of death for teens. Risk-taking behavior is known to contribute to fatal crashes in young drivers and occupants. The objective of this study was to analyze behaviors that influence the risk of crash injury in Latino adolescent males.

**Method:** The Youth Risk Behavior Survey (YRBS) is a multistage cluster sample of students in U.S. public and private high schools, with oversampling of Hispanics. Among other risk behavior topics, three questions are directly related to motor vehicle occupant crash injuries: use of seat belts, riding with a driver who had been drinking, and driving when drinking. Analysis was restricted to Hispanic and non-Hispanic Whites age  $\geq 15$  ( $n=8,520$ ). Data were analyzed using Stata survey procedures that account for survey weights and clustering. Differences between groups were tested using linear regression, controlling for age, with post-estimation tests to compare Hispanic males to Hispanic females and to non-Hispanic White males.

**Results:** Thirteen percent of male Hispanics in this age group reported that they rarely or never wore a seat belt. The percentage of those who rarely or never wore a seat belt was 4% higher for male Hispanics than for female Hispanics.