

# Lawrence Berkeley National Laboratory

## Recent Work

### Title

Transcriptomic Studies of the Response to Exogenous Exposure and Endogenous Production of Biofuel Candidates in E. Coli

### Permalink

<https://escholarship.org/uc/item/9x50c30r>

### Authors

Dahl, R.  
Rutherford, Becky J.  
Mukhopadhyay, Aindrila  
et al.

### Publication Date

2009-12-18

## Transcriptomic Studies of the Response to Exogenous Exposure and Endogenous Production of Biofuel Candidates in E. Coli

R. Dahl<sup>1,3</sup>, B. Rutherford<sup>1,3</sup>, A. Mukhopadhyay<sup>2,3</sup>, J. D. Keasling<sup>1,2,3,4\*</sup> Department of Chemical Engineering, University of California, Berkeley, CA 94720<sup>2</sup> Physical Biosciences Division, Lawrence Berkeley National Laboratory, Berkeley, CA 94720<sup>3</sup> Joint BioEnergy Institute, Emeryville, CA 95608 (www.jbei.org)<sup>4</sup> Department of Bioengineering, University of California, Berkeley, CA 94720 \* robdahl@berkeley.edu

The production of biofuels in microbial systems presents a unique challenge to the host cell. Not only is the cell exposed to the solventogenic fuel molecule itself, the coordinated overexpression of an exogenous pathway presents a large burden on the cell's physiology, both in depleting nutrients and introducing foreign intermediates which could have toxic side-effects. At JBEI, we have focused on the impacts of short-chain alcohols (e.g. butanol) and isoprenoid-based fuels (e.g. isopentenol). Microarrays (in complementary studies with other system-wide "Omics" studies) were used to characterize E. coli's response to these challenges. The general response has been one of a combination of oxidative, hyperosmotic (chaotrophic/ desiccation) and heat shock. In this poster, we present some of these results as well as the unique challenge to engineer the cell to better cope with these stresses.