

UC Davis

UC Davis Previously Published Works

Title

National guideline for the field triage of injured patients: Recommendations of the National Expert Panel on Field Triage, 2021.

Permalink

<https://escholarship.org/uc/item/9z36j7g3>

Journal

The Journal of Trauma and Acute Care Surgery, 93(2)

Authors

Newgard, Craig

Fischer, Peter

Gestring, Mark

et al.

Publication Date

2022-08-01

DOI

10.1097/TA.00000000000003627

Copyright Information

This work is made available under the terms of a Creative Commons Attribution-NonCommercial-NoDerivatives License, available at

<https://creativecommons.org/licenses/by-nc-nd/4.0/>

Peer reviewed

OPEN

National guideline for the field triage of injured patients: Recommendations of the National Expert Panel on Field Triage, 2021

Craig D. Newgard, MD, MPH, FACEP, Peter E. Fischer, MD, Mark Gestring, MD, Holly N. Michaels, MPH, Gregory J. Jurkovich, MD, FACS, E. Brooke Lerner, PhD, FAEMS, Mary E. Fallat, MD, Theodore R. Delbridge, MD, MPH, Joshua B. Brown, MD, MSc, FACS, Eileen M. Bulger, MD, and the Writing Group for the 2021 National Expert Panel on Field Triage, Portland, Oregon

In the United States, unintentional injury remains the leading cause of death and years of potential life lost among children and young adults, and the third most common cause of death overall.^{1,2} Injury is the most common reason for use of 9-1-1 emergency medical services (EMS) in the United States,³ with EMS playing a critical role in the early evaluation and care of injured patients.⁴ An important aspect of EMS care is field triage—the process of identifying seriously injured patients in need of care in specialized trauma centers from among the larger number of patients with minor to moderate injuries who can be cared for in nontrauma hospitals. To accomplish this task quickly and efficiently, EMS clinicians use specific prehospital criteria known as the field triage guideline. The triage guideline was originally developed in 1976, with periodic revisions every 5 to 10 years.⁵ The most recent evidence-based revisions to the field triage guideline were completed in 2011.⁶

Concentrating the most seriously injured patients in trauma centers through field triage is predicated on the principle that patients have better outcomes in these hospitals. A landmark study demonstrated 20% lower in-hospital mortality and 25% lower 1-year mortality among seriously injured adults treated in Level I trauma centers compared with nontrauma hospitals.⁷ Other studies have shown that regionalized trauma systems are associated with reductions in mortality,^{8–11} with the benefit driven primarily by Level I trauma centers.^{8,9} The benefits are similar for children, particularly when treated in pediatric trauma centers^{12–14} and in trauma centers with high emergency department (ED) pediatric readiness.^{15,16} For older adults, the benefit

of tertiary trauma centers is less clear, with some studies showing reduced mortality^{17,18} and others no effect.^{7,19} Until the evidence becomes clearer, the prevailing view is that seriously injured older adults should be managed in trauma centers.

The effectiveness of field triage is measured at the system level using metrics termed *undertriage* and *overtriage*. Undertriage is the percentage of seriously injured patients missed by field triage processes and transported to nontrauma hospitals, which is associated with increased mortality in adults and children.^{20–23} Overtriage is the percentage of patients with minor to moderate injuries identified by field triage criteria as having serious injuries and transported to trauma centers unnecessarily, representing overuse of limited resources and inefficiency in the system. Undertriage and overtriage are inversely related.²⁴ Trauma systems have prioritized the goal of minimizing undertriage and accepting a higher level of overtriage to avoid increased mortality, with targets set at ≤5% and ≤35%, respectively.⁴ A systematic review of field triage performance across all ages showed 14% to 34% undertriage and 12% to 31% overtriage.²⁵ Undertriage of children is up to 51%²⁶ and has increased with recent triage guidelines.²⁷ Undertriage is highest among older adults,^{28–30} with half of seriously injured older adults treated in nontrauma centers nationally.³¹ Reducing undertriage was an important goal of the panel for the current guideline revision.

The purpose of this report is to present the final 2021 field triage guideline and to describe the process of guideline development and the supporting evidence. The guideline is intended for use in civilian 9-1-1 EMS systems and is not intended to guide

Submitted: February 25, 2022, Revised: March 9, 2022, Accepted: March 15, 2022, Published online: April 27, 2022.

From the Department of Emergency Medicine (C.D.N., J.R.L.), Center for Policy and Research in Emergency Medicine, Oregon Health and Science University, Portland, Oregon; Department of Surgery (P.E.F.), University of Tennessee Health Science Center, Memphis, Tennessee; Department of Surgery (M.G.), University of Rochester, Rochester, New York; Committee on Trauma (H.N.M., M.N., M.D., J.D.), American College of Surgeons, Chicago, Illinois; Department of Surgery (G.J.J.), UC Davis Health, Sacramento, California; Department of Emergency Medicine (E.B.L.), University at Buffalo, Buffalo, New York; Department of Surgery (M.E.F.), University of Louisville School of Medicine, Norton Children's Hospital, Louisville, Kentucky; Maryland Institute for Emergency Medical Services Systems (T.R.D.), Baltimore, Maryland; Division of Trauma and General Surgery, Department of Surgery (J.B.B.), University of Pittsburgh Medical Center, Pittsburgh, Pennsylvania; Department of Surgery (E.M.B.), University of Washington, Seattle, Washington.

Supplemental digital content is available for this article. Direct URL citations appear in the printed text, and links to the digital files are provided in the HTML text of this article on the journal's Web site (www.jtrauma.com).

Address for reprints: Craig D. Newgard, MD, MPH, Department of Emergency Medicine, Center for Policy and Research in Emergency Medicine, Oregon Health and Science University, 3181 SW Sam Jackson Park Rd, mail code CR-114, Portland, OR 97239-3098; email: newgardc@ohsu.edu.

Copyright © 2022 The Author(s). Published by Wolters Kluwer Health, Inc. on behalf of the American Association for the Surgery of Trauma. This is an open-access article distributed under the terms of the Creative Commons Attribution-Non Commercial-No Derivatives License 4.0 (CCBY-NC-ND), where it is permissible to download and share the work provided it is properly cited. The work cannot be changed in any way or used commercially without permission from the journal.

DOI: 10.1097/TA.0000000000003627

mass casualty events or in-hospital trauma team responses. The evidence to support the current guideline is based on civilian trauma systems. The guideline is intended for patients in whom maximal resuscitative care is appropriate and does not apply to patients with limited goals of care.

METHODS

Study Design

We conducted an evidence-based revision of the field triage guideline using an interdisciplinary national expert panel and systematic reviews of the field triage literature. We used the Reporting Tool for Practice Guidelines in Health Care³² to report the 2021 revision to the field triage guideline. A complete Reporting Tool for Practice Guidelines in Health Care checklist is included as supplemental online content (Supplemental Digital Content, Supplementary Data 1, <http://links.lww.com/TA/C515>).

Panel Participant Recruitment

The expert panel included EMS clinicians, EMS physicians, emergency physicians, trauma surgeons, pediatric surgeons, nurses, EMS medical directors, experts in EMS training and education, EMS and trauma system administrators, researchers, and representatives from stakeholder organizations. The function of the panel was to review the evidence base, provide stakeholder feedback, assess usability and feasibility, and make informed decisions about revisions to the triage guideline.

Systematic Reviews and Evidence Base

We organized multiple systematic reviews in advance of the guideline revision. The reviews were targeted to controversial aspects of the guideline, opportunities for new or modified criteria, and to identify relevant literature published since the 2011 guideline, including assessment of the quality of evidence and risk of bias. The systematic reviews included the predictive utility of out-of-hospital motor Glasgow Coma Scale (GCS) score versus total GCS,³³ circulatory measures,³⁴ respiratory measures,³⁵ mechanism of injury and special considerations criteria,³⁶ and the overall performance of the triage guideline.²⁵

An inherent challenge in field triage is defining a “seriously injured” patient, which has varied widely across studies. Most triage research has used one of the following categories to define “serious injury”: (1) anatomic injury severity (e.g., Injury Severity Score ≥ 16), (2) critical resource use (e.g., blood transfusion requirements, certain operative interventions, and specific “life-saving” procedures), (3) in-hospital mortality, or (4) a combination of categories.^{25,33–36} We considered any of these definitions to represent “serious injury.”

Criteria for addition and removal of triage criteria

For the 2011 guideline, the threshold to add new triage criteria was a positive predictive value of 20% or greater, with removal of criteria when predictive evidence was lacking.⁶ Because the positive predictive value is dependent on the prevalence of disease (e.g., serious injury) and therefore not readily comparable across studies, we worked with experts in predictive analytics to identify rigorous statistical criteria to guide the addition and removal of triage criteria. Ultimately, we opted to use positive likelihood ratios (+LRs) and area under the receiver operating characteristic curves (AUROCs) because they combine

sensitivity and specificity, are not influenced by disease prevalence, and provide more balanced metrics (Table 1). We used +LRs because individual triage criteria generally favor specificity over sensitivity. When all triage criteria are combined, the collective sensitivity of the guideline is raised. We also considered ease of use in the field and quality of the evidence.

Process for Generating the Updated Guidelines

We assembled a steering committee years in advance to develop key questions for the systematic reviews, organize, plan, and orchestrate the revision process. The expert panel met for 2 days in April 2021 to review the evidence base and discuss potential revisions to the guideline. Following the meeting, the steering committee drafted proposed revisions to the guideline and compiled additional data to address questions from the panel. A second meeting with the panel was held 2 months later to discuss the draft revisions, present additional data, and reach consensus on recommendations for the new guideline. Following the second meeting, the steering committee integrated the additional revisions and sent the draft guideline to stakeholder organizations for feedback. The steering committee integrated feedback from each of these organizations and again returned the updated guideline to the expert panel for review. This process was repeated until all comments, suggestions, and feedback had been addressed.

In parallel with preparations for revisions to the guideline, the EMS Subcommittee of the American College of Surgeons Committee on Trauma developed and piloted a 40-question electronic end-user feedback tool in the fall of 2020. The tool was distributed to 29 national organizations to gather information about use of the field triage guideline directly from EMS clinicians. Responses from 3,958 EMS clinicians³⁷ were shared with the expert panel and integrated into the guideline revision process.

RESULTS

Overview

The 2021 field triage guideline includes important clarifications regarding nomenclature and terminology. The name has been revised to “National Guideline for the Field Triage of Injured Patients,” reflecting the goal and intended function of the document. The name can be shortened to “Field Triage Guideline,” as needed.

Format and Structure

There are substantive changes to the format and structure of the guideline. Because stakeholder feedback and research indicated that the step-wise algorithmic format of prior versions was

TABLE 1. Statistical Criteria Used to Add and Remove Individual Triage Criteria

- To *add* a new field triage criterion: $+LR \geq 2$ or $AUROC \geq 0.60$
Magnitude of predictive utility:
 - Large effect: $+LR \geq 10$, $AUROC \geq 0.80$
 - Moderate effect: $+LR 5-9$, $AUROC 0.7-0.79$
 - Small effect: $+LR 2-4$, $AUROC 0.6-0.69$
- To *remove* a field triage criterion: *no evidence* or $+LR 1.0-1.5$ or $AUROC 0.50-0.55$ across multiple studies (triage criteria were not removed based on a single study)

overly complex for field use, the expert panel modified the structure to align with the flow of information to EMS and actual use of the guideline.^{37–39} The redesigned structure consolidates triage criteria into two main categories based on risk of serious injury: (1) high-risk criteria (red box), including Injury Patterns (previously “Anatomic Criteria”) and Mental Status and Vital Signs (previously “Physiologic Criteria”), and (2) moderate-risk criteria (yellow box), including Mechanism of Injury and EMS Judgment (previously “Special Considerations”). Each risk category is aligned with recommendations for a destination hospital. The guideline is intended to be read from top-to-bottom (risk) and left-to-right (flow of information to EMS).

Specific Field Triage Guideline Recommendations

The 2021 guideline is detailed in Figure 1. Changes from the 2011 field triage guideline are summarized in Table 2. The 2011 guideline is included in the online supplement for reference (Supplemental Digital Content, Supplementary Data 2, <http://links.lww.com/TA/C516>).

Injury Patterns (Previously Step 2 Anatomic Criteria)

Injury patterns are highly specific, yet insensitive for identifying seriously injured patients.⁴⁰ We added one new criterion and revised six criteria for clarity. Two criteria remain unchanged. We also revised the order of criteria to align with a head-to-toe field-based rapid physical assessment.

New and Modified Criteria

- New criterion: Active bleeding requiring a tourniquet or wound packing with continuous pressure
Rationale: Research in military settings has shown that early field application of tourniquets is associated with improved survival and few complications.^{41–43} Tourniquet use was not included in the 2011 guideline because of insufficient evidence in the civilian setting. Multiple civilian studies have since been published on the appropriate application of field tourniquets, safety, effectiveness, and specificity for serious injury. Among 306 civilian trauma patients with tourniquet application, 92% required surgical intervention within 8 hours and field application was associated with lower transfusion requirements and higher survival.⁴⁴ Additional studies showed similar results.^{45–47} The panel added “wound packing with continuous pressure” to capture external bleeding requiring operative intervention in anatomic locations not amenable to tourniquet placement.

- Criterion clarified: Penetrating injuries to the head, neck, torso, and proximal extremities
Rationale: This criterion was revised from “proximal to elbow or knee”⁶ to “proximal extremities” to simplify the criterion based on EMS feedback. This criterion includes impalement.

- Criterion clarified: Skull deformity, suspected skull fracture
Rationale: This criterion was revised from “Open or depressed skull fracture” based on EMS feedback.

- Criterion clarified: Suspected spinal injury with new motor or sensory loss
Rationale: This criterion was revised from “Paralysis” based on EMS feedback.

- Criterion clarified: Chest wall instability, deformity, or suspected flail chest

Rationale: This criterion was revised from “Chest wall instability or deformity (e.g., flail chest)” based on EMS feedback.

- Criterion clarified: Suspected pelvic fracture
Rationale: This criterion was revised from “Pelvic fractures” based on EMS feedback. While field use of this criterion has shown lower predictive utility than other anatomic criteria, having a pelvic fracture by *International Classification of Diseases, Ninth Revision*, diagnosis codes increased the +LR to 6.2.⁴⁰ The panel felt that this criterion should be retained, noting an opportunity for EMS training.

- Criterion clarified: Suspected fracture of two or more proximal long bones

Rationale: This criterion was revised from “Two or more proximal long-bone fractures” based on EMS feedback.

Retained Criteria (No Changes)

- Retained criterion: Crushed, degloved, mangled, or pulseless extremity

- Retained criterion: Amputation proximal to wrist or ankle
Rationale: While the panel debated changes to the level of amputation (e.g., hand, digit) based on the limited availability of hand surgeons in many regions, they ultimately decided to retain the criterion without changes.

Mental Status and Vital Signs (Previously Step 1 Physiologic Criteria)

These criteria are highly specific, but insensitive for identifying seriously injured patients. The panel focused on expanding this category based on new evidence, with attention to feasibility of use in the field. There are five new criteria, three retained criteria, and three criteria removed.

New and Modified Criteria

- New criterion: “Unable to follow commands (motor GCS <6)” replaces total “GCS ≤13.”

Rationale: Twelve head-to-head studies compared total GCS to motor GCS, with high AUROCs for both measures (0.8–0.9).³³ The AUROC difference in predictive performance between the two measures was small and unlikely to have clinical impact,³³ particularly considering ease of use and the dichotomized cut point used for field triage. Feedback from EMS and the expert panel indicated strong preference for simplifying the criterion for feasibility and EMS training. This measure also applies to young children,^{48,49} as lacking spontaneous or purposeful movements. Patients with language barriers who are unable to understand commands is a potential limitation of this criterion.

- New criterion: heart rate (HR) > systolic blood pressure (SBP) (adults and older adults)

Rationale: The systematic review of circulatory predictors identified 29 studies evaluating shock index (HR/SBP), most of which used a value of 1.0.³⁴ Among out-of-hospital studies, pooled estimates showed a sensitivity of 37%, a specificity of

National Guideline for the Field Triage of Injured Patients

RED CRITERIA

High Risk for Serious Injury

| Injury Patterns | Mental Status & Vital Signs |
|--|--|
| <ul style="list-style-type: none"> Penetrating injuries to head, neck, torso, and proximal extremities Skull deformity, suspected skull fracture Suspected spinal injury with new motor or sensory loss Chest wall instability, deformity, or suspected flail chest Suspected pelvic fracture Suspected fracture of two or more proximal long bones Crushed, degloved, mangled, or pulseless extremity Amputation proximal to wrist or ankle Active bleeding requiring a tourniquet or wound packing with continuous pressure | <p>All Patients</p> <ul style="list-style-type: none"> Unable to follow commands (motor GCS < 6) RR < 10 or > 29 breaths/min Respiratory distress or need for respiratory support Room-air pulse oximetry < 90% <p>Age 0–9 years</p> <ul style="list-style-type: none"> SBP < 70mm Hg + (2 x age in years) <p>Age 10–64 years</p> <ul style="list-style-type: none"> SBP < 90 mmHg or HR > SBP <p>Age ≥ 65 years</p> <ul style="list-style-type: none"> SBP < 110 mmHg or HR > SBP |

Patients meeting any one of the above RED criteria should be transported to the highest-level trauma center available within the geographic constraints of the regional trauma system

YELLOW CRITERIA

Moderate Risk for Serious Injury

| Mechanism of Injury | EMS Judgment |
|---|--|
| <ul style="list-style-type: none"> High-Risk Auto Crash <ul style="list-style-type: none"> Partial or complete ejection Significant intrusion (including roof) <ul style="list-style-type: none"> >12 inches occupant site OR >18 inches any site OR Need for extrication for entrapped patient Death in passenger compartment Child (age 0–9 years) unrestrained or in unsecured child safety seat Vehicle telemetry data consistent with severe injury Rider separated from transport vehicle with significant impact (eg, motorcycle, ATV, horse, etc.) Pedestrian/bicycle rider thrown, run over, or with significant impact Fall from height > 10 feet (all ages) | <p>Consider risk factors, including:</p> <ul style="list-style-type: none"> Low-level falls in young children (age ≤ 5 years) or older adults (age ≥ 65 years) with significant head impact Anticoagulant use Suspicion of child abuse Special, high-resource healthcare needs Pregnancy > 20 weeks Burns in conjunction with trauma Children should be triaged preferentially to pediatric capable centers <p>If concerned, take to a trauma center</p> |

Patients meeting any one of the YELLOW CRITERIA WHO DO NOT MEET RED CRITERIA should be preferentially transported to a trauma center, as available within the geographic constraints of the regional trauma system (need not be the highest-level trauma center)

Figure 1. 2021 National Guideline for the Field Triage of Injured Patients. *For the red criteria transport recommendations, patients in extremis (e.g., unstable airway, severe shock, or traumatic arrest) may require transport to the closest hospital for initial stabilization, before transport to a Level I or II trauma center for definitive care. Pediatric patients meeting the red criteria should be preferentially triaged to pediatric-capable trauma centers. The EMS Judgment criteria should be considered in the context of resources available in the regional trauma system, including consideration of online medical control for further direction. Examples of patients with special, high-resource health care needs include tracheostomy with ventilator dependence and cardiac assist devices, among others. Patients with combined burns and trauma should be preferentially transported to a trauma center with burn care capability. If not available, then a trauma center takes precedence over a burn center. Specific age used to define “children” is based on local system resources and practice patterns.

85%, and an AUROC of 0.72 for identifying seriously injured patients.³⁴ Among five head-to-head studies comparing shock index to SBP, all favored shock index, although the quality of evidence was low.³⁴ Assessing if HR is greater than SBP

achieves the goal of identifying patients with a shock index of >1.0 and facilitates EMS training. Pediatric studies have used an age-adjusted shock index to predict serious injury,^{50,51} but the panel felt that calculating an age-adjusted shock index

TABLE 2. Summary of Changes to the Field Triage Guidelines

| Type of Change | Changes in 2021 Field Triage Guidelines | Age Range | 2011 Field Triage Guidelines |
|---------------------------|---|-------------------------------|---|
| Format and structure | Two categories of triage criteria, based on risk of serious injury — high risk versus moderate risk (from top to bottom organization) Within each risk category, the groups of criteria are listed from left to right to follow the flow of information to EMS Injury patterns criteria are organized from head-to-toe to align with rapid field assessment | All ages | 4 Categories of triage criteria, classified as “steps” No alignment with flow of information to EMS No specific order |
| | <i>Injury Patterns</i> | All ages | Anatomic criteria (step 2) |
| New criterion | Active bleeding requiring a tourniquet or wound packing with continuous pressure | All ages | None |
| Clarified criteria | Skull deformity, suspected skull fracture | All ages | Open or depressed skull fracture |
| | Suspected spinal injury with new motor or sensory loss | All ages | Paralysis |
| | Chest wall instability, deformity or suspected flail chest | All ages | Chest wall instability or deformity (e.g., flail chest) |
| | Suspected pelvic fracture | All ages | Pelvic fractures |
| | Suspected fracture of two or more proximal long bones | All ages | Two or more proximal long-bone fractures |
| | <i>Mental Status and Vital Signs</i> | All ages | Physiologic criteria (step 1) |
| New criteria | Motor GCS <6 (unable to follow commands) | All ages | GCS ≤13 |
| | Heart rate >SBP | ≥10 y | None |
| | SBP <70 mm Hg + (2 × age in years) | 0–9 y | None |
| | Respiratory distress or need for respiratory support | All ages | Respiratory rate <20 in infant aged <1 y; ventilatory support |
| Relocated criteria | Room air pulse oximetry <90% | All ages | None |
| | SBP <110 mm Hg for older adults | ≥65 y | SBP <110 might represent shock after age 65 y (Special Considerations section) |
| | <i>Mechanism of Injury Criteria</i> | All ages | Mechanism criteria (step 3) |
| New criterion | Child (age 0–9 y) unrestrained or in unsecured child safety seat | 0–9 y | None |
| Modified criteria | Rider separated from transport vehicle with significant impact (e.g., motorcycle, ATV, horse, etc.) | All ages | Motor cycle crash >20 mph |
| | Fall from height >10 ft (all ages) | All ages | Adults: >20 ft (one story is equal to 10 ft) Children: >10 ft or two to three times the height of the child |
| Modified criterion | Pedestrian/bicycle rider thrown, run over, or with significant impact | | Auto vs. pedestrian/bicyclist thrown, run over, or with significant (>20 mph) impact |
| | <i>EMS Judgment</i> | All ages | Special considerations criteria (step 4) |
| New criteria | Low level falls in young children (≤ 5 y) or older adults (≥ 65 y) with significant head impact | 0–5 y, ≥65 y | Older adults — low impact mechanisms (e.g., ground level falls) might result in severe injury |
| | Suspicion of child abuse | Any child, with focus on ≤5 y | None |
| Modified criteria | Special, high resource health care needs | All ages | None |
| | Anticoagulation use | All ages | Anticoagulants and bleeding disorders — patients with head injury are at high risk for rapid deterioration |
| Transport recommendations | Patients meeting any of the high risk criteria (Injury Patterns and Mental Status and Vital Signs) “should be preferentially transported to the highest level trauma center available within the geographic constraints of the regional trauma system” | | Patients meeting any of the Step 1 (physiologic) or Step 2 (anatomic) criteria “should be transported preferentially to the highest level of care within the defined trauma system” |

Continued next page

TABLE 2. (Continued)

| Type of Change | Changes in 2021 Field Triage Guidelines | Age Range | 2011 Field Triage Guidelines |
|----------------|---|-----------|---|
| | Patients not meeting high risk criteria, but meeting any of the moderate risk criteria “should be preferentially transported to a trauma center, as available within the geographic constraints of the regional trauma system (need not be the highest level trauma center)”. | | Patients not meeting Step 1 or 2 criteria but meeting Step 3 criteria, “transport to a trauma center, which, depending upon the defined trauma system, need not be the highest level trauma center” Patients not meeting Steps 1, 2, or 3, but meeting Step 4 criteria, “transport to a trauma center or hospital capable of timely and thorough evaluation and initial management of potentially serious injuries. Consider consultation with medical control.” |

would be cumbersome and nonfeasible for field use. Therefore, this criterion only applies to adults and older adults.

- New criterion: $SBP < 70 \text{ mm Hg} + (2 \times \text{age in years})$ (children 0–9 years)

Rationale: Two studies showed that age-adjusted hypotension (calculated using this formula) in the ED is an independent predictor of mortality among injured children.^{12,16} Inclusion of this criterion aligns the triage guideline with Advanced Trauma Life Support training⁵² and was viewed by the panel as a pediatric-specific training opportunity for EMS. Children older than 9 years reach the adult threshold of $SBP < 90 \text{ mm Hg}$ using the formula. Because hypotension is a late finding of pediatric shock (decompensated shock), EMS training on the use of visual cues (e.g., pallor, mottling, cyanosis) is encouraged, as represented in the Pediatric Assessment Triangle.^{53–55}

- New criterion: “Respiratory distress or need for respiratory support” replaces “need for ventilatory support” and “respiratory rate < 20 in infant aged < 1 year.”

Rationale: The criterion “need for ventilatory support” was added in 2011⁶ based on three studies showing that need for airway management and assisted ventilation was highly predictive of serious injury and death.^{56–58} There have since been four studies evaluating the need for respiratory support (variably defined as assisted ventilation, intubation, or need for mechanical ventilation), which showed a sensitivity of 8% to 53% and a specificity of 61% to 100% for identifying patients with serious injury.³⁵ The panel revised the wording to “need for respiratory support” based on EMS feedback. Because there is not a specific respiratory rate threshold for injured infants,⁵⁸ the “respiratory rate < 20 in infants” criterion was removed. The panel included “respiratory distress” to facilitate EMS training on important examination findings that precede the need for respiratory support, particularly in children.^{53–55}

- New criterion: Room-air pulse oximetry $< 90\%$

Rationale: Pulse oximetry is widely available on portable monitors used by EMS and has been evaluated in five studies, with most using a cut point of $< 90\%$.³⁵ Pulse oximetry had AUROC values of 0.59 to 0.76 for identifying patients with serious injury, similar to the respiratory rate criterion.³⁵ While most studies were conducted in adults, one study demonstrated the predictive utility of pulse oximetry in injured children⁵⁷ and another study showed the benefit of respiratory support and correction of hypoxia among young children with traumatic brain injury.⁵⁹ Therefore, this criterion applies to patients of all ages.

Retained Criteria (No Changes)

- Retained criterion: $SBP < 90 \text{ mm Hg}$

Rationale: The predictive utility of hypotension is supported by 49 studies, most of which evaluated a cut point of $SBP < 90 \text{ mm Hg}$.³⁴ A meta-analysis of 17 studies showed that prehospital $SBP < 90 \text{ mm Hg}$ had a pooled sensitivity of 19%, a specificity of 95%, and an AUROC of 0.67 for patients with serious injuries.³⁴ Higher thresholds for SBP modestly raised sensitivity, but lowered specificity,³⁴ and the panel sought to preserve the specificity of this measure. This criterion applies to patients 10 years and older, with use of a higher threshold for older adults.

- Retained criterion: $SBP < 110 \text{ mm Hg}$ for older adults

Rationale: The criterion “ $SBP < 110 \text{ mm Hg}$ might represent shock after age 65 years” was added to the “Special Considerations” section in 2011 to address the issue of undertriage among older adults.⁶ Because $SBP < 90 \text{ mm Hg}$ has a sensitivity of 4% to 5% for identifying seriously injured older adults, a higher SBP threshold improves sensitivity (13–29%) while preserving specificity (83–93%) in this population.³⁴ This criterion was moved from the Special Considerations section to Mental Status and Vital Signs for clarity and consistency.

- Retained criterion: Respiratory rate of < 10 or > 29 breaths per minute

Rationale: Respiratory rate is the most commonly studied respiratory triage criterion (25 studies), with respiratory rate of < 10 or > 29 breaths per minute being the most studied parameters.³⁵ This criterion had a pooled sensitivity of 13% and a specificity of 96% for identifying seriously injured patients, with an AUROC of 0.70.³⁵ While most studies were conducted in adults, a respiratory rate < 10 or > 29 breaths/minute demonstrated good predictive utility in children and older adults, yet with more variability in the accuracy estimates.³⁵ This criterion applies to patients of all ages.

Mechanism of Injury Criteria

Because anatomic and physiologic criteria identify less than half of patients with serious injuries,^{28,60,61} the mechanism criteria are important in the triage process. However, the mechanism criteria are less specific for serious injuries (lower +LR) and therefore are included in the “moderate risk” category. Based on high undertriage associated with previous versions of the guideline,^{25–27,29} the panel considered changes to reduce

undertriage, particularly in children. There is one new criterion, three modified criteria, and four unchanged criteria.

New and Modified Criteria

- New criterion: Child (age 0–9 years) unrestrained or in unsecured child safety seat

Rationale: Motor vehicle crashes are a common cause of pediatric injury. Lack of appropriate restraints is a consistent factor among seriously injured children.^{62–66} Unrestrained children have higher injury severity, greater trauma resource needs and are more likely to die than restrained children.^{62–64} Lack of restraint use also has been shown to predict seriously injured children involved in motor vehicle crashes.⁶⁶ The panel felt that this criterion was most pertinent for children 0 to 9 years, which provided consistency with the age range for pediatric blood pressure to simplify EMS training.

- Modified criterion: Significant intrusion (including roof) >12 in occupant site or >18 in any site or need for extrication of the entrapped patient

Rationale: As criteria already present in the guideline,^{6,67} additional studies have confirmed the predictive utility of these criteria in adults and children.^{36,60,61,68} Extrication >20 minutes was removed from the 2006 guideline based on varying definitions of “prolonged extrication” in the literature and the belief that the intrusion criteria would capture patients requiring extrication.^{6,67} However, a systematic review showed that extrication of any duration was a significant predictor of serious injury in adults and children³⁶ and that predictive utility was retained down to ≥5 minutes.⁶¹ Based on these studies, the panel added the extrication criterion back to the guideline, without a specific time requirement. Because different studies used “extrication” and “entrapment” interchangeably, the panel integrated these terms for the criterion.

- Modified criterion: Rider separated from transport vehicle with significant impact (e.g., motorcycle, ATV, horse, etc.)

Rationale: Different versions of the motorcycle crash criterion have been present since the 1990 guideline,⁵ despite limited data. A study of adults not meeting physiologic or anatomic criteria showed that motorcycle crash >20 mph or with rider separation had poor overall predictive utility (+LR, 1.0–1.2).⁶¹ With only a single study evaluating the motorcycle criterion in the past 10 years³⁶ and the speed component offering little predictive yield,⁶¹ the panel removed the speed requirement and broadened the type of transport vehicle for greater application to children.⁶⁰

- Modified criterion: fall from height >10 ft (all ages)

Rationale: The 2011 guideline specified falls >20 ft in adults and >10 ft in children (or two to three times the height of the child).⁶ However, the >10 ft criterion has good predictive utility for children⁶⁰ and adults.⁶¹ The criterion specifying two to three times the height of the child was based on research in young children falling from bunk beds⁶⁹ but has not demonstrated improved prediction compared with a >10 ft criterion.⁶⁰ For consistency and simplicity, the panel opted to use the same fall height

for children and adults and to remove the age-based height for children.

- Modified criterion: Pedestrian/bicycle rider thrown, run over, or with significant impact

Rationale: This triage criterion was included in the 1987 guideline, with slight modifications over time.^{5,6} Six studies published since 2011 showed mixed results (+LR, 0.4–2.8).³⁶ In a study of children not meeting physiologic or anatomic criteria, the pedestrian criterion was predictive for patients run over and with significant impact (>20 mph).⁶⁰ Among adults, this criterion demonstrated predictive utility with higher speed of impact (+LR ≥2.2).⁶¹ Because this criterion has long existed in the triage guideline with reasonable predictive utility for children, the panel retained the criterion and simplified the wording.

Retained Criteria (No Changes)

- Retained criterion: Ejection (partial or complete) from automobile

Rationale: Among multiple studies published since 2011, most showed that ejection remains a significant predictor of serious injury and death in adults and children.^{36,60,61}

- Retained criterion: Death in passenger compartment

Rationale: In several studies published since 2011, death of another passenger in the same vehicle predicted serious injury in adults and children.^{36,60,61}

- Retained criterion: Vehicle telemetry data consistent with severe injury

Rationale: This criterion was added to the 2006 guideline based on promising developments in automated collision notification systems and retained in 2011 based on six studies demonstrating predictive utility and the potential for transmission to 9-1-1 dispatch centers.⁶ In five recent studies, crash algorithms had good predictive utility (+LR, 4.7–22.2),³⁶ yet studies evaluating real-time use of vehicle telemetry for field triage are lacking.

Emergency Medical Services Judgment (Previously Step 4 Special Considerations)

The “Special Considerations” step has changed over time to include special populations, unique triage factors, and EMS provider judgment.⁶ While some studies of EMS provider judgment have had mixed results,^{70,71} others have shown judgment to be independently associated with serious injury.⁷² The panel felt that EMS judgment plays an important role in field triage, but is dependent on training and experience. For the current guideline, the panel created a category for “EMS Judgment” to replace “Special Considerations” and provided structured guidance on factors to consider in the decision-making process. The criteria in this section generally have less evidence and lower predictive utility, but remain important considerations in field triage. There are three new criteria, one modified criterion, and three unchanged criteria.

New and Modified Criteria

- New criterion: Suspicion of child abuse

Rationale: Child abuse can be difficult to diagnose and have subtle presentations, yet with potentially devastating consequences.

Mortality is elevated among abused children, especially with recurrent episodes of abuse.⁷³ Abused children frequently require specialty care to address their injuries and the complex legal, logistical, social, and investigative aspects of these incidents. Trauma centers are required to have protocols in place to provide comprehensive evaluation of such children, with guidelines and best practices published by several national trauma organizations.^{74–76} Training EMS clinicians to recognize the signs of child abuse and integration of prehospital information into ED-based clinical decision support systems^{77,78} are supported by multiple national organizations. Based on these considerations, the panel added suspicion of child abuse.

- New criterion: Special, high-resource health care needs

Rationale: Various comorbid conditions were in the triage guidelines from 1987 to 2006, but were removed in 2011 because of lack of evidence.^{5,6} Among five recent studies evaluating the use of comorbidities for field triage,³⁶ some showed that comorbidities were independently associated with death and could reduce undertriage among older adults.³⁶ However, the predictive utility of comorbidities varied across studies (+LR, 0.8–3.1).³⁶ The panel recognized that injured patients with special health care needs related to comorbidities (e.g., ventilator dependence or ventricular assist devices) may require the resources and expertise of trauma centers.

- Modified criterion: Low level falls in young children (age ≤5 years) or older adults (age ≥65 years) with significant head impact.

Rationale: The panel moved these criteria from the Mechanism and Special Considerations sections to EMS Judgment. Research has shown that some children incur serious injuries from low-height falls, including falls from standing,⁶⁰ and that such falls are a common cause of traumatic brain injury in young children.⁷⁹ For older adults, ground-level falls can cause serious injury and death,^{19,80–82} which were the reasons for inclusion in the 2011 guideline.⁶ However, ground-level falls are common among older persons and therefore are relatively nonspecific for serious injury (+LR, 1.2–1.9).²⁵ Based on concerns that these criteria could result in overtriage, the panel included these factors under EMS Judgment and added “with significant head impact.”

- Modified criterion: Anticoagulation use

Rationale: “Coagulopathy” was added to the triage guideline in 1990⁵ and included in the 2011 guideline as “Anticoagulants and bleeding disorders — patients with head injury are at high risk for rapid deterioration.”³⁶ Five recent studies evaluating anticoagulant use for triage showed relatively low predictive utility (+LR, 0.73–1.8).³⁶ Some research suggests that such a criterion could help identify older adults with intracranial hemorrhage,⁸³ but other studies show otherwise.⁸⁴ In a prospective study of older adults transported by EMS, the incidence of brain hemorrhage was similar between patients taking versus not taking anticoagulants.⁸⁵ Based on these data, the panel felt that use of anticoagulants (including antiplatelet agents) was best considered in the context of EMS Judgment.

Retained Criteria (No Changes)

- Retained criterion: Pregnancy >20 weeks

Rationale: The pregnancy criterion was added to the 1999 guideline⁵ and refined to “pregnancy >20 weeks” in 2006.⁶⁷ While pregnancy does not necessarily increase the likelihood of serious injury, simultaneous management of the mother and unborn child can create complex clinical scenarios requiring trauma centers with obstetrics capabilities. Therefore, the panel felt that this factor should be part of EMS Judgment.

- Retained criterion: Burns in conjunction with trauma

Rationale: Consistent with Advanced Trauma Life Support teaching, when a burn patient has other injuries, the injuries should be evaluated and potentially prioritized over the burn. Trauma centers have the capability to quickly evaluate these patients to expedite care for both clinical conditions.

- Retained criterion: Children should be triaged preferentially to pediatric capable trauma centers

Rationale: For injured children, research has demonstrated higher survival in pediatric trauma centers compared with adult or mixed trauma centers.¹² However, many regions do not have access to pediatric trauma centers.⁸⁶ While transport to a pediatric trauma center is preferable, the panel felt that transport to pediatric versus adult trauma centers should be determined by local protocols and proximity. Based on stakeholder and expert feedback, the panel chose not to use a specific age to define children, as there is insufficient evidence for a specific age limit and systems have established varying age limits based on local resources and practice patterns. Because high ED pediatric readiness has been associated with improved short- and long-term survival of children in US trauma centers,^{15,16} all trauma centers are strongly encouraged to meet such criteria.

TRANSPORT RECOMMENDATIONS

Emergency medical services systems vary by geography, organization, resources, service levels, staffing, training, access to air medical services, travel times, oversight, and governance. Trauma centers are hospitals that are prepared to provide emergent care for seriously injured patients through resources, personnel, expertise, education, and quality improvement programs. There are national standards for adult and pediatric trauma centers, with trauma center designation (Levels I to V) typically made at the state level (Table 3). State trauma systems may be inclusive or exclusive, with inclusive systems categorizing most hospitals and demonstrating lower injury-related mortality.⁸⁷

Recognizing the variability in EMS and trauma systems, transport recommendations in the guideline allow local flexibility. There is not a “one size fits all” that will work for all systems. While the survival benefit of regionalized trauma care is driven primarily by Level I hospitals,^{7–9} there are large regions across the United States that do not have immediate access to such trauma centers. Although 84% to 88% of US residents have access to a Level I or II trauma center within 60 minutes, these proportions are substantially lower when limited to ground travel and shorter time windows.^{88,89} Access to pediatric trauma centers is even lower,⁸⁶ with widely variable proximity by state.⁹⁰ Rural regions have the most limited access to Levels I and II trauma centers,^{86,88,89} resulting in higher undertriage, more interhospital

TABLE 3. Characteristics of Trauma Centers

| Level | Criteria |
|-------|---|
| I | <ul style="list-style-type: none"> • Regional resource center expected to manage large numbers of seriously injured patients • Admit $\geq 1,200$ trauma patients or have ≥ 240 admissions with ISS ≥ 16 per year • Attending trauma surgeon participates in major resuscitations in ED, present at operative procedures, and actively involved in critical care of all seriously injured patients (24-h in-house availability) • Immediate availability of board-certified emergency physicians, general surgeons, anesthesiologists, neurosurgeons, and orthopedic surgeons • Maintain a surgically directed critical care service • Participate in resident training • Be a leader in education and outreach activities • Conduct trauma research |
| II | <ul style="list-style-type: none"> • Regional resource center expected to manage large numbers of seriously injured patients • Same standards for provision of clinical care without the volume requirements • No requirement for resident training, education, outreach, trauma research, or surgically directed critical care service |
| III | <ul style="list-style-type: none"> • Capability to initially manage the majority of injured patients • Transfer agreements with Level I or II trauma centers for seriously injured patients • Continuous general surgical coverage |
| IV | <ul style="list-style-type: none"> • Often serve rural regions and supplement care within a larger trauma system • Initial evaluation and assessment of injured patients, with expected transfer of many patients to higher-level trauma centers • Transfer agreements with higher-level trauma centers • 24-h emergency coverage by a physician or midlevel provider • Frequently lack continuous surgical coverage |

From Resources for the Optimal Care of the Injured Patient, Committee on Trauma, American College of Surgeons, 2014. There is variation in state-to-state definitions and designations of trauma centers. There are separate processes and criteria for pediatric trauma centers.

ISS, Injury Severity Score.

transfers, and longer transfer distances compared with urban settings.⁹¹ The triage guideline is intended to provide a template that can be adapted for use in all systems.

When feasible, patients meeting the “high risk” criteria should be triaged to the highest-level trauma center within the region, including consideration of air medical services. Injured patients meeting the physiologic criteria have lower mortality when cared for in Level I trauma centers.⁹² Air medical services may offer advanced care clinicians, access to additional interventions, and more rapid transport. Emergency medical services medical directors and trauma system managers are encouraged to evaluate the resources relevant to their systems to guide implementation of the field triage guideline. Because time is known to be crucial for certain trauma patients,⁹³ field triage favors short time intervals. However, the current evidence is insufficient to make specific recommendations regarding transport times and when air medical services should be activated.⁹⁴ Some EMS systems may opt to implement a closest hospital approach for patients with an unstable airway, severe shock, traumatic arrest, or other “extremis” conditions for initial stabilization, before higher level transport for definitive care.

IMPLEMENTATION AND ADHERENCE TO THE FIELD TRIAGE GUIDELINE

The triage guideline is not useful if not fully implemented into trauma systems and adopted by EMS clinicians. Following the 2006 triage guideline, only 17% of states had full adoption of the new guideline, with 61% using an older version or a different protocol altogether.⁹⁵ In a study of six metropolitan regions, only one region had adopted the most recent triage guideline within 2 years and 36% of triage criteria in use had been previously removed or never included.⁹⁶ Compliance with the field triage

guideline varies widely, with lowest adherence for the physiologic criteria.⁹⁷ Strict adherence would reduce undertriage.⁹⁸ While there are many hurdles to implementing an updated guideline, translating the science into practice is arguably the most important step of realizing effective field triage practices. The 2021 guideline is organized to facilitate ease of use, increase speed of decision-making, and promote adherence. We recommend adoption at the state level (similar to the trauma center designation process), allowing regional and local EMS and trauma systems to determine system-specific adaptations for hospital selection.

FUTURE RESEARCH

There is substantial need for future research to inform the triage guideline. Noninvasive monitor technology and point-of-care testing hold promise for field triage, particularly for seriously injured patients not meeting the high-risk criteria. Systematic reviews of circulatory and respiratory criteria identified several promising measures (e.g., point-of-care lactate, end-tidal CO₂, and heart rate variability), but more research and technology are needed to facilitate field use.^{34,35} Research on new criteria added to the 2021 guideline will be particularly important, as well as studies on the real-time use of automated collision notification systems for field triage.

The 2021 guideline includes changes in format, structure, and content. Research is needed to evaluate the usability, performance, adherence, and application of the new guideline (including the impact on health outcomes), particularly compared with the 2011 guideline. Research on efficient and effective training methods, including training frequency, is also needed. Understanding how and why EMS clinicians make triage decisions, including concordance versus discordance with the guideline, will be important in optimizing triage performance. Based on the slow and variable uptake of previous triage guidelines,^{95,96}

creating new ways of disseminating, implementing, and monitoring adherence will be important to realizing the true potential of the guideline. Finally, there is a need for more system-based research to inform transport times, when air medical services should be activated, and the role of different provider levels.

CONCLUSION

The 2021 field triage guideline is based on the most current science, a national panel of interdisciplinary experts, direct feedback from EMS clinicians, and input from many stakeholders. This guideline presents an opportunity to improve the prehospital care of injured patients across the United States. Effective field triage is foundational to trauma systems, concentrating the most seriously injured patients in trauma centers to improve survival after injury.

AUTHORSHIP

E.M.B., C.D.N., P.E.F., M.G., and H.N.M. contributed in the study conception and design. R.C. and J.R.L. contributed in the systematic reviews. R.C., J.R.L., E.M.B., and C.D.N. contributed to the literature search (systematic reviews and other). P.E.F., M.G., H.N.M., and E.M.B. contributed to the EMS feedback. E.M.B., C.D.N., P.E.F., M.G., and H.N.M. participated as the steering committee. H.N.M., M.D., M.N., and J.D. participated as administrative support. All authors participated as panel members. E.M.B. was the panel leader. All authors contributed to the interpretation of results. E.M.B. obtained funding. C.D.N. performed the drafting of manuscript. All authors contributed in the critical revision.

ACKNOWLEDGMENTS

We thank the Office of Emergency Medical Services at the National Highway Traffic Safety Administration for their support, guidance, and involvement in this project.

Additional Authors Included in the Writing Group:

Jeffrey M. Goodloe, MD, FACEP, FAEMS, Department of Emergency Medicine, University of Oklahoma School of Community Medicine Tulsa, Oklahoma; John H. Armstrong, MD, University of South Florida Morsani College of Medicine, Tampa, Florida; John M. Gallagher, MD, FAEMS, FACEP, Board of Directors, National Association of EMS Physicians, Overland Park, Kansas; Stewart C. Wang, MD PhD FACS, Department of Surgery, University of Michigan, Ann Arbor, Michigan; Brian J. Eastridge, MD, FACS, Division of Trauma and Emergency General Surgery, Department of Surgery, University of Texas Health Science Center at San Antonio, San Antonio, Texas; N. Clay Mann, PhD, MS, MBA, Department of Pediatrics, University of Utah School of Medicine, Salt Lake City, Utah; Ron R. Lawler, BUS, NRP, Sanford Ambulance, Fargo, North Dakota; Jeffrey P. Salomone, MD, FACS, Banner Desert Medical Center, Mesa, Arizona; Roger Chou, MD, FACP, Departments of Medicine and Medical Informatics and Clinical Epidemiology, Oregon Health & Science University, Portland, Oregon; Nathan A.M. Christopherson, DNP, MBA, MSN, RN, EMT-P, Department of Surgery, Donald and Barbara Zucker School of Medicine at Hofstra/Northwell, Northwell Health, Manhasset, New York; Jorie Klein, MSN, MBA, BSN, RN, Texas Department of State Health Services, Austin, Texas; Scott M. Sasser MD, FACEP, Prisma Health Medical Group, Department of Emergency Medicine, University of South Carolina School of Medicine Greenville, Greenville, South Carolina; Laura N. Godat, MD FACS, Division of Trauma, Surgical Critical Care, Burns and Acute Care Surgery, Department of Surgery, UC San Diego Health, San Diego, California; Jeff Gilchrist, MHA, BA, RN, CEN, CPEN, NREMT-P, CCERT-P, UnityPoint Health, Marshalltown, Iowa; Joshua R. Lupton, MD, MPH, MPhil, Center for Policy and Research in Emergency Medicine, Department of Emergency Medicine, Oregon Health & Science University, Portland, Oregon; Robert T. Russell, MD, MPH, Pediatric Surgery, Department of Surgery, Children's of Alabama, University of Alabama at Birmingham, Birmingham, Alabama; Dennis Rowe, EMT-P, Government and Industry Relations, Priority OnDemand, Priority Ambulance, Knoxville, Tennessee; Melanie Neal, MS, Committee on Trauma, American College of Surgeons, Chicago, Illinois; Mackenzie Dafferner, MPH, Committee on Trauma, American College of Surgeons, Chicago, Illinois; Jimm Dodd, MS, MA; Committee on Trauma, American College of Surgeons, Chicago, Illinois.

Expert Panel

Eileen Bulger, ACS Steering Committee; Craig Newgard, ACS Steering Committee + expertise based; Mark Gestring, ACS Steering Committee; Greg Jurkovich, ACS Steering Committee; Joshua Brown, ACS Steering Committee + expertise based; Peter Fischer, ACS Steering Committee; E. Brooke Lerner, ACS Steering Committee; Mary Fallat, ACS Steering Committee + American Academy of Pediatrics; Clay Mann, ACS Steering Committee + expertise based; Brian Eastridge, ACS Steering Committee; Bellal Joseph, expertise based; Laura Godat, expertise based; John Armstrong, past guidelines; Jorie Klein, past guidelines; Scott Sasser, past guidelines; Stewart C. Wang, past guidelines; Jeff Goodloe, American College of Emergency Physicians; Lisa Gray, Emergency Medical Services for Children; Jeffrey Gilchrist, Emergency Nurses Association; Ron Lawler, National Association of EMS Educators; Dennis Rowe, National Association of Emergency Medical Technicians; Theodore Delbridge, National Association of State EMS Officials; Jon Krohmer, National Highway Traffic Safety Administration; Robert Russell, Pediatric Trauma Society; Nathan Christopherson, Society of Trauma Nurses; Jeffrey Salomone, National Registry of EMTs; and John M. Gallagher, National Association of EMS Physicians.

List of Organizations and Federal Agencies Endorsing the Field Triage Guidelines

American College of Emergency Physicians, Emergency Medical Services for Children, Emergency Nurses Association, National Association of EMS Educators, National Association of Emergency Medical Technicians, National Association of State EMS Officials, Pediatric Trauma Society, Society of Trauma Nurses, National Registry of EMTs, National Association of EMS Physicians, American Academy of Pediatrics, and American College of Surgeons Committee on Trauma.

DISCLOSURE

The authors declare no conflicts of interest.

The American College of Surgeons was funded to perform an evidence-based revision of the Field Triage Guidelines as part of Cooperative Agreement number 693J91950007 between the American College of Surgeons and the National Highway Traffic Safety Administration, Office of Emergency Medical Services funded in part by the Health Resources and Services Administration, Maternal and Child Health Bureau, and Emergency Medical Services for Children Program. The contents are those of the authors and do not necessarily represent the official views of, nor an endorsement, by the US Government. For more information, please visit EMS.gov and HRSA.gov.

REFERENCES

1. *10 Leading Causes of Death, United States 2019, Both Sexes, All Ages, All Races*. Atlanta, GA: Centers for Disease Control and Prevention; 2021. Available at: <https://wisqars-viz.cdc.gov:8006/lcd/home>. Accessed February 18, 2022.
2. Borse NN, Rudd RA, Dellinger AM, Sleet DA. Years of potential life lost from unintentional child and adolescent injuries—United States, 2000–2009. *J Safety Res*. 2013;45:127–131.
3. Wang HE, Mann NC, Jacobson KE, Ms MD, Mears G, Smyrski K, et al. National characteristics of emergency medical services responses in the United States. *Prehosp Emerg Care*. 2013;17(1):8–14.
4. *Resources for Optimal Care of the Injured Patient*. 6th. ed. American College of Surgeons; 2014.
5. Mackersie RC. History of trauma field triage development and the American College of Surgeons criteria. *Prehosp Emerg Care*. 2006;10(3):287–294.
6. Sasser SM, Hunt RC, Faul M, Sugerman D, Pearson WS, Dulski T, et al. Guidelines for field triage of injured patients: recommendations of the National Expert Panel on Field Triage, 2011. *MMWR Recomm Rep*. 2012;61(RR-1):1–20.
7. MacKenzie EJ, Rivara FP, Jurkovich GJ, Nathens AB, Frey KP, Egleston BL, et al. A national evaluation of the effect of trauma-center care on mortality. *N Engl J Med*. 2006;354(4):366–378.
8. Sampalis JS, Denis R, Lavoie A, Frechette P, Boukas S, Nikolis A, et al. Trauma care regionalization: a process-outcome evaluation. *J Trauma*. 1999;46(4):565–579 discussion 79–81.
9. Mullins RJ, Veum-Stone J, Helfand M, Zimmer-Gembeck M, Hedges JR, Southard PA, et al. Outcome of hospitalized injured patients after institution of a trauma system in an urban area. *JAMA*. 1994;271(24):1919–1924.
10. Nathens AB, Jurkovich GJ, Cummings P, Rivara FP, Maier RV. The effect of organized systems of trauma care on motor vehicle crash mortality. *JAMA*. 2000;283(15):1990–1994.

11. Nathens AB, Jurkovich GJ, Rivara FP, Maier RV. Effectiveness of state trauma systems in reducing injury-related mortality: a national evaluation. *J Trauma*. 2000;48(1):25–30 discussion-1.
12. Sathya C, Alali AS, Wales PW, Scales DC, Karanickolas PJ, Burd RS, et al. Mortality among injured children treated at different trauma center types. *JAMA Surg*. 2015;150(9):874–881.
13. Pracht EE, Tepas JJ 3rd, Langland-Orban B, Simpson L, Pieper P, Flint LM. Do pediatric patients with trauma in Florida have reduced mortality rates when treated in designated trauma centers? *J Pediatr Surg*. 2008;43(1):212–221.
14. Wang NE, Saynina O, Vogel LD, Newgard CD, Bhattacharya J, Phibbs CS. The effect of trauma center care on pediatric injury mortality in California, 1999 to 2011. *J Trauma Acute Care*. 2013;75(4):704–716.
15. Newgard CD, Lin A, Goldhaber-Fiebert JD, Marin JR, Smith M, Cook JNB, et al. Association of emergency department pediatric readiness with mortality to 1 year among injured children treated at trauma centers. *JAMA Surg*. 2022;e217419.
16. Newgard CD, Lin A, Olson LM, Cook JNB, Gausche-Hill M, Kuppermann N, et al. Evaluation of emergency department pediatric readiness and outcomes among US trauma centers. *JAMA Pediatr*. 2021.
17. Pracht EE, Langland-Orban B, Flint L. Survival advantage for elderly trauma patients treated in a designated trauma center. *J Trauma*. 2011;71(1):69–77.
18. Garwe T, Stewart K, Newgard CD, Stoner J, Sacra JC, Cody P, et al. Survival benefit of treatment at or transfer to a tertiary trauma center among injured older adults. *Prehosp Emerg Care*. 2019;1:18.
19. Staudenmayer KL, Hsia RY, Mann NC, Spain DA, Newgard CD. Triage of elderly trauma patients: a population-based perspective. *J Am Coll Surg*. 2013;217(4):569–576.
20. Haas B, Gomez D, Zagorski B, Stukel TA, Rubenfeld GD, Nathens AB. Survival of the fittest: the hidden cost of undertriage of major trauma. *J Am Coll Surg*. 2010;211(6):804–811.
21. Yoder A, Bradburn EH, Morgan ME, Vernon TM, Bresz KE, Gross BW, et al. An analysis of overtriage and undertriage by advanced life support transport in a mature trauma system. *J Trauma Acute Care*. 2020;88(5):704–709.
22. Haas B, Stukel TA, Gomez D, Zagorski B, De Mestral C, Sharma SV, et al. The mortality benefit of direct trauma center transport in a regional trauma system: a population-based analysis. *J Trauma Acute Care*. 2012;72(6):1510–1515 discussion 5-7.
23. Hewes HA, Christensen M, Taillac PP, Mann NC, Jacobsen KK, Fenton SJ. Consequences of pediatric undertriage and overtriage in a statewide trauma system. *J Trauma Acute Care*. 2017;83(4):662–667.
24. Newgard CD, Hsia RY, Mann NC, Schmidt T, Sahni R, Bulger EM, et al. The trade-offs in field trauma triage: a multiregion assessment of accuracy metrics and volume shifts associated with different triage strategies. *J Trauma Acute Care*. 2013;74(5):1298–1306 discussion 306.
25. Lupton JR, Davis-O'Reilly C, Jungbauer RM, Newgard CD, Fallat ME, Brown JB, et al. Under-triage and over-triage using the field triage guidelines for injured patients: a systematic review. *Prehosp Emerg Care*. 2022; [epub before print].
26. van der Sluijs R, van Rein EAJ, Wijnand JGJ, Leenen LPH, van Heijl M. Accuracy of pediatric trauma field triage: a systematic review. *JAMA Surg*. 2018;153(7):671–676.
27. Lerner EB, Cushman JT, Drendel AL, Badawy M, Shah MN, Guse CE, et al. Effect of the 2011 revisions to the field triage guidelines on under- and over-triage rates for pediatric trauma patients. *Prehosp Emerg Care*. 2017;21(4):456–460.
28. Newgard CD, Zive D, Holmes JF, Bulger EM, Staudenmayer K, Liao M, et al. A multisite assessment of the American College of Surgeons Committee on Trauma field triage decision scheme for identifying seriously injured children and adults. *J Am Coll Surg*. 2011;213(6):709–721.
29. Newgard CD, Fu R, Zive D, Rea T, Malveau S, Daya M, et al. Prospective validation of the national field triage guidelines for identifying seriously injured persons. *J Am Coll Surg*. 2016;222(2):146–158.
30. Garwe T, Stewart K, Stoner J, Newgard CD, Scott M, Zhang Y, et al. Out-of-hospital and inter-hospital under-triage to designated tertiary trauma centers among injured older adults: a 10-year statewide geospatial-adjusted analysis. *Prehosp Emerg Care*. 2017;21(6):734–743.
31. Uribe-Leitz T, Jarman MP, Sturgeon DJ, Harlow AF, Lipsitz SR, Cooper Z, et al. National study of triage and access to trauma centers for older adults. *Ann Emerg Med*. 2020;75(2):125–135.
32. Chen Y, Yang K, Marusic A, Qaseem A, Meerpohl JJ, Flottorp S, et al. A reporting tool for practice guidelines in health care: the RIGHT statement. *Ann Intern Med*. 2017;166(2):128–132.
33. Chou R, Totten AM, Carney N, Dandy S, Fu R, Grusing S, et al. Predictive utility of the total Glasgow Coma Scale versus the motor component of the Glasgow Coma Scale for identification of patients with serious traumatic injuries. *Ann Emerg Med*. 2017;70(2):143–57 e6.
34. Newgard CD, Cheney TP, Chou R, Fu R, Daya MR, O'Neil ME, et al. Out-of-hospital circulatory measures to identify patients with serious injury: a systematic review. *Acad Emerg Med*. 2020;27(12):1323–1339.
35. Daya MR, Cheney TP, Chou R, Fu R, Newgard CD, O'Neil ME, et al. Out-of-hospital respiratory measures to identify patients with serious injury: a systematic review. *Acad Emerg Med*. 2020;27(12):1312–1322.
36. Lupton JR, Davis-O'Reilly C, Jungbauer RM, Newgard CD, Fallat ME, Brown JB, et al. Mechanism of injury and special considerations as predictive of serious injury: a systematic review. *Acad Emerg Med*. 2022; (In Press).
37. Fischer PEGM, Sagraves SG, Michaels HN, Patel B, Dodd J, Campion EM, Vander Kolk WE, Bulger EM. The national trauma triage protocol: how EMS perspective can inform the guideline revision. *Trauma Surg Acute Care Open*. 2022;7(1).
38. Jones CM, Cushman JT, Lerner EB, Fisher SG, Seplaki CL, Veazie PJ, et al. Prehospital trauma triage decision-making: a model of what happens between the 9-1-1 call and the hospital. *Prehosp Emerg Care*. 2016;20(1):6–14.
39. Newgard CD, Nelson MJ, Kampp M, Saha S, Zive D, Schmidt T, et al. Out-of-hospital decision making and factors influencing the regional distribution of injured patients in a trauma system. *J Trauma*. 2011;70(6):1345–1353.
40. Lerner EB, Roberts J, Guse CE, Shah MN, Swor R, Cushman JT, et al. Does EMS perceived anatomic injury predict trauma center need? *Prehosp Emerg Care*. 2013;17(3):312–316.
41. Kragh JF Jr., Littrel ML, Jones JA, Walters TJ, Baer DG, Wade CE, et al. Battle casualty survival with emergency tourniquet use to stop limb bleeding. *J Emerg Med*. 2011;41(6):590–597.
42. Kragh JF Jr., Walters TJ, Baer DG, Fox CJ, Wade CE, Salinas J, et al. Survival with emergency tourniquet use to stop bleeding in major limb trauma. *Ann Surg*. 2009;249(1):1–7.
43. Beekley AC, Sebesta JA, Blackburne LH, Herbert GS, Kauvar DS, Baer DG, et al. Prehospital tourniquet use in Operation Iraqi Freedom: effect on hemorrhage control and outcomes. *J Trauma*. 2008;64(2 Suppl):S28–S37 discussion S.
44. Scerbo MH, Holcomb JB, Taub E, Gates K, Love JD, Wade CE, et al. The trauma center is too late: major limb trauma without a pre-hospital tourniquet has increased death from hemorrhagic shock. *J Trauma Acute Care*. 2017;83(6):1165–1172.
45. Inaba K, Siboni S, Resnick S, Zhu J, Wong MD, Haltmeier T, et al. Tourniquet use for civilian extremity trauma. *J Trauma Acute Care*. 2015;79(2):232–237 quiz 332–3.
46. Ode G, Studnek J, Seymour R, Bosse MJ, Hsu JR. Emergency tourniquets for civilians: can military lessons in extremity hemorrhage be translated? *J Trauma Acute Care*. 2015;79(4):586–591.
47. Schroll R, Smith A, McSwain NE Jr., Myers J, Rocchi K, Inaba K, et al. A multi-institutional analysis of prehospital tourniquet use. *J Trauma Acute Care*. 2015;79(1):10–14 discussion 4.
48. Acker SN, Ross JT, Partrick DA, Nadlonek NA, Bronsert M, Bensard DD. Glasgow motor scale alone is equivalent to Glasgow Coma Scale at identifying children at risk for serious traumatic brain injury. *J Trauma Acute Care*. 2014;77(2):304–309.
49. Cicero MX, Cross KP. Predictive value of initial Glasgow Coma Scale score in pediatric trauma patients. *Pediatr Emerg Care*. 2013;29(1):43–48.
50. Acker SN, Ross JT, Partrick DA, Tong S, Bensard DD. Pediatric specific shock index accurately identifies severely injured children. *J Pediatr Surg*. 2015;50(2):331–334.
51. Strutt J, Flood A, Kharbada AB. Shock index as a predictor of morbidity and mortality in pediatric trauma patients. *Pediatr Emerg Care*. 2019;35(2):132–137.
52. *ATLS Advanced Trauma Life Support 10th edition Student Course Manual*. 10th ed. Chicago, Illinois: American College of Surgeons; 2018.
53. Dieckmann RA, Brownstein D, Gausche-Hill M. The pediatric assessment triangle: a novel approach for the rapid evaluation of children. *Pediatr Emerg Care*. 2010;26(4):312–315.
54. Gausche-Hill M, Eckstein M, Horeczko T, McGrath N, Kurobe A, Ullum L, et al. Paramedics accurately apply the pediatric assessment triangle to drive management. *Prehosp Emerg Care*. 2014;18(4):520–530.

55. Fuchs S, Terry M, Adelgais K, Bokholdt M, Brice J, Brown KM, et al. Definitions and assessment approaches for emergency medical services for children. *Pediatrics*. 2016;138(6).
56. Newgard CD, Rudser K, Hedges JR, Kerby JD, Stiell IG, Davis DP, et al. A critical assessment of the out-of-hospital trauma triage guidelines for physiologic abnormality. *J Trauma*. 2010;68(2):452–462.
57. Newgard CD, Rudser K, Atkins DL, Berg R, Osmond MH, Bulger EM, et al. The availability and use of out-of-hospital physiologic information to identify high-risk injured children in a multisite, population-based cohort. *Prehosp Emerg Care*. 2009;13(4):420–431.
58. Newgard CD, Cudnik M, Warden CR, Hedges JR. The predictive value and appropriate ranges of prehospital physiological parameters for high-risk injured children. *Pediatr Emerg Care*. 2007;23(7):450–456.
59. Gaither JB, Spaite DW, Bobrow BJ, Keim SM, Barnhart BJ, Chikani V, et al. Effect of implementing the out-of-hospital traumatic brain injury treatment guidelines: the Excellence in Prehospital Injury Care for Children Study (EPIC4Kids). *Ann Emerg Med*. 2021;77(2):139–153.
60. Lerner EB, Badawy M, Cushman JT, Drendel AL, Fumo N, Jones CMC, et al. Does mechanism of injury predict trauma center need for children? *Prehosp Emerg Care*. 2021;25(1):95–102.
61. Lerner EB, Shah MN, Cushman JT, Swor RA, Guse CE, Brasel K, et al. Does mechanism of injury predict trauma center need? *Prehosp Emerg Care*. 2011;15(4):518–525.
62. Starnes M. Child passenger fatalities and injuries, based on restraint use, vehicle type, seat position, and number of vehicles in the crash, Department of Transportation #HS 809 784 Technical Report. Washington, DC: National Center for Statistics and Analysis; 2005. Available at: <https://crashstats.nhtsa.dot.gov/Api/Public/ViewPublication/809784>. Accessed May 16, 2022.
63. Chan L, Reilly KM, Telfer J. Odds of critical injuries in unrestrained pediatric victims of motor vehicle collision. *Pediatr Emerg Care*. 2006;22(9):626–629.
64. Valent F, McGwin G Jr., Hardin W, Johnston C, Rue LW 3rd. Restraint use and injury patterns among children involved in motor vehicle collisions. *J Trauma*. 2002;52(4):745–751.
65. Benedetti M, Klinich KD, Manary MA, Flannagan CAC. Factors affecting child injury risk in motor-vehicle crashes. *Stapp Car Crash J*. 2019;63:195–211.
66. Newgard CD, Lewis RJ, Jolly BT. Use of out-of-hospital variables to predict severity of injury in pediatric patients involved in motor vehicle crashes. *Ann Emerg Med*. 2002;39(5):481–491.
67. Sasser SM, Hunt RC, Sullivent EE, Wald MM, Mitchko J, Jurkovich GJ, et al. Guidelines for field triage of injured patients. Recommendations of the National Expert Panel on Field Triage. In: *MMWR Recomm Rep*. vol. 58 (RR-1). 2009:1–35.
68. Isenberg D, Cone DC, Vaca FE. Motor vehicle intrusion alone does not predict trauma center admission or use of trauma center resources. *Prehosp Emerg Care*. 2011;15(2):203–207.
69. Macgregor DM. Injuries associated with falls from beds. *Inj Prev*. 2000;6(4):291–292.
70. Mulholland SA, Cameron PA, Gabbe BJ, Williamson OD, Young K, Smith KL, et al. Prehospital prediction of the severity of blunt anatomic injury. *J Trauma*. 2008;64(3):754–760.
71. Mulholland SA, Gabbe BJ, Cameron P, Victorian State Trauma Outcomes Registry and Monitoring Group. Is paramedic judgement useful in prehospital trauma triage? *Injury*. 2005;36(11):1298–1305.
72. Newgard CD, Kampp M, Nelson M, Holmes JF, Zive D, Rea T, et al. Deciphering the use and predictive value of “emergency medical services provider judgment” in out-of-hospital trauma triage: a multisite, mixed methods assessment. *J Trauma Acute Care*. 2012;72(5):1239–1248.
73. Deans KJ, Thackeray J, Askegard-Giesmann JR, Earley E, Groner JJ, Minneci PC. Mortality increases with recurrent episodes of nonaccidental trauma in children. *J Trauma Acute Care*. 2013;75(1):161–165.
74. Rosen NG, Escobar MA Jr., Brown CV, Moore EE, Sava JA, Peck K, et al. Child physical abuse trauma evaluation and management: a Western Trauma Association and Pediatric Trauma Society critical decisions algorithm. *J Trauma Acute Care*. 2021;90(4):641–651.
75. Escobar MA Jr., Flynn-O'Brien KT, Auerbach M, Tiyyagura G, Borgman MA, Duffy SJ, et al. The association of nonaccidental trauma with historical factors, examination findings, and diagnostic testing during the initial trauma evaluation. *J Trauma Acute Care*. 2017;82(6):1147–1157.
76. ACS Trauma Quality Programs Best Practices Guidelines for Trauma Center Recognition of Child Abuse, Elder Abuse, and Intimate Partner Violence; 2019, American College of Surgeons Committee on Trauma, Chicago, Illinois. Available at: https://www.facs.org/media/o0wdimys/abuse_guidelines.pdf. Accessed May 16, 2022.
77. Berger RP, Saladino RA, Fromkin J, Heineman E, Suresh S, McGinn T. Development of an electronic medical record-based child physical abuse alert system. *J Am Med Inform Assoc*. 2018;25(2):142–149.
78. Rosenthal B, Skrbn J, Fromkin J, Heineman E, McGinn T, Richichi R, et al. Integration of physical abuse clinical decision support at 2 general emergency departments. *J Am Med Inform Assoc*. 2019;26(10):1020–1029.
79. Haarbauer-Krupa J, Haileyesus T, Gilchrist J, Mack KA, Law CS, Joseph A. Fall-related traumatic brain injury in children ages 0–4 years. *J Safety Res*. 2019;70:127–133.
80. Chisholm KM, Harruff RC. Elderly deaths due to ground-level falls. *Am J Forensic Med Pathol*. 2010;31(4):350–354.
81. Spaniolas K, Cheng JD, Gestring ML, Sangosanya A, Stassen NA, Bankey PE. Ground level falls are associated with significant mortality in elderly patients. *J Trauma*. 2010;69(4):821–825.
82. Newgard CD, Lin A, Yanez ND, Bulger E, Malveau S, Caughey A, et al. Long-term outcomes among injured older adults transported by emergency medical services. *Injury*. 2019;50(6):1175–1185.
83. Nishijima DK, Gaona SD, Waechter T, Maloney R, Bair T, Blitz A, et al. Out-of-hospital triage of older adults with head injury: a retrospective study of the effect of adding “anticoagulation or antiplatelet medication use” as a criterion. *Ann Emerg Med*. 2017.
84. Newgard CD, Lin A, Eckstrom E, Caughey A, Malveau S, Griffiths D, et al. Comorbidities, anticoagulants, and geriatric-specific physiology for the field triage of injured older adults. *J Trauma Acute Care*. 2019;86(5):829–837.
85. Nishijima DK, Gaona SD, Waechter T, Maloney R, Blitz A, Elms AR, et al. The incidence of traumatic intracranial hemorrhage in head-injured older adults transported by EMS with and without anticoagulant or antiplatelet use. *J Neurotrauma*. 2018;35(5):750–759.
86. Nance ML, Carr BG, Branas CC. Access to pediatric trauma care in the United States. *Arch Pediatr Adolesc Med*. 2009;163(6):512–518.
87. Utter GH, Maier RV, Rivara FP, Mock CN, Jurkovich GJ, Nathens AB. Inclusive trauma systems: do they improve triage or outcomes of the severely injured? *J Trauma*. 2006;60(3):529–535 discussion 35–37.
88. Branas CC, MacKenzie EJ, Williams JC, Schwab CW, Teter HM, Flanagan MC, et al. Access to trauma centers in the United States. *JAMA*. 2005;293(21):2626–2633.
89. Carr BG, Bowman AJ, Wolff CS, Mullen MT, Holena DN, Branas CC, et al. Disparities in access to trauma care in the United States: a population-based analysis. *Injury*. 2017;48(2):332–338.
90. *Pediatric Trauma Centers — Availability, Outcomes, and Federal Support Related to Pediatric Trauma Care*. Washington, DC: U.S. Government Accountability Office; 2017: Contract No.: GAO-17-334. Available at: <https://www.gao.gov/products/gao-17-334>. Accessed May 16, 2022.
91. Newgard CD, Fu R, Bulger E, Hedges JR, Mann NC, Wright DA, et al. Evaluation of rural vs urban trauma patients served by 9-1-1 emergency medical services. *JAMA Surg*. 2017;152(1):11–18.
92. Hannan EL, Farrell LS, Cooper A, Henry M, Simon B, Simon R. Physiologic trauma triage criteria in adult trauma patients: are they effective in saving lives by transporting patients to trauma centers? *J Am Coll Surg*. 2005;200(4):584–592.
93. Newgard CD, Meier EN, Bulger EM, Buick J, Sheehan K, Lin S, et al. Revisiting the “golden hour”: an evaluation of out-of-hospital time in shock and traumatic brain injury. *Ann Emerg Med*. 2015;66(1):30–41 e1–3.
94. Newgard CD, Braverman MA, Phuong J, Shipper ES, Price MA, Bixby PJ, et al. Developing a National Trauma Research Action Plan: Results from the prehospital and mass casualty research Delphi survey. *J Trauma Acute Care*. 2022;92(2):398–406.
95. Sasser SM, Ossmann E, Wald MM, Lerner EB, Hunt RC. Adoption of the 2006 field triage decision scheme for injured patients. *West J Emerg Med*. 2011;12(3):275–283.
96. Barnett AS, Wang NE, Sahni R, Hsia RY, Haukoos JS, Barton ED, et al. Variation in prehospital use and uptake of the national Field Triage Decision Scheme. *Prehosp Emerg Care*. 2013;17(2):135–148.
97. van Rein EAJ, van der Sluijs R, Raaijmakers AMR, Leenen LPH, van Heijl M. Compliance to prehospital trauma triage protocols worldwide: a systematic review. *Injury*. 2018;49(8):1373–1380.
98. Newgard CD, Fu R, Lerner EB, Daya M, Jui J, Wittwer L, et al. Role of guideline adherence in improving field triage. *Prehosp Emerg Care*. 2017;1–11.