UC Irvine

UC Irvine Previously Published Works

Title

Feasibility and acceptability of cognitive behavioral therapy for insomnia (CBT-I) or acupuncture for insomnia and related distress among cancer caregivers.

Permalink

<https://escholarship.org/uc/item/0094083k>

Journal

Palliative & supportive care, 18(6)

ISSN

1478-9515

Authors

Applebaum, Allison J
Buda, Kara
Hoyt, Michael A
et al.

Publication Date

2020-12-01

DOI

10.1017/s1478951520001017

Peer reviewed

Original Article

Cite this article: Applebaum AJ, Buda K, Hoyt MA, Shaffer K, Garland S, Mao JJ (2020). Feasibility and acceptability of cognitive behavioral therapy for insomnia (CBT-I) or acupuncture for insomnia and related distress among cancer caregivers. *Palliative and Supportive Care* **18**, 644–647. <https://doi.org/10.1017/S1478951520001017>

Received: 20 July 2020

Revised: 5 September 2020

Accepted: 10 September 2020

Key words:

Acupuncture; Cancer caregivers; Cognitive Behavioral Therapy for Insomnia (CBT-I); Insomnia

Author for correspondence:

Allison J. Applebaum, Assistant Attending Psychologist, Director, Caregivers Clinic, Department of Psychiatry and Behavioral Science, Memorial Sloan Kettering Cancer Center, 641 Lexington Avenue, 7th Floor, New York, NY 10022, USA.
E-mail: applebaa@mskcc.org

Feasibility and acceptability of cognitive behavioral therapy for insomnia (CBT-I) or acupuncture for insomnia and related distress among cancer caregivers

Allison J. Applebaum, PH.D.¹ , Kara Buda, M.A.¹, Michael A. Hoyt, PH.D.², Kelly Shaffer, PH.D.³, Sheila Garland, PH.D.⁴ and Jun J. Mao, M.D.⁵

¹Department of Psychiatry and Behavioral Sciences, Memorial Sloan Kettering Cancer Center, New York, NY;

²Department of Population Health and Disease Prevention and the Chao Family Comprehensive Cancer Center, University of California, Irvine, CA; ³Center for Behavioral Health and Technology, University of Virginia School of Medicine, Charlottesville, VA; ⁴Department of Psychology, Memorial University, St. John's, Newfoundland, Canada and ⁵Integrative Medicine Service, Memorial Sloan Kettering Cancer Center, New York, NY

Abstract

Objective. Insomnia is a common, distressing, and impairing psychological outcome experienced by informal caregivers (ICs) of patients with cancer. Cognitive behavioral therapy for insomnia (CBT-I) and acupuncture both have known benefits for patients with cancer, but such benefits have yet to be evaluated among ICs. The purpose of the present study was to evaluate the feasibility, acceptability and preliminary effects of CBT-I and acupuncture among ICs with moderate or greater levels of insomnia.

Method. Participants were randomized to eight sessions of CBT-I or ten sessions of acupuncture.

Results. Results highlighted challenges of identifying interested and eligible ICs and the impact of perception of intervention on retention and likely ultimately outcome.

Significance of the results. Findings suggest preliminary support for non-pharmacological interventions to treat insomnia in ICs and emphasize the importance of matching treatment modality to the preferences and needs of ICs.

Introduction

Insomnia is a common, distressing, and impairing outcome informal cancer caregivers (ICs) (Carter, 2003). Despite evidence supporting behavioral and integrative interventions for insomnia in psycho-oncology (Garland et al., 2019), none have been tested among cancer ICs. While cognitive behavioral therapy for insomnia (CBT-I) is recommended as a first line treatment (Rossman, 2019), it is not clear whether it will be acceptable and effective for ICs given the challenges of adhering to sleep restriction and stimulus control in the context of caregiving. Similarly, there is evidence for the use of acupuncture for insomnia (Yin et al., 2017), but it is not clear whether acupuncture will be acceptable or effective for ICs.

There is a clear need to test effects of existing evidenced-based non-pharmacological therapeutic interventions for insomnia among cancer ICs. The purpose of this feasibility study was to determine whether a future randomized controlled trial can be done, should be done and if so, how it should be done (Eldridge et al., 2016). This involves evaluation of the acceptability of CBT-I and acupuncture for insomnia and related distress among cancer ICs, and considerations of measurement, recruitment, retention, and elements of future study design.

Methods

Participants

Participants were recruited from Memorial Sloan Kettering Cancer Center (MSK) over 15 months and were (i) ≥ 18 years of age; (ii) a self-reported IC to a patient with any site/stage of cancer; (iii) able to read English and provide informed consent; and (iv) scored ≥ 7 on the Insomnia Severity Index (Bastien et al., 2001) and met DSM-5 criteria for Insomnia Disorder. Participants were excluded if they reported: (i) presence of sleep disorder other than insomnia; (ii) alcohol or drug dependence that compromises comprehension of assessments; (iii) engaging in night shift work; or (iv) being a daily smoker.

Procedure

ICs were randomized to seven sessions of CBT-I or ten sessions of acupuncture. Questionnaires were completed at baseline (T1), after intervention completion (T2), and at 3 months (T3).

Interventions

CBT-I included sleep restriction, stimulus control, cognitive restructuring, relaxation training, and sleep hygiene (Rossman, 2019). CBT-I consisted of four 60-minute weekly sessions, followed by three sessions over four weeks. Our manual incorporated the caregiving context as a key contributor to the onset and maintenance of insomnia. Participants kept nightly sleep logs and records of caregiving-related activities interfering with sleep.

Acupuncture is a component of Traditional Chinese Medicine in which an acupuncturist inserts needles at specific locations on the body. Our manualized acupuncture protocol included standardized points commonly used to address sleep problems with additional points to treat comorbid symptoms like pain and anxiety if needed. Licensed acupuncturists inserted 8–16 Seirin needles (0.16–0.25 mm × 15–40 mm) and manipulated the needles to achieve De Qi (a sensation of aching and soreness) (Mao et al., 2007b). Participants received acupuncture twice weekly for 2 weeks, then weekly for six more weeks, for a total of ten treatments for eight weeks. The first acupuncture visit involved a detailed history and examination lasting 60 min, with each subsequent session lasting 30 min, for a total time of 330.

Measures

Questionnaire measures included the *Causal Attributions of my Insomnia Questionnaire* (CAM-I; Harvey et al., 2013) and the *Acupuncture and CBT-I Expectancy Scales* (AES; Mao et al., 2007a).

Subjective measures of insomnia included the *Insomnia Severity Index* (ISI; Bastien et al., 2001), the *Pittsburgh Sleep Quality Index* (PSQI; Buysse et al., 1989), the *Multidimensional Fatigue Inventory-Short Form* (MFSI-SF; Stein et al., 2004), and the *Hyperarousal Scale* (HAS; Pavlova et al., 2001).

Objective sleep and stress measures included: *Wrist Actigraphy* (Morgenthaler et al., 2006), and *Diurnal Cortisol*, which were measured over two days prior to T1 and after T2. Additional assessments included the *Hospital Anxiety and Depression Scale* (HADS; Snaith and Zigmond, 1986), and a *Qualitative Interview* at T1 and T2 that assessed perceptions of causes of insomnia and strategies used to improve sleep. The interview explored participants' experiences with the treatment.

Data analyses

Feasibility was informed by the number of eligible ICs, the percentage who participated, and completion rates. Acceptability was assessed as the percentage of participants completing ≥ 4 of 7 sessions of CBT-I and ≥ 8 of 10 sessions of acupuncture. Acceptability was also observed through interviews. Descriptive statistics and mean change scores examined person-level patterns of change. The study was approved by the MSK Institutional Review Board (protocol 17-001).

Results

Among 32 prospective participants, 41% ($N = 13$ ICs) enrolled and were randomized, 7 to CBT-I and 6 to acupuncture. Among non-enrollees, one was not eligible, 14 were not interested, and three were unreachable (Figure 1). All participants

were female and had a college degree. The majority ($N = 11$, 92%) were white, married/cohabitating ($N = 8$, 62%), and provided care since the patient's diagnosis ($N = 9$, 70%). Most lived with the patient ($N = 10$, 77%) and were his/her spouse/partner or child ($N = 10$, 77%). Participants were, on average, 51 years old (range 25–73).

Acceptability

All participants randomized to CBT-I completed T1, three completed T2 and two completed the T3 assessment. Two participants randomized to acupuncture dropped before T1 and one did not complete the acupuncture protocol; the remaining three completed all acupuncture sessions and assessments. Of the 11 participants completing T1, only six agreed to provide salivary cortisol and actigraph data. Reasons for attrition included bereavement, preference for CBT-I when randomized to acupuncture, and loss of interest in treatment.

Semi-structured interviews were completed by 11 participants at T1 and five at T2. At T1, four participants preferred CBT-I and five preferred acupuncture. Those preferring CBT-I desired skills they could use over the long-term (e.g., "I can't put the needles in me after treatment"). Those preferring acupuncture anticipated it would "provide relaxation" and expressed interest in a modality that "required very little effort." At T2, participants described CBT-I as "demanding and emotionally challenging" but ultimately "incredibly effective" in reducing insomnia. Sleep restriction was identified as a powerful but challenging technique, and the accountability built into sessions particularly helpful. Regarding acupuncture, participants reported satisfaction with the limited effort required, and two believed that it helped with non-sleep related symptoms.

Perceived causes of insomnia and expectancy of treatment

The most frequently endorsed category of causes of insomnia was *thinking patterns* (e.g., *can't shut off thoughts*), endorsed by 11 participants, followed by *sleep-related thoughts* (e.g., *thinking about falling asleep*; endorsed by five participants) and *sleep-related emotions* (*stress, anxiety*; endorsed by four participants). Participants endorsed relatively similar expectation of acupuncture and CBT-I (13.75 versus 12.43) in addressing insomnia and their ability to cope with related distress.

Impact of CBT-I and acupuncture on insomnia

At T2, CBT-I participants had lower mean scores for insomnia severity (CBT-I 11.67–6.33, acupuncture 10.33–9.00, $d = 0.59$), hyperarousal (CBT-I 42–35, acupuncture 53.0–47.0, $d = 1.39$), anxiety (CBT-I 11.83–7.67, acupuncture 15.60–15.00, $d = 1.37$), depression (CBT-I 7.14–6.00, acupuncture 10.25–10.00, $d = -1.57$), and fatigue (CBT-I 15.00–6.67, acupuncture 19.00–11.67, $d = -0.96$) than acupuncture participants.

When given the option only six participants (three in each arm) provided salivary cortisol data at T1 and T2 and six participants (four randomized to CBT-I and two to acupuncture) completed data collection via wrist actigraphy.

Discussion

Results provide only modest evidence of feasibility of both non-pharmacological modalities for the treatment of insomnia

Fig. 1. Trend data.

among ICs and highlight key challenges in recruitment and feasibility of objective assessments. Only 32 potential participants were identified over 15 months despite multipronged recruitment (e.g., direct referral by treating physician, recruitment in-person at clinics, fliers). Though study staffing was limited, this may reflect ICs'

need for treatment that addresses multiple facets of quality-of-life (e.g., three potential participants reported desire for psychotherapy and did not want to risk randomization to acupuncture). Framing insomnia as an element of IC burden that is related to other facets of functioning may have made the trial more appealing.

The most frequent reason for non-enrollment was lack of interest; only one IC did not meet insomnia symptom threshold criteria. Both CBT-I and acupuncture are time-intensive and require a commitment from ICs that may not be possible during intense periods of caregiving. Future studies are needed to identify optimal timing of recruitment to accommodate more ICs who are willing and capable of participation.

Treatment preference emerged as an important indicator of potential impact of these approaches; interviews highlighted preferences for CBT-I among participants desiring skills they could independently use post-treatment, and preference for acupuncture among those preferencing minimal effort. Data also underscored the acceptability of these modalities, with CBT-I participants characterizing the intervention as demanding but incredibly effective, and those to acupuncture reporting great satisfaction with the limited effort required to participate with perception of benefit beyond insomnia management. At baseline, most participants attributed their insomnia to thinking patterns, which highlights the appropriateness of CBT-I in this particular population.

Th challenges of collecting objective measures of insomnia and related distress (i.e., salivary cortisol and actigraphy) are evident. Adherence to these protocols may be difficult and future studies will need to consider alternate strategies that provide similar data with more limited effort.

Conclusions and future directions

Our study highlighted the challenges of recruitment and retention of ICs. Data provide preliminary evidence for CBT-I and acupuncture to positively impact insomnia and related distress, though suggest these approaches may fit unique IC needs. Future studies should consider matching treatment modality to IC preference and stated needs and offer telehealth-delivered CBT-I to ameliorate the burden associated with participation in-person.

Funding sources

This research was supported by funding from the National Institutes of Health (T32CA009461, PI: Jaime Ostroff, PhD; P30CA008748, PI: Craig Thompson, MD).

Data sharing

The data that support the findings of this study are available on request from the corresponding author; the data are not publicly available due to privacy or ethical restrictions.

Conflicts of interest

The authors have no conflicts of interest to report.

References

- Bastien CH, Vallieres A and Morin CM (2001) Validation of the insomnia severity index as an outcome measure for insomnia research. *Sleep Medicine* 2(4), 297–307.
- Buysse DJ, Reynolds 3rd CF, Monk TH, *et al.* (1989) The Pittsburgh Sleep Quality Index: A new instrument for psychiatric practice and research. *Psychiatry Research* 28(2), 193–213.
- Carter PA (2003) Family caregivers' sleep loss and depression over time. *Cancer Nursing* 26(4), 253–259.
- Eldridge SM, Chan CL, Campbell MJ, *et al.* (2016) CONSORT 2010 statement: Extension to randomised pilot and feasibility trials. *Pilot and Feasibility Studies* 2, 64.
- Garland SN, Xie SX, DuHamel K, *et al.* (2019) Acupuncture versus cognitive behavioral therapy for insomnia in cancer survivors: A randomized clinical trial. *JNCI: Journal of the National Cancer Institute* 111(12), 1323–1331.
- Harvey AG, Soehner A, Lombrozo T, *et al.* (2013) 'Folk theories' about the causes of insomnia. *Cognitive Therapy and Research* 37(5). doi: 10.1007/s10608-013-9543-2.
- Mao JJ, Armstrong K, Farrar JT, *et al.* (2007a) Acupuncture expectancy scale: Development and preliminary validation in China. *Explore (NY)* 3(4), 372–377.
- Mao JJ, Farrar JT, Armstrong K, *et al.* (2007b) De qi: Chinese acupuncture patients' experiences and beliefs regarding acupuncture needling sensation - An exploratory survey. *Acupuncture in Medicine* 25(4), 158–165.
- Morgenthaler T, Kramer M, Alessi C, *et al.* (2006) Practice parameters for the psychological and behavioral treatment of insomnia: An update. An American Academy of Sleep Medicine report. *Sleep* 29(11), 1415–1419.
- Pavlova M, Berg O, Gleason R, *et al.* (2001) Self-reported hyperarousal traits among insomnia patients. *Journal of Psychosomatic Research* 51(2), 435–441.
- Rossman J (2019) Cognitive-behavioral therapy for Insomnia: An effective and underutilized treatment for Insomnia. *American Journal of Lifestyle Medicine* 13(6), 544–547.
- Snaith RP and Zigmond AS (1986) The hospital anxiety and depression scale. *British Medical Journal (Clinical Research Edition)* 292(6516), 344.
- Stein KD, Jacobsen PB, Blanchard CM, *et al.* (2004) Further validation of the multidimensional fatigue symptom inventory-short form. *Journal of Pain and Symptom Management* 27(1), 14–23.
- Yin X, Gou M, Xu J, *et al.* (2017) Efficacy and safety of acupuncture treatment on primary insomnia: A randomized controlled trial. *Sleep Medicine* 37, 193–200.