

UCSF

UC San Francisco Previously Published Works

Title

Toddlers at Elevated Likelihood for Autism: Exploring Sensory and Language Treatment Predictors.

Permalink

<https://escholarship.org/uc/item/00f2n4tz>

Journal

Journal of Early Intervention, 45(1)

ISSN

1053-8151

Authors

Jatkar, Alapika
Garrido, Dunia
Silverman, Greyson
[et al.](#)

Publication Date

2023-03-01

DOI

10.1177/10538151211067227

Peer reviewed



Published in final edited form as:

J Early Interv. 2023 March ; 45(1): 39–62. doi:10.1177/10538151211067227.

Toddlers at Elevated Likelihood for Autism: Exploring Sensory and Language Treatment Predictors

Alapika Jatkar¹, Dunia Garrido², Shuting Zheng³, Greyson Silverman⁴, Heba Elsayed^{4,5}, Paige Huguely Davis¹, Helen Lee⁶, Elizabeth R. Crais⁴, John Sideris⁶, Lauren Turner-Brown⁷, Grace T. Baranek⁶, Linda R. Watson⁴, Rebecca Grzadzinski¹

¹The University of North Carolina at Chapel Hill, Carolina Institute for Developmental Disabilities, Chapel Hill, NC, USA

²The University of Granada, Mind, Brain, and Behaviors Research Center, Granada, Spain

³The University of California at San Francisco, San Francisco, CA, USA

⁴The University of North Carolina at Chapel Hill, Division of Speech and Hearing Sciences, Chapel Hill, NC, USA

⁵Alexandria University, Phoniatics Unit, Otolaryngology Department, Egypt

⁶The University of Southern California, Mrs. T.H. Chan Division of Occupational Science and Occupational Therapy, Los Angeles, CA, USA

⁷The University of North Carolina at Chapel Hill, Department of Psychiatry, Chapel Hill, NC, USA

Abstract

Baseline child characteristics may predict treatment outcomes in children with or at elevated likelihood of developing autism (EL-ASD). Little is known about the role of child sensory and language features on treatment outcome. Participants were randomly assigned to a parent-mediated intervention or control condition. Analyses explored the relationship between baseline child sensory and language characteristics and changes in ASD symptoms over approximately 9 months. Higher baseline sensory hyporeactivity was significantly related to less improvement in social communication (SC) for the treatment group only. More baseline atypical vocalizations were significantly related to less improvement on SC across treatment and control groups. This work provides an initial framework to encourage the tailoring of interventions for EL-ASD children, suggesting sensory reactivity and atypical vocalizations may be useful behaviors to consider in treatment planning.

Article reuse guidelines: sagepub.com/journals-permissions

Corresponding Authors: Alapika Jatkar, Carolina Institute for Developmental Disabilities, The University of North Carolina at Chapel Hill, 101 Renee Lynne Ct, NC 27599-6305, USA. alapikajatkar@gmail.com, Rebecca Grzadzinski, Carolina Institute for Developmental Disabilities, The University of North Carolina at Chapel Hill, 101 Renee Lynne Ct, NC, 27599-6305, USA. rebecca_grzadzinski@med.unc.edu.

Heba Elsayed, Paige Huguely Davis, Elizabeth R. Crais, John Sideris, Grace T. Baranek, Linda R. Watson, Rebecca Grzadzinski, these authors were now affiliated to The University of North Carolina at Chapel Hill, Program for Early Autism Research Leadership and Service (PEARLS), Chapel Hill, NC, USA.

Declaration of Conflicting Interests

The author(s) declared no potential conflicts of interest with respect to the research, authorship, and/or publication of this article.

Keywords

autism spectrum disorders; disabilities and development delays; language and communication; child development; infants and toddlers; young children

Background

Autism spectrum disorder (referred to throughout as autism) is a neurodevelopmental difference that is characterized by delays or differences in developing social communication (SC) skills and the presence of restricted and repetitive behaviors (RRBs). Symptoms of autism usually consolidate into a diagnosis when a child is between 2 and 3 years of age; however, the median age of diagnosis is older than 4 years in the United States (Shaw et al., 2020). Significant research efforts have identified early screening and symptom presentations that may precede a formal diagnosis.

As research in early identification improves accuracy of a later diagnosis, recent studies have run parallel to explore child characteristics that may predict later child outcomes. This is in an effort to better understand child trajectories and improve tailored treatments for children with, or at elevated likelihood for neurodevelopmental difference such as autism (EL-ASD; Bearss et al., 2015; Garrido et al., 2017; Pellecchia et al., 2016; Schreibman et al., 2009; Siller et al., 2013; Tamis-Lemonda et al., 2001; Warren et al., 2010). EL-ASD children, identified due to having a sibling with autism or through early screening, can begin early intervention (EI) services by 18 months of age (or earlier), if sufficient symptomology of autism or other developmental delay can be identified. EI studies have focused on the efficacy of applied behavior analysis (ABA)-based treatment strategies, including naturalistic developmental behavioral interventions (NDBI; Landa, 2018). These studies have found modest effects on improving functional outcomes of EL-ASD children (Vivanti et al., 2018). NDBIs are correlated with increased adaptive behavior and intelligence quotient (IQ) in children with autism (Vivanti & Rogers, 2014). EI-related improvements have been related to treatment intensity in some studies, with treatments of more than 30 hours a week associated with maximal gains (Eldevik et al., 2009), although this needs further exploration (Crank et al., 2021). Research has suggested that NDBIs involving caregiver and clinician components may be most effective (Hampton & Kaiser, 2016), although the relationship between provider (caregiver vs. clinician) and treatment outcome is unclear (Crank et al., 2021; Rogers et al., 2020). High-intensity interventions for young children with autism (up to 40 hr per week) and high caregiver and clinician involvement are both expensive and time-consuming (Leigh & Du, 2015; Ouyang et al., 2014; Pellicano et al., 2014). This can lead to substantial family and parental stress (Estes et al., 2019), as well as financial burden. Given the cost and burden of these interventions, identifying mechanisms of improvement or factors that contribute to better functional outcomes for children is a top priority to tailor EIs.

Certain child characteristics, such as high interest in toy contact, were related to better language outcomes in response to treatment (Schreibman et al., 2009; Yoder & Stone, 2006). Research has also suggested that higher baseline cognitive ability is a predictor of more

improvement over the course of intervention (Grzadzinski et al., under review; Harris & Handleman, 2000), though more studies are needed, as others have found that a lower baseline IQ is correlated with greater increases in cognitive scores (Ben-Itzhak et al., 2014) and some found no IQ effects (Sutera et al., 2007). In addition, nuanced exploration of child characteristics may reveal new information regarding mechanisms of treatment outcome, which could ultimately be incorporated into early intervention practices.

Though understudied, research has suggested that 70% to 90% of children diagnosed with autism display high levels of sensory features, and the *Diagnostic and Statistical Manual of Mental Disorders, 5th Edition (DSM-5)* now recognizes some sensory features under RRBs (Baranek et al., 2006; Ben-Sasson et al. 2009; López & Leekam, 2007). Sensory features are noted across modalities (e.g., auditory, tactile) and are often characterized into three different sensory response categories that regularly coexist. Sensory hyporeactivity is defined as either a decreased or absent reaction to sensory stimuli, hyperreactivity is an aversive or avoidant reaction to sensory stimuli, and sensory seeking behaviors extend, intensify, or modulate sensory experiences (Baranek et al., 2006; Ben-Sasson et al., 2008, 2009). Hyporeactivity has consistently been reported in young children with autism as well as in EL-ASD children (Baranek et al., 2006, 2013; Ben-Sasson et al. 2009). Hyperreactivity and sensory seeking behaviors are also documented (Baranek et al. 2019; Ben-Sasson et al., 2008, 2013; Brock et al., 2012; Wolff et al., 2019). Atypical sensory reactivity has been found in EL-ASD infants and toddlers in several studies, suggesting it as a potential precursor to an autism diagnosis. One infant sibling study found that at 12 months, children later diagnosed with autism showed elevated hyperreactivity on a parent report measure compared with children that did not go on to develop autism (Wolff et al., 2019). In another study, parents reported sensory differences among infant siblings, some as young as 6 months of age, in children who later met criteria for an autism diagnosis (Sacrey et al., 2015). Similarly, sensory reactivity reported by parents at 12 months was found to be a predictor of a later autism diagnosis in a community sample (Turner-Brown et al., 2013). Research has shown that the patterns of sensory reactivity also matter in child development of other skills. For instance, children with hyporeactivity presented with more communication deficits, including delays in spoken language (Baranek et al., 2013; Flippin & Watson, 2011; Patten et al., 2013; Philpott-Robinson et al., 2016). Although there is growing research that supports the relationship between early sensory reactivity and later autism diagnosis (Bizzell et al., 2020; Grzadzinski et al., 2020; Sacrey et al., 2014; Wolff et al., 2019), research has not explored if sensory reactivity patterns impact treatment outcomes.

EL-ASD toddlers who go on to be diagnosed with autism also show unique patterns of communication and language skills compared with typically developing peers. For example, between 18 and 24 months of age, children who go on to receive a diagnosis of autism have been found to have a lower proportion of speech-like vocalizations (Plumb & Wetherby, 2013). EL-ASD infants, identified due to ASD diagnosis in an older sibling, produce fewer total speech-like vocalizations at 6, 9, and 12 months (Paul et al., 2011), while typically developing infants were 17 times more likely to use canonical babbling (vocalizations that include at least one well-formed syllable, with each having at least one full vowel-like element and at least one consonant-like element with rapid transition between the two) at

9 to 12 months and 6 times more likely at 15 to 18 months (Patten et al., 2014). Parents of children with autism have indicated that their child's use of speech-like vocalizations is significantly different from reports of parents of typically developing children at 12 months of age, but did not consistently differ from those with other developmental disabilities (Watson et al., 2007). Global language skills, such as Verbal IQ, have been found to be a predictor of treatment outcomes (Grzadzinski et al., under review), but specific vocalization components, such as directedness of speech, frequency of canonical babbling, or atypical vocalizations, may be related to treatment outcomes as well. Further exploration of the impact of vocalization patterns on treatment outcomes in children at an elevated likelihood for or diagnosed with autism could provide new targets for treatment and tailor interventions.

Although the impacts of early sensory reactivity and vocalization patterns on a later autism diagnosis have been studied, their impact on outcomes is unknown. Thus, the aim of this study is to examine whether sensory reactivity and vocalization patterns impact changes in SC and RRB. We hypothesize that aspects of sensory reactivity (hyporeactivity) and vocalization patterns will predict SC and RRB behaviors about 9 months later, regardless of treatment status. By using multiple measures of sensory reactivity (parent report and direct observation), nuanced examinations of vocalization patterns, and a new treatment outcome measure (the Brief Observation of Social Communication Change; BOSCC; Grzadzinski et al., 2016), the results of this study will maximize the potential of data collected through a completed randomized control trial (Watson et al., 2017) to provide the field with novel intervention targets for EL-ASD toddlers.

Method

Participants

This study included a sample of 87 EL-ASD toddlers (60 boys) identified at 12 months (± 2 weeks) based on community screenings with the *First Year Inventory version 2.0* (FYI; Baranek et al., 2003). Participants were recruited by mailing the FYI to parents of children in the study catchment area, based on state birth records, just prior to the child's first birthday. The children were randomized to a parent-mediated early intervention (Watson et al., 2017; Adapted Responsive Teaching; ART; $n = 45$) or Referral to Early Intervention and Monitoring (REIM; $n = 42$); referred to as Group in analyses. Participants were seen at baseline (prior to treatment onset) and ~9 months later, post-treatment. Time 1 (14 months, ± 0.77 months) and Time 2 (23 months, ± 0.86), respectively. Additional eligibility criteria included birthweight $> 2,500$ g and English spoken as a primary language at home. All participants' parents provided their written consent before evaluation began, as in accordance with protocols approved by the Institutional Review Board (see Table 1).

Measures

Cognitive Abilities

Mullen Scales of Early Learning (MSEL).—At Time 1 and Time 2, children were administered the MSEL (see Table 1; Mullen, 1995). The MSEL yields standard *T*-scores

in the domains of fine motor, visual reception (VR), receptive language, and expressive language (EL) at Time 1. VR and EL, as proxies of nonverbal and verbal developmental level, respectively, were covariates in analyses.

Sensory Reactivity

Sensory Processing Assessment (SPA).—This behavioral measure assesses a child's sensory reactivity to a variety of stimuli in a 15-minute play-based observation designed for children aged 6 months to 9 years of age (Baranek, 1999; Baranek et al., 2007, 2013). Results yield scores in hyporeactivity (a decreased orienting response to visual, tactile, and auditory stimuli; SPA-Hypo), hyperactivity (an aversive or avoidant response to visual, tactile, and auditory stimuli; SPA-Hyper), and sensory seeking behaviors (an intense craving, repetition or fascination of sensory experiences with objects or body; SPA-Seek). The SPA was administered at Time 1 (14 months, ± 0.77 months), and administrators were blind to group status. Of the 87 children, three were lost to follow-up at Time 2, and for one additional child, only parent report assessments were available at Time 2.

Sensory Experiences Questionnaire (SEQ).—This questionnaire consists of 43 items appropriate for children 5 months to 12 years of age, and assesses the type and frequency of a child's sensory reactions to everyday stimuli (Baranek, 1999; Baranek et al., 2006, 2013; Little et al., 2011). Responses are rated on a 5-point Likert-type scale ranging from 1 (*almost never*) to 5 (*almost always*); higher scores indicate more sensory symptoms. The SEQ, a companion to the SPA, yields scores on hyporeactivity, hyperreactivity, and sensory seeking behaviors and has high test–retest reliability (Little et al., 2011) and internal consistency (Baranek et al., 2006). At Time 1, parents completed the SEQ about their child.

Vocalizations

At Time 1, child vocalizations were coded as described in Garrido et al. (2017). Vocalizations were coded from 30 min of video observation of the child in naturalistic and semi-naturalistic social/play contexts. The context of the first 10 min was a parent–child free-play with a standard set of developmentally appropriate toys. Video recordings of the administration of the Communication and Symbolic Behavior Scales Developmental Profile (CSBS-DP; Wetherby & Prizant, 2002) were used to code child vocalizations for an additional 20 min. With vocalization data missing for one child at Time 1, a minimum of 75 children were available for all analyses.

As described in Garrido et al. (2017), two types of speech vocalizations (canonical and non-canonical) and three types of nonspeech vocalizations (atypical, distress, and pleasure) were coded; see Figure 1. Vocalizations were coded as discrete events with an onset occurring when the child initiated a sound and an ending if there was at least 1 second break in the child's vocalization. All vocalizations were coded as either directed to a communication partner or non-directed (see Garrido et al., 2017 for coding criteria). Frequencies of each vocalization were summed. Three coders who were unaware of child information were trained on non-study videos to identify the types of vocalizations described in Figure 1 and attained at least 90% event-by-event agreement with the established training video codes before coding study videos. To assess interrater reliability, 20% of the study videos were

randomly selected to be independently coded by two coders, both of whom were unaware of which videos would be double-coded. Given the time-consuming and therefore costly process required for coding, double coding 20% of the study was deemed feasible within the resources available for the study as well as sufficient to evaluate reliability of the coded data. Reliability was estimated from intraclass correlation coefficients (ICCs), using SPSS statistics version 22.0, using a two-way random effects model for absolute agreement. Interrater agreement was excellent across vocalization type, with intraclass correlation coefficients ranging from 0.91 to 0.995 (see Garrido et al. (2017) for additional details). One participant was missing language coding, and was therefore removed from analyses.

Autism Symptoms

The BOSCC (Grzadzinski et al., 2016) is a treatment response measure that quantifies changes in autism symptoms. The BOSCC coding scheme was applied to parent–child free-play videos at Time 1 (14 months, ± 0.77) and Time 2 (23 months, ± 0.86) to specifically examine changes in child autism behaviors in the SC (BOSCC-SC) and RRB (BOSCC-RRB) domains. The 10-min parent–child free-play video was the same as was coded for the child vocalizations described above (though the addition of the first 20 min of the CSBS was specific to vocalization coding). Prior to coding independently, coders participated in training that involved review of the BOSCC coding scheme, practice watching and coding video observations, and participation in discussions with already reliable coders. New coders obtained interrater agreement standards that the authors deemed adequate: no more than three items with more than one point disagreement AND within four points across summed totals for all items, across three consecutive videos (Grzadzinski et al., 2016). Codes from coders that had not met the above interrater agreement standards were never included in data sets. Regular consensus meetings were held to confirm ongoing reliability on the BOSCC across coders. During the coding process, to maintain reliability standards, a primary coder was randomly chosen and used for data analysis, while a secondary coder was randomly chosen and used for reliability analysis. This method was chosen, rather than a fully crossed model, due to the time-consuming nature of coding and the availability of coders (Grzadzinski et al., 2016). Coders were blind to group and timepoint.

Twenty-seven videos, representing 15% of the coded samples, were randomly chosen for independent coding by two coders to assess inter-rater reliability (IRR). The coders were blind to which videos were being double-coded. The IRR for the SC and RRB domain was high (intra-class correlation coefficient = 0.85, 95% confidence interval [CI] = [0.69, 0.93] and 0.88, 95% CI = [0.75, 0.94]), respectively. Change scores in BOSCC-SC and BOSCC-RRB domains were calculated by obtaining ratio scores to take into account Time 1 values. The use of change scores was chosen due to the goals of the BOSCC as a metric used specifically for quantifying change over time. T1 BOSCC scores were used as a covariate in analyses.

Correlations.—See Table 2 for correlation analyses.

Groups.—Forty-five children were randomly assigned to the ART group, an adaptation of the Responsive Teaching Curriculum (Mahoney & Macdonald, 2007) built on the idea

that pivotal behaviors play key roles in positive outcomes in infants both EL-ASD and with ASD. Forty-two children were randomly assigned to the REIM group, where the children and their families received no direct intervention services, but did receive contact from the project coordinator every 5 to 6 weeks. In the event that a parent indicated a need for help in accessing EI services, the project coordinator provided contact information and offered to initiate a referral (Watson et al., 2017).

Analyses.—Multicollinearity tests revealed no multicollinearity between variables ($1 < \text{variance inflation factor [VIF]} < 10$). Moderation analyses (controlling for Time 1 MSEL-EL and MSEL-VR) were conducted with 10,000 bootstraps to assess whether the child characteristics (e.g., vocalization, sensory) significantly moderated the relationship between group (ART vs. REIM; binary dummy coded) and ratio changes in child SC (BOSCC-SC-ratio) and RRBs (BOSCC-RRB-ratio). In the absence of significant moderation effects, linear regressions were conducted to test main effects, without covariates (Model 1) and with covariates (Model 2; Time 1 verbal MSEL-EL and nonverbal MSEL-VR). An alpha level of .05 was used to determine significance. All analyses were conducted using IBM SPSS Statistics, and the moderation effect was tested using PROCESS v3.5 for SPSS (Hayes, 2017).

Results

Social Communication

See Table 3.

Sensory hyporeactivity.—A moderation analysis revealed that there was a statistically significant interaction between SPA hyporeactivity and Group ($p = .02$) such that higher SPA Hyporeactivity at Time 1 was associated with fewer improvements in SC within the treatment group only. See Figure 2. Using the SEQ, moderation analyses testing the interaction between Group and Time 1 SEQ hyporeactivity were not statistically significant; subsequent main effect regression analyses also revealed that SEQ Hyporeactivity at Time 1 was not a significant predictor of BOSCC-SC-ratio.

Sensory hyperreactivity.—A moderation analysis revealed that there was not a statistically significant interaction between SPA hyperreactivity and Group. Subsequent main effect regression models indicated that Time 1 SPA hyperreactivity was a significant predictor of BOSCC-SC-ratio— $\beta = 0.136, t = 2.17, p = .033$). However, when controlling for baseline VR and EL, SPA hyperreactivity fell just below the statistical threshold of significance— $\beta = 0.119, t(79) = 1.917, p = .06$. Moderation analyses revealed that Time 1 SEQ hyperreactivity did not significantly moderate the relationship between Group and SC outcome; in addition, subsequent main effect regression models indicated that SEQ hyperreactivity at Time 1 was not a significant predictor of SC outcome.

Sensory seeking.—Moderation analyses revealed that SPA and SEQ seeking at Time 1 were not significantly moderating the relationship between Group and SC outcome. Furthermore, main effect regression models indicated that Time 1 SPA Seeking and Time 1 SEQ Seeking were not significant predictors of BOSCC-SC-ratio.

Vocalizations.—Moderation and subsequent main effect models analyses indicated that none of the vocalization variables, with the exception of atypical vocalizations, were significant moderators or main effect variables on BOSCC-SC-ratio. A moderation analysis revealed that there was not a statistically significant interaction between atypical vocalizations and treatment group ($p = .408$), suggesting that atypical vocalizations did not differentially impact BOSCC-SC-ratio based on treatment group. Main effect regression analyses indicated that atypical vocalizations significantly predicted BOSCC-SC-ratio, $\beta = 0.007$, $t(81) = 3.486$, $p = .001$, even when controlling for baseline VR and EL, $\beta = 0.008$, $t(81) = 3.769$, $p < .0001$; see Figure 3.

Restricted and Repetitive Behaviors

See Table 4.

Sensory hyporeactivity.—Moderation analyses indicated that there was not a significant interaction between SPA hyporeactivity and Group on BOSCC-RRB-ratio; subsequent main effect regression analyses revealed that SPA hyporeactivity at Time 1 was not a significant predictor of BOSCC-RRB-ratio. Moderation analyses indicated that there was not a significant interaction between SEQ hyporeactivity and Group on BOSCC-RRB-ratio; subsequent main effect regression analyses revealed that SEQ hyporeactivity at Time 1 was not a significant predictor of BOSCC-RRB-ratio.

Sensory hyperreactivity.—Moderation analyses indicated that there was not a significant interaction between SPA hyperreactivity and Group on BOSCC-RRB-ratio; subsequent main effect regression analyses revealed that SPA hyperreactivity at Time 1 was not a significant predictor of BOSCC-RRB-ratio. Moderation analyses indicated that there was not a significant interaction between SEQ hyperreactivity and Group on BOSCC-RRB-ratio; subsequent main effect regression analyses revealed that SEQ hyperreactivity at Time 1 was not a significant predictor of BOSCC-RRB-ratio.

Sensory seeking.—Moderation analyses indicated that there was not a significant interaction between SPA seeking and Group on BOSCC-RRB-ratio; subsequent main effect regression analyses revealed that SPA seeking at Time 1 was not a significant predictor of BOSCC-RRB-ratio. Moderation analyses indicated that there was not a significant interaction between SEQ seeking and Group on BOSCC-RRB-ratio; subsequent main effect regression analyses revealed that SEQ seeking at Time 1 was not a significant predictor of BOSCC-RRB-ratio.

Vocalizations.—Moderation and subsequent main effect models analyses indicated that none of the vocalization variables, with the exception of atypical vocalizations, were significant moderators or main effect variables on BOSCC-RRB-ratio. Moderation analyses indicated that atypical vocalization was not a significant moderator of the relationship between group and BOSCC-RRB-ratio. Subsequent main effect regression analyses indicated that Time 1 atypical vocalizations were a significant predictor of BOSCC-RRB-ratio ($\beta = -0.007$, $t = -2.087$, $p = .04$). When controlling for VR and EL, atypical

vocalizations fell just below statistical significance ($\beta = -0.006$, $t = -19.981$, $p = .052$); see Figure 4.

Discussion

Research is limited on child characteristics that impact treatment outcomes. This exploratory study evaluated how child sensory and vocalization patterns predict response to treatment in EL-ASD toddlers. Our results suggest that baseline sensory reactivity and the number of atypical vocalizations predict how much improvement a child will make in social communication skills over the course of 9 months. Child characteristics before the start of treatment, including sensory reactivity and vocalization patterns, may provide early indicators for a child's trajectory in autism symptoms as well as novel treatment targets that can be tailored to a child's baseline presentation.

Research indicates that children with autism display atypical sensory response patterns (Baranek et al., 2006; Ben-Sasson et al. 2009) and these patterns may be present in infancy before a definitive autism diagnosis is currently possible (Sacrey et al., 2015; Turner-Brown et al., 2013; Wolff et al., 2019). This study found that SPA hyporeactivity (e.g., reduced reactivity to sensory stimuli when it is expected) was a significant predictor of BOSCC-SC ratio in the treatment group, and this relationship held true when controlling for baseline VR and EL. This suggests that intervention outcomes for some infants at EL-ASD might be improved by focusing first on addressing sensory reactivity, specifically behavioral orienting to sensory stimuli, before introducing new social communication skills, in at least a subset of children at EL-ASD. Such interventions might include: teaching parents to identify their children's sensory preferences and reactivity patterns, assist their child in tuning to the appropriate sensory stimuli, and ensure that sensory stimuli are optimally salient and meaningful in daily routines. These suggestions may have implications for tailoring the child's sensory environment to encourage optimal levels of engagement as appropriate to the situation, or for tailoring intervention strategies to specifically address sensory hyporeactivity to improve treatment outcome.

Similar results were found when analyzing SPA hyperreactivity results; however, when controlling for baseline VR and EL, SPA hyperreactivity was no longer a significant predictor of BOSCC-SC-ratio (alpha value was just under significance). This may suggest that verbal and nonverbal skills are important to understanding the progress of a child during treatment and/or there may be a relationship between verbal and nonverbal skills and sensory hyperreactivity that impacts later outcome. Alternatively, our limited sample size may limit the power of our results, particularly since the results fell just above our significance threshold. Future research should continue to explore the relationship between sensory hyperreactivity and child SC outcomes, particularly in larger samples and while taking into account verbal and nonverbal skills.

Research has shown that increased atypical vocalizations are associated with diagnostic status, above and beyond baseline verbal and nonverbal skills. The results of this exploratory study suggest that Time 1 atypical vocalizations predicted BOSCC-SC change for children, regardless of group assignment. Other components of vocalizations including canonical

(Oller et al., 1999; Paul et al., 2011) and noncanonical (Oller et al., 1994), directed or nondirected, were not significant predictors of changes in social communication symptom change. Although this finding suggests that atypical vocalizations warrant consideration in early intervention for EL-ASD infants and toddlers, we are not aware of any existing intervention research that addresses strategies for, or outcomes of, including these as early intervention targets. Research on the functions of these vocalizations for the children and strategies that adults can use to respond in supportive ways to atypical vocalizations may be useful in further clarifying the clinical implications of this finding.

Early intervention trials of toddlers with ASD diagnoses have begun to examine baseline child and parent characteristics that impact changes in response to treatment (Beglinger & Smith, 2005; Carter et al., 2011; Kasari et al., 2010; Oosterling et al., 2010; Rogers et al., 2012; Zachor et al., 2007) (joint attention; Olafsen et al., 2006; Watson et al., 2013). For example, child object interest (Fossum et al., 2018) and IQ (Grzadzinski et al., under review; Harris & Handleman, 2000) at baseline have been linked with an amount of improvement over the course of an intervention. These results add to this growing literature suggesting that in a sample of children with early increased likelihood of later ASD behaviors that may present presymptomatically, such as sensory hyporeactivity and atypical vocalizations, could be crucial to tailoring early intervention practices such that future treatments are more effective. Identifying these prediagnostic markers provides targets for a potential very early (before 12 months) presymptomatic intervention for EL-ASD children. Future research should continue to evaluate child and family characteristics that may be present in infancy and could be targeted in a presymptomatic intervention to promote optimal developmental trajectories.

As these results suggest that behaviors such as sensory hyporeactivity and atypical vocalizations may be present presymptomatically (or at least prior to symptom consolidation into a diagnosable condition), further exploration of presymptomatic treatment and intervention practices is necessary. Currently, clinical practices do not focus on presymptomatic intervention for autism. Implementing presymptomatic intervention that is specifically tailored to certain features, such as sensory hyporeactivity and atypical vocalizations, could be crucial to promoting better adaptive outcomes in infants and toddlers who are particularly likely to make slow progress in social communication and language skills. Presymptomatic intervention could improve future functional behavior, and lessen the severity of impairments.

The current study has several limitations, and the results should be considered in light of these. After careful consideration, given the vastly exploratory nature of the original research question, a decision was made to not control for multiple comparisons to ensure that possible significant findings were not missed. If the decision had been made to control for multiple comparisons by using a more stringent alpha level, for example, the primary results of this manuscript would remain unchanged (impact of atypical vocalizations on amount of SC change). That said, future studies with larger samples should examine these variables with more robust alpha criteria. The sample size is relatively small, and the majority of the sample consisted of white male children with highly educated parents despite attempting to be representative of the larger local community (See Table 1). These results

may not be successfully extrapolated to more diverse populations; thus, future work would benefit from larger, more diverse samples. The results surrounding the atypical vocalizations may be particularly limited due to the use of only one measure of nuanced vocalizations; however, given the intricacy of coding 30 min of vocalizations, examining the predictive value of early vocalizations for outcomes in this population is a valuable contribution. The observational measures used in this work (SPA, vocalizations coding, and BOSCC) use brief samples of child behavior that intend to represent broader, pervasive characteristics, although these limited snapshots may not be fully representative of a particular child's skills in these areas. It is also important to note that we only have two timepoints to evaluate treatment-related change (pre- and post-treatment); this limits the interpretation of results as we cannot evaluate trends over time or regression to the mean. Future studies would benefit from the use of multiple measurements across more than 2 time points. The results of the work focused on direct observation (SPA) showed a relationship between Sensory Hyporeactivity and BOSCC-SC Ratio; however, this same relationship was not present for the parent-reported hyporeactivity (SEQ). These results could be limited due to the obstacles parents face when attempting to characterize hyporeactive behavior—that is, the absence of a response may be more difficult to recognize than the presence of an “over-response.” Clinicians may also observe certain behaviors when administering specific stimuli under more controlled conditions, but the same behaviors may be less evident outside of a clinical environment if, for example, the child functions more adaptively within predictable home routines. While parent reports on the SEQ questionnaire may be a broader and more ecologically-valid way to capture some aspects of sensory reactivity in daily routines, it is also possible that there is variability in the parent's ability to accurately report these behaviors in such a young sample. For example, interpretation or understanding of these behaviors may be based on parental experience such as experience raising older siblings or opportunities to observe their own child in relation to same-age peers. Research has shown that parents who are reporting on the later-born sibling of a child with autism indicate the sibling has fewer atypical sensory behaviors on the SEQ than parents who are reporting on the sibling of a typically developing child (Wolff et al., 2019), suggesting that parents of high familial likelihood siblings interpret the questions on the SEQ differently and in light of (or in comparison with) their experience of a child with autism. Although this sample was a community sample, and not a familial likelihood group, these types of discrepancies in parental experiences could be considered in future studies. Using a variety of measures, these analyses examined the impact of child characteristics, specifically sensory reactivity and atypical vocalizations, on treatment outcome in children with or at elevated likelihood for autism. These results of this exploratory study indicate that sensory hyporeactivity and atypical vocalizations should be further studied as potentially important characteristics to focus on when tailoring interventions to ensure that they are maximally effective for children with elevated likelihood for autism.

Overall, this initial work provides a stepping stone for future studies to examine baseline child characteristics as predictors of early intervention outcomes. Specifically, a child's baseline vocalization patterns (e.g., number of atypical vocalizations) as well as sensory reactivity may have significant implications for the amount of improvement to expect over the course of an intervention. As such, determinations about utility of an intervention may

need to be adjusted based on each child's baseline behaviors. Similarly, treatment goals and behavior targets may need to be determined based on a child's profile; for example, a child who often is hyporeactive to the social and nonsocial stimuli in the environment may benefit from caregivers or interventionists increasing the salience of stimuli or selecting activities and stimuli that have been previously observed to engage the child's interest and attention. . This work introduces novel questions to the field of early intervention research, highlighting the need for additional examinations of baseline skills/characteristics as researchers interpret the value of intervention practices.

Funding

The author(s) disclosed receipt of the following financial support for the research, authorship, and/or publication of this article: The research reported here was supported by the Institute of Education Sciences (IES), US Department of Education through grant #R324A100305 and by the National Institutes of Health (NIH) through grants #5T32HD040127-17 and #KL2TR002490 to the University of North Carolina at Chapel Hill. This article's contents are solely the responsibility of the authors and do not necessarily represent the official views of the funding agencies.

References

- Baranek GT (1999). Autism during infancy: A retrospective video analysis of sensory-motor and social behaviors at 9–12 months of age. *Journal of Autism and Developmental Disorders*, 29, 213–224. 10.1023/A:1023080005650 [PubMed: 10425584]
- Baranek GT, Watson LR, Crais E, & Reznick S (2003). *First Year Inventory (FYI) 2.0* Chapel Hill, NC: University of North Carolina.
- Baranek GT, Boyd BA, Poe MD, David FJ, & Watson LR (2007). Hyperresponsive sensory patterns in young children with autism, developmental delay, and typical development. *American Journal on Mental Retardation*, 112(4), 233–245. 10.1352/0895-8017(2007)112[233:HSPIYC]2.0.CO;2 [PubMed: 17559291]
- Baranek GT, Carlson M, Sideris J, Kirby AV, Watson LR, Williams KL, & Bulluck J (2019). Longitudinal assessment of stability of sensory features in children with autism spectrum disorder or other developmental disabilities. *Autism Research*, 12, 100–111. 10.1002/aur.2008 [PubMed: 30194913]
- Baranek GT, David FJ, Poe MD, Stone WL, & Watson LR (2006). Sensory Experiences Questionnaire: Discriminating sensory features in young children with autism, developmental delays, and typical development. *Journal of Child Psychology and Psychiatry and Allied Disciplines*, 47, 591–601. 10.1111/j.1469-7610.2005.01546.x [PubMed: 16712636]
- Baranek GT, Watson LR, Boyd BA, Poe MD, David FJ, & McGuire L (2013). Hyporesponsiveness to social and nonsocial sensory stimuli in children with autism, children with developmental delays, and typically developing children. *Development and Psychopathology*, 25(2), 307–320. 10.1017/S0954579412001071 [PubMed: 23627946]
- Bearss K, Johnson C, Smith T, Lecavalier L, Swiezy N, Aman M, McAdam DB, Butter E, Stillitano C, Minshawi N, Sukhodolsky DG, Mruzek DW, Turner K, Neal T, Hallett V, Mulick JA, Green B, Handen B, Deng Y, & Scahill L (2015). Effect of parent training vs parent education on behavioral problems in children with autism spectrum disorder: A randomized clinical trial. *JAMA—Journal of the American Medical Association*, 313(15), 1524–1533. 10.1001/jama.2015.3150
- Beglinger L, & Smith T (2005). Concurrent validity of social subtype and IQ after early intensive behavioral intervention in children with autism: A preliminary investigation. *Journal of Autism and Developmental Disorders*, 35, 295–303. 10.1007/s10803-005-3292-3 [PubMed: 16119470]
- Ben-Itzhak E, Watson LR, & Zachor DA (2014). Cognitive ability is associated with different outcome trajectories in autism spectrum disorders. *Journal of Autism and Developmental Disorders*, 44(9), 2221–2229. 10.1007/s10803-014-2091-0 [PubMed: 24710810]

- Ben-Sasson A, Carter AS, & Briggs-Gowan MJ (2009). Sensory over-responsivity in elementary school: Prevalence and social-emotional correlates. *Journal of Abnormal Child Psychology*, 37, 705–716. 10.1007/s10802-008-9295-8 [PubMed: 19153827]
- Ben-Sasson A, Cermak SA, Orsmond GI, Tager-Flusberg H, Kadlec MB, & Carter AS (2008). Sensory clusters of toddlers with autism spectrum disorders: Differences in affective symptoms. *Journal of Child Psychology and Psychiatry and Allied Disciplines*, 49, 817–825. 10.1111/j.1469-7610.2008.01899.x [PubMed: 18498344]
- Ben-Sasson A, Soto TW, Martínez-Pedraza F, & Carter AS (2013). Early sensory over-responsivity in toddlers with autism spectrum disorders as a predictor of family impairment and parenting stress. *Journal of Child Psychology and Psychiatry and Allied Disciplines*, 54, 846–853. 10.1111/jcpp.12035 [PubMed: 23336424]
- Bizzell E, Ross J, Rosenthal C, Dumont R, & Schaaf R (2020). Sensory features as a marker of autism spectrum disorders. *Journal of Autism and Developmental Disorders*, 50, 2240–2246. 10.1007/s10803-019-03948-8 [PubMed: 30825080]
- Brock ME, Freuler A, Baranek GT, Watson LR, Poe MD, & Sabatino A (2012). Temperament and sensory features of children with autism. *Journal of Autism and Developmental Disorders*, 42, 2271–2284. 10.1007/s10803-012-1472-5 [PubMed: 22366913]
- Carter AS, Messinger DS, Stone WL, Celimli S, Nahmias AS, & Yoder P (2011). A randomized controlled trial of Hanen’s “More Than Words” in toddlers with early autism symptoms. *Journal of Child Psychology and Psychiatry and Allied Disciplines*, 52, 741–752. 10.1111/j.1469-7610.2011.02395.x [PubMed: 21418212]
- Crank JE, Sandbank M, Dunham K, Crowley S, Bottema-Beutel K, Feldman J, & Woynarowski TG (2021). Understanding the effects of naturalistic developmental behavioral interventions: A project AIM meta-analysis. *Autism Research*, 14, 817–834. 10.1002/aur.2471 [PubMed: 33480474]
- Eldevik S, Hastings RP, Hughes JC, Jahr E, Eikeseth S, & Cross S (2009). Meta-analysis of early intensive behavioral intervention for children with autism. *Journal of Clinical Child and Adolescent Psychology*, 38(3), 439–450. 10.1080/15374410902851739 [PubMed: 19437303]
- Estes A, Swain DM, & MacDuffie KE (2019). The effects of early autism intervention on parents and family adaptive functioning. *Pediatric Medicine*, 2, 21. 10.21037/pm.2019.05.05 [PubMed: 31583390]
- Flippin M, & Watson LR (2011). Relationships between the responsiveness of fathers and mothers and the object play skills of children with autism spectrum disorders. *Journal of Early Intervention*, 33(3), 220–234. 10.1177/1053815111427445
- Fossum KL, Williams L, Garon N, Bryson SE, & Smith IM (2018). Pivotal response treatment for preschoolers with autism spectrum disorder: Defining a predictor profile. *Autism Research*, 11(1), 153–165. 10.1002/aur.1859 [PubMed: 28861936]
- Garrido D, Watson LR, Carballo G, Garcia-Retamero R, & Crais ER (2017). Infants at-risk for autism spectrum disorder: Patterns of vocalizations at 14 months. *Autism Research*, 10, 1372–1383. 10.1002/aur.1788 [PubMed: 28371531]
- Grzadzinski R, Carr T, Colombi C, McGuire K, Dufek S, Pickles A, & Lord C (2016). Measuring changes in social communication behaviors: Preliminary development of the Brief Observation of Social Communication Change (BOSCC). *Journal of Autism and Developmental Disorders*, 46, 2464–2479. 10.1007/s10803-016-2782-9 [PubMed: 27062034]
- Grzadzinski R, Donovan K, Truong K, Nowell S, Lee H, Sideris J, Turner-Brown L, Baranek GT, & Watson LR (2020). Sensory reactivity at 1 and 2 years old is associated with ASD severity during the preschool years. *Journal of Autism and Developmental Disorders*, 50, 3895–3904. 10.1007/s10803-020-04432-4 [PubMed: 32157566]
- Hampton LH, & Kaiser AP (2016). Intervention effects on spoken-language outcomes for children with autism: A systematic review and meta-analysis. *Journal of Intellectual Disability Research*, 60, 444–463. 10.1111/jir.12283 [PubMed: 27120988]
- Harris SL, & Handleman JS (2000). Age and IQ at intake as predictors of placement for young children with autism: A four- to six-year follow-up. *Journal of Autism and Developmental Disorders*, 30(2), 137–142. 10.1023/A:1005459606120 [PubMed: 10832778]

- Hayes AF (2017). Introduction to mediation, moderation, and conditional process analysis: A regression-based approach. Guilford Publications.
- Kasari C, Gulsrud AC, Wong C, Kwon S, & Locke J (2010). Randomized controlled caregiver mediated joint engagement intervention for toddlers with autism. *Journal of Autism and Developmental Disorders*, 40, 1045–1056. 10.1007/s10803-010-0955-5 [PubMed: 20145986]
- Landa RJ (2018). Efficacy of early interventions for infants and young children with, and at risk for, autism spectrum disorders. *International Review of Psychiatry*, 30(1), 25–39. 10.1080/09540261.2018.1432574 [PubMed: 29537331]
- Leigh JP, & Du J (2015). Brief report: Forecasting the economic burden of autism in 2015 and 2025 in the United States. *Journal of Autism and Developmental Disorders*, 45, 4135–4139. 10.1007/s10803-015-2521-7 [PubMed: 26183723]
- Little LM, Freuler AC, Houser MB, Guckian L, Carbine K, David FJ, & Baranek GT (2011). Psychometric validation of the Sensory Experiences Questionnaire. *American Journal of Occupational Therapy*, 65(2), 207–210. 10.5014/ajot.2011.000844
- López B, & Leekam SR (2007). Central coherence theory: A review of theoretical assumptions. *Childhood and Learning*, 30(3), 439–457. 10.1174/021037007781787462
- Mahoney G, & Macdonald J (2007). Autism and developmental delays in young children: The responsive teaching curriculum for parents and professionals. PRO-ED.
- Mullen EM (1995). Mullen Scales of Early Learning, AGS Edition: Manual and Item Administrative Books. American Guidance Services.
- Olafsen KS, Rønning JA, Kaaresen PI, Ulvund SE, Handegård BH, & Dahl LB (2006). Joint attention in term and preterm infants at 12 months corrected age: The significance of gender and intervention based on a randomized controlled trial. *Infant Behavior and Development*, 29(4), 554–563. 10.1016/j.infbeh.2006.07.004 [PubMed: 17138308]
- Oller DK, Eilers RE, Neal AR, & Schwartz HK (1999). Precursors to speech in infancy: The prediction of speech and language disorders. *Journal of Communication Disorders*, 32(4), 223–245. 10.1016/S0021-9924(99)00013-1 [PubMed: 10466095]
- Oller DK, Eilers RE, Steffens ML, Urbano R, & Lynch MP (1994). Speech-like vocalizations in infancy: An evaluation of potential risk factors. *Journal of Child Language*, 21(1), 33–58. 10.1017/S0305000900008667 [PubMed: 8006094]
- Oosterling I, Visser J, Swinkels S, Rommelse N, Donders R, Woudenberg T, Roos S, Van Der Gaag RJ, & Buitelaar J (2010). Randomized controlled trial of the focus parent training for toddlers with autism: 1-Year outcome. *Journal of Autism and Developmental Disorders*, 40, 1447–1458. 10.1007/s10803-010-1004-0 [PubMed: 20440639]
- Ouyang L, Grosse SD, Riley C, Bolen J, Bishop E, Raspa M, & Bailey DB (2014). A comparison of family financial and employment impacts of fragile X syndrome, autism spectrum disorders, and intellectual disability. *Research in Developmental Disabilities*, 35(7), 1518–1527. 10.1016/j.ridd.2014.04.009 [PubMed: 24755230]
- Patten E, Ausderau KK, Watson LR, & Baranek GT (2013). Sensory response patterns in nonverbal children with ASD. *Autism Research and Treatment*, 2013, Article 436286. 10.1155/2013/436286 [PubMed: 23956859]
- Patten E, Belardi K, Baranek GT, Watson LR, Labban JD, & Oller DK (2014). Vocal patterns in infants with autism spectrum disorder: Canonical babbling status and vocalization frequency. *Journal of Autism and Developmental Disorders*, 44(10), 2413–2428. 10.1007/s10803-014-2047-4 [PubMed: 24482292]
- Paul R, Fuerst Y, Ramsay G, Chawarska K, & Klin A (2011). Out of the mouths of babes: Vocal production in infant siblings of children with ASD. *Journal of Child Psychology and Psychiatry and Allied Disciplines*, 52, 588–598. 10.1111/j.1469-7610.2010.02332.x [PubMed: 21039489]
- Pellecchia M, Beidas RS, Marcus SC, Fishman J, Kimberly JR, Cannuscio CC, Reisinger EM, Rump K, & Mandell DS (2016). Study protocol: Implementation of a computer-assisted intervention for autism in schools: A hybrid type II cluster randomized effectiveness-implementation trial. *Implementation Science*, 11, Article 154. 10.1186/s13012-016-0513-4 [PubMed: 27884169]

- Pellicano E, Dinsmore A, & Charman T (2014). What should autism research focus upon? Community views and priorities from the United Kingdom. *Autism*, 18(7), 756–770. 10.1177/1362361314529627 [PubMed: 24789871]
- Philpott-Robinson K, Lane AE, & Harpster K (2016). Sensory features of toddlers at risk for autism spectrum disorder. *American Journal of Occupational Therapy*, 70(4), 7004220010p1–7004220010p8. 10.5014/ajot.2016.019497
- Plumb AM, & Wetherby AM (2013). Vocalization development in toddlers with autism spectrum disorder. *Journal of Speech, Language, and Hearing Research*, 56(2), 721–734. 10.1044/1092-4388(2012/11-0104)
- Rogers SJ, Estes A, Lord C, Vismara L, Winter J, Fitzpatrick A, Guo M, & Dawson G (2012). Effects of a brief early start Denver model (ESDM)-based parent intervention on toddlers at risk for autism spectrum disorders: A randomized controlled trial. *Journal of the American Academy of Child and Adolescent Psychiatry*, 51(10), 1052–1065. 10.1016/j.jaac.2012.08.003 [PubMed: 23021480]
- Rogers SJ, Yoder P, Estes A, Warren Z, McEachin J, Munson J, Rocha M, Greenson J, Wallace L, Gardner E, Dawson G, Sugar CA, Helleman G, & Whelan F (2020). A multisite randomized controlled trial comparing the effects of intervention intensity and intervention style on outcomes for young children with autism. *Journal of the American Academy of Child & Adolescent Psychiatry*, 60(6), 710–722. 10.1016/j.jaac.2020.06.013 [PubMed: 32853704]
- Sacrey LAR, Germani T, Bryson SE, & Zwaigenbaum L (2014). Reaching and grasping in autism spectrum disorder: A review of recent literature. *Frontiers in Neurology*, 5, Article 6. 10.3389/fneur.2014.00006 [PubMed: 24478753]
- Sacrey LAR, Zwaigenbaum L, Bryson S, Brian J, Smith IM, Roberts W, Szatmari P, Roncadin C, Garon N, Novak C, Vaillancourt T, McCormick T, Mackinnon B, Jilderda S, & Armstrong V (2015). Can parents' concerns predict autism spectrum disorder? A prospective study of high-risk siblings from 6 to 36 months of age. *Journal of the American Academy of Child and Adolescent Psychiatry*, 54(6), 470–478. 10.1016/j.jaac.2015.03.014 [PubMed: 26004662]
- Schreibman L, Stahmer AC, Barlett VC, & Dufek S (2009). Brief report: Toward refinement of a predictive behavioral profile for treatment outcome in children with autism. *Research in Autism Spectrum Disorders*, 3(1), 163–172. 10.1016/j.rasd.2008.04.008 [PubMed: 20046210]
- Shaw KA, Maenner MJ, Baio J, Washington A, Christensen DL, Wiggins LD, Pettygrove S, Andrews JG, White T, Rosenberg CR, Constantino JN, Fitzgerald RT, Zahorodny W, Shenouda J, Daniels JL, Salinas A, Durkin MS, & Dietz PM (2020). Early identification of autism spectrum disorder among children aged 4 years—Early autism and developmental disabilities monitoring network, six sites, United States, 2016. *MMWR Surveillance Summaries*, 69(3), 1–11. 10.15585/MMWR.SS6903A1
- Siller M, Hutman T, & Sigman M (2013). A parent-mediated intervention to increase responsive parental behaviors and child communication in children with ASD: A randomized clinical trial. *Journal of Autism and Developmental Disorders*, 43, 540–555. 10.1007/s10803-012-1584-y [PubMed: 22825926]
- Sutera S, Pandey J, Esser EL, Rosenthal MA, Wilson LB, Barton M, Green J, Hodgson S, Robins DL, Dumont-Mathieu T, & Fein D (2007). Predictors of optimal outcome in toddlers diagnosed with autism spectrum disorders. *Journal of Autism and Developmental Disorders*, 37(1), 98–107. 10.1007/s10803-006-0340-6 [PubMed: 17206522]
- Tamis-LeMonda CS, Bornstein MH, & Baumwell L (2001). Maternal responsiveness and children's achievement of language milestones. *Child Development*, 72, 748–767. 10.1111/1467-8624.00313 [PubMed: 11405580]
- Turner-Brown LM, Baranek GT, Reznick JS, Watson LR, & Crais ER (2013). The first year inventory: A longitudinal follow-up of 12-month-old to 3-year-old children. *Autism*, 17(5), 527–540. 10.1177/1362361312439633 [PubMed: 22781058]
- Vivanti G, Kasari C, Green J, Mandell D, Maye M, & Hudry K (2018). Implementing and evaluating early intervention for children with autism: Where are the gaps and what should we do? *Autism Research*, 11, 16–23. 10.1002/aur.1900 [PubMed: 29206358]

- Vivanti G, & Rogers SJ (2014). Autism and the mirror neuron system: Insights from learning and teaching. *Philosophical Transactions of the Royal Society B: Biological Sciences*, 369, 20130184. 10.1098/rstb.2013.0184
- Warren SF, Brady N, Sterling A, Fleming K, & Marquis J (2010). Maternal responsivity predicts language development in young children with fragile X syndrome. *American Journal on Intellectual and Developmental Disabilities*, 115(1), 54–75. 10.1352/1944-7558-115.1.54 [PubMed: 20025359]
- Watson LR, Baranek GT, Crais ER, Steven Reznick J, Dykstra J, & Perryman T (2007). The First Year Inventory: Retrospective parent responses to a questionnaire designed to identify one-year-olds at risk for autism. *Journal of Autism and Developmental Disorders*, 37, 49–61. 10.1007/s10803-006-0334-4 [PubMed: 17219058]
- Watson LR, Crais ER, Baranek GT, Dykstra JR, & Wilson KP (2013). Communicative gesture use in infants with and without autism: A retrospective home video study. *American Journal of Speech-language Pathology*, 22(1), 25–39. 10.1044/1058-0360(2012/11-0145) [PubMed: 22846878]
- Watson LR, Crais ER, Baranek GT, Turner-Brown L, Sideris J, Wakeford L, Kinard J, Reznick JS, Martin KL, & Nowell SW (2017). Parent-mediated intervention for one-year-olds screened as at-risk for autism spectrum disorder: A randomized controlled trial. *Journal of Autism and Developmental Disorders*, 47, 3520–3540. 10.1007/s10803-017-3268-0 [PubMed: 28861651]
- Wetherby AM, & Prizant BM (2002). *Communication and symbolic behavior scales: developmental profile*. Paul H Brookes Publishing Co.
- Wolff JJ, Dimian AF, Botteron KN, Dager SR, Elison JT, Estes AM, Hazlett HC, Schultz RT, Zwaigenbaum L, Piven J, Chappell C, Shaw D, McKinstry R, Constantino J, Pruett J, Pandey J, Paterson S, Elison J, Evans AC, & Gu H (2019). A longitudinal study of parent-reported sensory responsiveness in toddlers at-risk for autism. *Journal of Child Psychology and Psychiatry and Allied Disciplines*, 60, 314–324. 10.1111/jcpp.12978 [PubMed: 30350375]
- Yoder P, & Stone WL (2006). A randomized comparison of the effect of two prelinguistic communication interventions on the acquisition of spoken communication in preschoolers with ASD. *Journal of Speech, Language, and Hearing Research*, 49(4), 698–711. 10.1044/1092-4388(2006/051)
- Zachor DA, Ben-Itzhak E, Rabinovich AL, & Lahat E (2007). Change in autism core symptoms with intervention. *Research in Autism Spectrum Disorders*, 1(4), 304–317. 10.1016/j.rasd.2006.12.001




Figure 1.
Language variables.




Figure 2.
BOSCC SC change ratio versus SPA hyporeactivity.




Figure 3. BOSCC SC change ratio versus number of atypical vocalizations.




Figure 4.
BOSCC RRB change ratio versus number of atypical vocalizations.

Table 1.Demographics ($n = 87$).

Demographic Variables	REIM (42)		ART (45)	
	Timepoint 1 mean (SD)	Timepoint 2 mean (SD)	Timepoint 1 mean (SD)	Timepoint 2 mean (SD)
Age (in months)	14 (0.78)	22 (0.79)	14 (0.77)	23 (0.99)
Mullen Scales of Early Learning Domains				
Fine motor	48 (9.21)	43 (11.09)	47 (10.44)	40 (13.67)
Visual reception	47 (11.17)	47 (12.30)	43 (10.04)	45 (13.80)
Receptive language	33 (12.33)	47 (15.76)	33 (9.47)	42 (16.95)
Expressive language,	36 (12.08)	41 (13.53)	34 (10.98)	41 (11.86)
<i>n</i> (%)				
Sex (males)		32 (76)		28 (62)
Child race				
White		31 (74)		29 (64)
African-American		6 (14)		12 (27)
Mixed race/other		5 (12)		4 (9)
Primary caregiver education level				
Less than high school		0 (0)		3 (7)
High-school diploma/High-school equivalency diploma		7 (16.7)		4 (9)
Vocational/associates/some college		8 (19)		7 (16)
4-year college degree		11 (26)		13 (29)
Graduate/professional degree		16 (38)		18 (40)

Note. REIM = Referral to Early Intervention and Monitoring; ART = Adapted Responsive Teaching.

Table 2.

Correlations.

Variables of Interest		BOSCC RRB ratio change	BOSCC SC ratio change	Canonical Directed	Canonical Non-Directed	Non-Canonical Directed	Non-Canonical Non-Directed	Atypical vocalizations	Distress vocalizations	Pleasure vocalizations	SPA hyporeactivity	SPA seeking	SEQ hyporeactivity	SEQ seeking
BOSCC RRB ratio change	Pearson's correlation	1												
	Sig. (2-tailed)													
	<i>N</i>	75												
BOSCC SC ratio change	Pearson's correlation	-.286*	1											
	Sig. (2-tailed)	0.013												
	<i>N</i>	75	83											
Canonical Directed	Pearson's correlation	0.157	-0.051	1										
	Sig. (2-tailed)	0.181	0.65											
	<i>N</i>	74	82	82										
Canonical Non-Directed	Pearson's correlation	-0.021	0.2	.283*	1									
	Sig. (2-tailed)	0.859	0.072	0.01										
	<i>N</i>	74	82	82	82									
Non-Canonical Directed	Pearson's correlation	0.16	-0.06	.320**	0.181	1								
	Sig. (2-tailed)	0.174	0.59	0.003	0.103									
	<i>N</i>	74	82	82	82	82								
Non-Canonical Non-Directed	Pearson's correlation	0.09	0.105	0.018	.665**	.273*	1							
	Sig. (2-tailed)	0.447	0.346	0.87	0	0.013								
	<i>N</i>	74	82	82	82	82	82							
Atypical vocalizations	Pearson's correlation	-.239*	.363**	-0.049	0.139	0.211	0.178	1						
	Sig. (2-tailed)	0.04	0.001	0.664	0.213	0.057	0.109							
	<i>N</i>	74	82	82	82	82	82	82						
Distress vocalizations	Pearson's correlation	-0.094	-0.013	0.103	-0.022	0.098	0.114	0.024	1					
	Sig. (2-tailed)	0.426	0.907	0.559	0.843	0.379	0.308	0.831						
	<i>N</i>	74	82	82	82	82	82	82	82					
Pleasure vocalizations	Pearson's correlation	0.154	-0.006	0.102	0.105	-0.022	-0.036	0.196	-0.115	1				
	Sig. (2-tailed)	0.189	0.959	0.562	0.348	0.842	0.747	0.078	0.305					
	<i>N</i>	74	82	82	82	82	82	82	82	82				
SPA hyporeactivity	Pearson's correlation	-0.041	.272*	-0.19	0.036	-.273*	0.025	0.097	0.019	-0.004	1			
	Sig. (2-tailed)	0.726	0.013	0.088	0.745	0.013	0.822	0.384	0.865	0.974				
	<i>N</i>	75	83	82	82	82	82	82	82	82	99			

Variables of Interest		BOSCC RRB ratio change	BOSCC SC ratio change	Canonical Directed	Canonical Non-Directed	Non-Canonical Directed	Non-Canonical Non-Directed	Atypical vocalizations	Distress vocalizations	Pleasure vocalizations	SFA hyporeactivity	SFA hyperactivity	SFA seeking	SEQ hyporeactivity	SEQ hyperactivity	SEQ seeking
SPA hyporeactivity	Pearson's correlation	-0.031	.234*	0.02	-0.083	0.013	0.033	0.042	.237*	-0.146	0.029	1				
	Sig. (2-tailed)	0.79	0.033	0.857	0.457	0.907	0.765	0.71	0.032	0.191	0.774					
	N	75	83	82	82	82	82	82	82	82	99	99				
SPA seeking	Pearson's correlation	0.114	-0.148	-0.084	-0.02	-0.052	0.122	-0.127	-0.049	-0.188	0.104	0.136	1			
	Sig. (2-tailed)	0.329	0.181	0.453	0.86	0.64	0.275	0.257	0.662	0.091	0.305	0.18				
	N	75	83	82	82	82	82	82	82	82	99	99	99			
SEQ hyporeactivity	Pearson's correlation	-0.153	0.175	-0.076	0.041	-0.387***	-0.022	-0.09	0.076	-0.067	.311**	-0.024	0.088	1		
	Sig. (2-tailed)	0.191	0.113	0.499	0.712	0	0.847	0.423	0.497	0.551	0.002	0.813	0.387			
	N	75	83	82	82	82	82	82	82	82	99	99	99	100		
SEQ hyperactivity	Pearson's correlation	0.055	-0.014	0.202	0.034	-0.007	-0.026	-0.076	0.146	0.008	0.083	0.115	-0.181	.346***	1	
	Sig. (2-tailed)	0.638	0.898	0.068	0.763	0.949	0.816	0.497	0.19	0.942	0.414	0.258	0.074	0		
	N	75	83	82	82	82	82	82	82	82	99	99	99	100	100	
SEQ seeking	Pearson's correlation	0.009	0.028	-0.111	0.033	0.132	0.01	-0.049	-0.031	0.085	-0.028	0.133	-0.005	-0.084	0.113	1
	Sig. (2-tailed)	0.939	0.804	0.322	0.771	0.237	0.928	0.664	0.783	0.446	0.781	0.188	0.963	0.406	0.265	
	N	75	83	82	82	82	82	82	82	82	99	99	99	100	100	100

Note. BOSCC = Brief Observation of Social Communication Change; RRB = restricted and repetitive behaviors; SC = social communication; SFA = Sensory Processing Assessment; SEQ = Sensory Experiences Questionnaire.

* Correlation is significant at the .05 level (2-tailed).

*** Correlation is significant at the .01 level (2-tailed).

Table 3.

Moderation and main effects: BOSCC SC change ratio.

Variables	Moderation Effects							Main Effects					
	Coefficient	SE	t	Significance	Model	Variables	B	t	Sig.	R	R Square	R Square Change	
SPA (Observational)	Hyporeactivity	0.02	0.04	0.42	0.67	1							
	Group	-0.26	0.12	-2.23	0.03								
	Group × Hyporeactivity	0.12	0.05	2.45	0.02								
	Visual Reception	0.00	0.00	-1.54	0.13								
	Expressive Language	0.01	0.00	2.83	0.01								
	Hyperreactivity	0.11	0.13	0.85	0.40	1	Hyperreactivity	0.14	2.17	0.03	0.23	0.05	
	Group	-0.06	0.22	-0.28	0.78								
	Group × Hyperreactivity	0.02	0.15	0.15	0.88	2	Hyperreactivity	0.12	1.92	0.06	0.34	0.12	0.06
	Visual Reception	0.00	0.00	-2.05	0.04		Visual Reception	0.00	-1.96	0.05			
	Expressive Language	0.00	0.00	1.87	0.06		Expressive Language	0.00	1.89	0.06			
SEQ (Parent Report)	Seeking	-0.03	0.05	-0.73	0.47	1	Seeking	-0.04	-1.35	0.18	0.15	0.02	
	Group	-0.02	0.16	-0.10	0.92								
	Group × Seeking	0.00	0.06	-0.03	0.97	2	Seeking	-0.03	-1.25	0.22	0.31	0.09	0.07
	Visual Reception	-0.01	0.00	-2.28	0.03		Visual Reception	0.00	-2.29	0.02			
	Expressive Language	0.00	0.00	1.52	0.13		Expressive Language	0.00	1.58	0.12			
	Hyporeactivity	0.03	0.06	0.53	0.60	1	Hyporeactivity	0.05	1.60	0.11	0.18	0.03	
	Group	-0.12	0.15	-0.83	0.41								
	Group × Hyporeactivity	0.05	0.07	0.78	0.44	2	Hyporeactivity	0.06	1.92	0.06	0.34	0.12	0.09
	Visual Reception	0.00	0.00	-1.93	0.06		Visual Reception	0.00	-1.99	0.05			
	Expressive Language	0.00	0.00	2.47	0.02		Expressive Language	0.00	2.45	0.02			
Hyperreactivity	Hyperreactivity	-0.05	0.07	-0.75	0.46	1	Hyperreactivity	-0.01	-0.13	0.90	0.01	0.00	
	Group	-0.15	0.18	-0.85	0.40								

Variables	Moderation Effects							Main Effects					R Square Change
	Coefficient	SE	t	Significance	Model	Variables	B	t	Sig.	R	R Square		
Group × Hyperreactivity	0.07	0.09	0.79	0.43	2	Hyperreactivity	-0.01	-0.27	0.79	0.28	0.08	0.08	
Visual Reception	0.00	0.00	-1.97	0.05		Visual Reception	0.00	-2.12	0.04				
Expressive Language	0.00	0.00	2.00	0.05		Expressive Language	0.00	2.04	0.04				
Seeking	0.02	0.07	0.26	0.79	1	Seeking	0.01	0.25	0.80	0.03	0.00	—	
Group	0.04	0.21	0.20	0.84									
Group × Seeking	-0.02	0.08	-0.29	0.77	2	Seeking	0.00	0.01	0.99	0.28	0.08	0.08	
Visual Reception	0.00	0.00	-2.10	0.04		Visual Reception	0.00	-2.12	0.04				
Expressive Language	0.00	0.00	1.93	0.06		Expressive Language	0.00	2.01	0.05				
Canonical Directed	0.00	0.00	-0.63	0.53	1	Canonical Directed	0.00	-0.46	0.65	0.05	0.00	—	
Group	0.00	0.07	-0.03	0.97									
Group × Canonical Directed	0.00	0.00	-0.14	0.89	2	Canonical Directed	0.00	-1.09	0.28	0.30	0.09	0.09	
Visual Reception	0.00	0.00	-2.20	0.03		Expressive Language	0.00	2.01	0.05				
Expressive Language	0.00	0.00	1.99	0.05		Visual Reception	0.00	-2.23	0.03				
Canonical Non-directed	0.00	0.00	1.15	0.25	1	Canonical Non-directed	0.01	1.82	0.07	0.20	0.04	—	
Group	-0.02	0.05	-0.36	0.72									
Group × Canonical Non-directed	0.01	0.01	0.64	0.52	2	Canonical Non-directed	0.01	1.59	0.12	0.32	0.10	0.06	
Visual Reception	0.00	0.00	-2.05	0.04		Visual Reception	0.00	-2.26	0.03				
Expressive Language	0.00	0.00	1.20	0.23		Expressive Language	0.00	1.22	0.23				
Non-canonical Directed	0.00	0.00	-0.91	0.37	1	Non-canonical Directed	0.00	-0.54	0.59	0.06	0.00	—	
Group	-0.02	0.08	-0.23	0.82									
Group × Non-Canonical Directed	0.00	0.00	0.16	0.88	2	Non-canonical Directed	0.00	-1.16	0.25	0.30	0.09	0.09	

Variables	Moderation Effects						Main Effects					
	Coefficient	SE	t	Significance	Model	Variables	B	t	Sig.	R	R Square	R Square Change
Visual Reception	0.00	0.00	-2.30	0.02		Expressive Language	0.00	1.98	0.05			
Expressive Language	0.00	0.00	1.96	0.05		Visual Reception	0.00	-2.37	0.02			
Non-Canonical Non-directed	0.00	0.01	0.79	0.43	1	Non-canonical Non-directed	0.00	0.95	0.35	0.11	0.01	
Group	0.00	0.06	-0.03	0.98								
Group × Non-Canonical Non-Directed	0.00	0.01	-0.14	0.89	2	Non-canonical Non-directed	0.00	0.94	0.35	0.29	0.08	0.07
Visual Reception	0.00	0.00	-2.22	0.03		Visual Reception	0.00	-2.26	0.03			
Expressive Language	0.00	0.00	1.64	0.11		Expressive Language	0.00	1.66	0.10			
Atypical	0.01	0.00	1.52	0.13	1	Atypical	0.01	3.49	0.00	0.36	0.13	
Group	-0.08	0.06	-1.25	0.22								
Group × Atypical	0.00	0.00	0.83	0.41	2	Atypical	0.01	3.77	0.00	0.47	0.22	0.09
Visual Reception	-0.01	0.00	-2.57	0.01		Visual Reception	0.00	-2.50	0.01			
Expressive Language	0.00	0.00	2.28	0.03		Expressive Language	0.00	2.13	0.04			
Distress	0.00	0.00	-0.88	0.38	1	Distress	0.00	-0.12	0.91	0.01	0.00	
Group	-0.07	0.07	-1.07	0.29								
Group × Distress	0.00	0.00	1.18	0.24	2	Distress	0.00	-0.21	0.83	0.27	0.07	0.07
Visual Reception	-0.01	0.00	-2.40	0.02		Visual Reception	0.00	-2.25	0.03			
Expressive Language	0.00	0.00	1.81	0.08		Expressive Language	0.00	1.67	0.10			
Pleasure	-0.01	0.01	-0.80	0.43	1	Pleasure	0.00	-0.05	0.96	0.01	0.00	
Group	-0.05	0.06	-0.84	0.40								
Group × Pleasure	0.01	0.01	1.02	0.31	2	Pleasure	0.00	-0.17	0.86	0.27	0.07	0.07
Visual Reception	0.00	0.00	-2.16	0.03		Visual Reception	0.00	-2.24	0.03			
Expressive Language	0.00	0.00	1.74	0.09		Expressive Language	0.00	1.70	0.09			

Note. Models 1 and 2 represent effects before and while controlling for visual reception and expressive language. SPA = Sensory Processing Assessment; SEQ = Sensory Experiences Questionnaire. Bolded entries highlight significant results.

^aSPA Hyporeactivity main effects were not run due to significant moderation effects, as described in the analyses section.

Author Manuscript

Author Manuscript

Author Manuscript

Author Manuscript

Table 4.

Moderation and main effects: BOSCC RRB change ratio.

Variables	Moderation Effects										Main Effects				
	Coefficient	SE	t	Significance	Model	Variables	B	t	Sig.	R	R	R	Square	Square	Change
SPA (Observational)	Hyporeactivity	0.06	0.06	1.04	0.30	1	Hyporeactivity	-0.01	-0.35	0.73	0.04	0.00	0.00	0.00	—
	Group	0.28	0.19	1.45	0.15										
	Group × Hyporeactivity	-0.10	0.08	-1.32	0.19	2	Hyporeactivity	0.00	0.09	0.93	0.16	0.03	0.03	0.03	0.03
	Visual Reception	0.00	0.00	0.97	0.33		Visual Reception	0.00	0.79	0.43					
	Expressive Language	0.00	0.00	0.91	0.37		Expressive Language	0.00	0.84	0.40					
	Hyperreactivity	0.02	0.20	0.10	0.92	1	Hyperreactivity	-0.03	-0.27	0.79	0.03	0.00	0.00	0.00	—
	Group	0.13	0.34	0.39	0.69										
	Group × Hyperreactivity	-0.06	0.23	-0.28	0.78	2	Hyperreactivity	-0.02	-0.22	0.83	0.17	0.03	0.03	0.03	0.03
	Visual Reception	0.00	0.00	0.89	0.38		Visual Reception	0.00	0.76	0.45					
	Expressive Language	0.00	0.00	0.87	0.39		Expressive Language	0.00	0.85	0.40					
SEQ (Parent Report)	Seeking	0.15	0.07	2.16	0.03	1	Seeking	0.04	0.98	0.33	0.11	0.01	0.01	0.01	—
	Group	0.35	0.23	1.52	0.13										
	Group × Seeking	-0.11	0.08	-1.39	0.17	2	Seeking	0.07	1.73	0.09	0.26	0.07	0.07	0.05	
	Visual Reception	0.01	0.00	1.54	0.13		Visual Reception	0.00	1.15	0.25					
	Expressive Language	0.01	0.00	1.70	0.09		Expressive Language	0.00	1.32	0.19					
	Hyporeactivity	0.05	0.09	0.56	0.57	1	Hyporeactivity	-0.07	-1.32	0.19	0.15	0.02	0.02	0.02	—
	Group	0.34	0.22	1.54	0.13										
	Group × Hyporeactivity	-0.15	0.10	-1.46	0.15	2	Hyporeactivity	-0.05	-1.04	0.30	0.20	0.04	0.04	0.04	0.02
	Visual Reception	0.00	0.00	0.76	0.45		Visual Reception	0.00	0.76	0.45					
	Expressive Language	0.00	0.00	0.58	0.56		Expressive Language	0.00	0.60	0.55					
Hyperreactivity	-0.05	0.10	-0.51	0.61	1	Hyperreactivity	0.03	0.47	0.64	0.06	0.00	0.00	0.00	—	
Group	-0.21	0.27	-0.79	0.44											

Variables	Moderation Effects						Main Effects						
	Coefficient	SE	t	Significance	Model	Variables	B	t	Sig.	R	R Square	R Square Change	
Canonical	Group × Hyperreactivity	0.12	0.13	0.95	0.35	2	Hyperreactivity	0.03	0.40	0.69	0.17	0.03	0.03
	Visual Reception	0.00	0.00	0.98	0.33		Visual Reception	0.00	0.76	0.45			
	Expressive Language	0.00	0.00	0.84	0.41		Expressive Language	0.00	0.84	0.40			
	Seeking	-0.07	0.10	-0.75	0.45	1	Seeking	0.01	0.08	0.94	0.01	0.00	—
	Group	-0.24	0.32	-0.75	0.46								
	Group × Seeking	0.11	0.12	0.89	0.38	2	Seeking	0.00	-0.03	0.98	0.16	0.03	0.03
	Visual Reception	0.00	0.00	0.81	0.42		Visual Reception	0.00	0.79	0.43			
	Expressive Language	0.00	0.00	1.00	0.32		Expressive Language	0.00	0.84	0.41			
	Canonical Directed	0.00	0.00	-0.08	0.93	1	Canonical Directed	0.00	1.35	0.18	0.16	0.03	—
	Group	-0.03	0.10	-0.27	0.79								
Canonical Directed	Group × Canonical Directed	0.00	0.00	0.73	0.47	2	Canonical Directed	0.00	0.70	0.49	0.21	0.05	0.02
	Visual Reception	0.00	0.00	0.96	0.34		Visual Reception	0.00	0.86	0.39			
	Expressive Language	0.00	0.00	0.71	0.48		Expressive Language	0.00	0.69	0.50			
	Canonical Non-directed	0.00	0.01	-0.44	0.66	1	Canonical Non-directed	0.00	-0.18	0.86	0.02	0.00	—
	Group	0.03	0.08	0.39	0.70								
	Group × Canonical Non-directed	0.00	0.01	-0.10	0.92	2	Canonical Non-directed	0.00	-0.60	0.55	0.21	0.04	0.04
	Visual Reception	0.00	0.00	0.96	0.34		Visual Reception	0.00	0.92	0.36			
	Expressive Language	0.00	0.00	1.18	0.24		Expressive Language	0.00	1.22	0.23			
	Non-canonical Directed	0.00	0.00	1.33	0.19	1	Non-canonical Directed	0.00	1.37	0.17	0.16	0.03	—
	Group	0.12	0.13	0.96	0.34								
Non-Canonical Directed	Group × Non-canonical Directed	0.00	0.00	-0.91	0.36	2	Non-canonical Directed	0.00	1.02	0.31	0.23	0.05	0.03
	Visual Reception	0.00	0.00	0.98	0.33		Visual Reception	0.00	0.97	0.33			

Variables	Moderation Effects							Main Effects				
	Coefficient	SE	t	Significance	Model	Variables	B	t	Sig.	R	R Square	R Square Change
Non-Canonical Non-Directed	0.00	0.00	0.75	0.45		Expressive Language	0.00	0.73	0.47			
	0.00	0.01	0.37	0.71	1	Non-canonical Non-directed	0.01	0.77	0.45	0.09	0.01	—
	0.02	0.09	0.20	0.84		Group						
Visual Reception	0.00	0.01	0.30	0.76	2	Non-canonical Non-directed	0.00	0.70	0.49	0.22	0.05	0.04
	0.00	0.00	0.95	0.35		Visual Reception	0.00	0.87	0.39			
	0.00	0.00	1.13	0.26		Expressive Language	0.00	1.11	0.27			
Other Vocalizations	-0.01	0.01	-1.69	0.10	1	Atypical	-0.01	-2.09	0.04	0.24	0.06	—
	0.01	0.10	0.06	0.95		Group						
	0.00	0.01	0.62	0.54	2	Atypical	-0.01	-1.98	0.05	0.30	0.09	0.03
Distress Vocalizations	0.00	0.00	1.13	0.26		Visual Reception	0.00	0.89	0.38			
	0.00	0.00	1.10	0.28		Expressive Language	0.00	1.02	0.31			
	0.00	0.00	-0.30	0.76	1	Distress	0.00	-0.80	0.43	0.09	0.01	—
Pleasure Vocalizations	0.04	0.10	0.43	0.67		Group						
	0.00	0.00	-0.22	0.83	2	Distress	0.00	-0.59	0.55	0.21	0.04	0.04
	0.00	0.00	0.95	0.35		Visual Reception	0.00	0.86	0.39			
Group × Pleasure	0.00	0.00	1.01	0.32		Expressive Language	0.00	1.06	0.29			
	0.00	0.01	-0.02	0.98	1	Pleasure	0.01	1.33	0.19	0.15	0.02	—
	-0.03	0.08	-0.35	0.73		Group						
Visual Reception	0.02	0.02	1.33	0.19	2	Pleasure	0.01	1.04	0.30	0.23	0.05	0.03
	0.00	0.00	0.99	0.33		Visual Reception	0.00	0.82	0.42			
	0.00	0.00	1.07	0.29		Expressive Language	0.00	0.98	0.33			

Note. Models 1 and 2 represent effects before and while controlling for visual reception and expressive language. SPA = Sensory Processing Assessment; SEQ = Sensory Experiences Questionnaire. Bolded entries highlight significant results.