

UC Davis

UC Davis Previously Published Works

Title

The Role of Health in the Technology Acceptance Model Among Low-Income Asian American Older Adults: Cross-Sectional Survey Analysis.

Permalink

<https://escholarship.org/uc/item/00k1h9dc>

Authors

DeLange Martinez, Pauline

Tancredi, Daniel

Pavel, Misha

et al.

Publication Date

2024-12-03

DOI

10.2196/57009

Peer reviewed

Original Paper

The Role of Health in the Technology Acceptance Model Among Low-Income Asian American Older Adults: Cross-Sectional Survey Analysis

Pauline DeLange Martinez¹, PhD; Daniel Tancredi², PhD; Misha Pavel³, PhD; Lorena Garcia⁴, DrPH; Heather M Young¹, PhD

¹Betty Irene Moore School of Nursing, University of California, Davis, Sacramento, CA, United States

²Department of Pediatrics, University of California, Davis, Sacramento, CA, United States

³Khoury College of Computer Science, Northeastern University, Boston, MA, United States

⁴Department of Public Health Sciences, University of California, Davis, Sacramento, CA, United States

Corresponding Author:

Pauline DeLange Martinez, PhD
Betty Irene Moore School of Nursing
University of California, Davis
2750 48th St
Sacramento, CA, 95817
United States
Phone: 1 916 426 2862
Email: pdmartinez@ucdavis.edu

Abstract

Background: Self-rated health is associated with information and communications technology (ICT) use among older adults. Non-US born, older Asian American individuals are more inclined to rate their health as fair or poor compared to individuals from other racial and ethnic backgrounds. This population is also less likely to use ICTs as compared to White older Americans. Furthermore, cognitive decline may impact technology acceptance. In a previous adaptation of the technology acceptance model for low-income, Asian American older adults, perceived usefulness (PU), perceived ease of use (PEOU), age, educational attainment, ethnicity, and English proficiency were significant predictors of ICT use. However, the association between health and technology acceptance has not been explored among Asian American older adults.

Objective: This study examined the role of self-rated health and subjective cognitive decline in the acceptance and use of ICTs among low-income, Asian American older adults.

Methods: This cross-sectional survey included Asian American individuals aged ≥ 62 years living in affordable housing for older adults ($N=392$). Using hierarchical multiple regression, we explored the association between self-rated health and ICT use and technology acceptance model mediators (PU and PEOU) while adjusting for demographics, English proficiency, and subjective cognitive decline. Contrast statements were used to estimate contrasts of interest. To further examine the separate and joint association between age and subjective cognitive decline and the dependent variables, we examined scatterplots with locally estimated scatterplot smoothing lines, revealing that the relationship between subjective cognitive decline and ICT use varied in 3 age segments, which led to updating our analysis to estimate differences in ICT use among age categories with and without subjective cognitive decline.

Results: Self-rated health was not significantly associated with ICT use ($\beta=.087$; $P=.13$), PU ($\beta=.106$; $P=.10$), or PEOU ($\beta=.062$; $P=.31$). However, the interaction terms of subjective cognitive decline and age significantly improved the model fit for ICT use ($\Delta R^2=0.011$; $P=.04$). In reviewing scatterplots, we determined that, in the youngest age group (62-74 years), ICT use increased with subjective cognitive decline, whereas in the older age groups (75-84 and ≥ 85 years), ICT use decreased with subjective cognitive decline, more so in the oldest age category. Through regression analysis, among participants with subjective cognitive decline, ICT use significantly decreased in the middle and older age groups as compared to the youngest age group. However, among participants without subjective cognitive decline, the difference in use among age groups was not significant.

Conclusions: This study contributes to the understanding of the complex relationship between health and ICT acceptance among low-income, Asian American older adults and suggests the need for tailored interventions to promote digital engagement and quality of life for this population.

(*JMIR Form Res* 2024;8:e57009) doi: [10.2196/57009](https://doi.org/10.2196/57009)

KEYWORDS

aged; older adults; Asian American; immigrant; vulnerable populations; internet; information and communications technology; ICT; digital divide; technology acceptance model; mobile phone

Introduction

Background

Asian Americans are the fastest-growing ethnic group aged ≥ 65 years in the United States, with 85% of this group being non-US born [1-4]. Despite their rapid population growth, Asian American older adults remain under-studied and under-served, attributed in part to their heterogeneity, encompassing >40 ethnic subgroups characterized by diverse languages, cultural backgrounds, and immigration histories [1,2,5-7]. Asian American older adults are less likely to use information and communications technologies (ICTs) such as smartphones, tablets, computers, the internet, social media, or other applications as compared to White older Americans [8,9]. Intersectional factors such as socioeconomic status and English proficiency play a significant role in diminishing the likelihood of ICT use among this population [8-10]. According to data from the California Health Interview Survey (CHIS), non-US born Asian American older adults are more inclined to rate their health as fair or poor compared to individuals from other racial and ethnic backgrounds [11]. While there is a significant association between self-rated health and ICT use among older adults [12-15], this association has not been explored among Asian American older adults.

Self-rated health is a subjective measure that encompasses multiple dimensions of an individual's well-being, including physical, mental, social, and functional aspects. Self-rated health is influenced by personal and cultural beliefs as well as health behaviors [16]. Self-rated health has an independent effect on health outcomes even when adjusting for other measures of medical, physical, cognitive, emotional, and social health [17]. Among Asian American individuals, self-rated health is influenced by factors such as gender; age; and level of acculturation, including English proficiency [18].

Poor self-rated health among Asian American older adults is also closely linked to mental well-being. Analysis of survey data from a large sample of Korean American and Chinese American older adults collected from 2011 to 2019 uncovered significant and positive associations among poor self-rated health, depression, number of chronic diseases diagnosed, and functional disabilities [19]. Furthermore, CHIS data revealed that Asian American older adults with limited English proficiency (LEP) were 3 times more likely to have a perceived mental health need as compared to their English-speaking counterparts [20]. According to the National Asian Pacific Center on Aging, >60% of Asian American individuals aged ≥ 65 years have LEP, and these individuals are more likely to rate their health as poor [1,20]. Among Asian American older

adults, poor self-rated health is attributed to a variety of factors, including low educational attainment, low health literacy, racism and discrimination, disparities in quality of care accumulated over time, and delayed use of health care [11]. Self-rated health among Asian American older adults may be further impacted by challenges such as living far from relatives and experiencing cultural differences and limited social networks in their new environment [1,21,22].

Asian American older adults have higher poverty rates (9.3%) than the general older adult population in the United States (8.9%) [23]. Low-income older adults are more likely to rate their health as poor and report lower ICT use [24,25]. In an analysis of the CHIS, the combination of Asian ethnicity and low income had an interactive, negative effect on ICT use [10]. In fact, Asian American older adults in the lowest income category were 14 times less likely to use the internet for health information compared to non-Hispanic White older adults in the highest income category [10]. More research is needed to understand the factors that impact ICT adoption among low-income, ethnic minority older adults to inform interventions to address this digital divide.

Theoretical Framework

The technology acceptance model (TAM) is a widely accepted framework used to understand the adoption of new technologies [26]. The model suggests that attitudes toward technology, including perceived usefulness (PU) and perceived ease of use (PEOU), significantly influence an individual's intention to use and actual use of technology.

Nayak et al [14] and Chen and Chan [15] each independently adapted the TAM, adding self-rated health as a covariate predicting ICT use among older adults. In a study of adults aged ≥ 60 years in the United Kingdom, poor self-rated health was significantly, positively associated with hours per week spent using the internet when adjusting for attitudes toward technology, including PU and PEOU, and demographics [14]. Similarly, in a study of adults aged ≥ 55 years in Hong Kong, poor self-rated health was positively associated with use of gerontechnologies, a broad term encompassing ICTs, housing and daily living technology, health technology, and education and recreation technologies, when adjusting for attitudes toward technology and demographics. In this study, other health variables were incorporated as covariates, including number of health conditions, physical functioning, cognitive ability, social relationships, and attitude toward life and satisfaction [15].

While Nayak et al [14] and Chen and Chan [15] reported an inverse relationship between self-rated health and technology use, other studies have found them to be positively associated.

Werner et al [12] conducted a cross-sectional survey analysis of ethnically and income-diverse adults aged ≥60 years in the United States, demonstrating that better ratings of general physical health and role-related emotional health were independently and positively associated with email and computer use, respectively, when adjusting for demographic factors. In another cross-sectional survey analysis engaging adults aged ≥80 years, Sims et al [13] found that participants with better self-rated health were more likely to report using a greater variety of ICT devices and applications.

Overall, studies support a significant association between self-rated health and ICT use among older adults. However, some have found this relationship to be positive [12,13], whereas others have found it to be negative [14,15]. This association has not been explored specifically among older Asian American individuals, who are more likely to rate their health as fair or poor as compared to older adults of other ethnicities [11].

Cognitive Ability and ICT Use

Beyond self-rated health, cognitive ability is another important predictor of ICT use. When adjusting for self-rated health and demographic factors, cognitive ability was significantly, positively associated with gerontechnology use and ICT use [15]. This relationship has been confirmed in multiple studies. For example, in a cross-sectional study of adults aged ≥65 years, subjective cognitive decline was significantly negatively associated with frequency of ICT use even when adjusting for

age, educational attainment, and depression [27]. In addition, in an analysis of the National Health and Aging Trends Study, self-reported memory problems were associated with decreased email, SMS text message, and internet use even when adjusting for age, sex, race and ethnicity, educational attainment, marital status, multimorbidity, and self-rated health [28]. In this study, we examine the influence of self-rated health on ICT acceptance among Asian American older adults adjusting for subjective cognitive decline.

Research Design

This cross-sectional study examined the relationships among self-rated health, attitudinal factors (PU and PEOU), and ICT use among low-income, Asian American older adults.

In this study, we tested three hypotheses:

1. Self-rated health will be significantly, positively associated with ICT use when adjusting for age, gender, educational attainment, ethnicity, English proficiency, and subjective cognitive decline (hypothesis 1, Figure 1).
2. Self-rated health will be significantly, positively associated with PU and PEOU when adjusting for age, gender, educational attainment, ethnicity, English proficiency, and subjective cognitive decline (hypothesis 2, Figure 2).
3. Adding health variables (self-rated health and subjective cognitive decline) as main effects or interaction terms to the TAM will significantly improve model fit (hypothesis 3, Figure 3).

Figure 1. Model examining the association between self-rated health and information and communications technology (ICT) use among Asian American older adults in low-income housing to test hypothesis 1.




Figure 2. Model examining the association among self-rated health, perceived usefulness, and perceived ease of use among Asian American older adults in low-income housing to test hypothesis 2. For each dependent variable shown on the right, a separate regression analysis was run.





Figure 3. Model examining self-rated health and subjective cognitive decline as potential moderating or effect-modifying factors among perceived usefulness, perceived ease of use, demographic factors, and information and communications technology (ICT) use among Asian American older adults in low-income housing to test hypothesis 3.



In examining hypotheses 1 and 2, we investigated PU and PEOU as potential mediators of the relationship between self-rated health and ICT use. Hypothesis 3 examines self-rated health and subjective cognitive decline as potential moderating or effect-modifying factors among PU; PEOU; demographic factors; and our dependent variable, ICT use. The exploration of effect modification is novel, but we felt that it was necessary to apply this model to an older adult population. This research builds on previous work [29].

Methods

Data and Sample

This study was a secondary data analysis of the Lighthouse for Older Adults program. The Lighthouse for Older Adults program provided high-speed broadband access, ICT devices, and a series of digital literacy trainings to residents living in 8 affordable housing communities for older adults located across California. All residents (N=1050) were invited to participate. To recruit participants, on-site staff distributed flyers about the Lighthouse program, set up informational tables in high traffic areas of the community, and held social events where residents could test out a variety of ICT devices to inform device selection for the project. All Lighthouse participants were required to complete pre- and postsurveys as well as attend at least one digital literacy training course. All participants were of low income; to qualify for affordable housing for older adults, residents' incomes must be <80% of the local area median income. The surveys were evidence based, and pilot-tested with residents and staff in 5 languages. The surveys were translated and then back translated by professional translators. The final surveys were self-administered in the residents' preferred language (eg,

Korean, Chinese, Vietnamese, Japanese, Hmong, and English), with staff available to assist as needed.

This study is a cross-sectional survey analysis of the presurveys from the Lighthouse program. Participants were included in this study if they were aged ≥ 62 years and identified as being of Asian ethnicity. We excluded participants who were missing survey responses to items incorporated into the outcome variable (ICT use). The final dataset included 392 participants.

Ethical Considerations

On the basis of the Human Research Protection-210 Determination Request, the University of California, Davis, Institutional Review Board determined that this research was exempt as it did not directly involve human participants and used deidentified secondary data (1938286-1). The original informed consent allowed for secondary analysis without additional consent, and the survey results were deidentified to maintain the privacy and confidentiality of the participants. As compensation for participating in the Lighthouse program, participants received a free ICT device along with a series of in-language digital literacy training classes. A description of Lighthouse program data collection and measures has been published previously [29].

Measures

Overview

The operationalization of ICT use, attitudinal variables (PU and PEOU), and demographic variables (age, gender, educational attainment, ethnicity, and English proficiency) has been previously reported [29] and is described in Table 1. We opted to combine smartphone, computer, and tablet use into a single outcome, referred to as ICT use, given that older adults often use these devices interchangeably for various tasks [30].

Table 1. Operationalization of outcome and mediator variables.

Variable and measures	Response options	Internal consistency, Cronbach α
ICT^a use		
<ul style="list-style-type: none"> How often do you use a desktop or laptop computer? How often do you use a tablet or iPad? How often do you use a smartphone (iPhone or Android)? How long have you been using technology, such as a computer, laptop, tablet, or smartphone? 	<ul style="list-style-type: none"> 0 (“never” or “I do not own”), 1 (“once or less than once per week”), 2 (“2 to 4 times per week”), or 3 (“about once per day”) 0 (“I’ve never used these”), 1 (“less than 1 year”), 2 (“1-2 years”), or 3 (“more than 2 years”) 	0.74
Perceived usefulness		
<ul style="list-style-type: none"> Technology helps me be connected with family and friends. Technology helps me learn new information and skills. 	Scale of 1 (“strongly disagree”) to 4 (“strongly agree”)	__ ^b
Perceived ease of use		
<ul style="list-style-type: none"> I feel comfortable with technology (reverse scored). Technology makes me nervous. I don’t feel confident about my ability to use technology. Technology is confusing. I feel apprehensive about using technology. I hesitate to use technology for fear of making mistakes I cannot correct. 	Scale of 1 (“strongly agree”) to 4 (“strongly disagree”)	0.89

^aICT: information and communications technology.

^bNot applicable.

The operationalization of ICT use was assessed by 4 researchers with expertise in survey methodology with older adults, and the data were reviewed by a biostatistician and a nurse researcher. Examination of a predicted probability plot indicated normal distributions of residuals aligning closely with the plot’s diagonal line. Furthermore, a scatterplot of residuals showed no obvious pattern, with points equally distributed above and below 0 on the x-axis and to the left and right of 0 on the y-axis.

Self-Rated Health

Self-rated health was operationalized as the average of 2 survey items: “In general, how would you rate your physical health?” and “In general, how would you rate your emotional health?” Responses were scored on a scale from 1 to 5, ranging from poor to excellent. These items were adapted from the validated Patient-Reported Outcomes Measurement Information System global physical health scale [31].

Subjective Cognitive Decline

We included subjective cognitive decline as a covariate; this was measured using 1 item: “During the past 12 months, have you experienced confusion or changes in memory that is happening more often or is getting worse?” Response options were binary. This item came from the cognitive decline module of the validated Behavioral Risk Factor Surveillance System [32]. On the basis of feedback from staff at affordable housing for older adults, the item was slightly adapted to increase cultural sensitivity and minimize participant stigma by modifying the phrase “memory loss” to “changes in memory.”

Analytic Strategy

First, we computed descriptive statistics for demographics, ICT use, self-rated health, and subjective cognitive decline. Next,

we used Pearson correlation analysis to examine relationships among all dependent and independent variables.

To test hypotheses 1 and 2, we conducted linear regression using self-rated health as the independent variable. To test hypothesis 1, ICT use was examined as the dependent variable; to test hypothesis 2, PU and PEOU were examined as dependent variables. We adjusted for age, gender, educational attainment, English proficiency, ethnicity, and subjective cognitive decline in all models.

To test hypothesis 3, we conducted hierarchical multiple regression using stepwise blocks [33]. Demographic variables, PU, and PEOU were added in the first step (model 1). Health variables (self-rated health and subjective cognitive decline) were added in the second step (model 2). Finally, in model 3, interaction terms were included in a forward stepwise regression. A total of 12 interaction terms were tested, including all pairwise combinations of health variables (self-rated health and subjective cognitive decline) with demographic and attitudinal variables (age, gender, educational attainment, English proficiency, PU, and PEOU). At each step, goodness of fit (adjusted R^2) was assessed, that is, the percentage of variability in the dependent variable that could be accounted for by the predictors. With each new set of terms, the change in R^2 was calculated to quantify the change in the predictive power. In case interactions were found, contrast statements were used to estimate contrasts of interest.

To further examine the separate and joint association between age and subjective cognitive decline and ICT use, we examined scatterplots with locally estimated scatterplot smoothing lines with age as the predictor. In total, 2 scatterplots were produced, including one with a single scatterplot of data from all

participants and another set stratified by presence of subjective cognitive decline. From there, we determined that the use of 3 major age categories captured the relationships well (ages of 62-74, 75-84, and ≥ 85 years). This led to us updating our analysis of hypothesis 3 and the associated contrast statements to estimate differences in ICT use among the age categories with and without subjective cognitive decline. For all analyses, the α level for testing significance was set to .05. All data analyses were conducted using SPSS Statistics (version 28; IBM Corp).

Results

Overview

Participant demographics are described in [Table 2](#). For the sample of 392 participants, the average age was 79.1 (SD 6.95) years.

In examining the 2 health variables, most participants (270/392, 68.9%) reported fair or poor self-rated health, and 27.9% (106/380) of the participants reported subjective cognitive decline. A total of 3.1% (12/392) of the participants declined to respond to the survey question about subjective cognitive decline. The prevalence of fair or poor self-rated health and subjective cognitive decline is detailed by demographic group in [Table 3](#). Next, we used correlation analysis to further examine relationships among ICT use, PU, PEOU, health, and demographic characteristics ([Table 4](#)).

Table 2. Participant demographics (N=392)^a.

	Participants, n (%)
Age (y)	
62-74	105 (27.6)
75-84	192 (50)
85-97	84 (22)
Gender	
Women	266 (68.2)
Men	124 (31.8)
Ethnicity	
Asian: other (eg, Japanese or Hmong)	14 (3.6)
Chinese	73 (18.6)
Filipino	12 (3.1)
Korean	274 (69.9)
Vietnamese	19 (4.8)
English proficiency	
Very good	8 (2.1)
Good	63 (16.3)
Not good	198 (51.3)
None at all	117 (30.3)
Educational attainment	
Never attended school	22 (5.9)
Some high school	133 (35.4)
Completed high school or GED ^b	76 (20.2)
Some college	62 (16.5)
College degree	60 (16)
Graduate degree	23 (6.1)
Years of experience using ICTs^c	
>2	227 (57.9)
1-2	45 (11.5)
<1	30 (7.7)
I have never used these	90 (23)
Computer use	
Approximately once per day	89 (22.7)
2-4 times per week	30 (7.7)
Once or less than once per week	27 (6.9)
Never	246 (62.8)
Tablet use	
Approximately once per day	106 (27)
2-4 times per week	35 (8.9)
Once or less than once per week	20 (5.1)
Never	231 (58.9)
Smartphone use	

	Participants, n (%)
Approximately once per day	243 (62)
2-4 times per week	40 (10.2)
Once or less than once per week	17 (4.3)
Never	92 (23.5)
Number of ICT device types used (computer, tablet, or smartphone)	
No devices	78 (19.9)
1 type	129 (32.9)
2 types	77 (19.6)
3 types	108 (27.6)

^aMissing data: 11 participants did not report their age, 2 participants did not report their gender, 6 participants did not report English proficiency, and 16 participants did not report their educational attainment.

^bGED: General Educational Development.

^cICT: information and communications technology.

Table 3. Prevalence of fair or poor self-rated health and subjective cognitive decline among Asian American older adults in low-income housing by demographic group.

	Fair or poor self-rated health, n/N (%)	Subjective cognitive decline, n/N (%)
All participants	270/392 (68.9)	106/380 (27.9)
Age (y)		
62-74	63/105 (60)	16/105 (15.2)
75-84	131/192 (68.2)	57/186 (30.6)
85-97	66/84 (78.6)	30/81 (37)
Gender		
Men	74/124 (59.7)	25/121 (20.7)
Women	194/266 (72.9)	81/257 (31.5)
Educational attainment		
High school degree or lower	173/231 (74.9)	63/226 (27.9)
Some college or higher	83/145 (57.2)	39/139 (28.1)
English proficiency		
Limited or no English	240/315 (76.2)	93/305 (30.5)
Speak English well or very well	27/71 (38)	11/69 (15.9)
Ethnicity		
Asian: other	9/14 (64.3)	2/12 (16.7)
Chinese	56/73 (76.7)	21/70 (30)
Filipino	4/12 (33.3)	1/11 (9.1)
Korean	190/274 (69.3)	76/268 (28.4)
Vietnamese	11/19 (57.9)	6/19 (31.6)

Table 4. Correlation analysis (Pearson r and 2-tailed P value) among the research variables (N=392).

	ICT ^a use	PU ^b	PEOU ^c	Self-rated health	Subjective cognitive decline	Age	Gender (men)	Educational attainment	English proficiency
ICT use									
r	1	0.218 ^d	0.327 ^d	0.248 ^d	-0.121 ^d	-0.225 ^d	0.187 ^d	0.321 ^d	0.260 ^d
P value	— ^e	<.001	<.001	<.001	.02	<.001	<.001	<.001	<.001
PU									
r	0.218 ^d	1	0.139 ^d	0.141 ^d	-0.055	-0.087	0.019	0.129 ^d	0.064
P value	<.001	—	.007	.007	.30	.10	.72	.01	.22
PEOU									
R	0.327 ^d	0.139 ^d	1	0.191 ^d	-0.126 ^d	-0.141 ^d	0.157 ^d	0.186 ^d	0.240 ^d
P value	<.001	.007	—	<.001	.02	.006	.002	<.001	<.001
Self-rated health									
r	0.248 ^d	0.141 ^d	0.191 ^d	1	-0.327 ^d	-0.225 ^d	0.117 ^d	0.297 ^d	0.406 ^d
P value	<.001	.007	<.001	—	<.001	<.001	.02	<.001	<.001
Subjective cognitive decline									
r	-0.121 ^d	-0.055	-0.126 ^d	-0.327 ^d	1	0.198 ^d	-0.113 ^d	-0.025	-0.183 ^d
P value	.02	.30	.02	<.001	—	<.001	.03	.64	<.001
Age									
r	-0.225 ^d	-0.087	-0.141 ^d	-0.225 ^d	0.198 ^d	1	-0.009	-0.106 ^d	-0.194 ^d
P value	<.001	.10	.006	<.001	<.001	—	.86	.04	<.001
Gender (men)									
r	0.187 ^d	0.019	0.157 ^d	0.117 ^d	-0.113 ^d	-0.009	1	0.331 ^d	0.189 ^d
P value	<.001	.72	.002	.02	.03	.86	—	<.001	<.001
Educational attainment									
r	0.321 ^d	0.129 ^d	0.186 ^d	0.297 ^d	-0.025	-0.106 ^d	0.331 ^d	1	0.403 ^d
P value	<.001	.01	<.001	<.001	.64	.04	<.001	—	<.001
English proficiency									
r	0.260 ^d	0.064	0.240 ^d	0.406 ^d	-0.183 ^d	-0.194 ^d	0.189 ^d	0.403 ^d	1
P value	<.001	.22	<.001	<.001	<.001	<.001	<.001	<.001	—
Korean									
r	-0.095	-0.203 ^d	-0.131 ^d	-0.063	0.016	0.003	0.014	-0.015	0.020
P value	.06	<.001	.01	.21	.76	.96	.78	.77	.70
Chinese									
r	0.051	0.171 ^d	0.058	-0.056	0.022	0.102 ^d	0.030	-0.048	-0.280 ^d
P value	.31	.001	.25	.27	.67	.047	.56	.36	<.001
Vietnamese									
r	0.100 ^d	0.049	-0.014	0.007	0.019	-0.113 ^d	-0.001	0.002	0.079
P value	.049	.35	.79	.89	.71	.03	.98	.97	.12
Filipino									
R	0.104 ^d	0.091	0.220 ^d	0.252 ^d	-0.072	-0.077	-0.090	0.172 ^d	0.329 ^d

	ICT ^a use	PU ^b	PEOU ^c	Self-rated health	Subjective cognitive decline	Age	Gender (men)	Educational attainment	English proficiency
<i>P</i> value	.04	.08	<.001	<.001	.16	.13	.08	.001	<.001
Asian—other									
<i>r</i>	−0.084	−0.002	0.011	0.032	−0.045	−0.012	−0.013	−0.028	0.142 ^d
<i>P</i> value	.10	.97	.83	.52	.38	.81	.79	.59	.005

^aICT: information and communications technology.

^bPU: perceived usefulness.

^cPEOU: perceived ease of use.

^dThe correlation is significant at a significance level of .01 (2-tailed).

^eNot applicable.

Testing Hypothesis 1: Linear Regression Examining the Association Between Self-Rated Health and ICT Use

Self-rated health was not a significant predictor of ICT use when adjusting for age, gender, educational attainment, English

proficiency, ethnicity, and subjective cognitive decline; the standardized coefficient of self-rated health was 0.087 (95% CI −0.131 to 0.978; $P=.13$; Table 5). Therefore, our findings do not support hypothesis 1 (self-rated health will be significantly, positively associated with ICT use).

Table 5. Linear regression to examine the predictive value of self-rated health on information and communications technology (ICT) use among Asian American older adults in low-income housing adjusting for control variables (N=392)^a.

Independent variable	ICT use	
	β^b (95% CI)	<i>P</i> value
Age	−.154 (−0.137 to −0.029)	.003
Gender (men)	.070 (−0.281 to 1.412)	.19
Educational attainment	.214 (0.276-0.903)	<.001
English proficiency	.110 (−0.073 to 1.218)	.08
Korean	Reference group	Reference group
Chinese	.120 (0.184-2.198)	.02
Vietnamese	.087 (−0.176 to 3.102)	.08
Filipino	−.001 (−2.327 to 2.296)	.99
Asian—other	−.083 (−4.132 to .331)	.10
Self-rated health	.087 (−.131 to .978)	.13
Subjective cognitive decline	−.055 (−1.346 to .413)	.30

^a β values are standardized regression coefficients.

^bAdjusted $R^2=0.171$.

Testing Hypothesis 2: Linear Regression Examining Outcomes of PU and PEOU

Using multiple regression, we determined that self-rated health was not a significant predictor of PU ($\beta=.106$, 95% CI −0.022 to 0.274; $P=.10$) or PEOU ($\beta=.062$, 95% CI −0.073 to 0.231;

$P=.31$) when adjusting for age, gender, educational attainment, English proficiency, ethnicity, and subjective cognitive decline (Table 6). Our findings do not support hypothesis 2 (self-rated health will be significantly, positively associated with PU and PEOU).

Table 6. Linear regression with perceived usefulness (PU) and perceived ease of use (PEOU) as dependent variables among Asian American older adults in low-income housing (N=392)^a.

Independent variable	PU		PEOU	
	β^b (95% CI)	P value	β^c (95% CI)	P value
Age	-.049 (-0.021 to 0.008)	.38	-.059 (-0.023 to 0.006)	.27
Gender (men)	-.016 (-0.258 to 0.196)	.79	.120 (0.022-0.489)	.03
Educational attainment	.063 (-0.041 to 0.129)	.31	.037 (-0.060 to 0.114)	.54
English proficiency	.006 (-0.166 to 0.181)	.93	.120 (-0.012 to 0.343)	.07
Chinese	Reference group	Reference group	Reference group	Reference group
Korean	-.259 (-0.804 to -0.263)	<.001	-.159 (-0.629 to -0.075)	.01
Vietnamese	-.055 (-0.713 to 0.267)	.37	-.094 (-0.921 to 0.091)	.11
Filipino	.026 (-0.535 to 0.810)	.69	.151 (0.173 to 1.558)	.01
Asian: other	-.046 (-0.936 to 0.394)	.42	-.032 (-0.849 to 0.470)	.57
Self-rated health	.106 (-0.022 to 0.274)	.10	.062 (-0.073 to 0.231)	.31
Subjective cognitive decline	-.033 (-0.309 to 0.170)	.57	-.034 (-0.319 to 0.165)	.53

^a β values are standardized regression coefficients.

^bAdjusted $R^2=0.068$.

^cAdjusted $R^2=0.118$.

Testing Hypothesis 3: Hierarchical Multiple Regression

The results of the hierarchical multiple regression are shown in Table 7. In model 1, we found that PU and PEOU were significant predictors of ICT use (PU: $\beta=.560$, 95% CI 0.161-0.960; $P=.01$; and PEOU: $\beta=.888$, 95% CI 0.506-1.270; $P<.001$). In model 2, adding self-rated health and subjective cognitive decline did not significantly improve model fit for ICT use ($\Delta R^2=0.005$; $P=.15$). In model 3, of the 12 interaction terms that we examined, the interaction between subjective cognitive decline and age was the only one that was statistically significant. Therefore, contrast statements were used to estimate

contrasts of interest. Subsequently, we updated the single age variable to reflect 3 different age groups (62 to 74 years, 75 to 84 years, and ≥ 85 years) and reran the hierarchical multiple regression. In model 3, the interaction terms of subjective cognitive decline and age significantly improved model fit for ICT use ($\Delta R^2=0.011$; $P=.04$). Therefore, our findings support hypothesis 3 (adding self-rated health and subjective cognitive decline as main effects or interaction terms to the TAM will significantly improve model fit), and we reject the null hypothesis. Figure 4 depicts our final supported model of ICT acceptance.

Figure 4. Supported model of information and communications technology (ICT) acceptance among Asian American older adults in low-income housing.




Table 7. Hierarchical regression examining our adapted technology acceptance model among older Asian American individuals in low-income housing (N=392)^a.

Model and independent variable	ICT ^b use β (SE; 95% CI)	P value
Model 1^c		
Age of 62-74 years	Reference	Reference
Age of 75-84 years	-.205 (.435; -1.061 to 0.650)	.64
Age of ≥85 years	-1.215 (.541; -2.280 to -0.150)	.03
Gender (men)	.423 (.426; -0.415 to 1.261)	.32
Educational attainment	.539 (.157; 0.230-0.848)	.001
English proficiency	.579 (.308; -0.027 to 1.186)	.06
Korean	Reference	Reference
Chinese	.914 (.519; -0.108 to 1.936)	.08
Vietnamese	1.238 (.805; -0.347 to 2.823)	.13
Filipino	-1.250 (1.171; -3.553 to 1.053)	.29
Asian: other	-1.796 (1.146; -4.051 to .459)	.12
PU ^d	.560 (.203; 0.161-0.960)	.006
PEOU ^e	.888 (.194; 0.506-1.270)	<.001
Model 2 (adding self-rated health and subjective cognitive decline)^f		
Age of 62-74 years	Reference	Reference
Age of 75-84 years	-.081 (.438; -0.943 to 0.782)	.85
Age of ≥85 years	-1.063 (.545; -2.136 to 0.010)	.05
Gender (men)	.397 (.426; -0.441 to 1.235)	.35
Educational attainment	.525 (.159; 0.212-0.838)	.001
English proficiency	.390 (.323; -0.247 to 1.026)	.23
Korean	Reference	Reference
Chinese	.865 (.519; -0.156 to 1.886)	.10
Vietnamese	1.323 (.804; -0.260 to 2.906)	.10
Filipino	-1.324 (1.169; -3.624 to 0.976)	.26
Asian: other	-1.780 (1.143; -4.029 to 0.469)	.12
PU	.520 (.204; 0.120-0.921)	.01
PEOU	.855 (.194; 0.473-1.237)	<.001
Self-rated health	.374 (.275; -0.166 to 0.915)	.17
SCD ^g	-.438 (.442; -1.308 to 0.431)	.32
ΔR ²	0.005	0.15
Model 3 (adding interaction terms)^h		
Age of 62-74 years	Reference	Reference
Age of 75-84 years	.352 (.483; -0.599 to 1.303)	.47
Age of ≥85 years	-.458 (.623; -1.684 to 0.768)	.46
Gender (men)	.292 (.425; -0.545 to 1.128)	.49
Educational attainment	.566 (.159; 0.253-0.878)	<.001
English proficiency	.354 (.321; -0.278 to 0.987)	.27
Korean	Reference	Reference

Model and independent variable	ICT ^b use	
	β (SE; 95% CI)	P value
Chinese	.838 (.515; -0.176 to 1.852)	.11
Vietnamese	1.182 (.802; -0.397 to 2.761)	.14
Filipino	-1.187 (1.163; -3.474 to 1.101)	.31
Asian: other	-1.929 (1.136; -4.165 to .306)	.09
PU	.501 (.202; 0.102-0.899)	.01
PEOU	.856 (.193; 0.476-1.236)	<.001
Self-rated health	.356 (.273; -0.181 to 0.893)	.19
SCD	1.596 (.913; -0.201 to 3.393)	.08
SCD \times age of 62-74 years	Reference	Reference
SCD \times age of 75-84 years	-2.469 (1.074; -4.582 to -.357)	.02
SCD \times age of \geq 85 years	-2.920 (1.244; -5.367 to -.473)	.02
ΔR^2	0.011	0.04

^a β values are unstandardized regression coefficients.

^bICT: information and communications technology.

^cAdjusted $R^2=0.216$.

^dPU: perceived usefulness.

^ePEOU: perceived ease of use.

^fAdjusted $R^2=0.221$.

^gSCD: subjective cognitive decline.


^hAdjusted $R^2=0.232$.

Subanalysis of Interaction Between Subjective Cognitive Decline and Age

The relationship between subjective cognitive decline and ICT use varied in 3 age segments (Figure 5). In the youngest age

group (62 to 74 years), ICT use increased with the presence of subjective cognitive decline, whereas in the older age groups (75 to 84 years and \geq 85 years), ICT use decreased with the presence of subjective cognitive decline, more so in the oldest age category.

Figure 5. Association between information and communications technology (ICT) use and subjective cognitive decline by age group among Asian American older adults in low-income housing.



Using regression analysis, we determined that, among participants with subjective cognitive decline, ICT use significantly decreased in the middle and older age groups as

compared to the youngest age group. However, among participants without subjective cognitive decline, the difference in use among age groups was not significant (Table 8).

Table 8. Contrasts estimating differences in information and communications technology (ICT) use among age categories with and without subjective cognitive decline^a.

	ICT use	
	Estimate (95% CI)	P value
With subjective cognitive decline		
Age of 75-84 years vs age of 62-74 years	-2.509 (-4.906 to -0.111)	.04
Age of ≥85 years vs age of 62-74 years	-3.852 (-6.306 to -1.399)	.002
Without subjective cognitive decline		
Age of 75-84 years vs age of 62-74 years	0.022 (-0.939 to 0.984)	.96
Age of ≥85 years vs age of 62-74 years	-0.755 (-2.010 to 0.499)	.24

^aModel adjusted for gender, educational attainment, English proficiency, ethnicity, and self-rated health.

Discussion

Principal Findings

Overview

In this study, we aimed to investigate the relationship among self-rated health, ICT use, PU, and PEOU in a sample of low-income, Asian American older adults living in affordable housing for older adults. We also explored whether the addition of health variables (self-rated health and subjective cognitive decline) would improve TAM model fit. We discovered that the interaction between age and subjective cognitive decline significantly predicted ICT use, which prompted further subanalyses to better understand the direction and strength of these relationships.

Self-Rated Health Among Low-Income, Asian American Older Adults

Most of our participants (270/392, 68.9%) reported fair or poor health; this prevalence is nearly 3 times greater than that of all adults aged ≥65 years in California (23.5%) and more than double the prevalence among older Asian Californians (33.1%) [34]. However, according to the CHIS, among older Asian Californians living at or below the federal poverty line, the prevalence of fair or poor health mirrors that of our study population. In 2020 and 2021, a total of 56.9% of Asian American older adults and 75.3% of Korean, 63.7% of Vietnamese, and 51.3% of Chinese individuals aged ≥65 years in the low-income bracket in California reported fair or poor health [34].

In our study population, correlational analysis revealed that older age, lower educational attainment, LEP, and subjective cognitive decline were each significantly and positively associated with poor self-rated health. These findings are consistent with those of other larger studies of older adults in both China and the United States [18,35-37].

Associations Among Self-Rated Health, PU, PEOU, and ICT Use

In correlational analysis, self-rated health was also significantly positively associated with PU, PEOU, and ICT use (Table 4). However, significance was not observed when we adjusted for demographic and health factors (Tables 5-7). These findings indicate that factors outside of self-rated health may be greater drivers of technology attitudes and use. Our results are consistent with the findings of the cross-sectional survey by Werner et al [12] involving ethnically diverse, low-income adults aged ≥60 years in California, observing a significant positive association between general health and computer use. However, this relationship lost significance when adjustments were made for demographic and health factors [12]. Initial correlation analyses in other studies also revealed positive associations between self-rated health and ICT use, but these associations became mixed or even reversed directions (eg, changed to a negative association) when adjusting for a variety of covariates, including demographic factors; health factors; and TAM constructs such as PU, PEOU, relevance, and attitude [13,15,36].

Significant Interaction Among Age, Subjective Cognitive Decline, and Technology Use

Despite these findings, adding health covariates significantly improved the TAM's ability to predict ICT use. In particular, the interaction between subjective cognitive decline and age significantly improved model fit. This interaction emerged as an important aspect of our analysis.

Our findings for the youngest age group contradict those of prevailing studies that indicate a negative association between cognitive decline and ICT use [38-41]. In previous studies, mild cognitive impairment has been associated with greater computer-related challenges, reduced frequency and duration of computer use, less accurate mouse use, and longer pauses between mouse movements [39,40].

One possible explanation for our findings is that younger participants, having more lifetime technology exposure, found using ICT devices less cognitively challenging compared to

older participants with less previous technology exposure [42-44]. Therefore, the presence of subjective cognitive decline among younger participants may have less of an impact on their actual ICT use. There is evidence suggesting that younger individuals with dementia or mild cognitive impairment have more positive attitudes toward technology as compared to their older counterparts [45].

In addition, it is possible that younger participants with subjective cognitive decline may use ICTs to compensate for cognitive changes. The compensatory use of ICTs is a growing area of research, with a recent systematic review indicating that ICTs can effectively improve prospective memory, task execution, and time management among individuals with impaired cognition [46].

Our contrast analysis indicated that ICT use significantly decreased with age among participants with subjective cognitive decline but not among those without subjective cognitive decline. Our dataset did not indicate the level of severity of cognitive decline, although it is possible that older participants that reported subjective cognitive decline had worse cognition than younger participants who reported subjective cognitive decline. All participants lived independently, although staff at their housing communities described observing cognitive decline among some of the oldest residents. On the basis of previous studies [38-41], it is plausible that older participants with worse cognition would use ICTs less as compared to younger participants with less severe cognitive changes.

As our study was cross-sectional, we cannot determine whether decreasing cognition led to less ICT use or whether more frequent ICT use led to a decrease in cognitive decline. Longitudinal studies suggest that ICT use reduces cognitive decline over time or even improves cognitive function [47], including executive function [48], memory performance [38,48], and the ability to switch between tasks [49]. One study even identified positive associations between daily computer use, hippocampal volume, and memory and executive function [41]. In addition, intervention studies suggest that training older adults to use tablet computers can improve cognitive capacities such as processing speed and episodic memory [50,51].

Of note, within our study population, subjective cognitive decline was significantly associated with English proficiency. In addition, both subjective cognitive decline and English proficiency were significantly associated with age, ICT use, and PEOU. While few studies have delved into the association between subjective cognitive decline and LEP, one study suggests that older Chinese American individuals with LEP are more likely to report subjective cognitive decline than their English-speaking counterparts [52]. Further research is needed to understand the intricate relationships among age, subjective cognitive decline, English proficiency, and ICT use. Cognitive decline experienced concurrently with language barriers may pose unique challenges to technology use among older adults.

Complexity of the Relationship Between Health and ICT Use

The relationship between health and ICT use is complex. Cognitive abilities or physical health can be both a precursor

and outcome of ICT use. Increasingly, ICTs modulate access to health care, services, information, and social interaction, thereby influencing overall health outcomes. Individuals who do not use ICTs face limitations in accessing telemedicine, web-based prescription refills, web-based health information, and web-based management of Medicare and other types of health insurance. Moreover, ICT use may have adverse effects on health, particularly if it exposes older adults to scams, fraud, or identity theft. The internet offers unlimited access to information—some of it reliable, but much of it being false. In cases in which individuals struggle to discern trustworthy information, the internet can become a source of confusion or, even worse, misinformation.

Study Limitations

Our study has several limitations. The data represent a specific population of Asian American older adults; most participants were Korean, and all participants lived in an age-limited setting (affordable housing for older adults), limiting the generalizability of our findings. However, few studies exist targeting this significant Asian population group, enhancing within-group understanding. The use of self-reported measures, including self-rated health and subjective cognitive decline, could introduce response bias.

Due to low educational attainment among study participants, validated survey measures were shortened and adapted to the population. In this study, we operationalized TAM constructs based on elements analogous to the original TAM measures. Nevertheless, the comparability of results across studies is challenging due to variations in outcomes, the operationalization of constructs, and the inclusion of diverse covariates in regression analyses.

Despite these limitations, our sample was unique in that all participants were of low income and represented various Asian ethnicities and most participants had LEP. Our findings have implications for promoting digital engagement among low-income Asian American older adults. Our findings indicate that health-related factors, particularly subjective cognitive decline, can play a role in shaping technology adoption and use patterns. Tailored interventions and support strategies should consider the age-related nuances in the relationship between cognitive decline and technology use.

Enhancing factors such as PEOU and PU through digital literacy training and user interface customization can be particularly beneficial for increasing ICT acceptance among this population [53-58]. Demonstrating culturally relevant use cases for technology, such as connecting with overseas relatives, accessing health information, or enjoying entertainment in one's native language, can make technology more meaningful for Asian American older adults. For example, the Lighthouse for Older Adults program, which provided the data for this study, improved technology adoption in affordable housing for older adults by offering equipment, training, and support, thereby fostering a community of learners who benefit from shared experiences [59].

The study population had very low English proficiency, and most technology uses Roman characters and is English based.

Addressing language barriers is crucial for promoting ICT acceptance and use among low-income, Asian American older adults. Expanding the language capabilities of voice interfaces such as Google Voice and Alexa can make these technologies more user-friendly for those who do not write, read, or speak English [60,61].

Furthermore, the costs of internet access and devices can hinder ICT use among low-income older adults. During the COVID-19 pandemic, several policies were implemented to address these financial barriers. The Accessible, Affordable Internet for All Act of 2021 allocated US \$94 billion for broadband infrastructure and subsidies [62]; the Infrastructure Investment and Jobs Act of 2021 set aside US \$65 billion for internet access, digital literacy, and rural expansion [63]; and the Affordable Connectivity Program offered low-income families a US \$30 monthly subsidy toward their internet costs and required internet service providers to be transparent about pricing [63]. Although the Affordable Connectivity Program ended in February 2024, the Lifeline support program now offers families a smaller

subsidy. New state programs are emerging, but continued advocacy is necessary to improve the affordability of digital access.

Future research could further explore the reasons behind the age-dependent effects of cognitive decline on technology use and examine potential interventions to enhance technology accessibility as well as digital literacy and technology adoption among older adults with varying health profiles. Longitudinal studies and intervention designs could provide a deeper understanding of these dynamics.

Conclusions

This study contributes to our understanding of the intricate relationship among self-rated health, subjective cognitive decline, and the acceptance and use of ICTs among low-income, Asian American older adults. By highlighting the age-specific effects of cognitive decline on technology adoption, our findings can inform targeted interventions to promote digital engagement and enhance the quality of life of aging populations.

Acknowledgments

This work was funded by the Healthy Aging in a Digital World initiative at the University of California, Davis, and the Center for Information Technology Research in the Interest of Society and the Banatao Institute. The University of California, Davis, School of Medicine Office of Research Evaluation Unit, including Rebeca Giacinto, Pauline DeLange Martinez, Stuart Henderson, and Julie Rainwater, designed the Lighthouse surveys and provided oversight for data collection and entry. The authors thank all Lighthouse program participants for their time and effort to complete the evaluation surveys. They also thank the staff at each site for their support with participant recruitment and data collection and entry. Lighthouse project leadership included the Front Porch Center for Innovation and Wellbeing (Davis Park, Kari Olson, and Jennifer Lee), Eskaton (Therese ten Brinke and Sheri Peifer), and the Center for Information Technology Research in the Interest of Society (David Lindeman and Gale Berkowitz). Finally, the authors thank Lisa Miller and Ladson Hinton for contributing to the conceptualization of their research design.

Data Availability

The datasets generated during and analyzed during this study are available from the corresponding author on reasonable request.

Authors' Contributions

PDM led conceptual design, data analysis, and authorship. HMY made substantial conceptual and design contributions, providing invaluable input along all stages of manuscript preparation. DT provided oversight for operationalization of key variables, data analysis, reporting results, and interpretation of the data. MP and LG helped critically revise the manuscript and agreed to be accountable for the accuracy of the work.

Conflicts of Interest

None declared.

References

1. Wu B, Qi X. Addressing health disparities among older Asian American populations: research, data, and policy. *Public Policy Aging Rep.* 2022;32(3):105-111. [FREE Full text] [doi: [10.1093/ppar/prac015](https://doi.org/10.1093/ppar/prac015)] [Medline: [35992733](https://pubmed.ncbi.nlm.nih.gov/35992733/)]
2. Budiman A, Ruiz NG. Key facts about Asian Americans, a diverse and growing population. Pew Research Center. Apr 19, 2021. URL: <https://www.pewresearch.org/short-reads/2021/04/29/key-facts-about-asian-americans/> [accessed 2021-12-30]
3. Budiman A, Ruiz NG. Asian Americans are the fastest-growing racial or ethnic group in the U.S. Pew Research Center. Apr 9, 2021. URL: <https://www.pewresearch.org/short-reads/2021/04/09/asian-americans-are-the-fastest-growing-racial-or-ethnic-group-in-the-u-s/> [accessed 2024-11-14]
4. Vespa J, Armstrong DM, Medina L. Demographic turning points for the United States: population projections for 2020 to 2060. United States Census Bureau. Feb 2020. URL: <https://www.census.gov/library/publications/2020/demo/p25-1144.html> [accessed 2024-11-25]

5. Doãn LN, Takata Y, Sakuma KL, Irvin VL. Trends in clinical research including Asian American, Native Hawaiian, and Pacific Islander participants funded by the US National Institutes of Health, 1992 to 2018. *JAMA Netw Open*. Jul 03, 2019;2(7):e197432. [FREE Full text] [doi: [10.1001/jamanetworkopen.2019.7432](https://doi.org/10.1001/jamanetworkopen.2019.7432)] [Medline: [31339543](https://pubmed.ncbi.nlm.nih.gov/31339543/)]
6. Ghosh C. Asian American health research: baseline data and funding. In: Trinh-Shevrin C, Islam NS, Rey MJ, editors. *Asian American Communities and Health: Context, Research, Policy, and Action*. Hoboken, NJ. John Wiley & Sons; 2009:73-106.
7. Jang Y, Kim MT. Limited English proficiency and health service use in Asian Americans. *J Immigr Minor Health*. Apr 24, 2019;21(2):264-270. [FREE Full text] [doi: [10.1007/s10903-018-0763-0](https://doi.org/10.1007/s10903-018-0763-0)] [Medline: [29797103](https://pubmed.ncbi.nlm.nih.gov/29797103/)]
8. Yoon H, Jang Y, Kim S, Speasmaker A, Nam I. Trends in internet use among older adults in the United States, 2011-2016. *J Appl Gerontol*. May 2021;40(5):466-470. [doi: [10.1177/0733464820908427](https://doi.org/10.1177/0733464820908427)] [Medline: [32131670](https://pubmed.ncbi.nlm.nih.gov/32131670/)]
9. Kim J, Lee HY. Older minority immigrants' digital technology use: does English proficiency matter? *Gerontologist*. Nov 2016;56(Suppl_3):139. [doi: [10.1093/geront/gnw162.557](https://doi.org/10.1093/geront/gnw162.557)]
10. Yoon H, Jang Y, Vaughan PW, Garcia M. Older adults' internet use for health information: digital divide by race/ethnicity and socioeconomic status. *J Appl Gerontol*. Jan 2020;39(1):105-110. [doi: [10.1177/0733464818770772](https://doi.org/10.1177/0733464818770772)] [Medline: [29661052](https://pubmed.ncbi.nlm.nih.gov/29661052/)]
11. Du Y, Xu Q. Health disparities and delayed health care among older adults in California: a perspective from race, ethnicity, and immigration. *Public Health Nurs*. Sep 16, 2016;33(5):383-394. [doi: [10.1111/phn.12260](https://doi.org/10.1111/phn.12260)] [Medline: [26990795](https://pubmed.ncbi.nlm.nih.gov/26990795/)]
12. Werner JM, Carlson M, Jordan-Marsh M, Clark F. Predictors of computer use in community-dwelling, ethnically diverse older adults. *Hum Factors*. Oct 2011;53(5):431-447. [FREE Full text] [doi: [10.1177/0018720811420840](https://doi.org/10.1177/0018720811420840)] [Medline: [22046718](https://pubmed.ncbi.nlm.nih.gov/22046718/)]
13. Sims T, Reed AE, Carr DC. Information and communication technology use is related to higher well-being among the oldest-old. *J Gerontol B Psychol Sci Soc Sci*. Sep 01, 2017;72(5):761-770. [doi: [10.1093/geronb/gbw130](https://doi.org/10.1093/geronb/gbw130)] [Medline: [27702839](https://pubmed.ncbi.nlm.nih.gov/27702839/)]
14. Nayak LU, Priest L, White AP. An application of the technology acceptance model to the level of internet usage by older adults. *Univ Access Inf Soc*. Feb 16, 2010;9(4):367-374. [doi: [10.1007/s10209-009-0178-8](https://doi.org/10.1007/s10209-009-0178-8)]
15. Chen K, Chan AH. Gerontechnology acceptance by elderly Hong Kong Chinese: a senior technology acceptance model (STAM). *Ergonomics*. Mar 24, 2014;57(5):635-652. [doi: [10.1080/00140139.2014.895855](https://doi.org/10.1080/00140139.2014.895855)] [Medline: [24655221](https://pubmed.ncbi.nlm.nih.gov/24655221/)]
16. Stanojevic Jerkovic O, Sauliune S, Šumskas L, Birt CA, Kersnik J. Determinants of self-rated health in elderly populations in urban areas in Slovenia, Lithuania and UK: findings of the EURO-URHIS 2 survey. *Eur J Public Health*. May 01, 2017;27(suppl_2):74-79. [doi: [10.1093/eurpub/ckv097](https://doi.org/10.1093/eurpub/ckv097)] [Medline: [26163468](https://pubmed.ncbi.nlm.nih.gov/26163468/)]
17. Benyamini Y. Why does self-rated health predict mortality? An update on current knowledge and a research agenda for psychologists. *Psychol Health*. Nov 2011;26(11):1407-1413. [doi: [10.1080/08870446.2011.621703](https://doi.org/10.1080/08870446.2011.621703)] [Medline: [22111660](https://pubmed.ncbi.nlm.nih.gov/22111660/)]
18. Kim MJ, Gorman BK. Acculturation and self-rated health among Asian immigrants: the role of gender and age. *Popul Res Policy Rev*. Feb 20, 2021;41(1):89-114. [doi: [10.1007/s11113-021-09642-x](https://doi.org/10.1007/s11113-021-09642-x)]
19. Jang Y, Yoon H, Li M, Park NS, Chiriboga DA, Wu B, et al. Self-rated health as a mediator between physical health conditions and depressive symptoms in older Chinese and Korean Americans. *PLoS One*. 2021;16(1):e0245136. [FREE Full text] [doi: [10.1371/journal.pone.0245136](https://doi.org/10.1371/journal.pone.0245136)] [Medline: [33417597](https://pubmed.ncbi.nlm.nih.gov/33417597/)]
20. Nguyen D. Acculturation and perceived mental health need among older Asian immigrants. *J Behav Health Serv Res*. Oct 20, 2011;38(4):526-533. [doi: [10.1007/s11414-011-9245-z](https://doi.org/10.1007/s11414-011-9245-z)] [Medline: [21598084](https://pubmed.ncbi.nlm.nih.gov/21598084/)]
21. National Academies of Sciences, Engineering, and Medicine. *Social Isolation and Loneliness in Older Adults: Opportunities for the Health Care System*. Washington, DC. National Academies Press; 2020.
22. Miyawaki CE. Association of social isolation and health across different racial and ethnic groups of older Americans. *Ageing Soc*. Nov 2015;35(10):2201-2228. [FREE Full text] [doi: [10.1017/S0144686X14000890](https://doi.org/10.1017/S0144686X14000890)] [Medline: [26494934](https://pubmed.ncbi.nlm.nih.gov/26494934/)]
23. Tran V. Asian American seniors are often left out of the national conversation on poverty. Urban Institute. May 31, 2017. URL: <https://www.urban.org/urban-wire/asian-american-seniors-are-often-left-out-national-conversation-poverty> [accessed 2024-01-21]
24. Byun M, Kim E, Ahn H. Factors contributing to poor self-rated health in older adults with lower income. *Healthcare (Basel)*. Nov 06, 2021;9(11):1515. [FREE Full text] [doi: [10.3390/healthcare9111515](https://doi.org/10.3390/healthcare9111515)] [Medline: [34828561](https://pubmed.ncbi.nlm.nih.gov/34828561/)]
25. Anderson M, Perrin A. Tech adoption climbs among older adults. Pew Research Center. May 17, 2017. URL: <https://www.pewresearch.org/internet/2017/05/17/tech-adoption-climbs-among-older-adults/> [accessed 2024-03-20]
26. Davis FD. Perceived usefulness, perceived ease of use, and user acceptance of information technology. *MIS Q*. Sep 1989;13(3):319-340. [doi: [10.2307/249008](https://doi.org/10.2307/249008)]
27. Bengte JF, Kiselica AM, Aguirre A, Hilsabeck RC, Douglas M, Paydarfar D, et al. Technology use and subjective cognitive concerns in older adults. *Arch Gerontol Geriatr*. Mar 2023;106:104877. [FREE Full text] [doi: [10.1016/j.archger.2022.104877](https://doi.org/10.1016/j.archger.2022.104877)] [Medline: [36459914](https://pubmed.ncbi.nlm.nih.gov/36459914/)]
28. Gell NM, Rosenberg DE, Demiris G, LaCroix AZ, Patel KV. Patterns of technology use among older adults with and without disabilities. *Gerontologist*. Jun 2015;55(3):412-421. [FREE Full text] [doi: [10.1093/geront/gnt166](https://doi.org/10.1093/geront/gnt166)] [Medline: [24379019](https://pubmed.ncbi.nlm.nih.gov/24379019/)]

29. DeLange Martinez P, Tancredi DJ, Pavel M, Garcia L, Young HM. Technology Acceptance Among Low-Income Asian American Older Adults: Cross-Sectional Survey Analysis. *J Med Internet Res*. Nov 22, 2024;26:e52498. [FREE Full text] [doi: [10.2196/52498](https://doi.org/10.2196/52498)] [Medline: [39576987](https://pubmed.ncbi.nlm.nih.gov/39576987/)]
30. Kononova A, Joshi P, Cotten S. Contrary to myth, older adults multitask with media and technologies, but studying their multitasking behaviors can be challenging. *Innov Aging*. Aug 2019;3(4):igz029. [FREE Full text] [doi: [10.1093/geroni/igz029](https://doi.org/10.1093/geroni/igz029)] [Medline: [31637313](https://pubmed.ncbi.nlm.nih.gov/31637313/)]
31. Hays RD, Schalet BD, Spritzer KL, Cella D. Two-item PROMIS® global physical and mental health scales. *J Patient Rep Outcomes*. 2017;1(1):2. [FREE Full text] [doi: [10.1186/s41687-017-0003-8](https://doi.org/10.1186/s41687-017-0003-8)] [Medline: [29757325](https://pubmed.ncbi.nlm.nih.gov/29757325/)]
32. Behavioral risk factor surveillance system: BRFSS questionnaires. Centers for Disease Control and Prevention. URL: <https://www.cdc.gov/brfss/questionnaires/index.htm> [accessed 2024-11-25]
33. Hierarchical regression in SPSS. IBM. URL: <https://www.ibm.com/support/pages/node/418249> [accessed 2024-05-24]
34. California Health Interview Survey (CHIS): questionnaires and survey topics. UCLA Center for Health Policy Research. URL: <https://healthpolicy.ucla.edu/our-work/california-health-interview-survey-chis/chis-design-and-methods/survey-topics-and-questionnaires> [accessed 2024-11-25]
35. Xu D, Arling G, Wang K. A cross-sectional study of self-rated health among older adults: a comparison of China and the United States. *BMJ Open*. Jul 31, 2019;9(7):e027895. [FREE Full text] [doi: [10.1136/bmjopen-2018-027895](https://doi.org/10.1136/bmjopen-2018-027895)] [Medline: [31371288](https://pubmed.ncbi.nlm.nih.gov/31371288/)]
36. Liang X, Xiong F, Xie F. The effect of smartphones on the self-rated health levels of the elderly. *BMC Public Health*. Mar 15, 2022;22(1):508. [FREE Full text] [doi: [10.1186/s12889-022-12952-0](https://doi.org/10.1186/s12889-022-12952-0)] [Medline: [35292010](https://pubmed.ncbi.nlm.nih.gov/35292010/)]
37. Lee J, Jang SN. Have changes in internet use during the COVID-19 pandemic affected older adults' self-rated health? A cross-sectional study of young-old and old-old populations in Korea. *Geriatr Nurs*. 2022;48:145-149. [FREE Full text] [doi: [10.1016/j.gerinurse.2022.09.012](https://doi.org/10.1016/j.gerinurse.2022.09.012)] [Medline: [36219933](https://pubmed.ncbi.nlm.nih.gov/36219933/)]
38. Choi EY, Wisniewski KM, Zelinski EM. Information and communication technology use in older adults: a unidirectional or bi-directional association with cognitive function? *Comput Human Behav*. Aug 2021;121:106813. [FREE Full text] [doi: [10.1016/j.chb.2021.106813](https://doi.org/10.1016/j.chb.2021.106813)] [Medline: [33986562](https://pubmed.ncbi.nlm.nih.gov/33986562/)]
39. Kaye J, Mattek N, Dodge HH, Campbell I, Hayes T, Austin D, et al. Unobtrusive measurement of daily computer use to detect mild cognitive impairment. *Alzheimers Dement*. Jan 2014;10(1):10-17. [FREE Full text] [doi: [10.1016/j.jalz.2013.01.011](https://doi.org/10.1016/j.jalz.2013.01.011)] [Medline: [23688576](https://pubmed.ncbi.nlm.nih.gov/23688576/)]
40. Seelye A, Hagler S, Mattek N, Howieson DB, Wild K, Dodge HH, et al. Computer mouse movement patterns: a potential marker of mild cognitive impairment. *Alzheimers Dement (Amst)*. Dec 01, 2015;1(4):472-480. [FREE Full text] [doi: [10.1016/j.dadm.2015.09.006](https://doi.org/10.1016/j.dadm.2015.09.006)] [Medline: [26878035](https://pubmed.ncbi.nlm.nih.gov/26878035/)]
41. Silbert LC, Dodge HH, Lahna D, Promjunyakul NO, Austin D, Mattek N, et al. Less daily computer use is related to smaller hippocampal volumes in cognitively intact elderly. *J Alzheimers Dis*. 2016;52(2):713-717. [FREE Full text] [doi: [10.3233/JAD-160079](https://doi.org/10.3233/JAD-160079)] [Medline: [26967228](https://pubmed.ncbi.nlm.nih.gov/26967228/)]
42. Berkowsky RW, Rikard RV, Cotten SR. Signing off: predicting discontinued ICT usage among older adults in assisted and independent living. In: *Proceedings of the First International Conference on Human Aspects of IT for the Aged Population*. 2015. Presented at: ITAP 2015; August 2-7, 2015; Los Angeles, CA. [doi: [10.1007/978-3-319-20913-5_36](https://doi.org/10.1007/978-3-319-20913-5_36)]
43. Cotten SR. Examining the roles of technology in aging and quality of life. *J Gerontol B Psychol Sci Soc Sci*. Sep 01, 2017;72(5):823-826. [doi: [10.1093/geronb/gbx109](https://doi.org/10.1093/geronb/gbx109)] [Medline: [28817928](https://pubmed.ncbi.nlm.nih.gov/28817928/)]
44. Rikard RV, Berkowsky RW, Cotten SR. Discontinued information and communication technology usage among older adults in continuing care retirement communities in the United States. *Gerontology*. 2018;64(2):188-200. [FREE Full text] [doi: [10.1159/000482017](https://doi.org/10.1159/000482017)] [Medline: [29130976](https://pubmed.ncbi.nlm.nih.gov/29130976/)]
45. Guzman-Parra J, Barnestein-Fonseca P, Guerrero-Pertiñez G, Anderberg P, Jimenez-Fernandez L, Valero-Moreno E, et al. Attitudes and use of information and communication technologies in older adults with mild cognitive impairment or early stages of dementia and their caregivers: cross-sectional study. *J Med Internet Res*. Jun 01, 2020;22(6):e17253. [FREE Full text] [doi: [10.2196/17253](https://doi.org/10.2196/17253)] [Medline: [32442136](https://pubmed.ncbi.nlm.nih.gov/32442136/)]
46. Brandt Å, Jensen MP, Sørberg MS, Andersen SD, Sund T. Information and communication technology-based assistive technology to compensate for impaired cognition in everyday life: a systematic review. *Disabil Rehabil Assist Technol*. Oct 14, 2020;15(7):810-824. [doi: [10.1080/17483107.2020.1765032](https://doi.org/10.1080/17483107.2020.1765032)] [Medline: [32407217](https://pubmed.ncbi.nlm.nih.gov/32407217/)]
47. Kamin ST, Lang FR. Cognitive functions buffer age differences in technology ownership. *Gerontology*. Aug 19, 2016;62(2):238-246. [doi: [10.1159/000437322](https://doi.org/10.1159/000437322)] [Medline: [26304629](https://pubmed.ncbi.nlm.nih.gov/26304629/)]
48. Kesse-Guyot E, Charreire H, Andreeva VA, Touvier M, Hercberg S, Galan P, et al. Cross-sectional and longitudinal associations of different sedentary behaviors with cognitive performance in older adults. *PLoS One*. Oct 17, 2012;7(10):e47831. [FREE Full text] [doi: [10.1371/journal.pone.0047831](https://doi.org/10.1371/journal.pone.0047831)] [Medline: [23082222](https://pubmed.ncbi.nlm.nih.gov/23082222/)]
49. Tun PA, Lachman ME. The association between computer use and cognition across adulthood: use it so you won't lose it? *Psychol Aging*. Sep 2010;25(3):560-568. [FREE Full text] [doi: [10.1037/a0019543](https://doi.org/10.1037/a0019543)] [Medline: [20677884](https://pubmed.ncbi.nlm.nih.gov/20677884/)]
50. Chan MY, Haber S, Drew LM, Park DC. Training older adults to use tablet computers: does it enhance cognitive function? *Gerontologist*. Jun 2016;56(3):475-484. [FREE Full text] [doi: [10.1093/geront/gnu057](https://doi.org/10.1093/geront/gnu057)] [Medline: [24928557](https://pubmed.ncbi.nlm.nih.gov/24928557/)]

51. Vaportzis E, Martin M, Gow AJ. A tablet for healthy ageing: the effect of a tablet computer training intervention on cognitive abilities in older adults. *Am J Geriatr Psychiatry*. Aug 2017;25(8):841-851. [FREE Full text] [doi: [10.1016/j.jagp.2016.11.015](https://doi.org/10.1016/j.jagp.2016.11.015)] [Medline: [28082016](https://pubmed.ncbi.nlm.nih.gov/28082016/)]
52. Li C, Zeng X, Yang X, Zhu CW, Neugroschl JA, Aloysi A, et al. Subjective cognitive complaints in Chinese - and English - speaking older adults. *Alzheimers Dement*. Dec 31, 2021;17(S6):e054658. [doi: [10.1002/alz.054658](https://doi.org/10.1002/alz.054658)]
53. Arthanat S. Promoting information communication technology adoption and acceptance for aging-in-place: a randomized controlled trial. *J Appl Gerontol*. May 29, 2021;40(5):471-480. [FREE Full text] [doi: [10.1177/0733464819891045](https://doi.org/10.1177/0733464819891045)] [Medline: [31782347](https://pubmed.ncbi.nlm.nih.gov/31782347/)]
54. Lagana L, Garcia JJ. The mental health impact of computer and internet training on a multi-ethnic sample of community-dwelling older adults: results of a pilot randomised controlled trial. *Int J Biomed Sci*. Sep 2013;9(3):135-147. [FREE Full text] [Medline: [24151452](https://pubmed.ncbi.nlm.nih.gov/24151452/)]
55. González A, Ramírez MP, Viadel V. ICT learning by older adults and their attitudes toward computer use. *Curr Gerontol Geriatr Res*. 2015;2015:849308. [FREE Full text] [doi: [10.1155/2015/849308](https://doi.org/10.1155/2015/849308)] [Medline: [26346158](https://pubmed.ncbi.nlm.nih.gov/26346158/)]
56. Senteio CR. Investigating the enduring impact of a community-based health education program to promote African American elders' use of technology designed to support chronic disease self-management. *Geriatrics (Basel)*. Oct 13, 2018;3(4):70. [FREE Full text] [doi: [10.3390/geriatrics3040070](https://doi.org/10.3390/geriatrics3040070)] [Medline: [31011105](https://pubmed.ncbi.nlm.nih.gov/31011105/)]
57. Czaja SJ, Boot WR, Charness N, Rogers WA, Sharit J. Improving social support for older adults through technology: findings from the PRISM randomized controlled trial. *Gerontologist*. May 08, 2018;58(3):467-477. [FREE Full text] [doi: [10.1093/geront/gnw249](https://doi.org/10.1093/geront/gnw249)] [Medline: [28201730](https://pubmed.ncbi.nlm.nih.gov/28201730/)]
58. Lor M, Bowers BJ. Feasibility of audio-computer-assisted self-interviewing with color-coding and helper assistance (ACASI-H) for Hmong older adults. *Res Nurs Health*. Aug 21, 2017;40(4):360-371. [FREE Full text] [doi: [10.1002/nur.21796](https://doi.org/10.1002/nur.21796)] [Medline: [28431187](https://pubmed.ncbi.nlm.nih.gov/28431187/)]
59. Giacinto RE, Martinez P, Neves S, Sullivan M, Kipke R, Henderson S. Lighthouse for older adults evaluation report. University of California, Davis, School of Medicine Office of Research Evaluation Unit. 2023. URL: <https://health.citris-uc.org/wp-content/uploads/2023/08/Lighthouse-UCD-SOMOR-Evaluation-Summary-Report-FINAL-PDF.pdf> [accessed 2024-11-25]
60. Kinsella B. Google Assistant now supports Cantonese on android and within actions on Google SDK. Voicebot.ai. Oct 23, 2019. URL: <https://voicebot.ai/2019/10/23/google-assistant-now-support-cantonese-on-android-and-within-actions-on-google-sdk/> [accessed 2024-03-18]
61. Pyae A, Scifleet P. Investigating the role of user's English language proficiency in using a voice user interface: a case of Google home smart speaker. In: Proceedings of the Extended Abstracts of the 2019 CHI Conference on Human Factors in Computing Systems. 2019. Presented at: CHI EA '19; May 4-9, 2019; Glasgow, Scotland. [doi: [10.1145/3290607.3313038](https://doi.org/10.1145/3290607.3313038)]
62. H.R.1783 - Accessible, Affordable Internet for All Act. Congress.Gov. 2021. URL: <https://www.congress.gov/bills/117/congress-117/house/bills/1783/text?q=%7B%22search%22%3A%5B%22internet+for+all%22%5D%7D&r=1&s=1> [accessed 2024-11-25]
63. High-speed internet discount program becomes permanent. AARP. URL: <https://states.aarp.org/west-virginia/high-speed-internet-discount-program-becomes-permanent> [accessed 2024-01-16]

Abbreviations

- CHIS:** California Health Interview Survey
ICT: information and communications technology
LEP: limited English proficiency
PEOU: perceived ease of use
PU: perceived usefulness
TAM: technology acceptance model

Edited by A Mavragani; submitted 01.02.24; peer-reviewed by K-C Wu, H Shin, J Freeman; comments to author 21.05.24; revised version received 06.08.24; accepted 29.10.24; published 03.12.24

Please cite as:

DeLange Martinez P, Tancredi D, Pavel M, Garcia L, Young HM

The Role of Health in the Technology Acceptance Model Among Low-Income Asian American Older Adults: Cross-Sectional Survey Analysis

JMIR Form Res 2024;8:e57009

URL: <https://formative.jmir.org/2024/1/e57009>

doi: [10.2196/57009](https://doi.org/10.2196/57009)

PMID: [39625744](https://pubmed.ncbi.nlm.nih.gov/39625744/)

©Pauline DeLange Martinez, Daniel Tancredi, Misha Pavel, Lorena Garcia, Heather M Young. Originally published in JMIR Formative Research (<https://formative.jmir.org>), 03.12.2024. This is an open-access article distributed under the terms of the Creative Commons Attribution License (<https://creativecommons.org/licenses/by/4.0/>), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work, first published in JMIR Formative Research, is properly cited. The complete bibliographic information, a link to the original publication on <https://formative.jmir.org>, as well as this copyright and license information must be included.