

# Lawrence Berkeley National Laboratory

## Lawrence Berkeley National Laboratory

### **Title**

Use of bromodeoxyuridine immunocapture to identify psychrotolerant phenanthrene-degrading bacteria in phenanthrene-enriched polluted Baltic Sea sediments

### **Permalink**

<https://escholarship.org/uc/item/0179x7qs>

### **Author**

Edlund, A.

### **Publication Date**

2009-03-04

# Use of bromodeoxyuridine immunocapture to identify psychrotolerant phenanthrene-degrading bacteria in phenanthrene-enriched polluted Baltic Sea sediments

Anna Edlund<sup>1,2</sup> & Janet K. Jansson<sup>2,3</sup>

<sup>1</sup>University College, School of Life Sciences, Huddinge, Sweden;

<sup>2</sup>Swedish University of Agricultural Sciences, Department of Microbiology, Uppsala, Sweden;

<sup>3</sup>Lawrence Berkeley National Laboratory, Earth Sciences Division, Berkeley, CA, USA

Keywords: phenanthrene; bromodeoxyuridine; terminalrestriction fragment length polymorphism; Baltic Sea sediment; Exiguobacterium

## Abstract

The aim of this study was to enrich and identify psychrotolerant phenanthrene-degrading bacteria from polluted Baltic Sea sediments. Polyaromatic hydrocarbon (PAH)-contaminated sediments were spiked with phenanthrene and incubated for 2 months in the presence of bromodeoxyuridine that is incorporated into the DNA of replicating cells. The bromodeoxyuridine-incorporated DNA was extracted by immunocapture and analyzed by terminal-restriction fragment length polymorphism and 16S rRNA gene cloning and sequencing to identify bacterial populations that were growing. In addition, degradation genes were quantified in the bromodeoxyuridine-incorporated DNA by real-time PCR. Phenanthrene concentrations decreased after 2 months of incubation in the phenanthrene-enriched sediments and this reduction correlated to increases in copy numbers of xylE and phnAc dioxygenase genes. Representatives of Exiguobacterium, Schewanella, Methylobacter, Pseudomonas, Bacteroides and an uncultured Deltaproteobacterium and a Gammaproteobacterium dominated the growing community in the phenanthrene-spiked sediments. Isolates that were closely related to three of these bacteria (two pseudomonads and an Exiguobacterium sp.) could reduce phenanthrene concentrations in pure cultures and they all harbored phnAc dioxygenase genes. These results confirm that this combination of culture-based and molecular approaches was useful for identification of actively growing bacterial species with a high potential for phenanthrene degradation.

## Introduction

High concentrations of polyaromatic hydrocarbons (PAHs) have accumulated in sediments of some coastal regions of the Baltic Sea as a result of industrial activities. For example, sediments from the bay of Örsörumsviken, located on the southeast coast of Sweden, were highly contaminated with PAHs, polychlorinated biphenyls (PCBs) and mercury (Hg) from paper mill operations between 1915 and 1980 (Palm et al., 2001). Örsörumsviken was therefore the focus of a large Swedish remediation initiative from 2001 to 2003, whereby the contaminated sediments were removed by dredging. The dredging project was successful, and only low levels of pollutants could be detected in the sediments following dredging (Edlund & Jansson, 2006). In a previous study, we found that the total and actively growing bacterial communities in

the polluted sediments in the bay before dredging were significantly different from those present in the bay after the polluted surface sediment layers were removed (Edlund & Jansson, 2006). A few bacterial isolates that could transform phenanthrene to a colored intermediate on agar medium were obtained from the polluted sediments (Edlund & Jansson, 2006).

Our aim here was to use a combination of molecular and cultivation-based approaches to further evaluate the intrinsic microbial potential for phenanthrene degradation in polluted Baltic Sea sediments. First, we enriched phenanthrene-degrading bacterial populations by spiking PAH-contaminated sediments that were collected previously from Örsörumsviken with phenanthrene. To specifically identify metabolically active bacterial populations stimulated to grow in the presence of

phenanthrene, we used bromodeoxyuridine immunocapture as a molecular tool that does not rely on cultivation for identification of growing bacteria and that was also successfully used previously for identification of active bacterial populations in sediments (Edlund & Jansson, 2006; Edlund *et al.*, 2008). The reason for using a cultivation-independent method was to overcome the known biases of cultivation-based approaches that normally detect only a fraction of the microorganisms inhabiting natural communities. Also, our recent studies using molecular approaches have shown that the microbial diversity of Baltic Sea sediments is still largely unexplored, with relatively few 16S rRNA genes representative of cultured microorganisms in existing databases (Edlund *et al.*, 2006, 2008). The bromodeoxyuridine immunocapture approach relies on the addition of bromodeoxyuridine as a thymidine analog that is incorporated into the DNA of replicating cells. The bromodeoxyuridine-incorporated DNA is extracted from bulk DNA extracts by immunocapture onto magnetic beads (Urbach *et al.*, 1999; Artursson & Jansson, 2003; Edlund & Jansson, 2006; Edlund *et al.*, 2008) and the bacterial community composition of the bromodeoxyuridine-incorporated DNA can then be determined by terminal-restriction fragment length polymorphism (T-RFLP), and cloning and sequencing of 16S rRNA genes (Edlund & Jansson, 2006; Edlund *et al.*, 2008). In addition, we aimed to quantify levels of bromodeoxyuridine-labeled functional genes reported to be involved in PAH degradation (Simon *et al.*, 1993; Kim & Zylstra, 1995; Lauri & Lloyd-Jones, 1999) by real-time PCR in the spiked sediments.

The second approach was to study isolates obtained previously from the contaminated sediments to evaluate their ability to degrade phenanthrene as a sole carbon and energy source and for the presence of PAH degradation genes in their genomes. In particular, we focused on isolates that, based on their 16S rRNA gene sequences, were represented as potentially growing members of the sediment microbiota in the phenanthrene-enriched sediments using the molecular approaches described above.

The experiments were carried out at temperatures typical of the Baltic Sea site of origin, i.e. 5 °C. The results of this study therefore provide information about the identities of psychrotolerant bacteria that have potential for phenan-

threne degradation in cold Baltic Sea sediments. We also obtained information regarding which genes and degradation pathways were present after long-term exposure to PAH compounds in cold aquatic habitats. This information is valuable for optimization of remediation of other PAH-contaminated environments in the Baltic Sea area and other cold regions.

## Materials and methods

### Sediment collection and microcosm setup

Polluted surface sediments were collected on October 5, 2002 from Örserumsviken (Edlund & Jansson, 2006) and stored frozen at –80 °C after collection. Three sediment cores from four separate sampling stations in the bay (Edlund & Jansson, 2006) were thawed at 4 °C overnight. The most dominant bacterial community members in all of these sediment cores were previously shown to be similar using a molecular fingerprinting approach (Edlund & Jansson, 2006). One hundred and thirty milliliters of sediment were taken from each of the 12 sediment cores, pooled to a volume of 1.56 L and thoroughly mixed with a spoon. Fifty-milliliter portions of the pooled sediments were aliquoted into 200-mL Pyrex flasks with tightened screw caps. Phenanthrene was dissolved in acetone and added to a subsample of sediment (25% of the total sediment volume). The acetone was evaporated before mixing with the remaining sediment, according to Brinch *et al.* (2002) to avoid impact of the solvent on the subsequent experiments. Phenanthrene was added to a final concentration of 10 µg mL<sup>-1</sup> and 1 mL of 2 mM Bromodeoxyuridine (Roche, Mannheim, Germany) was added to relevant treatments. Details of treatments and experimental replication are shown in Table 1. The sediment microcosms were incubated in the dark at 5 °C (the *in situ* temperature of the sediments at the sampling location) for a period of 60 days. Sampling of microcosms (see details below) was performed on days 0, 1, 2, 7 and 60 of incubation. All microcosms were gently stirred with a sterile spatula at each sampling period and also when bromodeoxyuridine was added.

**Table 1.** Sediment microcosm treatments

Treatment number (replicates)	Phenanthrene addition (day added)	Bromodeoxyuridine addition (day added)	Comments
1 (3)	–	–	Negative control, used for total community DNA extraction
2 (6)	–	Day 0	Control to identify active bacteria without added phenanthrene
3 (6)	Day 0	Day 0	Both phenanthrene and bromodeoxyuridine added at start of experiment
4 (3)	Day 0	Days 0, 1, 6	Test repeated doses of bromodeoxyuridine in presence of phenanthrene
5 (3)	Day 0	Day 58	Test effect of bromodeoxyuridine addition 2 days prior to last sampling date

### Quantification of phenanthrene by GC-MS

The total phenanthrene concentration was determined using the protocol developed by Pizzul *et al.* (2006) with the following modifications for pure cultures: 1 mL toluene was added to a 1-mL sample in 3-mL glass tubes with Teflon-lined caps. The tubes were shaken vigorously for 20 h at 200 r.p.m. and the extracts were centrifuged for 10 min at 492 g. After centrifugation, the supernatants were analyzed directly by GC-MS on an HP-5MS capillary column (250- $\mu\text{m}$  inner diameter, 30-m length, 0.25- $\mu\text{m}$  nominal) (Agilent Technologies, Palo Alto, CA) using an HP 6890 Mass Selective Detector. The oven program was: 150 °C for 2 min, followed by ramping at 5 °C min<sup>-1</sup> up to 250 °C maintained for 5 min. The injector temperature was 250 °C. The same procedure was used for sediment samples, except that 5 mL toluene was added to 15-mL glass tubes with Teflon-lined caps containing a 5-mL sediment sample, the supernatants were diluted 10 times in toluene before analysis and GC-MS was performed on an HP 19091S-433 capillary column (0.25 mm inner diameter, 30 m length, 0.25 cm thickness) (Agilent Technologies) using an HP 5971 Mass Selective Detector, and the oven program was: 80 °C for 4 min, followed by ramping at 7 °C min<sup>-1</sup> up to 310 °C maintained for 4 min.

Identification and quantification of phenanthrene was performed by comparisons with external standards ranging from 0.5 to 10  $\mu\text{g}$  phenanthrene mL<sup>-1</sup>. The recovery of phenanthrene in freshly spiked uninoculated controls was > 95%. The background level of phenanthrene in the sediment was 0.2  $\mu\text{g}$  mL<sup>-1</sup> sediment slurry.

### DNA extraction and bromodeoxyuridine-immunocapture

DNA from metabolically active bacteria (i.e. replicating bacteria) was extracted using the bromodeoxyuridine immunocapture approach (Urbach *et al.*, 1999; Artursson & Jansson, 2003; Edlund & Jansson, 2006; Edlund *et al.*, 2006, 2008). This method relies on incorporation of bromodeoxyuridine as a thymidine analog into growing cells during DNA replication. Total DNA was extracted from 0.5-mL sediment samples using a FastDNA<sup>®</sup>SPIN<sup>®</sup> Kit (for soil) following the manufacturer's instructions (Q-BIOgene, Carlsbad, CA). DNA with incorporated bromodeoxyuridine was purified by immunocapture using antibodies against bromodeoxyuridine (Urbach *et al.*, 1999; Artursson & Jansson, 2003; Edlund & Jansson, 2006; Edlund *et al.*, 2008). Monoclonal anti-bromodeoxyuridine antibodies (3  $\mu\text{L}$ ; Roche, Basel, Switzerland) were mixed at a 1 : 9 ratio with sheared and denatured herring sperm DNA (1.25 mg mL<sup>-1</sup> in PBS; Promega, Madison, WI) to block nonspecific binding. The mixture was incubated for 30 min at room temperature. Magnetic beads (Dynabeads) coated

with goat anti-mouse immunoglobulin G (DYNAL, Oslo, Norway) were washed three times with 1 mg mL<sup>-1</sup> acetylated bovine serum albumin (BSA) in phosphate-buffered saline (PBS) buffer (PBS-BSA) using a magnetic particle concentrator (DYNAL) and re-suspended in PBS-BSA to the initial concentration. A 25- $\mu\text{L}$  portion of sediment DNA (c. 1  $\mu\text{g}$ ) was denatured (heated for 5 min at 100 °C, quickly transferred to ice and kept on ice for 5 min), mixed with 30  $\mu\text{L}$  of the herring sperm DNA antibody mixture and incubated for 1 h in the dark at room temperature with constant agitation. The samples were mixed with 25- $\mu\text{L}$  portions of Dynabeads and the incubation was continued for an additional 1 h. After incubation, the samples containing the Dynabeads, bromodeoxyuridine-labeled DNA and unlabeled DNA were washed five times in 0.5 mL PBS-BSA. After every wash, unlabeled DNA (i.e. DNA that did not bind to the beads) was discarded. To elute the remaining bromodeoxyuridine-containing DNA fraction, 100  $\mu\text{L}$  of 1.7 mM bromodeoxyuridine (in PBS-BSA) was added, and the samples were incubated for 1 h in the dark at room temperature with constant agitation. Two microliters of glycogen (20 mg mL<sup>-1</sup>; GE Healthcare, Bio-Sciences, NJ) were added to the bead supernatants and DNA was isolated by two rounds of ethanol precipitation and dissolved in 20  $\mu\text{L}$  of sterile-distilled water. The DNA was washed using a QIAamp<sup>®</sup> DNA Micro Kit (QIAGEN, Hilden, Germany) according to the manufacturer's instructions, before PCR amplification.

The efficacy of bromodeoxyuridine immunocapture was confirmed by comparing visible bands on agarose gels after DNA extraction, immunocapture and PCR amplification with control sediment samples to which no bromodeoxyuridine was added. No PCR amplified product was visible in the controls without bromodeoxyuridine, demonstrating that the extraction was successful.

### T-RFLP

Bacterial 16S rRNA gene sequences were PCR amplified from the bromodeoxyuridine-immunocaptured DNA using the following bacterial primers: fd1-Hex AGAGTTTGAT CMTGGCTCAG (Weisburg *et al.*, 1991) labeled at the 5'-end with hexachlorofluorescein, and 926r CCGTCA ATTCCTTTRAGTTT (Muyzer *et al.*, 1995). The PCR products (12  $\mu\text{L}$ , c. 20 ng DNA) were digested for 2 h (50  $\mu\text{L}$  total volume) using HhaI restriction enzyme (Promega). The lengths of the fluorescent terminal restriction fragments (TRFs) were determined using an ABI capillary sequencer (Applied Biosystems, Foster City, CA) as described previously (Hjort *et al.*, 2007). The sizes of the TRFs were determined by comparison with the internal GS ROX-500 size standard (Applied Biosystems). Before injection, 1.4  $\mu\text{L}$  of the DNA was denatured in the presence of 10  $\mu\text{L}$  Hi-Di<sup>™</sup> formamide and 0.04  $\mu\text{L}$  GS ROX-500 size standard at 95 °C

for 5 min. Injection was performed electrokinetically at 10 kV for 50 s and the electrophoresis was run at 7.5 kV for 179 min. Relative peak areas for TRFs were determined by dividing the individual TRF area by the total area of peaks within the thresholds of 60–490 bp. Only peaks with relative fluorescence intensities over 50 U and that contributed to the total community abundance with 1.5% or more were included in the later analyses.

Diversity, defined as evenness and richness of the bacterial community members detected as TRFs by T-RFLP analysis, was calculated using Shannon's diversity and equitability index (Margalef, 1958).

### Cloning and sequencing

Bacteria that were stimulated to grow in the phenanthrene-enriched sediments after 60 days of incubation were identified by cloning and sequencing of 16S rRNA genes that were amplified from bromodeoxyuridine-immunocaptured DNA from microcosms treated with phenanthrene and bromodeoxyuridine (treatment 3; Table 1) using the same primers as in T-RFLP. The PCR amplified products from bromodeoxyuridine-immunocaptured DNA extracted from the three replicate microcosms were pooled together before library construction. Ten nanograms of the amplified, c. 900-bp, bromodeoxyuridine-incorporated 16S rRNA gene fragments were cloned using a TOPO TA cloning kit (Invitrogen, Carlsbad, CA) following the manufacturer's instructions. Cloned inserts were screened by RFLP after amplification using the M13f and M13r primers (supplied with the TOPO cloning kit) using amplification conditions according to the manufacturer's instructions. The PCR products were digested with HaeIII and RsaI (Promega) in individual reactions and then pooled together. The resulting restriction fragments were separated by gel electrophoresis in 2.2% agarose gels and the gels were stained with ethidium bromide ( $1 \text{ mg L}^{-1}$ ). 16S rRNA genes from representative clones of each RFLP group were partially sequenced (c. 900 bp) on an Applied Biosystems International PRISM

377 DNA sequencer using the M13f and M13r primers (Invitrogen). Sequences were compared with sequences in the GenBank database using BLASTN software (Altschul *et al.*, 1997). All 16S rRNA gene sequences were also tested for chimeras in Chimera Check, version 2.7, Ribosomal Database project II (Cole *et al.*, 2003). Chimeric 16S rRNA gene sequences that were detected within the data set were discarded. Cloned 16S rRNA gene sequences have been deposited in the EMBL nucleotide sequence database under the following accession numbers: AM489700–AM489706.

### Real-time (q) PCR

PCR primers used for specific detection of the following dioxygenase gene fragments: *phnAc* (144 bp), *nahAc* (212 bp), *pahAc* (139 bp), *pdoA1* (358 bp) and *xylE* (197 bp), were designed using the software BEACON DESIGNER, v. 6.0 (Premier Biosoft, International, Palo Alto, CA) (Table 2). Positive controls: *Burkholderia* sp., *Pseudomonas putida* G7, *P. putida* OUS82 and *Mycobacterium* sp. (Table 3) (Johnsen *et al.*, 2006, 2007) were a gift from Anders Johnsen, Department of Geochemistry, Geological Survey of Denmark and Greenland. The pUC19:*xylE* vector was a gift from Victor de Lorenzo, National Center for Biotechnology, Madrid, Spain. PCR reactions were individually optimized using diluted PCR products amplified from the control strains (Table 2).

Genes that gave a positive signal when amplified from environmental DNA (*xylE* and *phnAc*) were quantified by real-time qPCR. For this purpose, cleaned PCR products of *xylE* and *phnAc* genes were used as DNA standards. Clean PCR products for positive controls were obtained using the ExoSAP-IT kit (GE Healthcare) according to the manufacturer's instructions. The quality and concentration of PCR products were determined using an ND-1000 Spectrophotometer (Nanodrop Technologies, Wilmington, DE). qPCR was performed on a Light Cycler 2.0 instrument (LC, Roche). For each treatment, triplicate qPCR reactions were prepared from the bromodeoxyuridine-labeled DNA representing three separate sediment microcosms. A master mix

**Table 2.** PCR primers (constructed using software BEACON DESIGNER, v. 6.0), positive control strains and their corresponding dioxygenase genes

Primer pairs (5'–3')	Target gene and host (accession no, NCBI)	Sources
Phn_fwd: ATATATCAGGACCAGACATTATTC Phn_rev: GATGAGGTGATTGTGATGCG	Partial <i>phnAc</i> dioxygenase gene, 144 bp, <i>Burkholderia</i> sp., AF061751	Johnsen <i>et al.</i> (2006)
Nah_fwd: GCAAATGACCTCCAATACG Nah_rev: CCGAATGCCATCAACTCC	Partial <i>nahAc</i> naphthalene dioxygenase gene, 212 bp, <i>Pseudomonas putida</i> G7, AB237655	Johnsen <i>et al.</i> (2007)
Pah_fwd: AAAACTTTGTGGGTGATG Pah_rev: TATTTGGAGGTCATTTGC	Partial <i>pahAc</i> naphthalene dioxygenase gene, 139 bp, <i>Pseudomonas putida</i> OUS82, AB004059	Johnsen <i>et al.</i> (2007)
Pdo_fwd: TCATTGCGAAGAAGACAG Pdo_rev: CCGTGAATAAAGGCTGAG	Partial <i>pdoA1</i> PAH ring-hydroxylating dioxygenase Gene, 358 bp, <i>Mycobacterium</i> sp., AJ494745	Johnsen <i>et al.</i> (2007)
<i>xylE</i> _fwd: CAGCTGCGTGTACTGGACAT <i>xylE</i> _rev: AAGGTTGTGGATGAGGATG	pUC19: <i>xylE</i> vector, Partial catechol-2,3-dioxygenase gene, 197 bp	*de Lorenzo

\*The *xylE* vector was a gift from Victor de Lorenzo, National Center for Biotechnology, Madrid, Spain.

**Table 3.** Relative abundances of TRFs and 16S rRNA gene clones amplified from bromodeoxyuridine-immunocaptured DNA extracted from phenanthrene-spiked sediment after 60 days incubation

TRF length bp (Hhal)	Closest match (NCBI) and accession number	Matching isolate* and accession number	Relative abundance (%)	
			TRF <sup>†</sup>	Clone <sup>‡</sup>
60	<i>Exiguobacterium oxidotolerans</i> 99% sequence similarity to AY489701 §	<i>Exiguobacterium</i> sp. AE3 99% sequence similarity to EF530574	5.9	7.3
76	<i>Schewanella baltica</i> 99% sequence similarity to AJ000214		31.1	34.3
79	<i>Methylobacterium methanica</i> 100% sequence similarity to AF150806		30.3	24.3
97	<i>Deltaproteobacterium</i> 99% sequence similarity to DQ079636		8.7	11.4
101	<i>Bacteroidetes</i> bacterium 98% sequence similarity to DQ185611		9.4	12.8
128	Unidentified bacterium		5.0	X
205	Unclassified <i>Gammaproteobacterium</i> , 100% sequence similarity to AM167991§	<i>Pseudomonas</i> sp. AE2 98% sequence similarity to EF528261	1.8	2.8
218	<i>Pseudomonas</i> sp. 99% sequence similarity to AM167976§	<i>Pseudomonas</i> sp. AE1 97% sequence similarity to EF528260	.8	7.1

\*Strain designations given to isolates based on closest matches to their full-length 16S rRNA gene sequences.

<sup>†</sup>Data represent the average relative abundance values of TRFs from three replicate sediment microcosms.

<sup>‡</sup>16S rRNA gene sequences were digested *in silico* using the recognition site for restriction enzyme Hhal.

§Based on partial 16S rRNA gene sequences (Edlund & Jansson, 2006).

X, 16S rRNA gene sequence not obtained in clone library.

containing 3 mM MgCl<sub>2</sub>, 0.5 μM of each primer and 1 μL bovine serum albumin (BSA; 1 mg mL<sup>-1</sup>) was prepared on ice and added to a Platinum<sup>®</sup> SYBR<sup>®</sup> Green qPCR Super-Mix-UGD kit (Invitrogen). Eighteen microliters of master mix were combined with 2 μL of serially diluted template in capillaries, which were sealed and placed in the LightCycler. The following temperature profile was used for qPCR: after initial uracil DNA glycosylase (UGD) incubation at 50 °C for 120 s and at 95 °C for 120 s, amplification was carried out for 45 cycles with denaturation at 94 °C for 5 s, annealing at 60 °C for 5 s and extension at 72 °C for 8 s. The temperature transition rate was 20 °C s<sup>-1</sup>. Melting curve analysis was conducted in one cycle with three segments: 95 °C for 60 s, 60 °C for 15 s and then increased to 95 °C. The temperature transition rate was 20 °C s<sup>-1</sup> for the first and second segments and 0.1 °C for the third segment. Fluorescence measurements were taken during the final temperature ramp. Melting curves were routinely checked to confirm quantification of the desired product. To confirm that the correct genes were amplified from environmental DNA, direct sequencing of qPCR products was performed with qPCR primers (Table 2).

The numbers of gene copies in the positive controls were calculated from the formula developed previously by Whelan *et al.* (2003).

$$\text{DNA (copy)} = \frac{6.02 \times 10^{23}(\text{copy/mol}) \times \text{DNA amount (g)}}{\text{DNA length (dp)} \times (660(\text{g mol}^{-1} \text{ dp}^{-1}))}$$

DNA concentration was determined using an ND-1000 Spectrophotometer and this value was then inserted into the formula above to calculate the numbers of copies for each gene. The threshold cycle ( $C_t$ ) values for each replicate dilution were measured to generate standard curves. The  $C_t$  value for each standard was plotted against the log<sub>10</sub> of the starting DNA quantity to generate a standard curve. The standard curve for *xylE* ranged from 1 × 10<sup>3</sup> to 1 × 10<sup>7</sup> gene copies and the standard curve for *phnAc* ranged from 1 × 10<sup>3</sup> to 1 × 10<sup>9</sup> gene copies. The resulting straight-line equations that were used for calculations of copies were  $C_t = -3.03 \times \log_{10}(\text{copy number}) + 44.49$  and  $C_t = -3.06 \times \log_{10}(\text{copy number}) + 35.95$  for *xylE* and *phnAc*, respectively. The *phnAc* and *xylE* genes that were amplified and sequenced from the sediments were deposited in the NCBI nucleotide sequence database under the following accession numbers: EU251476 and EU251477, respectively.

### Phenanthrene degradation by isolates

Three bacterial strains that were previously isolated from the polluted Baltic Sea sediments after 7 days of incubation in phenanthrene-spiked microcosms (Edlund & Jansson, 2006) were assigned to the genus level based on their 16S rRNA gene sequences and given candidate strain numbers (Table 3). The full-length 16S rRNA gene sequences for the isolates were deposited in GenBank under the following accession numbers: EF528260, EF528261 and EF530574.

The isolates were tested for their ability to grow on and degrade phenanthrene in liquid cultures. Phenanthrene concentrations in the cultures were monitored using GC-MS as described above. Degradation experiments were performed in sediment extract medium to mimic nutrient conditions typical of the area of isolation. Sediment extract medium was prepared by filtering Baltic Sea surface sediments through 0.2- $\mu\text{m}$  filters (Sarstedt, Nümbrecht, Germany) and collecting the water fraction. The cells were pregrown in 10% tryptic soy broth medium (Difco Laboratories, Detroit, MI), collected by centrifugation (10 min, 970 g), washed once with sterile water and resuspended in 100 mL sediment extract medium containing 10  $\mu\text{g}$  phenanthrene  $\text{mL}^{-1}$  to  $c. 5 \times 10^7$  CFU  $\text{mL}^{-1}$  sediment medium in 500-mL glass flasks. Subsequent  $\text{OD}_{600\text{nm}}$  values were diluted to 0.12 for all bacterial strains to ensure that all cultures started with approximately the same amount of cells. The cultures were incubated in the dark at 5 °C with shaking (150 r.p.m.) and triplicate cultures were established for each isolate. Uninoculated controls containing only phenanthrene-spiked sediment extract medium were also established in triplicate. Flasks were covered with parafilm and wrapped in an aluminum foil to avoid evaporation of phenanthrene and exposure to light. Cell growth was monitored by enumeration of CFU on 10% tryptic soy broth agar plates, incubated in the dark at 5 °C.

### Screening isolates for degradation genes

Selected bacterial isolates were screened by PCR for the presence of *phnAc* and *xylE* genes using the same primer pairs as those used for qPCR of DNA directly extracted from sediment (Table 2). PCR conditions were as described previously (Edlund *et al.*, 2006) with the following modifications: each step in a cycle was 40 s, annealing temperature was 60 °C and 35 cycles of amplification were performed. The amplified DNA was verified by gel electrophoresis of PCR products in 2.2% agarose gels (containing 2.2% ethidium bromide). To determine sequence identities, the PCR products were cloned and sequenced as described above. The *phnAc* genes were deposited in the EMBL nucleotide sequence database under the following accession numbers: AM491127–AM491129.

## Results

### Phenanthrene transformation in sediment microcosms

Phenanthrene levels in phenanthrene-spiked sediment slurries were significantly reduced after 60 days of incubation. The sediments that were spiked with 10  $\mu\text{g}$   $\text{mL}^{-1}$  phenanthrene (treatments 3, 4 and 5, Table 1) contained on average


3.7  $\mu\text{g}$   $\text{mL}^{-1}$  phenanthrene on initial sampling dates, suggesting that a substantial part of the phenanthrene was immediately absorbed to the flasks, parafilm and/or sediment particles. However, after 60 days, the phenanthrene concentration had decreased to 0.8  $\mu\text{g}$   $\text{mL}^{-1}$ , suggesting that microorganisms in the sediment might be degrading the compound.

### Actively growing bacterial community profiles and bacterial identification

Bromodeoxyuridine-immunocapture was used to selectively extract DNA corresponding to actively growing bacterial populations in the sediments enriched with phenanthrene. To ensure that sufficient bromodeoxyuridine was available throughout the 60-day incubation period, bromodeoxyuridine was added repeatedly to some treatments and compared with treatments where bromodeoxyuridine was only added at the beginning of the experiment. The bacterial community structure of the bromodeoxyuridine-immunocaptured DNA was profiled by T-RFLP and the resulting TRFs were compared between treatments.

When bromodeoxyuridine was added three times during the first week of incubation (treatment 4, Fig. 1), there was a decrease in the number of TRFs detected at days 2 and 7, compared with treatments with only one addition (treatment 2, Fig. 1), suggesting that some bacteria may either have been inhibited or may have been favored by the increased bromodeoxyuridine concentration and grew to the expense of other populations (e.g. those represented by TRFs 101 and 218). However, after 2 months of incubation, the community with repeated bromodeoxyuridine additions (treatment 4, Fig. 1) resembled that where bromodeoxyuridine was added only at day 0 (treatment 3, Fig. 1). In the microcosms where bromodeoxyuridine was added at day 58 (2 days before the last sampling; treatment 5, Fig. 1), two novel TRFs appeared at 338 and 369 bp, indicating that these bacterial populations had sufficiently high growth rates to incorporate bromodeoxyuridine. Their appearance may also imply that new niches were available at the end of the incubation period.

The numbers and relative abundances of TRFs from the T-RFLP data were used to determine the impact of the different incubation conditions on the diversity of the actively growing members of the bacterial community in the sediments. In addition, the total dominant bacterial members of the community were monitored during the incubation period (treatment 1). There was a significant decrease in the phylotype diversity of the total community after 60 days of incubation, from  $c. 40$  initial TRFs ( $H'$ : 3.2;  $E$ : 0.9) to 10 TRFs ( $H'$ : 1.7;  $E$ : 0.5) (Fig. 1, treatment 1). The numbers of TRFs in the bromodeoxyuridine-immunocaptured DNA representing the active community were relatively similar over the entire incubation period in the



**Fig. 1.** Representative electropherograms from TRF analysis (T-RFLP) of 16S rRNA gene fragments amplified from total DNA extracts (treatment 1) and from bromodeoxyuridine-labeled DNA (treatments 2–5) representing sediment microcosm treatments (see Table 1 for a more detailed description of treatments). 16S rRNA gene community profiles from total DNA extracts were analyzed at all sampling dates, including day 0 (before incubation started), and all other profiles were analyzed at days 1, 2, 7 and 60 of incubation. The x-axis in each individual electropherogram represents TRF sizes with a minimum of 60 and a maximum of 500 bp in each case. The y-axis represents fluorescence intensity and the maximum values varied between samples. For the sake of comparison, the y-axis of the electropherograms was adjusted so that the abundant peaks could all be visualized within the same area. The SD for mean relative abundance values (%) of TRFs from three replicate sediment microcosms ranged from SD:  $\pm 0.9$  to 1.6%.

phenanthrene-spiked microcosms, with *c.* 10 TRFs (2 days:  $H'$ : 1.7 and  $E$ : 0.7; 2 months:  $H'$ : 1.5 and  $E$ : 0.6). However, in the unspiked microcosms without added phenanthrene, the numbers of TRFs from actively growing bacteria decreased from 10 to 6 and the diversity decreased from an  $H'$  of 1.7 to 1.0 after 60 days of incubation (Fig. 1).

A clear shift in the T-RFLP profiles of the actively growing bacterial community was observed in sediments that were spiked with phenanthrene after 60 days of incubation compared with the TRFs in unspiked microcosms (Fig. 1). Several TRFs (e.g. TRFs 60, 76, 79, 97 and 128 bp) increased in relative abundance in phenanthrene-spiked sediments, suggesting that they were positively influenced by phenanthrene addition. By contrast, some of the TRFs detected in the control microcosms without phenanthrene (TRFs 210, 213 and 242 bp) had a higher relative abundance compared with the microcosms spiked with phenanthrene. This suggests that the corresponding populations were negatively impacted when phenanthrene was added or that phenanthrene had a below-average positive effect on these populations. However, some TRFs (e.g. 201, 205 and 218 bp) were dominant in both phenanthrene-spiked and -unspiked microcosms, indicating that the corresponding populations were neither effected positively nor negatively by addition of phenanthrene.

Bacteria that were growing and possibly enriched in the phenanthrene-spiked sediments were further identified by cloning and sequencing of 16S rRNA genes amplified from bromodeoxyuridine-immunocaptured DNA (treatment 3) in order to match sequence identities to specific TRFs. A subset of 70 clones (out of *c.* 2000 total) was screened by RFLP using HhaI and RsaI. Seven unique restriction patterns were obtained, indicating a low diversity of metaboli-

cally active community members in polluted Baltic Sea sediments, similar to the results of a previous study of sediments collected in the same sampling area (Edlund & Jansson, 2006). All of the clones could be matched to specific TRFs in the T-RFLP profiles (Table 3). This enabled us to estimate the relative abundances of the corresponding populations in the sediment community. Both the T-RFLP and the cloning approaches resulted in similar relative abundance values (Table 3), confirming the validity of these molecular tools.

Populations with 16S rRNA gene sequences corresponding to *Schewanella baltica* (TRF 76) and *Methylomonas methanica* (TRF 79) were the most dominant metabolically active populations (each *c.* 30% relative abundance), using both approaches (Table 3). These bacteria were only identified in the phenanthrene-spiked microcosms, in addition to *Exiguobacterium oxidotolerans* (TRF 60) and an unidentified bacterium (TRF 128) (Tables 3 and 4). Therefore, it is likely that bacteria representative of *S. baltica*, *M. methanica*, *E. oxidotolerans* and the unidentified bacterium played a role in phenanthrene degradation or at least were stimulated to grow in its presence.

A *Pseudomonas* sp. (TRF 218) was present in all samples, including those to which bromodeoxyuridine had not been added, suggesting that it was dominant and metabolically active irrespective of the incubation conditions tested (Tables 3 and 4). TRFs 97, 101 and 205, corresponding to an uncultured *Deltaproteobacterium*, a *Bacteroidetes* sp. and a *Pseudomonas* strain, respectively, were also abundant and metabolically active irrespective of phenanthrene addition (Table 4).

TRFs 101 and 205, corresponding to the *Bacteroidetes* sp. and the *Pseudomonas* sp. mentioned above, were not present



**Table 4.** Representative TRFs present in community profiles from total DNA extracts and in bromodeoxyuridine-immunocaptured DNA after 60 days of incubation in sediment microcosms

TRF (bp) treatment*	Total DNA		Bromodeoxyuridine-labeled DNA		
	1	2	3	4	5
60	–	–	+	+	+
76	–	–	+	+	+
79	–	–	+	+	+
97	+	–	+	+	+
101	+	+	+	–	–
–	–	–	–	–	–
128	–	–	+	+	+
205	+	+	+	–	–
–	–	–	–	–	–
218	+	+	+	+	+


±, TRF present/not present in community profiles from triplicate sediment microcosms.

\*See Table 1 for description of treatments.

in treatment 4, where bromodeoxyuridine was added several times, suggesting that they were inhibited by high levels of bromodeoxyuridine addition or outcompeted by other more fast-growing bacteria (Table 4).

### Quantification of dioxygenase genes in actively growing bacteria


Gene copies of the *xylE* and *phnAc* genes encoding catechol-2,3-dioxygenase and the iron–sulfur protein (large) subunit of PAH dioxygenase, respectively, were monitored in bromodeoxyuridine-immunocaptured DNA by real-time PCR over the sampling period. None of the other functional genes tested (Table 2) were detected in any samples. In the phenanthrene-spiked sediments, *xylE* gene copy numbers increased from *c.* 40 000 copies mL<sup>-1</sup> sediment on day 1 to *c.* 260 000 copies mL<sup>-1</sup> sediment on day 7 (Fig. 2). Between days 7 and 60, the gene copy numbers decreased to approximately original levels at 45 000 copies mL<sup>-1</sup> sediment. There was also an increase in the number of *xylE* gene copies in the unspiked microcosms containing background levels of phenanthrene from *c.* 25 000 copies mL<sup>-1</sup> sediment on day 1 to *c.* 82 000 on day 7, followed by a decrease to *c.* 30 000 copies mL<sup>-1</sup> sediment by the 60-day sampling date. The *phnAc* genes were considerably more abundant in the phenanthrene-enriched sediments, with copy numbers that increased from *c.* 1.5 × 10<sup>5</sup> copies mL<sup>-1</sup> sediment on day 1, to *c.* 3 × 10<sup>6</sup> copies mL<sup>-1</sup> sediment on day 7, followed by a decrease to *c.* 3.7 × 10<sup>4</sup> copies mL<sup>-1</sup> sediment after 2 months of incubation (Fig. 2). By contrast, in the sediments without added phenanthrene *phnAc* gene copy numbers remained relatively low and stable, ranging from 2500 to 5000 copies mL<sup>-1</sup> throughout the incubation period.



**Fig. 2.** Quantification of dioxygenase genes *xylE* (a) and *phnAc* (b) in phenanthrene-enriched sediment (black bars) and unspiked sediment (white bars) during 60 days of incubation (treatments 2 and 3, Table 1, respectively). The data represent the average of triplicate sediment microcosms and the error bars represents the SD of the mean.

### Characterization of isolates

Three of the 16S rRNA gene sequences that were identified as representative of abundant and active bacteria in the bromodeoxyuridine-extracted DNA from the sediment microcosms were closely related (97–99%) to isolates that were obtained previously from the same sediments (Table 3). These isolates had 16S rRNA gene sequences that were related to 16S rRNA gene sequences of a pseudomonad (TRF 218; i.e. one of those that was abundant and active in all samples, Table 4), another pseudomonad (TRF 205; i.e. abundant and active irrespective of phenanthrene addition, but inhibited by repeated doses of bromodeoxyuridine, Table 4) and *E. oxidotolerans* (TRF 60; abundant and active in the presence of phenanthrene, Table 4). Based on their full-length 16S rRNA gene sequences, we provided each of the isolates with a strain designation as follows: pseudomonad corresponding to TRF 218: *Pseudomonas* sp. AE1 pseudomonad corresponding to TRF 205: *Pseudomonas* sp.



**Fig. 3.** Phenanthrene degradation (■) and log CFU (●) in sediment extract medium spiked with 10 µg phenanthrene mL<sup>-1</sup>. (a) *Exiguobacterium* sp. AE3, (b) *Pseudomonas* sp. AE1, (c) *Pseudomonas* sp. AE2 and (d) uninoculated control. Error bars represent the SD of the mean of triplicate cultures. The error bars for the growth curves are within the area of the data points.

AE2, and *Exiguobacterium* species corresponding to TRF 60: *Exiguobacterium* sp. AE3 (Table 3).

The isolates were tested for their ability to degrade phenanthrene and were also screened for the presence of dioxygenase genes that could express enzymes functioning in phenanthrene degradation. Phenanthrene levels were reduced to background levels in sediment extract medium incubated with all three bacterial cultures, compared with negligible reduction of the compound in uninoculated controls (Fig. 3). The rates of phenanthrene removal were different for the different bacterial cultures. *Exiguobacterium* sp. AE3 reduced phenanthrene to background levels after 18 days of incubation, whereas cultures of the two pseudomonads, *Pseudomonas* sp. AE1 and *Pseudomonas* sp. AE2, depleted phenanthrene after 30 and 34 days, respectively (Fig. 3). The numbers of CFUs increased for all of the tested bacteria during the incubation period, suggesting that they used phenanthrene as a carbon source. All isolates carried an identical *phnAc* gene (99% sequence similarity to AF061751, NCBI) but none of them harbored *xylE* genes. This suggests that the *phnAc* dioxygenase enzyme was used for phenanthrene degradation by the isolates. Moreover, the 144-bp

long *phnAc* gene fragments that were sequenced from isolates and sediments were 100% identical when multiple sequence alignment was performed with the MUSCLE software, version 3.5, available at <http://drive5.com/muscle> (Edgar, 2004).

## Discussion

Molecular approaches, such as those applied in this study, are useful for identification of metabolically active bacteria, including not yet described species (i.e. not yet cultured) that may play significant roles in pollutant degradation or in other community processes (Nogales *et al.*, 1999). Specifically, we identified bacterial populations that grew in Baltic Sea sediments that were spiked with phenanthrene and incubated for 60 days. The incubation conditions were cold (5 °C) and the community was exposed to oxygen during sampling and incubation, thereby also selecting for populations that were both psychrotolerant and oxygen tolerant. The combination of T-RFLP, qPCR, cloning and sequencing of bromodeoxyuridine-labeled DNA enabled the composition of the dominant and active members of the sediment

bacterial community to be identified in parallel with identification and quantification of PAH degradation genes that were present during the enrichment process.

The source of material in this study was sediment that had been collected previously from a polluted site in the Baltic Sea subsequent to remediation by dredging (Edlund *et al.*, 2006) and it was stored frozen before setup of the phenanthrene enrichments. Although it was not possible to resample the original site to obtain fresh polluted sediments, because these had been removed during the dredging process, the frozen sediment did contain pollutants and populations representative of the original sampling site. We are aware that some microorganisms originally present in the sediment might have died during sampling and storage. However, it was encouraging to find that several of the 16S rRNA gene sequences representative of growing populations that were detected initially were still detected as growing members of the community in the present study.

Some of the 16S rRNA gene sequences that we detected in the phenanthrene-spiked microcosms were related to isolates obtained previously from the same PAH-contaminated Baltic Sea sediments (Edlund & Jansson, 2006). In this study, we found that three of the isolates, given the strain designations *Exiguobacterium* sp. AE3, *Pseudomonas* sp. AE1 and *Pseudomonas* sp. AE2, were capable of growth on and degradation of phenanthrene. Although *Exiguobacterium* is a relatively unstudied genus, a couple of strains that derive from extreme environments have been described previously (Collins *et al.*, 1983; Vishnivetskaya *et al.*, 2000). These strains have shown interesting features such as high survival during long-term freezing, rapid growth at low temperatures and a high osmotic tolerance (Collins *et al.*, 1983; Vishnivetskaya *et al.*, 2000; Isao *et al.*, 2004), similar to the conditions of the sediments used in our study. In addition, pseudomonads related to the *Pseudomonas* isolates in our study, *Pseudomonas fluorescens* and *Pseudomonas migulae*, were isolated previously from phenanthrene-polluted environments and these had a high capacity for phenanthrene degradation (Sanseverino *et al.*, 1993; Muñoz *et al.*, 2003).

Although one might expect anaerobic bacterial populations to carry out degradation functions in the largely anaerobic sediments *in situ*, aerobic degrading isolates and aerobic PAH degradation genes were found. Therefore, the question remains as to whether these populations play a key role in bioremediation of Baltic Sea sediments. We know from recent studies of a relatively clean area of the Baltic Sea that the microbial community structure varies at different vertical layers in the sediment and this is correlated to redox potential, with the upper sediment layers being oxic (Edlund *et al.*, 2008). In the present investigation, we collected surface sediments, including the upper presumably oxic layers, and this could be the source of our isolates and aerobic degradation genes that were subsequently enriched.

Other studies have also reported high numbers of aerobic aromatic degrading organisms and genes in apparent anaerobic aquifers (Hendrickx *et al.*, 2005; Fahy *et al.*, 2006). Therefore, the relative contributions of aerobic and anaerobic processes in these environments remain to be elucidated.

All of our phenanthrene-degrading isolates harbored the *phnAc* gene, which is involved in the initial hydroxylation step of phenanthrene degradation (Lauri & Lloyd-Jones, 1999). These results are in line with a previous study where enrichment of the *phn* genotype was observed in isolates from phenanthrene-amended PAH-contaminated soils from a broad geographical distribution (central Siberia, Antarctica, Ross Island and New Zealand) (Lauri & Lloyd-Jones, 2000).

In contrast to the isolates that only contained the *phnAc* gene, both *xylE* and *phnAc* genes were detected by qPCR of bromodeoxyuridine-labeled DNA extracted from the sediment (Fig. 2). Both genotypes were found to increase significantly after 7 days of incubation with phenanthrene. The *xylE* gene (EC 1.13.11.2 in Kyoto Encyclopedia of Genes and Genomes, KEGG available at <http://www.genome.jp/kegg/>) expresses *meta* cleavage of catechol in the lower pathways of PAH degradation. In line with our results, another study of PAH-contaminated soil showed that *xylE* was significantly expressed during phenanthrene degradation (Cunliffe *et al.*, 2006). However, we found a greater increase in expression levels of *phnAc* than of *xylE*. This difference may reflect that *phnAc* is critical for initial degradation of phenanthrene and that the dioxygenase encoded by *xylE* is one of several options for further degradation of the compound. The decrease in *xylE* and *phnAc* gene copy numbers at the end of the experiment may reflect the corresponding decrease in levels of phenanthrene. Moreover, all *phnAc* gene fragments that were sequenced from isolates and sediments were identical, indicating that this gene may have been recently transferred between members of the bacterial community. These results are in line with a previous study where highly conserved *phn* genes were shared by groups of taxonomically diverse hosts in naphthalene-contaminated seep sediments (Wilson *et al.*, 2003). However, we only analyzed 144 bp within the complete PAH-catabolic gene cluster that encompasses 11 451 bp in total (Lauri & Lloyd-Jones, 1999) and we do not know whether the remaining genes were present. Also, there could be other catabolic pathways involved in phenanthrene detection that we did not screen for, such as the *phd* genes described for phenanthrene degradation in river sediments (Goyal & Zylstra, 1996, 1997).

Our results indicated that *Exiguobacterium* sp. AE3 only grew in the microcosm experiments when phenanthrene was added. By contrast, the two pseudomonads, *Pseudomonas* sp. AE1 and *Pseudomonas* sp. AE2, were abundant and growing in the sediment microcosms in the presence and

absence of phenanthrene, suggesting that they could also utilize endogenous carbon sources for growth and that their growth might be independent of phenanthrene levels *in situ*. Previous studies applied stable isotope probing (SIP) to specifically study metabolically active phenanthrene-degrading bacteria in PAH-contaminated soil (Singleton *et al.*, 2005, 2006). In those studies, community members belonging to the genus *Pseudomonas* were responsible for pyrene degradation while some *Acidovorax* representatives were responsible for phenanthrene degradation. Taken together, these studies suggest that phenanthrene-degradation abilities are not species specific but that they are dependent on the local conditions and the genetic makeup of the indigenous bacterial flora.

In addition to the isolates mentioned above, bacteria related to *M. methanica* and *S. baltica* (Table 4) were found to be abundant and metabolically active in the phenanthrene-spiked sediments using the bromodeoxyuridine immunocapture approach, but we did not have isolates representative of these bacteria. The molecular data, however, provide a clue that these bacteria are interesting as potential phenanthrene degraders and they present target populations for specific isolation using selective cultivation conditions in the future. It is also possible that these bacteria use metabolites or excreted products from the phenanthrene-degrading community members for growth. In previous studies, it was shown that *Schewanella putrefaciens*, which is a close relative to *S. baltica*, has the potential to perform reductive dechlorination of carbon tetrachloride and some components of oil (Backhus *et al.*, 1997; Martín-Gil *et al.*, 2004) and is capable of growing both aerobically and anaerobically at the expense of many different electron acceptors (Nealson *et al.*, 1995; Kostka *et al.*, 1996). The roles of the *Methylobomonas* and *Schewanella* species in this study remain unknown. However, their presence in the spiked microcosms suggests that they may be involved in phenanthrene degradation but this hypothesis remains to be confirmed by a study of corresponding isolates.

After 2 months of incubation, the total community richness based on T-RFLP of directly extracted DNA decreased in all treatments by *c.* 75% (Fig. 1), which may be due to a 'bottle effect' (i.e. some bacterial populations were favored by the specific incubation conditions used and increased in growth while the growth of other populations was repressed, Pritchard *et al.*, 1979). In contrast to the totally extracted DNA, a decrease in the number of TRFs was not observed for 16S rRNA genes that were amplified from bromodeoxyuridine-labeled DNA, which would be expected if the incubation conditions alone were acting on the community. We previously found a low diversity of metabolically active bacteria in Baltic Sea sediments (Edlund & Jansson, 2006) and in soil (Hjort *et al.*, 2007) using the same approach. Therefore, these results suggest that only a small

fraction of the total bacterial community is active at any particular time, depending on the specific environmental conditions. However, it should be kept in mind that additional rare bacterial species that are below the detection limit of our method may also play important ecological roles.

In summary, we used a combination of approaches, to identify psychrotolerant bacteria and functional genes that were selectively enriched in the presence of phenanthrene. Some bacterial isolates, including a novel *Exiguobacterium* strain, were represented as growing members of the sediment community in the phenanthrene enrichments using molecular approaches and they could grow on and degrade phenanthrene at 5 °C. Therefore, these strains represent populations that are active in polluted Baltic Sea sediments and potential contributors to PAH transformations in oxic layers of the sediments *in situ*.

## Acknowledgements

We thank Anders R. Johnsen, Department of Geochemistry, Geological Survey of Denmark and Greenland, and Victor de Lorenzo, National Center for Biotechnology, Madrid, Spain, for supplying us with control bacterial strains for qPCR analysis; Elisabet Börjesson and Leticia Pizzul, Department of Microbiology, Swedish University of Agricultural Sciences, Uppsala, Sweden, and Tord Eriksson, Södertörn University college for assistance with GC-MS analyses and contribution towards cultivation of phenanthrene-transforming bacteria; and Sonja Löfmark and Benjamin Edvinsson, Department of Laboratory Medicine, Karolinska Institute, Sweden, for help with qPCR analysis and access to their equipment. We also thank Sara Sjöling, Södertörn University College, Huddinge, Sweden, and an anonymous reviewer for critically reviewing the manuscript. The study was funded by the Baltic Sea Foundation and in part by US Department of Energy Contract DE-AC02-05CH11231 with Lawrence Berkeley National Laboratory.

## References

- Altschul SE, Madden TL, Schaeffer AA, Zhang J, Zhang Z, Miller W & Lipman DJ (1997) Gapped BLAST and PSI-BLAST: a new generation of protein database search programs. *Nucleic Acids Res* **25**: 3389–3402.
- Artursson V & Jansson JK (2003) Use of bromodeoxyuridine immunocapture to identify bacteria associated with arbuscular mycorrhizal hyphae. *Appl Environ Microbiol* **69**: 6208–6215.
- Backhus DA, Picardal FW, Johnson S, Knowles T, Collins R, Radue A & Sanggou K (1997) Soil- and surfactant-enhanced reductive dechlorination of carbon tetrachloride in the presence of *Shewanella putrefaciens* 200. *J Contam Hydrol* **28**: 337–361.

- Brinch UC, Ekelund F & Jacobsen CS (2002) Method for spiking soil samples with organic compounds. *Appl Environ Microbiol* **68**: 1808–1816.
- Cole JR, Chai B, Farris RJ, Wang Q, Kulam SA & McGarrell DM (2003) The Ribosomal Database Project (RDP-II): previewing a new aligner that allows regular updates and the new prokaryotic taxonomy. *Nucleic Acids Res* **31**: 442–443.
- Collins MD, Lund BM, Farrow JAE & Schleifer KH (1983) Chemotaxonomic study of an alkaliphilic bacterium, *Exiguobacterium aurantiacum* gen. nov., sp. nov. *J Gen Microbiol* **129**: 2037–2042.
- Cunliffe M, Kawasaki A, Fellows E & Kertesz MA (2006) Effect of inoculum pretreatment on survival, activity and catabolic gene expression of *Sphingobium yanoikuyae* B1 in aged polycyclic aromatic hydrocarbon-contaminated soil. *FEMS Microbiol Ecol* **58**: 364–372.
- Edgar RC (2004) MUSCLE: multiple sequence alignment with high accuracy and high throughput. *Nucleic Acids Res* **32**: 1792–1797.
- Edlund A & Jansson JK (2006) Changes in dominant and active bacterial communities before and after dredging of highly polluted Baltic Sea sediments. *Appl Environ Microbiol* **72**: 6800–6807.
- Edlund A, Soule T, Sjöling S & Jansson JK (2006) Microbial community structure in polluted Baltic Sea sediments. *Environ Microbiol* **8**: 223–232.
- Edlund A, Hårdeman F, Jansson JK & Sjöling S (2008) Active bacterial community structure along vertical redox gradients in Baltic Sea sediment. *Environ Microbiol*, DOI: 10.1111/j.1462-2920.2008.01624.x.
- Fahy A, McGenity TJ, Timmis KN & Ball AS (2006) Heterogeneous aerobic benzene-degrading communities in oxygen-depleted groundwaters. *FEMS Microbiol Ecol* **58**: 260–270.
- Goyal AK & Zylstra GJ (1996) Molecular cloning of novel genes for polycyclic aromatic hydrocarbon degradation from *Comamonas testosteroni* GZ39. *Appl Environ Microbiol* **62**: 230–236.
- Goyal AK & Zylstra GJ (1997) Genetics of naphthalene and phenanthrene degradation by *Comamonas testosteroni*. *J Indust Microbiol Biotechnol* **19**: 401–407.
- Hendrickx B, Dejonghe W, Boënné W *et al.* (2005) Dynamics of an oligotrophic bacterial aquifer community during contact with a groundwater plume contaminated with benzene, toluene, ethylbenzene, and xylenes: an *in situ* mesocosm study. *Appl Environ Microbiol* **71**: 3815–3825.
- Hjort K, Lembke A, Speksnijder A, Smalla K & Jansson JK (2007) Community structure of actively growing bacterial populations in plant pathogen suppressive soil. *Microb Ecol* **53**: 399–413.
- Isao Y, Hishinuma-Narisawa M, Hirota K, Shingyo T, Takebe F, Nodasaka Y, Matsuyama H & Isao H (2004) *Exiguobacterium oxidotolerans* sp. nov., a novel alkaliphile exhibiting high catalase activity. *Int J Syst Evol Microbiol* **54**: 2013–2017.
- Johnsen A, de Liphay JR, Sørensen SJ, Ekelund F, Christensen P, Andersen O, Karlson U & Jacobsen CS (2006) Microbial degradation of street dust polycyclic aromatic hydrocarbons in microcosms simulating diffuse pollution of urban soil. *Environ Microbiol* **8**: 535–545.
- Johnsen A, Schmidt S, Hybholt TK, Henriksen S, Jacobsen CS & Andersen O (2007) Strong impact on the polyaromatic hydrocarbon (PAH) degrading community of a PAH-polluted soil but marginal effect on PAH degradation when priming with bioremediation soil dominated with *Mycobacteria*. *Appl Environ Microbiol* **73**: 1474–1480.
- Kim E & Zylstra GJ (1995) Molecular and biochemical characterization of two meta-cleavage dioxygenases involved in biphenyl and *m*-xylene degradation by *Beijerinckia* sp. strain B1. *J Bacteriol* **177**: 3095–3103.
- Kostka JE, Stucki JW, Neelson KH & Wu J (1996) Reduction of structural Fe(III) in smectite by a pure culture of *Schewanella putrefaciens* strain MR-1. *Clays Miner* **44**: 522–529.
- Lauri AD & Lloyd-Jones G (1999) The *phn* genes of *Burkholderia* sp. strain RP007 constitute a divergent gene cluster for polyaromatic hydrocarbon catabolism. *J Bacteriol* **181**: 531–540.
- Lauri AD & Lloyd-Jones G (2000) Quantification of *phnAc* and *nahAc* in contaminated New Zealand soils by competitive PCR. *Appl Environ Microbiol* **66**: 1814–1817.
- Margalef R (1958) Information theory in ecology. *Gen Sys* **3**: 36–71.
- Martin-Gil J, Ramos-Sanchez MC & Martin-Gil FJ (2004) *Shewanella putrefaciens* in a fuel-in-water emulsion from the Prestige oil spill. *Antonie van Leeuwenhoek* **86**: 283–285.
- Muñoz R, Guieysse B & Mattiasson B (2003) Phenanthrene biodegradation by an algal-bacterial consortium in two-phase partitioning bioreactors. *Appl Microbiol Biotechnol* **61**: 261–267.
- Muyzer G, Teske A, Wirsen CO & Jannasch HW (1995) Phylogenetic relationship of *Thiomicrospira* species and their identification in deep-sea hydrothermal vent samples by denaturing gradient gel electrophoresis of 16S rDNA fragments. *Arch Microbiol* **164**: 162–172.
- Neelson KH, Moseder DP & Saffarini A (1995) Anaerobic electron acceptor chemotaxis in *Shewanella putrefaciens*. *Appl Environ Microbiol* **61**: 1551–1554.
- Nogales B, Moore ERB, Abraham WR & Timmis KN (1999) Identification of the metabolically active members of a bacterial community in a polychlorinated biphenyl-polluted mooreland soil. *Environ Microbiol* **3**: 199–212.
- Palm A, Wängberg I & Broström-Lundén E (2001) Kvicksilver och organiska miljögifter i Örserumsviken, IVL Swedish Environmental Research institute Ltd. Report number B 1433.
- Pizzul L, del Pilar Castillo M & Stenström J (2006) Characterization of selected actinomycetes degrading polyaromatic hydrocarbons in liquid culture and spiked soil. *World J Microbiol Biotechnol* **22**: 745–752.
- Pritchard PH, Bourquin AW, Frederickson HL & Maziarz T (1979) System design factors affecting environmental fate

- studies in microcosms. *Workshop: Microbial degradation of pollutants in Marine environments* (Bourqin AW & Pritchard PM, eds), pp. 251–272, EPA-600/9-79-012. US Environmental Protection Agency, Washington, DC.
- Sanseverino J, Applegate BM, King JMH & Sayler GS (1993) Plasmid-mediated mineralization of naphthalene, phenanthrene, and anthracene. *Appl Environ Microbiol* **59**: 1931–1937.
- Simon MJ, Osslund TD, Saunders R, Ensley BD, Suggs S, Harcourt A, Suen WC, Cruden DL, Gibson DT & Zylstra GJ (1993) Sequences of genes encoding naphthalene dioxygenase in *Pseudomonas putida* G7 and NCIB 9816-4. *Gene* **127**: 31–37.
- Singleton DR, Powell SN, Sangaiah R, Gold A, Ball LM & Aitken MD (2005) Stable-isotope probing of bacteria capable of degrading salicylate, naphthalene or phenanthrene in a bioreactor treating contaminated soil. *Appl Environ Microbiol* **71**: 1202–1209.
- Singleton DR, Sangaiah R, Gold A, Ball LM & Aitken MD (2006) Identification and quantification of uncultivated proteobacteria associated with pyrene degradation in a bioreactor treating PAH-contaminated soil. *Environ Microbiol* **8**: 1736–1745.
- Urbach E, Vergin LL & Giovannoni SJ (1999) Immunochemical detection and isolation of DNA from metabolically active bacteria. *Appl Environ Microbiol* **65**: 1207–1213.
- Vishnivetskaya T, Kathariou S, McGrath J, Gilichinsky D & Tiedje JM (2000) Low-temperature recovery strategies for the isolation of bacteria from ancient permafrost sediments. *Extremophiles* **4**: 165–173.
- Weisburg WG, Barns SM, Pelletier DA & Lane DJ (1991) 16S Ribosomal DNA amplification for phylogenetic study. *J Bacteriol* **173**: 697–703.
- Whelan JA, Russel NB & Whelan E (2003) A method for the absolute quantification of cDNA using real-time PCR. *J Immunol Methods* **278**: 261–269.
- Wilson MS, Herrick JB, Jeon CO, Hinman DE & Madsen EL (2003) Horizontal transfer of *phnAc* dioxygenase genes within one of two phenotypically and genotypically distinctive naphthalene-degrading guilds from adjacent soil environments. *Appl Environ Microbiol* **69**: 2172–2181.