

UC Irvine

UC Irvine Previously Published Works

Title

Gauge-top unification

Permalink

<https://escholarship.org/uc/item/01g103c4>

Journal

Journal of High Energy Physics, 2009(07)

ISSN

1126-6708

Authors

Hosteins, Pierre

Kappl, Rolf

Ratz, Michael

et al.

Publication Date

2009

DOI

10.1088/1126-6708/2009/07/029

Copyright Information

This work is made available under the terms of a Creative Commons Attribution License, available at <https://creativecommons.org/licenses/by/4.0/>

Peer reviewed

Gauge-top unification

Pierre Hosteins, Rolf Kappl, Michael Ratz and Kai Schmidt-Hoberg

*Physik-Department T30, Technische Universität München,
James-Franck-Straße, 85748 Garching, Germany*

E-mail: pierre.hosteins@ph.tum.de, rolf.kappl@ph.tum.de,
mratz@ph.tum.de, kschmidt@ph.tum.de

ABSTRACT: Higher-dimensional models of grand unification allow us to relate the top Yukawa coupling y_t to the gauge coupling g . The tree level relation $y_t = g$ at the scale of grand unification implies, in the framework of the MSSM, a rather small ratio of Higgs expectation values $\tan\beta$. We find that, in the presence of localized Fayet-Iliopoulos terms, y_t is suppressed against g because the bulk fields acquire non-trivial profiles whose overlap is smaller than in the case of flat profiles. This increases the prediction for $\tan\beta$ to moderately large values. Thus $\tan\beta$ is related to the geometry of compact space. We also discuss explicit realizations of such settings in orbifold compactifications of the heterotic string. It turns out that anisotropic compactifications, allowing for an orbifold GUT interpretation, are favored.

KEYWORDS: Field Theories in Higher Dimensions, Superstrings and Heterotic Strings, Extended Supersymmetry

ARXIV EPRINT: [0905.3323](https://arxiv.org/abs/0905.3323)

Contents

1	Introduction	1
2	Gauge-top unification in extra dimensions	2
3	Corrections to the equality of y_t and g	5
3.1	Corrections from localized states	5
3.2	Diagonalization effects	5
3.3	Localization effects	6
4	Explicit string theory realization	8
4.1	Motivation of the orbifold GUT picture	8
4.2	A specific example	9
4.3	Mini-landscape survey	10
5	Phenomenological implications	11
6	Discussion	13

1 Introduction

The coupling strengths governing the interactions of the standard model (SM) exhibit a very peculiar pattern: on the one hand, the gauge and top Yukawa couplings are of order one, on the other hand all other Yukawa couplings are suppressed. This might tell us that couplings come in two classes with fundamentally different origin. In this study we shall investigate the scenario of “gauge-top unification” in which the top Yukawa coupling y_t and the gauge couplings g_a ($1 \leq a \leq 3$) unify at a high scale. Our analysis will be based on the minimal supersymmetric extension of the standard model (MSSM), as it appears to provide us with the most compelling scenario of gauge coupling unification, and therefore fits very nicely to the concept of grand unified theories (GUTs) [1, 2].

Arguably, the most compelling realizations of GUTs incorporate extra dimensions. In fact, string-theoretic orbifolds [3–10] and field-theoretic orbifold GUTs [11–18] allow us to retain the beautiful aspects of grand unification while avoiding the notorious problems. Merging both approaches [19–22] has lead us to explicit string-derived models which reproduce the MSSM in their low-energy limit and have a straightforward orbifold GUT interpretation [23–28] (for reviews see [29, 30]).

In this paper we shall study scenarios in which y_t arises from gauge interactions in more than four dimensions. We will discuss settings in which the Higgs fields arise from

extra components of the gauge multiplet, enforcing the tree-level relation

$$y_t = g \tag{1.1}$$

at the compactification scale [18, 31]. As we shall see, relation (1.1) implies, together with the updated mass of the top quark [32], uncomfortably small values for the ratio of Higgs expectation values $\tan\beta$. However, following earlier work by Lee, Nilles and Zucker [33] we find that localized Fayet-Iliopoulos terms, which are generically present in these compactifications, always reduce the value of y_t at the compactification scale, thus increasing the prediction for $\tan\beta$ to moderately large (or even large) values. This is because the bulk hypermultiplets attain non-trivial profiles whose overlap is always smaller than in the case of flat profiles. The precise value of $\tan\beta$ depends on the size and shape of compact space.

This paper is organized as follows. In section 2 we will discuss a class of simple orbifold GUT models in which $y_t = g$ at tree level. Section 3 is devoted to (quantum) corrections to this relation. In section 4 we will discuss explicit string theory realizations of these settings. Phenomenological implications are described in section 5. Finally, section 6 contains our conclusions.

2 Gauge-top unification in extra dimensions

Let us now consider field-theoretic orbifold GUT settings in which y_t arises from gauge interactions. We shall focus on models with two extra dimensions and an $SU(6)$ bulk gauge group. We will also consider further $U(1)$ factors, as their presence has important implications for the prediction of gauge and Yukawa couplings. The geometry of the model is $\mathbb{T}^2/\mathbb{Z}_2$. The resulting orbifold can be envisaged as a ravioli or pillow, whose corners correspond to the four fixed points. We label the fixed points by two integers n_2 and n'_2 which take the values 0 or 1 (see figure 1). The lengths of the edges of the pillow or the distances of the fixed points are given by πR_5 and πR_6 , respectively, where R_5 and R_6 denote the radii of the underlying 2-torus \mathbb{T}^2 . We restrict ourselves to rectangular tori, leaving the general case for future analysis.

The $SU(6)$ bulk gauge group gets broken to $SU(5) \times U(1)_X$ and $SU(4) \times SU(2)_L \times U(1)'$ at two inequivalent fixed points, i.e. fixed points with different n_2 . The low-energy gauge group emerges as the intersection of these local gauge groups, and is given by

$$G_{\text{SM}} = SU(3)_C \times SU(2)_L \times U(1)_Y \tag{2.1}$$

plus an additional $U(1)$ factor. In orbifold GUT language, this is a consequence of the \mathbb{Z}_2 boundary conditions

$$P = \text{diag}(1, 1, 1, 1, 1, -1) \quad \text{and} \quad P' = \text{diag}(1, 1, 1, -1, -1, 1) , \tag{2.2}$$

which are to be imposed at the fixed points with $n_2 = 0$ and $n_2 = 1$, respectively. In the description common to string theory, the difference between the two local boundary conditions can be ascribed to the presence of a discrete order 2 Wilson line in y_5 direction.




Figure 1. 6D orbifold GUT. We show the local gauge groups at the fixed points, labeled by the localization quantum numbers (n_2, n'_2) .

The emerging symmetry breaking pattern has been studied in the context of a 5D orbifold GUT [18], and happens to arise in a 6D orbifold GUT limit of promising string compactifications [26, 29, 30]. The bulk supersymmetry in six dimensions corresponds to $N = 2$ supersymmetry from a 4D perspective. The 6D gauge multiplet (V, Φ) contains the 4D vector field V as well as the chiral field Φ . The boundary conditions (2.2) result in orbifold parities for the components of Φ that are opposite to those of V ,

$$\begin{aligned} \Phi &= \Phi^a T_a \\ &= \begin{pmatrix} \Phi_{(8,1)_0}^{(--)} - \frac{1}{\sqrt{15}}\Phi_Y^{(--)} + \frac{1}{2\sqrt{15}}\Phi_X^{(--)} & \frac{1}{\sqrt{2}}\Phi_{(3,2)_{-5/6}}^{(-+)} & \frac{1}{\sqrt{2}}\Phi_{(3,1)_{-1/3}}^{(+)} \\ \frac{1}{\sqrt{2}}\Phi_{(3,2)_{5/6}}^{(-+)} & \Phi_{(1,3)}^{(--)} + \frac{3}{2\sqrt{15}}\Phi_Y^{(--)} + \frac{1}{2\sqrt{15}}\Phi_X^{(--)} & \frac{1}{\sqrt{2}}\Phi_{(1,2)_{1/2}}^{(++)} \\ \frac{1}{\sqrt{2}}\Phi_{(3,1)_{1/3}}^{(+)} & \frac{1}{\sqrt{2}}\Phi_{(1,2)_{-1/2}}^{(++)} & \frac{-5}{2\sqrt{15}}\Phi_{(1,1)_0}^{(--)} \end{pmatrix}. \end{aligned} \tag{2.3}$$

Here we decompose the generators of $SU(6)$ into the generators of $G_{SM} \times U(1)_\chi$ and elements of the coset, using an obvious notation. At the massless level, only the doubly even states

$$h_u = \Phi_{(1,2)_{1/2}}^{(++)} \quad \text{and} \quad h_d = \Phi_{(1,2)_{-1/2}}^{(++)} \tag{2.4}$$

are retained, which carry the quantum numbers of the MSSM Higgs doublets. The normalization factors in (2.3) are a consequence of the usual condition on the generators,

$$\text{tr}(T_a T_b) = \frac{1}{2}\delta_{ab}. \tag{2.5}$$

Both $\Phi = \Phi^a T_a$ and the various multiplets appearing on the right-hand side of (2.3) are canonically normalized.

The settings further contain a bulk hypermultiplet H that transforms as a **20**-plet under $SU(6)$. In $N = 1$ language

$$H = (\varphi, \varphi^c), \tag{2.6}$$

where φ transforms as **20** (3-index antisymmetric tensor) and φ^c as $\overline{\mathbf{20}}$ under SU(6). After orbifold projection, φ gives rise to one copy of quark doublets (q_3) and φ^c leads to one superfield transforming as a u -type quark (\bar{u}_3), as well as a lepton singlet \bar{e}_3 . We refrain from specifying the origin of the remaining SM matter here. Later, in section 4, we will clarify this issue in the context of string-derived models, where all anomaly constraints are automatically fulfilled.

In order to extract the top Yukawa coupling it is sufficient to work with component fields. We will then denote by ξ and η the two-component spinors contained respectively in φ and φ^c , and the scalar component of Φ is given by

$$\phi = \frac{A_5 + iA_6}{\sqrt{2}}. \quad (2.7)$$

The 4D Yukawa term

$$y_t \bar{u}_3 q_3 h_u \subset \mathcal{L}_{4D} \quad (2.8)$$

originates from the gauge interactions

$$\eta(\bar{\partial} + \sqrt{2}g_6 \phi)\xi \subset \mathcal{L}_{6D} \quad (2.9)$$

with g_6 denoting the 6D gauge coupling and $\bar{\partial} = \partial_5 + i\partial_6$ (cf. [34]).

In SU(n) notation (see e.g. [35]), one has specifically for the SU(6) case under consideration

$$\begin{aligned} & \frac{1}{3!} \eta_{i_1 i_2 i_3} \left[\delta_{j_1}^{i_1} \delta_{j_2}^{i_2} \delta_{j_3}^{i_3} \bar{\partial} + \sqrt{2} g_6 \left(\phi_{j_1}^{i_1} \delta_{j_2}^{i_2} \delta_{j_3}^{i_3} + \delta_{j_1}^{i_1} \phi_{j_2}^{i_2} \delta_{j_3}^{i_3} + \delta_{j_1}^{i_1} \delta_{j_2}^{i_2} \phi_{j_3}^{i_3} \right) \right] \xi^{j_1 j_2 j_3} \\ &= \frac{1}{3!} \eta_{i_1 i_2 i_3} \left[\delta_{j_3}^{i_3} \bar{\partial} + 3\sqrt{2} g_6 \phi_{j_3}^{i_3} \right] \xi^{i_1 i_2 i_3}. \end{aligned} \quad (2.10)$$

q_3 is contained in the SU(5) **10**-plet from the SU(6) **20**-plet ξ ,

$$\mathbf{10}^{ij} = \xi^{ij6}, \quad (2.11)$$

while \bar{u}_3 arises from the **10**-plet in the SU(6) $\overline{\mathbf{20}}$ -plet η ,

$$\frac{1}{2} \varepsilon_{ijklm} (\mathbf{10}^c)^{\ell m} = \eta_{ijk}. \quad (2.12)$$

Here the indices $i, j, k \dots$ run from 1 to 5; the above implies that q_3 and \bar{u}_3 do not stem from the same SU(5) **10**-plet.

We are now left with

$$\begin{aligned} \sqrt{2} g_6 \frac{1}{2} \eta_{ijk} \phi_6^k \xi^{ij6} &= \frac{\sqrt{2} g_6}{4} \varepsilon_{ijklm} (\mathbf{10}^c)^{ij} \mathbf{10}^{k\ell} \phi_6^m \\ &\supset \frac{g_6}{2} \varepsilon_{abc} \varepsilon_{\alpha\beta} \varepsilon^{abd} (\bar{u}_3)_d (q_3)^{c\alpha} (h_u)^\beta \\ &= g_6 \bar{u}_3 q_3 h_u. \end{aligned} \quad (2.13)$$

Here a, b, c, d are SU(3)_C indices and α, β are SU(2)_L indices. This calculation shows that $y_t = g$ at tree level, and confirms an earlier computation [26] (see also [18]).

3 Corrections to the equality of y_t and g

Having seen that $y_t = g$ at tree level, we now turn to discussing corrections to this relation. Such corrections are of different origin. First, there are radiative, logarithmic contributions to the wave function renormalization constants coming from fields localized at the fixed points (or ‘branes’) which do not respect the unified bulk gauge symmetry. Second, the top Yukawa coupling emerges from the 3×3 Yukawa matrix Y_u by diagonalization. Subdominant entries in Y_u can therefore shift this eigenvalue. Third, there are corrections coming from non-trivial localization properties of the bulk hypermultiplets. As we shall see, this effect yields generically the numerically dominant correction.

3.1 Corrections from localized states

The 6D β -function is comprised of a ‘power-law’ and logarithmic piece,

$$\beta_{6D} = b_6 \mu^2 R_5 R_6 + b_4 . \tag{3.1}$$

We have verified that the power-law pieces are universal for gauge and top couplings, as it should be. Assuming only the minimal matter content in the bulk to form the third family of the MSSM, one obtains

$$b_6^t = b_6^i = -4 \quad \text{for } 1 \leq i \leq 3 . \tag{3.2}$$

The logarithmic corrections are sensitive to fields sitting at the fixed points, where the $SU(6)$ bulk symmetry is broken. The b_4 coefficients might hence not be universal. One might think of these corrections as wave function renormalization constants localized at the fixed points. Their impact can be estimated as

$$|\Delta y_t - \Delta g_i|_{\log} \sim |\Delta g_i - \Delta g_j|_{\log} \sim |\Delta b_4 \ln(\Lambda/M_{GUT})| , \tag{3.3}$$

where Λ denotes the cut-off and Δb_4 denotes the respective difference of β -functions. These corrections are expected to be numerically similarly relevant as MSSM threshold corrections, which originate from the squarks and sleptons having masses that differ by an $\mathcal{O}(1 - 10)$ factor. Such effects will not be studied in detail in the present analysis.

3.2 Diagonalization effects

In general, we will expect the u -type Yukawa matrix to be of the form

$$Y_u = \begin{pmatrix} 0 & 0 & 0 \\ 0 & 0 & 0 \\ 0 & 0 & \mathcal{O}(g) \end{pmatrix} + \begin{pmatrix} s^{n_{11}} & s^{n_{12}} & s^{n_{13}} \\ s^{n_{21}} & s^{n_{22}} & s^{n_{23}} \\ s^{n_{31}} & s^{n_{32}} & s^{n_{33}} \end{pmatrix} , \tag{3.4}$$

where s denotes the typical expectation value of some singlet field, $n_{ij} \in \mathbb{N}$, and we suppressed coefficients. It is clear that, after a bi-unitary diagonalization, the top Yukawa coupling will be given by

$$y_t = \mathcal{O}(g) + \mathcal{O}(s^n) \tag{3.5}$$




Figure 2. Localization of two wavefunctions φ and φ^c with opposite charges in the extra dimensions.

with some $n \in \mathbb{N}$. However, if the singlet scale and the exponents n_{ij} in (3.4) are such as to give rise to realistic Yukawa couplings for the u and s quarks, this effect is expected to be negligible. In fact, as we shall see next, localization effects in the compact dimensions will generically have sizable impact on y_t , completely overwhelming the effect described in the present subsection.

3.3 Localization effects

Building on earlier analyses of the 5D case [36, 37], Lee, Nilles and Zucker (LNZ) [33] reported an interesting observation. Suppose there is a $U(1)$ symmetry, called $U(1)_{\text{LNZ}}$ in what follows, with localized Fayet-Iliopoulos (FI) terms, i.e.

$$\text{tr}(q_I) \neq 0, \tag{3.6}$$

where the trace extends over the states localized at the fixed points, labeled by $I = (n_2, n'_2)$. It turns out that then bulk states with non-zero $U(1)_{\text{LNZ}}$ charge develop a non-trivial profile in the extra dimensions, cf. figure 2. In such a situation, the effective four-dimensional top Yukawa coupling, which is given by the overlap of the corresponding wave functions, generically gets reduced. It is amusing to see that the presence of fixed points, which break $N = 2$ supersymmetry down to $N = 1$, also allows us to discriminate between the φ and φ^c components of the hypermultiplet.

In [33] it was shown that the profile for the zero modes on the 6D orbifold $\mathbb{T}^2/\mathbb{Z}_2$ is given by

$$\psi \simeq f \prod_I \left| \vartheta_1 \left(\frac{z - z_I}{2\pi} \middle| \tau \right) \right|^{\frac{1}{2\pi} g_6 q_\psi \xi_I} \exp \left(-\frac{1}{8\pi^2 \tau_2} g_6 q_\psi \xi_I (\text{Im}(z - z_I))^2 \right), \tag{3.7}$$

where we neglected a subleading logarithmic contribution, and f is a normalization factor. We explicitly checked that this profile is the same for bosonic and fermionic degrees of

freedom. The complex coordinate z is defined as

$$z = \frac{1}{R_5}x^5 + \frac{\tau}{R_6}x^6 \tag{3.8}$$

with τ denoting the modular parameter of the torus. In our settings we assume the torus to be rectangular, $\tau = i\tau_2 = iR_6/R_5$. The torus ϑ -functions can be written as [38]

$$\vartheta_1(z|\tau) = \sum_{n \in \mathbb{Z}} e^{i\pi\tau(n+\frac{1}{2})^2 + 2\pi i(n+\frac{1}{2})(z+\frac{1}{2})} . \tag{3.9}$$

Furthermore,

$$\xi_I = \frac{1}{16\pi^2}g_6 \Lambda^2 \left(\frac{1}{4} \text{tr}(q) + \text{tr}(q_I) \right) \tag{3.10}$$

is the quadratically divergent piece of the FI term with q the charges of the bulk fields under the corresponding $U(1)$, q_I the charges of fields localized at the fixed point I and Λ the cutoff of the theory. We will ignore the first term in the sum of the right-hand side of (3.10), i.e. we assume $\text{tr} q = 0$. We further take Λ to be the higher-dimensional Planck scale, $\Lambda^2 = M_P/\sqrt{V_{56}}$ with $M_P = 2.43 \cdot 10^{18}$ GeV and $V_{56} = 2\pi^2 R_5 R_6$ the volume of the extra dimensions. In addition we have the usual relation between the higher-dimensional gauge coupling and its four-dimensional counterpart, $g_6 = \sqrt{V_{56}}g$. The constant f can be determined through the normalization condition

$$\int_0^{\pi R_5} dx_5 \int_0^{2\pi R_6} dx_6 |\psi|^2 = 1 . \tag{3.11}$$

Let us note that the existence of the localization effect does not require $U(1)_{\text{LNZ}}$ to be unbroken down to low energies. Rather, the FI terms are dominated by loop momenta close to the cut-off scale such that $U(1)_{\text{LNZ}}$ broken somewhat below Λ will yield almost identical results.

In our case we consider the orbifold $\mathbb{T}^2/\mathbb{Z}_2$ with one Wilson line in the y_5 direction. This means that there are two pairs of equivalent fixed points, so all FI terms will have the same absolute value, given that the effective FI term in four dimensions vanishes. Then the lightest Kaluza-Klein masses are either $M = 1/(2R_5)$ or $M = 1/R_6$, due to the presence of the Wilson line. Since we are interested in the limit $R_5 \geq R_6$ and since the GUT group gets broken by the Wilson line, we identify the GUT scale with the corresponding Kaluza-Klein mass, $M_{\text{GUT}} = 1/(2R_5)$.

Let us now briefly compare the terms giving rise to the top Yukawa and gauge couplings, respectively. While the first is proportional to the overlap integral over $h_u q_3 \bar{u}_3$, the latter scales like the integral over $A \bar{u}_3 \bar{u}_3^\dagger$. The Higgs field itself retains a flat profile since it comes from a non-Abelian gauge multiplet. Taking further into account that the profile of \bar{u}_3^\dagger is identical to the one of \bar{u}_3 , while the profiles of q_3 and \bar{u}_3 are inverse to each other, it is clear that the top Yukawa coupling will always be reduced with respect to the gauge coupling, when the correct normalization is taken into account. This effect is illustrated in figure 3. As can be seen, the reduction depends on the given charges and is more pronounced for anisotropic compactifications. This behavior is to be expected,




Figure 3. Suppression of the top Yukawa coupling y_t relative to the gauge coupling g due to the localization effects as a function of the anisotropy. The inlay indicates that the region $R_5 > R_6$, where the distance between equivalent fixed points with coinciding traces is smaller than the one between inequivalent fixed points, is considered. In the other limit, i.e. for $R_6 \gg R_5$, there is no analogous suppression.

since in the limit $R_6 \rightarrow 0$ the 5D case should be recovered, where the bulk fields effectively become brane fields and hence the overlap should vanish. This is because in this limit the bulk fields get exponentially localized towards the opposite ends of the interval [36], such that the overlap becomes exponentially suppressed.

4 Explicit string theory realization

Our analysis is motivated by recent progress in string model building [23–28], where $\mathcal{O}(100)$ explicit (and globally consistent) models with the exact spectra of the MSSM, the so-called ‘heterotic mini-landscape’, have been derived. In a subclass of these models the top (but neither the bottom nor the τ) Yukawa coupling is related to the gauge coupling. This applies in particular to the two models that have been analyzed in some detail, [23, 24, 26, 28] and the ‘benchmark model 1A’ in [27]. In what follows, we will study gauge-top unification in these models. Our analysis should be viewed as the first step towards a full string theory calculation, where one computes the FI term in string theory, takes into account arbitrary values for the torus parameter τ and the full volume of the six compact dimensions. In the present study, we will restrict ourselves to the somewhat naive orbifold GUT picture, and will take the cutoff Λ to be the 6D Planck scale, as before.

4.1 Motivation of the orbifold GUT picture

Taking the orbifold GUT limit of a string compactification has a rather profound motivation. Witten proposed in a footnote [39, footnote 3] a possible way to explain the

discrepancy between the string and the GUT scales. This can be accomplished by considering highly anisotropic compactifications, where M_{GUT} is associated to the inverse of the largest radius, while all (or most of) the other radii are much smaller. In this case, the volume of compact space can be small enough to ensure that the perturbative description of the setting is still appropriate. This idea has been studied in some detail more recently [40]. The outcome of the analysis is that the above puzzle can be resolved if the largest radius is by a factor 50 or so larger than the other radii. The question of how to stabilize the largest radius has also been addressed in the framework of field theory, and it has been found that one can indeed obtain $R_5 \simeq 1/(2M_{\text{GUT}})$ [41–43]. Also here localized FI terms can play an important role. The question why one radius behaves so differently from the others will be addressed elsewhere [44]. In what follows, we will build on these results, and consider 6D orbifold GUT limits with particular emphasis on the highly anisotropic case $R_5 \gg R_6$. The transverse four dimensions will not be considered.

4.2 A specific example

For concreteness, let us first focus on the benchmark model 1A of [27]. In order to be in accord with the discussion in section 3.3 (and the analysis by LNZ [33]), here and below we take only U(1) factors orthogonal to the so-called ‘anomalous U(1)’ direction \mathbf{t}_{anom} into account. The general case will be explored elsewhere. One can choose the basis of these U(1)s such that the local FI terms at the fixed points are entirely in one U(1) direction, called U(1)_{LNZ} in what follows. The corresponding generator reads

$$\mathbf{t}_{\text{LNZ}} = \frac{1}{\sqrt{105}} \left(-\frac{3}{2}, 0, 0, 0, 0, 0, 0 \right) \left(-\frac{11}{4}, \frac{21}{4}, 0, \frac{11}{4}, 0, \frac{11}{4}, 0, 0 \right), \quad (4.1)$$

and is normalized such that $|\mathbf{t}_{\text{LNZ}}|^2 = 1/2$. In this normalization, the charges of φ and φ^c , containing q_3 and \bar{u}_3 , are

$$q_\varphi = -q_{\varphi^c} = \frac{1}{\sqrt{105}} \frac{3}{4}, \quad (4.2)$$

respectively. One further has

$$\text{tr}_{\text{SU}(5) \text{ brane}} q_{\text{LNZ}} = \frac{64}{\sqrt{105}} \quad \text{and} \quad \text{tr}_{\text{SU}(4) \times \text{SU}(2) \text{ brane}} q_{\text{LNZ}} = -\frac{64}{\sqrt{105}}, \quad (4.3)$$

where the SU(5) and SU(4) × SU(2) branes comprise the fixed points with $n_2 = 0$ and 1, respectively, with the fixed points carrying half of the trace. We can hence apply the results of section 3.3 in order to evaluate y_t . The ratio y_t/g for the present example is shown in figure 4. For this figure, we used our field theoretic estimate of the FI term, i.e. chose to identify Λ with the 6D Planck scale, as before. As discussed in section 3.3, the localization effect becomes more pronounced when the ratio R_5/R_6 increases. We also show what the suppression would be if we chose an increased cut-off, $\Lambda^2 = 4M_{\text{P}}/\sqrt{V_{56}}$, designed in such a way that for $R_5 = 50 R_{>5}$ Λ equals the heterotic string scale, $M_{\text{string}} \simeq 8 \cdot 10^{17}$ GeV, as in the proposal discussed by Hebecker and Trapletti [40].

It is amazing to see that the top Yukawa coupling gets only reduced if we take the ‘more appealing’ orbifold GUT limit $R_5 \gg R_6$. If we were to choose $R_6 \gg R_5$, which would




Figure 4. Estimate of y_t/g in the benchmark model 1A of [27]. For comparison, the dashed line shows the suppression which occurs with an increased cut-off, $\Lambda \rightarrow 2\Lambda$.

lead us to an orbifold GUT with $SU(3) \times SU(3)$ bulk group that gets broken to the SM at the two equivalent boundaries, the top Yukawa coupling would no longer be suppressed since $\text{tr } q_I$ vanishes at both boundaries.

4.3 Mini-landscape survey

We have repeated the analysis of section 4.2 for a subclass of the heterotic mini-landscape models [25, 27], in which

- there is an orbifold GUT limit as described in section 2;
- it has been explicitly verified that exotics decouple consistently with supersymmetry, i.e. with vanishing F - and D -terms.

All these models, including the model derived in [23, 24, 26, 28] and the ‘benchmark model 1A’ discussed before, are based on the same shift vector. They turn out to have the following family structure (up to vector-like states):

- 1st and 2nd families come from **16**-plets localized at $SO(10)$ fixed points, which correspond to the fixed points with $n_2 = 0$ in the orbifold GUT limit (figure 1);
- 3rd family \bar{d} and ℓ (i.e. the 3rd family $\bar{\mathbf{5}}$ in $SU(5)$ language) come from the $T_{2/4}$ twisted sectors and therefore are localized on two-dimensional submanifolds in compact 6D space;
- 3rd family \bar{u} , \bar{e} and q as well as the Higgs fields h_u and h_d are bulk fields, i.e. free to propagate everywhere in compact space.

Only 4 out of 56 candidate models do not have localized FI terms.

Yukawa couplings connecting the Higgs fields to matter may be written as overlap integrals. One could then expect that the couplings of the first two generations are suppressed by the total 6D volume while the τ and b Yukawas, y_τ and y_b , are suppressed by the volume of the 4D space transverse to the two-dimensional submanifold, while the top Yukawa y_t is unsuppressed, as discussed before. This leads us to the hierarchy

$$\text{Yukawa couplings of the first two generations} \ll y_\tau, y_b \ll y_t.$$

It should be mentioned that these localization properties are highly non-trivial (and come out ‘for free’). In fact, the cancellation of all 4D and higher-dimensional gauge anomalies, which is guaranteed by modular invariance [45], appears rather miraculous in field theory [26]. Also discrete anomalies have been shown to cancel [46].

Let us now come back to the question of the value of y_t at the compactification or GUT scale. As before in section 4.2, it is possible to rotate the basis of U(1) factors (orthogonal to $\mathfrak{t}_{\text{anom}}$) such that there is precisely one U(1), which we again call U(1)_{LNZ}, with the following properties:

- the chiral multiplets φ and φ^c have non-zero (and opposite) U(1)_{LNZ} charges;
- the traces $\text{tr}_{\text{SU}(5) \text{ brane}} q_{\text{LNZ}}$ and $\text{tr}_{\text{SU}(4) \times \text{SU}(2) \text{ brane}} q_{\text{LNZ}}$ have opposite sign (as in (4.3));
- $|\mathfrak{t}_{\text{LNZ}}|^2 = \frac{1}{2}$, i.e. the charges are in “GUT normalization”.

The statistics of the localized traces and the charges of the bulk hypermultiplets φ and φ^c are depicted in figure 5.

5 Phenomenological implications

Let us now come to some phenomenological implications of the gauge-top unification scenario. Given a certain pattern of soft terms as well as the top Yukawa coupling at the GUT scale, the value of $\tan \beta$ can be determined. Note that the value of $\tan \beta$ is also quite sensitive to the mass of the top quark, where the latest experimental value is given by $m_t = 173.1 \pm 1.3 \text{ GeV}$ [32].

It turns out that the resulting value for $\tan \beta$ mainly depends on $y_t(M_{\text{GUT}})$. To illustrate this dependence we plot the top Yukawa coupling at the GUT scale versus $\tan \beta$ for different values of the top quark mass in figure 6. We impose a specific set of boundary conditions at the GUT scale, the so-called mSUGRA set with $m_0 = m_{1/2} = -A_0 = 1 \text{ TeV}$, and use SOFTSUSY [47] for our numerical analysis.

The value of the unified gauge coupling on the other hand turns out to be only very weakly dependent on the soft parameters and is always close to $g \simeq 0.7$. This implies that, given the tree-level relation $y_t = g \simeq 0.7$, the resulting value for $\tan \beta$ is quite small, $\tan \beta \simeq 2$. An immediate question is whether such small values for $\tan \beta$ and hence the tree-level relation are still valid, since the small $\tan \beta$ region is probed experimentally by




Figure 5. Statistics of $|q_\varphi \text{tr } q_{\text{LNZ}}|_{n_2=n'_2=0}$ in a class of string-derived orbifold GUTs and suppression of the top Yukawa coupling for the average value.




Figure 6. (a) $\tan \beta$ as a function of the top Yukawa coupling at the GUT scale. The central line corresponds to the central value of the top quark mass, whereas the dashed and dotted lines correspond to the one and two sigma intervals, respectively. A larger top mass results in a larger $\tan \beta$. (b) shows for given $q_\varphi \text{tr } q_I$ the anisotropy that is needed to reduce y_t/g to 0.75.

searches for the light Higgs boson [48]. This tension can be seen by comparing the LEP bound on the SM Higgs mass, $m_h \geq 114.4 \text{ GeV}$, to the theoretical upper bound on the lightest Higgs mass as a function of $\tan \beta$. At tree-level the maximal mass for the lightest

Higgs is given by $m_h^2 \simeq M_Z^2 \cos^2 2\beta$, which vanishes for $\tan \beta = 1$. Radiative corrections can significantly increase the Higgs mass compared to the tree-level value, but still m_h is minimized for $\tan \beta$ around one in the MSSM. While the LEP Higgs bound does not apply for all of the MSSM parameter space, in the small $\tan \beta$ region it is applicable to a good approximation, since here the lightest CP-even Higgs boson couples to the Z with SM-like strength [49]. For the so called ‘ m_h^{\max} scenario’ within the MSSM, designed such that for fixed values of m_t and M_{SUSY} the predicted value of the lightest Higgs is maximized for each value of $\tan \beta$ and m_A , the lower bound on $\tan \beta$ is around 2 [49]. Although hence values of $\tan \beta$ around 2 are still possible, one should not forget that this is possible only when the parameters are tuned accordingly. In a large part of the parameter space $\tan \beta$ has to be larger. To see what is natural as the ‘smallest value of $\tan \beta$ without too much tuning’, we restrict ourselves to mSUGRA. Taking $m_{1/2} = m_0 = -A_0 = 1 \text{ TeV}$ and the top mass at its two sigma upper bound, the predicted Higgs mass is above the LEP bound for $\tan \beta \gtrsim 3.3$. This in turn implies a ‘natural range’ for the top Yukawa coupling at the GUT scale of $0.48 \lesssim y_t \lesssim 0.6$, which translates into $0.69 \lesssim y_t/g \lesssim 0.86$. Note that in the presence of vector-like matter (in complete $SU(5)$ representations) with masses below the GUT scale the relative suppression of y_t has to be stronger.

In conclusion we can say that the naive picture of gauge-top unification, although not excluded, seems to be possible for very special patterns of the soft terms only. However, we have seen in the previous sections that the top Yukawa coupling gets somewhat reduced with respect to the tree-level relation $y_t = g$, depending on the geometry of the extra-dimensional space. Turning this around means that, given a value for $\tan \beta$, we gain access to the geometry of the extra-dimensional space. In particular it seems that highly anisotropic compactifications are favored, in accord with [40].

6 Discussion

We have discussed orbifold GUT scenarios in which the top Yukawa coupling arises from gauge interactions. This leads to the tree-level relation $y_t = g$ at the GUT or compactification scale. It turns out that in scenarios with localized FI terms y_t is suppressed against g , which seems also to be required by data. The suppression depends on the size of the localized FI terms, the charge of the bulk hypermultiplets and, in particular, on the geometry of compact space. We have analyzed string-derived models which reproduce the MSSM below the compactification scale and which possess orbifold GUT limits with the above features. This allowed us to get a feeling for what the FI terms and charges ‘should be’. Using this as an input, we find that the observed value of the top Yukawa coupling seems to favor anisotropic geometries, which might also be a key ingredient for reconciling the scale of grand unification with the Planck or string scales. This result might be interpreted as further support for the idea of orbifold GUTs and, in particular, for their embedding in the heterotic string.

As mentioned, our analysis should be viewed as a step towards getting sufficiently accurate predictions from string theory. The precision of the present analysis is mainly limited by our ignorance of the cut-off Λ , for which we simply used the 6D Planck scale. One

can improve the accuracy significantly by directly computing the FI terms in string theory. Another important question is why, from a top-down perspective, one radius behaves so differently from the others. We plan to clarify these issues in the near future.

Acknowledgments

We would like to thank H.P. Nilles, P.K.S. Vaudrevange and M. Winkler for useful discussions and H.M. Lee for correspondence. This research was supported by the DFG cluster of excellence Origin and Structure of the Universe, and the SFB-Transregio 27 "Neutrinos and Beyond" by Deutsche Forschungsgemeinschaft (DFG). The work of P.H. was supported by the Alexander von Humboldt foundation.

References

- [1] H. Georgi and S.L. Glashow, *Unity of all elementary particle forces*, *Phys. Rev. Lett.* **32** (1974) 438 [SPIRES].
- [2] H. Fritzsch and P. Minkowski, *Unified interactions of leptons and hadrons*, *Ann. Phys.* **93** (1975) 193 [SPIRES].
- [3] L.J. Dixon, J.A. Harvey, C. Vafa and E. Witten, *Strings on orbifolds*, *Nucl. Phys. B* **261** (1985) 678 [SPIRES].
- [4] L.J. Dixon, J.A. Harvey, C. Vafa and E. Witten, *Strings on orbifolds. 2*, *Nucl. Phys. B* **274** (1986) 285 [SPIRES].
- [5] L.E. Ibáñez, H.P. Nilles and F. Quevedo, *Orbifolds and Wilson lines*, *Phys. Lett. B* **187** (1987) 25 [SPIRES].
- [6] L.E. Ibáñez, J.E. Kim, H.P. Nilles and F. Quevedo, *Orbifold compactifications with three families of $SU(3) \times SU(2) \times U(1)^n$* , *Phys. Lett. B* **191** (1987) 282 [SPIRES].
- [7] J.A. Casas, E.K. Katehou and C. Muñoz, *$U(1)$ charges in orbifolds: anomaly cancellation and phenomenological consequences*, *Nucl. Phys. B* **317** (1989) 171 [SPIRES].
- [8] J.A. Casas and C. Muñoz, *Three generation $SU(3) \times U(2) \times U(1)$ - Y models from orbifolds*, *Phys. Lett. B* **214** (1988) 63 [SPIRES].
- [9] A. Font, L.E. Ibáñez, H.P. Nilles and F. Quevedo, *Degenerate orbifolds*, *Nucl. Phys. B* **307** (1988) 109 [Erratum *ibid.* **B 310** (1988) 764]. [SPIRES].
- [10] A. Font, L.E. Ibáñez, F. Quevedo and A. Sierra, *The construction of 'realistic' four-dimensional strings through orbifolds*, *Nucl. Phys. B* **331** (1990) 421 [SPIRES].
- [11] Y. Kawamura, *Gauge symmetry reduction from the extra space S^1/Z_2* , *Prog. Theor. Phys.* **103** (2000) 613 [hep-ph/9902423] [SPIRES].
- [12] Y. Kawamura, *Triplet-doublet splitting, proton stability and extra dimension*, *Prog. Theor. Phys.* **105** (2001) 999 [hep-ph/0012125] [SPIRES].
- [13] G. Altarelli and F. Feruglio, *$SU(5)$ grand unification in extra dimensions and proton decay*, *Phys. Lett. B* **511** (2001) 257 [hep-ph/0102301] [SPIRES].
- [14] L.J. Hall and Y. Nomura, *Gauge unification in higher dimensions*, *Phys. Rev. D* **64** (2001) 055003 [hep-ph/0103125] [SPIRES].

- [15] A. Hebecker and J. March-Russell, *A minimal $S^1/(Z_2 \times Z_2)$ orbifold GUT*, *Nucl. Phys. B* **613** (2001) 3 [[hep-ph/0106166](#)] [[SPIRES](#)].
- [16] T. Asaka, W. Buchmüller and L. Covi, *Gauge unification in six dimensions*, *Phys. Lett. B* **523** (2001) 199 [[hep-ph/0108021](#)] [[SPIRES](#)].
- [17] L.J. Hall, Y. Nomura, T. Okui and D. Tucker-Smith, *SO(10) unified theories in six dimensions*, *Phys. Rev. D* **65** (2002) 035008 [[hep-ph/0108071](#)] [[SPIRES](#)].
- [18] G. Burdman and Y. Nomura, *Unification of higgs and gauge fields in five dimensions*, *Nucl. Phys. B* **656** (2003) 3 [[hep-ph/0210257](#)] [[SPIRES](#)].
- [19] T. Kobayashi, S. Raby and R.-J. Zhang, *Constructing 5D orbifold grand unified theories from heterotic strings*, *Phys. Lett. B* **593** (2004) 262 [[hep-ph/0403065](#)] [[SPIRES](#)].
- [20] S. Förste, H.P. Nilles, P.K.S. Vaudrevange and A. Wingarter, *Heterotic brane world*, *Phys. Rev. D* **70** (2004) 106008 [[hep-th/0406208](#)] [[SPIRES](#)].
- [21] T. Kobayashi, S. Raby and R.-J. Zhang, *Searching for realistic 4d string models with a Pati-Salam symmetry: orbifold grand unified theories from heterotic string compactification on a Z_6 orbifold*, *Nucl. Phys. B* **704** (2005) 3 [[hep-ph/0409098](#)] [[SPIRES](#)].
- [22] W. Buchmüller, K. Hamaguchi, O. Lebedev and M. Ratz, *Dual models of gauge unification in various dimensions*, *Nucl. Phys. B* **712** (2005) 139 [[hep-ph/0412318](#)] [[SPIRES](#)].
- [23] W. Buchmüller, K. Hamaguchi, O. Lebedev and M. Ratz, *Supersymmetric standard model from the heterotic string*, *Phys. Rev. Lett.* **96** (2006) 121602 [[hep-ph/0511035](#)] [[SPIRES](#)].
- [24] W. Buchmüller, K. Hamaguchi, O. Lebedev and M. Ratz, *Supersymmetric standard model from the heterotic string. II*, *Nucl. Phys. B* **785** (2007) 149 [[hep-th/0606187](#)] [[SPIRES](#)].
- [25] O. Lebedev et al., *A mini-landscape of exact MSSM spectra in heterotic orbifolds*, *Phys. Lett. B* **645** (2007) 88 [[hep-th/0611095](#)] [[SPIRES](#)].
- [26] W. Buchmüller, C. Lüdeling and J. Schmidt, *Local SU(5) unification from the heterotic string*, *JHEP* **09** (2007) 113 [[arXiv:0707.1651](#)] [[SPIRES](#)].
- [27] O. Lebedev et al., *The heterotic road to the MSSM with R parity*, *Phys. Rev. D* **77** (2008) 046013 [[arXiv:0708.2691](#)] [[SPIRES](#)].
- [28] W. Buchmüller and J. Schmidt, *Higgs versus matter in the heterotic landscape*, *Nucl. Phys. B* **807** (2009) 265 [[arXiv:0807.1046](#)] [[SPIRES](#)].
- [29] M. Ratz, *Notes on local grand unification*, [arXiv:0711.1582](#) [[SPIRES](#)].
- [30] H.P. Nilles, S. Ramos-Sanchez, M. Ratz and P.K.S. Vaudrevange, *From strings to the MSSM*, *Eur. Phys. J. C* **59** (2009) 249 [[arXiv:0806.3905](#)] [[SPIRES](#)].
- [31] I. Gogoladze, Y. Mimura and S. Nandi, *Unification of gauge, Higgs and matter in extra dimensions*, *Phys. Lett. B* **562** (2003) 307 [[hep-ph/0302176](#)] [[SPIRES](#)].
- [32] TEVATRON ELECTROWEAK WORKING GROUP collaboration, *Combination of CDF and D0 results on the mass of the top quark*, [arXiv:0903.2503](#) [[SPIRES](#)].
- [33] H.M. Lee, H.P. Nilles and M. Zucker, *Spontaneous localization of bulk fields: the six-dimensional case*, *Nucl. Phys. B* **680** (2004) 177 [[hep-th/0309195](#)] [[SPIRES](#)].
- [34] N. Arkani-Hamed, T. Gregoire and J.G. Wacker, *Higher dimensional supersymmetry in 4D superspace*, *JHEP* **03** (2002) 055 [[hep-th/0101233](#)] [[SPIRES](#)].
- [35] H. Georgi, *The state of the art-gauge theories*, in *Particles and Fields 1974*, ed. C.E. Carlson, AIP, New York U.S.A. (1975), pg. 575.

- [36] S. Groot Nibbelink, H.P. Nilles and M. Olechowski, *Instabilities of bulk fields and anomalies on orbifolds*, *Nucl. Phys. B* **640** (2002) 171 [[hep-th/0205012](#)] [[SPIRES](#)].
- [37] S. Groot Nibbelink, H.P. Nilles, M. Olechowski and M.G.A. Walter, *Localized tadpoles of anomalous heterotic U(1)'s*, *Nucl. Phys. B* **665** (2003) 236 [[hep-th/0303101](#)] [[SPIRES](#)].
- [38] D. Mumford, *Tata lectures on theta I*, first edition, Birkhäuser (1983).
- [39] E. Witten, *Strong coupling expansion of Calabi-Yau compactification*, *Nucl. Phys. B* **471** (1996) 135 [[hep-th/9602070](#)] [[SPIRES](#)].
- [40] A. Hebecker and M. Trapletti, *Gauge unification in highly anisotropic string compactifications*, *Nucl. Phys. B* **713** (2005) 173 [[hep-th/0411131](#)] [[SPIRES](#)].
- [41] W. Buchmüller, R. Catena and K. Schmidt-Hoberg, *Small extra dimensions from the interplay of gauge and supersymmetry breaking*, *Nucl. Phys. B* **804** (2008) 70 [[arXiv:0803.4501](#)] [[SPIRES](#)].
- [42] C. Gross and A. Hebecker, *A realistic unified gauge coupling from the micro-landscape of orbifold GUTs*, [arXiv:0812.4267](#) [[SPIRES](#)].
- [43] W. Buchmüller, R. Catena and K. Schmidt-Hoberg, *Enhanced symmetries of orbifolds from moduli stabilization*, [arXiv:0902.4512](#) [[SPIRES](#)].
- [44] R. Kappl, S. Ramos-Sánchez, M. Ratz and P.K.S. Vaudrevange, in preparation (2009).
- [45] C. Vafa, *Modular invariance and discrete torsion on orbifolds*, *Nucl. Phys. B* **273** (1986) 592 [[SPIRES](#)].
- [46] T. Araki et al., *(Non-)Abelian discrete anomalies*, *Nucl. Phys. B* **805** (2008) 124 [[arXiv:0805.0207](#)] [[SPIRES](#)].
- [47] B.C. Allanach, *SOFTSUSY: a C++ program for calculating supersymmetric spectra*, *Comput. Phys. Commun.* **143** (2002) 305 [[hep-ph/0104145](#)] [[SPIRES](#)].
- [48] S. Heinemeyer, W. Hollik and G. Weiglein, *Constraints on $\tan\beta$ in the MSSM from the upper bound on the mass of the lightest Higgs boson*, *JHEP* **06** (2000) 009 [[hep-ph/9909540](#)] [[SPIRES](#)].
- [49] G. Degrandi, S. Heinemeyer, W. Hollik, P. Slavich and G. Weiglein, *Towards high-precision predictions for the MSSM Higgs sector*, *Eur. Phys. J. C* **28** (2003) 133 [[hep-ph/0212020](#)] [[SPIRES](#)].