

UC Irvine

UC Irvine Previously Published Works

Title

Post-Ischemic Inhibition of the Mitochondrial Ca^{2+} Uniporter with Ru265 Protects in Ex Vivo and In Vivo Models of Cerebral Ischemia.

Permalink

<https://escholarship.org/uc/item/02n695fs>

Journal

Clinical Science

ISSN

0143-5221

Authors

Medvedeva, Yuliya V
Yin, Hong Z
Bazrafkan, Afsheen
[et al.](#)

Publication Date

2026-09-23

DOI

10.1042/cs20261842

Copyright Information

This work is made available under the terms of a Creative Commons Attribution License, available at <https://creativecommons.org/licenses/by/4.0/>

Peer reviewed

Post-Ischemic Inhibition of the Mitochondrial Ca^{2+} Uniporter with Ru265 Protects in Ex Vivo and In Vivo Models of Cerebral Ischemia.

Yuliya V. Medvedeva¹, Hong Z. Yin¹, Afsheen Bazrafkan¹, Masih A. Rafi¹, Evelyn Wang¹, Ivan A. Yeromin¹, Edward Sharman¹, Humza Ghaznavi¹, Sepehr Khatami¹, Kantapon Suayngam¹, Alexander L. Liu¹, Najwa W. Ellahib¹, Renee E. Lotfy¹, Justin J. Wilson⁵, Yama Akbari^{1,2,3,4}, and John H. Weiss^{1,2}

Affiliations:

¹ Department of Neurology, University of California Irvine, Irvine, CA 92697, United States.

² Department of Anatomy & Neurobiology, University of California Irvine, Irvine, CA 92697, United States.

³ Department of Neurological Surgery, University of California Irvine, Irvine, CA 92697, United States.

⁴ The Beckman Laser Institute & Medical Clinic, University of California Irvine, Irvine, CA 92697, United States.

⁵ Department of Chemistry and Biochemistry, University of California, Santa Barbara, 93106, Santa Barbara, CA, United States.

Abstract

Brain ischemia is a leading cause of morbidity and mortality in the aging population, yet no effective neuroprotective interventions currently exist. This therapeutic gap stems from two key challenges: the inability to deliver neuroprotective drugs before reperfusion and limited understanding of post-reperfusion events leading to delayed neurodegeneration. Zn^{2+} is an important contributor to ischemic brain injury, with mitochondrial Zn^{2+} uptake playing a central role in its neurotoxic effects. Ongoing Zn^{2+} accumulation occurs through the mitochondrial Ca^{2+} uniporter (MCU) in selectively vulnerable CA1 neurons after transient ischemia, contributing to delayed mitochondrial dysfunction. We previously demonstrated that post-ischemic administration of the non-selective MCU antagonist, ruthenium red attenuated neuronal damage in both an *ex vivo* oxygen glucose deprivation (OGD) mouse brain slice model and in an *in vivo* rat cardiac arrest (CA) model. In the present study, we test post-ischemic treatment with Ru265, a novel selective MCU blocker, in both ischemic models. In the slice model, OGD induced delayed Zn^{2+} accumulation in CA1 neuronal mitochondria (as previously reported), followed by late-onset persistent mitochondrial depolarization and sustained elevation of cytoplasmic Ca^{2+} levels, indicating loss of Ca^{2+} homeostasis. Application of Ru265 after OGD termination inhibited the mitochondrial Zn^{2+} accumulation and largely prevented subsequent mitochondrial depolarization and loss of cytosolic Ca^{2+} homeostasis. In the CA model, intravenous Ru265 administration during cardiopulmonary resuscitation largely mitigated mitochondrial damage, markedly reduced neuronal injury and improved behavioral outcomes 20-24 hours after ischemia. These findings suggest that post-ischemic MCU blockade may constitute a promising neuroprotective therapeutic strategy.

Corresponding Authors:

J. H. Weiss, MD, Ph.D.

Phone: (949) 824-6774

Email: jweiss@uci.edu

Yuliya V. Medvedeva, Ph.D.

Phone: (949) 824-8479

Email: yvmedved@uci.edu

Keywords: hippocampal slice, mitochondria, zinc, ischemia, mitochondrial Ca^{2+} uniporter, MCU, Ru265, oxygen glucose deprivation, cardiac arrest

Acknowledgements: Supported by NIH grant NS121227 (JHW) and Roneet Carmell Memorial Endowment Fund (YA). We wish to thank Joshua Woods for his contributions to the development of Ru265, and Miranda N. Cepkinian, Anushka Singhal, Catherine T. Vu, Dan T. Vo, Dylan T. Ngo, and Adelaide Hong for contributing to animal cardiac arrest experiments and data analysis. The authors declare no competing financial interests.

Introduction

Brain ischemia remains a leading cause of morbidity and mortality, particularly among the aging population. Despite decades of research, effective neuroprotective interventions have yet to be established in clinical practice [1]. To date, cardiopulmonary resuscitation (CPR) represents the only clinically proven treatment for improving neurological outcomes following cardiac arrest (CA). Although therapeutic hypothermia after resuscitation offered hope in early clinical studies, recent trials have questioned its benefits [2]. Similarly, prompt restoration of cerebral blood flow remains the sole evidence-based intervention for ischemic stroke. However, recent advances in cerebral revascularization techniques following stroke [3] have extended the therapeutic window and created new opportunities for adjunctive neuroprotective therapies.

Until recently, most treatments demonstrated to be effective in animal models were administered before the onset of ischemia. In clinical settings, however, opportunities for intervention typically arise only after the restoration of blood flow. Notably, following brief ischemia, select neuronal populations - particularly those in the hippocampal CA1 region and cortical layers 3, 5, and 6 - initially survive but subsequently undergo delayed neuronal death over the ensuing hours to days, a phenomenon well-documented in both rodents and humans [4-9]. These selectively vulnerable neurons represent potential targets for therapeutic salvage with timely interventions. Therefore, elucidating the sequence of post-ischemic molecular and cellular events that drive neurodegeneration in these vulnerable populations is essential for developing effective neuroprotective strategies.

Mitochondria have been consistently identified as primary targets of ischemic injury. Transient ischemia induces early mitochondrial damage and dysfunction, culminating in ATP depletion, reactive oxygen species (ROS) overproduction, and the release of pro-apoptotic mediators [10-12].

Accumulating evidence has implicated Zn^{2+} -mitochondria interactions as a key contributor to ischemic neuronal injury [13, 14]. Consistent with this, we recently observed prolonged Zn^{2+} accumulation within mitochondria of selectively vulnerable CA1 neurons - but not in the comparatively resistant CA3 neurons - occurring after transient ischemia [15]. Notably, delayed administration of either the Zn^{2+} chelator N,N,N',N'-tetrakis(2-pyridylmethyl)ethylenediamine (TPEN) or ruthenium red (RR), an antagonist of the mitochondrial Ca^{2+} uniporter (MCU; the major route for Zn^{2+} entry to mitochondria), attenuated both mitochondrial and neuronal damage at 4 hours post-ischemia in the *ex vivo* acute hippocampal slice model. RR was further shown to confer neuroprotection in an *in vivo* rat cardiac arrest model [16]. However, RR is highly non-selective, exerting numerous off-target effects beyond MCU blockade, and is unsuitable for human use [17]. Likewise, the alternative MCU inhibitor RU360 presents significant translational limitations, as it is membrane-impermeable and susceptible to redox transformations that render it inactive [18].

The present study investigated the neuroprotective potential of Ru265 [19], a recently synthesized MCU inhibitor that is substantially more potent, selective, and cell-permeable than previously available MCU blockers, offering considerable translational potential. Acute hippocampal slices were subjected to transient oxygen-glucose deprivation (OGD, 8 min) followed by reperfusion to model ischemia-reperfusion injury. Brief OGD induced persistent Zn^{2+} accumulation in CA1 neuronal mitochondria occurring after reperfusion, consistent with our previous observations [15], along with delayed injury at 4 hours post-reperfusion, including persistent mitochondrial depolarization and sustained elevation of cytoplasmic $[\text{Ca}^{2+}]$, indicative of disrupted Ca^{2+} homeostasis.

Delayed post-OGD application of Ru265 suppressed late mitochondrial Zn^{2+} accumulation and largely prevented both mitochondrial depolarization and loss of cytosolic Ca^{2+} homeostasis. In an *in vivo* rat CA model of transient global brain ischemia, intravenous Ru265 administration during CPR markedly attenuated ischemia-evoked mitochondrial swelling and fragmentation, diminished neuronal injury and improved behavioral outcome 20-24 hours post-ischemia.

Taken together, these findings provide compelling support for delayed administration of MCU antagonists as a therapeutic strategy to mitigate post-ischemic injury. Critically, such treatment can be administered following restoration of blood flow – a clinically relevant therapeutic window.

Materials and Methods

Reagents.

Ru265 is a novel, potent (IC₅₀ 2.6 nM), and membrane-permeable inhibitor of the MCU. Ru265 was synthesized as previously described [19, 20] by the laboratory of Justin Wilson. Ketamine was obtained from Sigma (St. Louis, MO). TOM20 antibodies were obtained from Santa Cruz Biotechnology (Santa Cruz, CA) and DyLight™ 488 antibodies were obtained from Jackson ImmunoResearch (West Grove, PA). All other reagents were purchased from Thermo Fisher Scientific (Waltham, MA).

Ethics Statement.

This study was carried out in accordance with the Guide for the Care and Use of Laboratory Animals of the National Institutes of Health and was approved by the Institutional Animal Care and Use Committee (IACUC) at the University of California, Irvine.

Animals.

Efforts were made to minimize animal numbers and suffering. Both male and female mice from 129S6/SvEvTac strain (Taconic Biosciences; 4-5 weeks old) were utilized for *ex vivo* experiments. All *in vivo* experiments were performed using adult male Wistar rats (Envigo, Philadelphia, PA), aged an average of approximately 66 days and weighing 363.35 ± 42.43 g. Rats were housed under standard laboratory conditions ($23 \pm 2^\circ\text{C}$, 60–70% humidity, 12 hour light/dark cycle) with ad libitum access to food and water. Animals were handled daily for two weeks prior to surgery and were calorically restricted to 25% of normal food intake for 14 hours prior to the cardiac arrest (CA) procedure, as standard with our prior publications [21, 22].

Preparation of acute brain slices.

Mice were deeply anesthetized with isoflurane, decapitated, and brains quickly removed and placed in ice-cold solution containing (in mM): 220 sucrose, 3 KCl, 1.25 NaH₂PO₄, 6 MgSO₄, 26 NaHCO₃, 0.2 CaCl₂, 10 Glucose and 0.42 ketamine (pH 7.35, 310 mOsm, equilibrated with 95% O₂ / 5% CO₂). Coronal brain slices (300 μm) were prepared as previously described [23] using a Leica VT1200 vibratome (Leica, Germany). After preparation, slices were incubated in artificial cerebrospinal fluid (ACSF) containing (in mM): 126 NaCl, 3 KCl, 1.25 NaH₂PO₄, 1 MgSO₄, 26 NaHCO₃, 2 CaCl₂, 10 Glucose (pH 7.35, 310 mOsm, equilibrated with 95% O₂ / 5% CO₂) for 1 hour at $34 \pm 0.5^\circ\text{C}$, and then kept at room temperature (20–23°C) in oxygenated ACSF. All experiments were performed at $32 \pm 0.5^\circ\text{C}$.

OGD/reperfusion in slices and Ru265 administration.

To induce short hypoxia-hypoglycemia, ACSF was changed to identical solution without glucose and equilibrated with 95% N₂ / 5% CO₂ (see Fig. 1A for experimental timeline). After 8 min, OGD was terminated by restoration of oxygenated ACSF containing glucose. The duration of OGD was determined by our previous experiments with the aim to withdraw OGD before Ca²⁺ deregulation occurs in the majority of CA1 neurons, while allowing sufficient time to evoke mitochondrial depolarization (loss of $\Delta\Psi_m$), Zn²⁺ accumulation and delayed (4 hours later) mitochondrial dysfunction [15, 23].

Ru265 (5 μM) was applied 3 min after the OGD termination for 30 min.

Fluorescent measurements.

Measurements of intracellular Zn²⁺ in individual hippocampal neurons. Slices were placed in a flow-through chamber (RC-27L, Warner Instruments) mounted on the stage of an upright microscope (BX51WI Olympus, Japan) and perfused with oxygenated ACSF at 2 ml/min. Membrane-impermeable Zn²⁺ indicator FluoZin-3 [24,

25] and Ca^{2+} indicator (either low affinity Fura-FF or high affinity Fura-2) were dissolved in pipette solution (containing, in mM: 125 K-Gluconate, 10 KCl, 3 Mg-ATP, 1 MgCl_2 , 10 HEPES, pH 7.25 with KOH, 290 mOsm with sucrose) at a concentration of 1 mM, and 1 μl placed in the tip of a patch-pipette (5–7 M Ω , borosilicate glass with filament), prior to backfilling with pipette solution. Neurons were loaded with indicator via patch-pipette which was slowly withdrawn after ~ 5 min, as previously described [23]. The intracellular Ca^{2+} level (as Fura-FF or Fura-2 ratio) was monitored during pipette withdrawal. If Fura ratio increased during withdrawal, even briefly, the cell was considered damaged and was not used. Fluorescence was excited at 340(20), 380(20) and 482(20) nm via a 40x water-immersion objective, using a Xenon light source (Sutter Instruments), and emitted fluorescence collected at 532 (40) nm using a cooled CCD camera (Hamamatsu, Japan) (All filters are bandpass with bandwidths indicated in parentheses.) Images were acquired every 15 or 30 sec. After background subtraction, images were analyzed using METAFLUOR 7.1 software (Universal Imaging). FluoZin-3 fluorescence changes are presented as $\Delta F/F_0 = (F - F_0)/F_0 \times 100\%$, where F is the current fluorescence intensity and F_0 is the average baseline fluorescence during the 10 min prior to OGD. The photobleaching-related decrease in FluoZin-3 fluorescence averaged 4–7 % per 10 min, depending on exposure time. If the rate of fluorescence decrease accelerated considerably during the experiment (more than 10%) the cell was withdrawn from the experiment. For background subtraction, we determined a fluorescence-free region for each recording located at least 100 μm from the loaded neuron(s). 1–2 neurons per slice were loaded, and only neurons that remained in sharp focus throughout the whole experiment were used.

To assess the level of mitochondrial Zn^{2+} accumulation following OGD, a high dose of carbonyl cyanide-4-(trifluoromethoxy)phenylhydrazone (FCCP, 2 μM , 5 min) was administered to the slice ~ 1 hour after OGD withdrawal. This treatment causes dissipation of the mitochondrial potential ($\Delta\Psi_m$) and release of ions accumulated within mitochondria to the cytoplasm, where they can be sensed by cytoplasmic fluorescent indicators. We assessed two parameters: the maximum amplitude of FluoZin-3 fluorescence increase within 15 min after the start of the FCCP application (see Fig. 2B); and the area under the FluoZin-3 curve, evaluated for 15 min starting from the onset of the of FCCP exposure (Fig. 2B).

Evaluation of changes in $\Delta\Psi_m$. To study changes in mitochondrial function we evaluated $\Delta\Psi_m$. Slices were bath loaded with the $\Delta\Psi_m$ -sensitive indicator, Rhodamine 123 (Rhod123, 26 μM , 30 min, 22–24 $^{\circ}\text{C}$). Rhod123 is positively charged and accumulates in negatively charged mitochondria where its fluorescence is quenched. Upon loss of $\Delta\Psi_m$ the indicator is released into the cytoplasm causing an increase in fluorescence [26]. FCCP (2 μM) was administered to control slices and 4 hours post-OGD to evoke full loss of $\Delta\Psi_m$ and a consequent increase in Rhod123 fluorescence, to evaluate the level of $\Delta\Psi_m$ before FCCP application. Rhod123 was excited at 540(25) nm, and emitted fluorescence collected at 605(55) nm. Images were acquired every 15 sec, background subtracted and analyzed. Data are presented as $\Delta F/F_0 = (F - F_0)/F_0 \times 100\%$. Regions of interest were monitored in the CA1 pyramidal cell layer. For background subtraction we used image of unloaded slices obtained using the same parameters as Rhod123-loaded slices.

Assessment of cytoplasmic Ca^{2+} levels using an expressible Ca^{2+} indicator. To evaluate the cytoplasmic Ca^{2+} concentration $[\text{Ca}^{2+}]_i$ in multiple CA1 neurons we used slices from Salsa6f/CaM mice. These mice express a protein-based ratiometric Ca^{2+} indicator in forebrain pyramidal neurons. ‘Salsa6f’ was produced by fusing a protein based Ca^{2+} indicator (GCaMP6f, Kd ~ 375 nM, green fluorescence) and a cell marker (tdTomato, red), permitting ratiometric measurement of cytoplasmic Ca^{2+} rises (as green/red fluorescence ratio). Transgenic mice with Salsa6f targeted to the Rosa26 locus were crossed with CaMKIIalpha-cre T29-1 transgenic mice, yielding expression of ‘Salsa6f’ to forebrain pyramidal neurons. GCaMP6f was excited at 482(20) nm and emitted fluorescence collected at 532(40) nm. TdTomato was excited at 540(25) nm, and emitted fluorescence collected at 605(55) nm. Changes in Ca^{2+} values are presented as the ratio of background subtracted GCaMP6f and tdTomato emission intensities. For background subtraction we used images of control, non-fluorescent slices, as for Rhod123 experiments.

CA/CPR and Ru265 administration.

The presented study utilizes an asphyxial CA+CPR model as previously described [27-29] (see Fig. 1B for experiment timeline). Shortly before experiment, rats were anesthetized with 2% isoflurane in a 50% O₂ / 50% N₂ mixture and intubated with a 14-gauge endotracheal tube (Braun Melsungen AG, Melsungen, Germany), connected to a mechanical ventilator and isoflurane vaporizer. The left femoral artery and vein were cannulated for continuous blood pressure monitoring and intravenous drug delivery. Invasive arterial blood pressure was measured continuously using a transducer (CWE Inc., Ardmore, PA). Baseline arterial blood gas (ABG) measurements (Abaxis, Union City, CA) were obtained prior to initiation of asphyxia to ensure that the arterial blood gases were at normal levels before CA.

After surgical preparation, the inhaled gas was switched to 100% O₂ for 2 minutes, after which the isoflurane was stopped to wash out anesthesia for 3 minutes, followed by the inlet disconnection from oxygen to allow room air to be mechanically delivered to the rat. Additionally, neuromuscular blockade was initiated by injection of 1 mL of intravenous Vecuronium (2 mg/kg) with 1 mL of heparinized saline. After 3 minutes of isoflurane washout, asphyxial CA was induced by turning the ventilator off and clamping the ventilator tubing. CA time was defined as systolic blood pressure <30 mm Hg and pulse pressure of 10 mm Hg or less. Duration of asphyxia was 8 min. In the last min of asphyxia, as the ventilator was being reconnected and turned on, epinephrine (0.01 mg/kg) and sodium bicarbonate (1 mmol/kg) were administered intravenously ahead of CPR to stimulate the sympathetic nervous system and manage acidosis, respectively [29]. CPR (manual sternal compressions at 180-240 per min) was performed after 8 minutes of asphyxia and continued until return of spontaneous circulation (ROSC). Our study used 3 groups of animals: 1) sham group (control, this group was subjected to all experimental procedure without asphyxia); 2) rats subjected to CA with placebo treatment; and 3) rats subjected to CA followed by intravenous administration of the MCU blocker Ru265 (1 mg/kg) at the time of CPR in order to reach the brain with ROSC (Fig. 1B), as we have demonstrated with other drugs administered during the CPR phase [28]. After obtaining post-CA ABG measurements 10 minutes after ROSC, vessels were decannulated and rats were extubated within the following 1-2 hours when weaned off the ventilator.

Post-CA care and behavioral assessment.

After ROSC, animals were monitored and maintained under standard conditions. At ~4.5-hour post-ROSC time point (after the behavioral testing), rats received subcutaneous injections of cefazolin (5 mg/kg), along with a mixture of lactated Ringer's solution and sterile saline (20 mL/kg) for hydration and infection prevention. Body weight for each rat was recorded after these injections and again at 24 hours post-ROSC to calculate overnight weight loss. After injections, rats were returned to the vivarium for recovery and continued monitoring. Neurological assessments using the Neurological Deficit Score (NDS) were conducted at ~4 and ~20 hours post-ROSC. All assessments were performed by lab members who were blinded to the experimental conditions. The NDS includes an evaluation of arousal, brainstem reflexes, motor and sensory function, behavior, and seizures, following the previously described evaluation criteria [28]. The rats were then perfused at 24 hour post-ROSC for brain tissue collection.

Histochemistry.

Cardiac perfusion. Twenty-four hours after CPR, animals were anesthetized with an intraperitoneal injection of sodium pentobarbital, and perfused transcardially with 200 ml of phosphate-buffered saline (PBS) for 2 min, followed by 500 ml of buffer containing 0.002% CaCl₂, 1.6% NaH₂PO₄ and 0.4% NaOH (pH 7.3), 4% PFA and 1% glutaraldehyde over the next 20 min. After perfusion, the brains were postfixed using 2% PFA / 2.5% glutaraldehyde at 4°C overnight, and 20 and 30 µm slices were cut using a vibratome (Leica VT1200).

Vanadium acid fuchsin (VAF) staining. To assess neuronal injury, 10 µm sections were stained with vanadium acid fuchsin (VAF) largely as previously described [30]. Slices were stained with VAF for 1–2 min, washed with

PBS, incubated in 0.01% borax solution for 20–30 seconds, and rinsed in distilled water. Finally, brain slices were cleared by acetate buffer (pH 3.3) for 30 sec and rinsed with distilled water. Stained slices were assessed and photographed using light microscopy. Cell counts of injured vs healthy neurons were carried out by an investigator blinded to experimental conditions.

Antibody labeling and confocal microscopy.

To assess mitochondrial morphology, 20 μm slices were stained with primary antibodies against the mitochondrial outer membrane protein TOM20 (1:200) and secondary anti-rabbit fluorescent antibodies (1:200, DyLight™ 488). The images of the hippocampal CA1 zone were taken using an inverted stage Olympus FV3000 confocal laser scanning microscope (Olympus, Japan) with a high-speed resonance scanner, IX3-ZDC2 Z-drift compensator and Olympus 40x silicone oil objective (UPLSAPO40XS, NA = 1.25, WD = 0.3 mm). A 488 nm diode laser was used for excitation and a high-sensitivity cooled GaAsP PMT was used for detection of signals.

An analysis of mitochondrial size and shape was performed blindly using coded images with ImageJ software as previously described [15]. To ensure that we evaluated only pyramidal neuronal mitochondria, only mitochondria in the immediate proximity of easily recognizable pyramidal neuronal nuclei were analyzed. Each single plane image was adjusted manually using brightness/contrast functions of ImageJ software to provide optimal discrimination of mitochondrial edges from background. We selected cells that showed clearly evident mitochondria in the perinuclear regions in a sharp focus. The mitochondria that were aligned with their long axes parallel to the nuclear membrane and round mitochondria in close nuclear proximity were assessed. Measures of long axes, parallel to, and short axes, perpendicular to nuclear membranes were obtained on all clearly demarcated mitochondria adjacent to and surrounding the nuclear circumference. Overlapping mitochondria were not analyzed. Average parameters were calculated for all measured mitochondria per cell.

Statistics and data analysis.

Statistical differences were assessed using two-sample t-tests for 2-group comparisons and One- or Two-Way ANOVA for 3 or more groups of data (Origin 9.1). All comparisons reflect sets of data substantially interleaved in time. Evaluations of mitochondrial and neuronal health and morphology were performed blindly using coded images.

In vitro experiments. Since our preliminary experiments did not find any differences in investigated parameters between slices of female and male animals, all further analysis included slices from animals of both sexes. We performed 1-2 slice experiments per animal per condition. All comparisons were made based on $n \geq 8$ slices. For evaluation of neuronal Ca^{2+} levels using Salsa6F/CaM mice, data were collected from single neurons (≥ 6 neurons per animal).

In vivo experiments. To analyze neuronal injury (using VAF staining), we evaluated 3–5 sections per animal with 50–100 cells examined per section. All data obtained from *in vivo* experiments were averaged per animal.

Behavioral data. Data were analyzed using IBM SPSS Statistics (Version 30.0; IBM Corp., Armonk, NY) and GraphPad Prism (Version 10.5; GraphPad Software Inc., La Jolla, CA). Normality was assessed with the Shapiro-Wilk test. Data for NDS and NDS sub-scores did not meet normality assumptions, thus the comparisons between the groups were done using the Mann–Whitney U test. For behavioral data analysis, significance was defined as $p < 0.05$.

Results.

Post-ischemic MCU blockade prevents delayed Zn^{2+} uptake into mitochondria

In order to evaluate the neuroprotective potential of the MCU inhibitor Ru265 [19] after brain ischemia, we first assessed its effect on post-ischemic mitochondrial Zn^{2+} accumulation. Experiments were conducted using the mouse hippocampal slice OGD model of transient ischemia developed in our laboratory [15, 16]. CA1

pyramidal neurons were loaded with the membrane-impermeable Zn^{2+} -sensitive indicator, FluoZin-3 [24, 25] and a Ca^{2+} -sensitive ratiometric indicator (Fura-FF or Fura-2) via patch-pipette prior to subjecting the slices to a short (8 min) OGD episode. This technique allows discrimination of Zn^{2+} vs Ca^{2+} signals in the cytoplasm of CA1 neurons [23] confirming the specificity of FluoZin-3 for the detection of cytoplasmic Zn^{2+} dynamics. OGD evoked a sharp rise in cytosolic Zn^{2+} levels, which largely recovered to baseline within 10-30 min following OGD termination and stayed low for the remainder of the 1-hour recording period (Fig. 2A). To assess whether this Zn^{2+} accumulates into mitochondria, slices were exposed to a high level of the protonophore, FCCP (2 μM), which rapidly depolarizes mitochondria, thereby releasing sequestered Zn^{2+} into the cytosol. Administration of FCCP at 60 min post-OGD resulted in a sharp rise in cytosolic Zn^{2+} levels, indicating substantial Zn^{2+} accumulation within mitochondria (Fig. 2A, B, D, E). In contrast, we did not observe any FCCP-evoked increase in intracellular Ca^{2+} levels, indicating an absence of significant mitochondrial Ca^{2+} accumulation at this delayed time point (Fig. 2A, B). These data correlate with our previous observations of long-lasting mitochondrial Zn^{2+} – but not Ca^{2+} – accumulation in CA1 neurons post-OGD [15]. We suggest that after ischemia termination, both Zn^{2+} and Ca^{2+} enter mitochondria via the MCU; however, Ca^{2+} is relatively rapidly extruded by the $\text{Na}^+/\text{Ca}^{2+}$ exchanger [31-33], whereas the Zn^{2+} remains sequestered within the mitochondria for a prolonged period.

In a subsequent set of experiments, the MCU antagonist Ru265 [19] was applied for 30 min beginning 3 min after OGD termination. Under these conditions, FCCP application at 60 min post-OGD no longer elicited a cytosolic Zn^{2+} rise, indicating that Ru265 effectively prevented Zn^{2+} entry into mitochondria (Fig. 2C, D, E).

MCU blockade following OGD attenuates delayed mitochondrial dysfunction

Previous studies have identified Zn^{2+} uptake through the MCU as a potent driver of mitochondrial damage, including persistent $\Delta\Psi_m$ depolarization, ROS generation, swelling, and fragmentation [15, 16, 34, 35]. Thus, to assess the ability of MCU inhibition to protect mitochondria from ischemia/reperfusion induced persistent dysfunction, we evaluated $\Delta\Psi_m$ at 4 h post-OGD. To measure the level of $\Delta\Psi_m$ depolarization, as an index of mitochondrial dysfunction, we employed the fluorescent indicator Rhodamine123 (Rhod123). This positively charged dye accumulates within negatively charged mitochondria, where its fluorescence is quenched, in proportion to $\Delta\Psi_m$. Application of the protonophore FCCP at high concentration (2 μM) dissipates the mitochondrial proton gradient, inducing complete $\Delta\Psi_m$ collapse, resulting in Rhod123 release and a corresponding rise in fluorescence (ΔF), thereby providing a measure of $\Delta\Psi_m$ prior to FCCP exposure. Using this approach, we evaluated $\Delta\Psi_m$ in the CA1 pyramidal cell layer. At 4 hours post-OGD, $\Delta\Psi_m$ was substantially reduced (by approximately 50%), indicative of severe mitochondrial dysfunction. Post-OGD application of Ru265 (5 μM) for 30 min (starting 3 min after OGD termination) markedly attenuated this delayed $\Delta\Psi_m$ loss, indicating improved mitochondrial functional recovery (Fig. 3A, B).

Delayed MCU blockade attenuates post-ischemic disruption of neuronal Ca^{2+} homeostasis

Disruption of neuronal Ca^{2+} homeostasis is a critical early event in the cascade leading to neurodegeneration after transient ischemia [36-38]. Thus, to further investigate neuroprotective effect of post-ischemic Ru265 treatment, we examined the ability of neurons to maintain physiological intracellular Ca^{2+} concentrations ($[\text{Ca}^{2+}]_i$).

Changes in $[\text{Ca}^{2+}]_i$ were monitored in freshly prepared slices from Salsa6f/CaM transgenic mice, which express the genetically encoded ratiometric Ca^{2+} indicator Salsa6f (GCaMP6f/tdTomato) [39] in forebrain pyramidal neurons, including those in the hippocampal CA1 region. Salsa6f is a fusion protein comprising the Ca^{2+} -sensitive green fluorescent indicator GCaMP6f (Kd ~375 nM), which exhibits increased green fluorescence upon cytosolic Ca^{2+} elevation, and the Ca^{2+} -insensitive red fluorescent protein tdTomato, which provides a stable reference signal enabling ratiometric Ca^{2+} imaging. Notably, at resting $[\text{Ca}^{2+}]_i$ levels, GCaMP6f fluorescence is barely distinguishable from background; therefore, tdTomato was used for cell identification and tracking

When slices from Salsa6f/CaM were exposed to elevated KCl (90 mM, 3 min), which depolarizes the plasma membrane triggering an activation of voltage gated Ca^{2+} channels (VGCC), a sharp increase in the GCaMP6f/tdTomato fluorescence ratio was observed, consistent with a sharp rise in $[\text{Ca}^{2+}]_i$ that recovered over the subsequent 5-8 minutes (Fig. 4A). Prolonged OGD induces a pronounced terminal $[\text{Ca}^{2+}]_i$ elevation (Fig. 4B), similar to our previous observations using synthetic Ca^{2+} indicators [15, 23]. Next, we applied transient (8 min) OGD and evaluated delayed changes in $[\text{Ca}^{2+}]_i$. We observed near-control $[\text{Ca}^{2+}]_i$ levels 1-hour after OGD termination (GCaMP6f/tdTomato ratio: 0.21 ± 0.01 vs. 0.16 ± 0.01 in control; $p = 0.1$). However, at later time points, $[\text{Ca}^{2+}]_i$ progressively increased, reaching a ratio of 0.41 ± 0.01 at 3 hours post-OGD ($p < 0.01$ vs. baseline), indicative of an impaired ability of neurons to maintain Ca^{2+} homeostasis. In contrast, post-OGD application of Ru265 (5 μM) markedly attenuated this delayed $[\text{Ca}^{2+}]_i$ elevation (0.25 ± 0.02 ; $p < 0.01$ vs. OGD alone) (Fig. 4C).

Delayed MCU blockade preserves mitochondrial morphology after in vivo CA

Subsequent experiments investigated the effects of Ru265 administration following global ischemia *in vivo*. Rats were randomly assigned to one of three experimental conditions: sham treatment, asphyxial CA (8 min) followed by CPR, or CA/CPR with intravenous administration of Ru265 (1 mg/kg) at the time of CPR. The dose employed in the present study is substantially below the 10 mg/kg previously reported to induce seizures in mice [40]. Additionally, no adverse effects were observed in any of the rats treated with 1 mg/kg of Ru265 ($n=11$).

After ~4 and ~20 h post-CPR, animals underwent behavioral testing. And ~24 h after CA, rats were perfused and brains isolated and processed for histological analysis (see Materials and Methods).

To assess mitochondrial morphology, 20 μm hippocampal slices were stained with antibody against the mitochondrial outer membrane protein, TOM-20. Confocal images (1000x) were obtained from the CA1 pyramidal cell layer (Fig. 5A). Mitochondrial morphological parameters were quantified using ImageJ software, as previously described [15, 16]. CA/CPR induced marked mitochondrial fragmentation and swelling, reflected by significant decreases in mitochondrial length, increases in width, and a substantially reduced length-to-width (L/W) ratio relative to sham-treated animals (Fig. 5B-E). Importantly, these morphological alterations were markedly attenuated by Ru265 treatment.

MCU inhibition after CA is neuroprotective

Our previous work demonstrated that brief episodes of global brain ischemia cause considerable injury of CA1 neurons 4 hours after CPR [41]. Here, we evaluated the extent of neuronal injury across multiple brain regions at 24 h post-ischemia and assessed the neuroprotective effect of post-ischemic MCU inhibition with Ru265. Neuronal injury was assessed using a modified vanadium acid fuchsin (VAF) staining procedure, which selectively labels acidophilic (injured) neurons. Stained sections were examined under light microscopy, and each identifiable neuron was classified as either healthy or injured. In the hippocampus, CA1 pyramidal neurons exhibited severe injury, whereas CA3 pyramidal neurons were considerably less affected, consistent with previously reported patterns of selective vulnerability [5-7]. Importantly, neuronal injury in both CA1 and CA3 was markedly reduced in Ru265-treated animals (Fig. 6).

In addition to the hippocampus, certain cortical neurons are known to be preferentially vulnerable to ischemic injury [8]. Consistent with this, pyramidal neurons in cortical layers III and V were substantially injured following CA/CPR. Notably, as observed in the hippocampus, Ru265 treatment significantly attenuated cortical neuronal injury (Fig. 7).

Delayed MCU blockade improves neurological recovery after CA

To assess neurological function following CA, we employed the Neurologic Deficit Scale (NDS). This rodent behavioral assessment is modeled after the Glasgow Coma Scale used in humans, with scores ranging from 0

(dead) to 80 (neurologically intact) [42, 43]. The NDS provides a cumulative score across multiple domains, including arousal, brainstem function, motor and sensory function, and behavioral responses. CA alone produced a substantial reduction in NDS scores relative to sham-treated controls. However, intravenous Ru265 administration after CA markedly improved NDS scores to near-control values at 20 h post-CPR (Fig. 8). Note that 4 h after CPR the NDS score was similar in Ru265- and placebo-treated ($p=0.37$), reflecting the slow recovery after CA-induced coma.

Discussion

Our previous and current studies have detected progressive, long-lasting Zn^{2+} accumulation in mitochondria of selectively vulnerable CA1 neurons following both *ex vivo* and *in vivo* ischemia [15, 41]. This delayed Zn^{2+} accumulation occurs predominantly via the MCU and contributes to ROS overproduction, mitochondrial dysfunction, and subsequent neuronal injury at 4 h post-ischemia [15, 16, 44]. As the limitations of previously available MCU inhibitors constrained both mechanistic conclusions and translational potential, the present study employed Ru265, a recently developed, highly potent, and relatively selective MCU inhibitor. In the *ex vivo* acute brain slice model, Ru265-mediated MCU blockade substantially decreased mitochondrial dysfunction and restored neuronal Ca^{2+} homeostasis at 3-4 h after OGD. In the *in vivo* rat CA model, intravenous Ru265 administration following CA markedly preserved mitochondrial morphology and considerably reduced neuronal injury in vulnerable hippocampal and cortical regions at 24 h post-CA. Furthermore, MCU inhibition reversed CA-induced behavioral deficits at 20 h post-CPR.

Thus, these findings identify mitochondrial Zn^{2+} accumulation as a viable therapeutic target in the post-ischemic period. The neuroprotective efficacy of Ru265 demonstrated across both *ex vivo* and *in vivo* ischemic models supports its potential as a foundation for future drug development targeting the MCU pathway.

Understanding mechanisms occurring after restoration of blood flow is key to effective neuroprotection

After transient cerebral ischemia, many vulnerable neurons, particularly CA1 hippocampal pyramidal neurons, initially appear to recover, but subsequently undergo delayed cell death hours to days later. The numerous deleterious cascades driving this neuronal death unfold after reperfusion, at a time when cerebral blood flow has been restored and therapeutic intervention is clinically feasible. Therefore, elucidating the precise sequence of events leading to this delayed neuronal death is essential for developing effective neuroprotective strategies.

Over decades of research, multiple treatments have been proposed and tested, yet most have failed to demonstrate meaningful neuroprotection in humans. This likely reflects the inability of many studies to distinguish between - and selectively target - pathological events occurring after ischemia versus those unfolding during ischemia itself.

One extensively studied mechanism is excitotoxicity, characterized by excessive activation of glutamate receptors (particularly NMDA receptors) leading to rapid neuronal Ca^{2+} overload. Although NMDA receptor blockers demonstrated high efficacy in animal models when administered prior to experimental ischemia, these agents consistently failed to provide neuroprotection in human clinical trials [45, 46]. This translational failure is likely attributable to the critical timing of drug administration: during acute ischemic events, massive NMDA-mediated Ca^{2+} influx occurs within minutes, causing neuronal damage well before therapeutic intervention is clinically feasible. Consequently, post-ischemic administration of NMDA receptor blockers likely had minimal impact on the progression of ischemic damage, as the primary excitotoxic insult had already occurred before treatment could be initiated.

Recent advances in reperfusion strategies following stroke opened new possibilities for neuroprotective interventions [47]. Identifying and targeting pathological events that occur specifically after the restoration of blood flow is crucial for the successful development of clinically translatable neuroprotective therapies.

Mitochondrial Zn²⁺ accumulation through the MCU is a key contributor to ischemic injury

Multiple studies support a critical role of Zn²⁺ in ischemic neuronal injury. Considerably elevated Zn²⁺ levels have been detected in vulnerable neurons after ischemia, including hippocampal CA1 neurons [48, 49]. Consistent with this, Zn²⁺ chelation confers neuroprotection in both *in vitro* and *in vivo* ischemia models [23, 48, 50, 51].

Although the brain contains substantial amounts of Zn²⁺, under physiological conditions it is sequestered in distinct Zn²⁺ pools. In neurons there are two major pools: (1) Zn²⁺ sequestered in synaptic vesicles, which is released upon strong synaptic stimulation and can enter neurons through certain Ca²⁺ permeable channels [14, 35]; and (2) Zn²⁺ bound to cytoplasmic Zn²⁺-binding proteins (in neurons, primarily metallothionein-III; MT3), from which it can be liberated by oxidative stress and acidosis which occur during ischemia/reperfusion [52-54]. Zn²⁺ release from MT3s appears to contribute substantially to post-ischemic CA1 neuronal injury [15, 55, 56].

While the mechanisms through which Zn²⁺ causes neuronal damage remain incompletely defined, multiple studies have identified mitochondria as the primary target of Zn²⁺ toxicity. Zn²⁺ enters mitochondria via the MCU and contributes to mitochondrial dysfunction, swelling and ROS generation [55, 57-60]. Additionally, Zn²⁺ has been found to induce irreversible inhibition of several mitochondrial enzymes involved in energy production and antioxidant defense, likely contributing to increased mitochondrial ROS production and mPTP induction [61, 62]. The mPTP induction, in turn, triggers release of apoptotic mediators including cytochrome C and apoptosis inducing factor [58], contributing to the activation of apoptotic injury pathways [63-66]. The ROS overproduction, in addition to damaging all cellular molecules, with particular impairment to mitochondrial DNA [67], triggers inflammatory cascades with immune cell infiltration, further exacerbating injury [68-73].

Our recent studies using *ex vivo* and *in vivo* ischemia models found that after short ischemic episodes Zn²⁺ accumulates in mitochondria via the MCU, where it persists for prolonged periods contributing to ROS overproduction, delayed mitochondrial dysfunction, structural damage, swelling and fragmentation [15, 16, 41, 44]. While Ca²⁺ also enters mitochondria following ischemia it is likely cleared relatively quickly via the Na⁺/Ca²⁺ exchanger, since we did not observe significant amount of Ca²⁺ in CA1 mitochondria 1 h after ischemia [15]. Importantly, while mitochondrial Zn²⁺ accumulation appears to exert a substantially more potent deleterious effect than cytoplasmic Zn²⁺ [51], cytoplasmic Ca²⁺ accumulation has also been shown to contribute to injury, in part through activation of NADPH oxidase [44, 74, 75].

The present study demonstrates that MCU inhibition with Ru265 following ischemia prevents mitochondrial Zn²⁺ accumulation, attenuates delayed mitochondrial injury, restores Ca²⁺ homeostasis, reduces neuronal damage and considerably improves behavioral recovery after CA. Taken together, these findings establish mitochondrial Zn²⁺ accumulation via the MCU as a critical and therapeutically targetable event in the post-ischemic cascade.

The novel MCU inhibitor Ru265 as a platform for future drug development

Until recently, pharmacological MCU blockade was not clinically achievable, as all available inhibitors were either highly non-selective and therefore unsuitable for human use (Ruthenium Red) [17] or membrane-impermeable and highly susceptible to oxidative degradation (RU360). The recent development of Ru265 has yielded an MCU antagonist with substantially greater selectivity and sufficient potency for use at relatively low doses, as we found that 1 mg/kg produced no overt adverse effects in rats, making it a promising foundation for future translational drug development.

The present study is an early investigation demonstrating that Ru265 represents a promising starting point for drug development for post-ischemic neuroprotective treatment. Future studies should more rigorously evaluate this compound, including assessment of its effectiveness across both sexes, as well as neuroprotective benefits measured at delayed time points – weeks and months post-ischemia. Furthermore, examining its efficacy when administered not only immediately following ischemia but also hours after the ischemic event would be particularly relevant for stroke patients, where timely interventions is often limited. Additionally, the development of improved Ru265 derivatives with reduced side effect profiles would further advance the therapeutic potential.

Ischemic injury is a complex process involving multiple deleterious events that collectively drive delayed neurodegeneration. Mitochondrial Zn^{2+} accumulation and ROS generation [44] appear to represent early initiating events in this pathological cascade, followed by - and likely contributing to - activation of apoptosis-triggering caspases, microglial activation, neutrophil infiltration, and broader immune system involvement, all of which perpetuate and amplify delayed tissue damage [7, 68, 69]. Ultimately, combined therapeutic strategies targeting distinct components of the downstream injury cascade at specific time points may provide substantial protection against delayed ischemic injury. Early post-ischemic MCU blockade could prove to be a highly effective component of such multimodal treatment strategies.

Clinical Perspectives

- Rationale: Despite being a leading cause of morbidity and mortality in the aging population, brain ischemia lacks effective neuroprotective interventions, largely due to the challenges of pre-reperfusion drug delivery and a limited understanding of the post-reperfusion mechanisms driving delayed neurodegeneration.
- Key findings: Following transient ischemia, zinc ions selectively accumulate within the mitochondria of vulnerable hippocampal CA1 pyramidal neurons, where they persist for prolonged periods and contribute to progressive mitochondrial dysfunction and delayed cellular injury. Notably, post-ischemic administration of Ru265 - a selective blocker of the mitochondrial calcium uniporter (MCU), which serves as the primary route for mitochondrial Zn^{2+} uptake - markedly attenuates delayed neuronal injury and improves neurological outcomes following brain ischemia.
- Clinical significance: These findings offer compelling evidence that administration of MCU antagonists, initiated after the restoration of cerebral blood flow – a clinically relevant therapeutic window - represents a novel and actionable therapeutic strategy for mitigating post-ischemic brain injury.

Data availability

Data that support our findings are available from the corresponding author upon reasonable request.

Funding

This work was supported by NIH grant NS121227 (JHW) and Roneet Carmell Memorial Endowment Fund (YA).

Abbreviations

AUC	area under the curve
ACSF	artificial cerebrospinal fluid
CA	cardiac arrest
CPR	cardiopulmonary resuscitation
FCCP	carbonyl cyanide-4-(trifluoromethoxy)phenylhydrazone
MCU	mitochondrial Ca^{2+} uniporter
OGD	oxygen-glucose deprivation

ROS	reactive oxygen species
Rhod123	Rhodamine 123
RR	ruthenium Red
Salsa6f/CaM mice	GCaMP6f/ CaMKIIalpha transgenic mice
TPEN	N,N,N',N'-tetrakis(2-pyridylmethyl)ethylenediamine

Figure legends

Figure 1. Experimental procedure. **A** – *Oxygen glucose deprivation (OGD) in brain slices.* Schematic show steps of experiment modeling ischemia/reperfusion in acute brain slices. Experiment was terminated at 4 h. Intramitochondrial Zn^{2+} levels were assessed at 1h, and cytoplasmic Ca^{2+} levels were evaluated at 1 h and 3 h after OGD withdrawal. **B** - *In vivo Cardiac arrest (CA) experimental procedure.* Schematic represents steps of an experiment modeling CA evoked transient global ischemia in rats. Ru265 was administered at the time of CPR along with standard intravenous epinephrine. All animals were intubated, attached to ventilator and cannulated before CA or sham procedure. Chest compressions were performed on sham animals to mimic CPR. Behavioral testing was performed at 4 and 20 h after CPR. At 24 h experiment was terminated, rats euthanized and brains processed for histological assessments.

Figure 2. MCU inhibition with Ru265 after OGD withdrawal prevents long-lasting Zn^{2+} accumulation into mitochondria. **A - C** – *Assessment of post-ischemic mitochondrial Zn^{2+} accumulation in individual CA1 neurons.* Neurons were pipette-loaded with FluoZin-3 and either Fura-FF or Fura-2 and subjected to 8 min OGD. At ~3 min after OGD withdrawal, Ru265 (5 μ M) or placebo treatment was applied for 30 min, and FCCP (2 μ M) administered at 1h post-OGD as indicated. **A** – Images show changes in FluoZin-3 fluorescence and in Fura-FF ratio in representative CA1 neuron subjected to OGD followed by FCCP after 1 h. The time marks on images indicate the time elapsed after OGD onset at which each image was acquired. Note that the second neuron visible in the field went out of focus during recording and was therefore excluded from analysis. **B** – *Upper traces* show FluoZin-3 fluorescence ($\Delta F/F_0 \times 100\%$, **blue**) and Fura-FF ratio (**black**) changes in the CA1 neuron marked with an arrow in (A). The region of interest used for recordings is denoted on the baseline (BL) FluoZin-3 image (A, upper left). The red vertical arrow illustrates the measurement of FCCP-induced FluoZin-3 amplitude increase; additionally, the red vertical and horizontal arrows together demarcate the area under the curve (AUC) presented in E. *Lower trace* represents background-subtracted FluoZin-3 fluorescent changes in the control CA1 neuron. **C** – Traces show FluoZin-3 fluorescence ($\Delta F/F_0 \times 100\%$, **blue**) and Fura-2 ratio (**black**) changes in the CA1 neuron subjected to OGD and treated with Ru265 after reperfusion; FCCP was administered 1 hour following OGD withdrawal. **D** - Bar graph shows mean FCCP-induced FluoZin-3 fluorescence increases ($\pm$ SE); individual symbols represent measurements from single neurons. **E** - Bar graph show mean AUC ($\pm$ SE) of FCCP-induced FluoZin-3 fluorescence changes calculated from the onset of FCCP administration over the subsequent 15 min; symbols show individual data points for each neuron. Neurons corresponding to representative traces shown in B and C are indicated by an X. * - $p < 0.01$, t-Test, control: n=8 neurons from 8 slices, 8 mice; Ru265: n=7 neurons from 6 slices, 6 mice.

Figure 3. Treatment with Ru265 post-OGD diminishes delayed loss of $\Delta\Psi_m$. **A** - *Assessment of $\Delta\Psi_m$.* Brain slices were subjected to short OGD (8 min)/reperfusion, followed by placebo treatment or Ru265 (5 μ M for 30 min). Next, slices were loaded with Rhod123, and 4 h after OGD termination, FCCP (2 μ M) was administered. *Upper* - pseudocolor Rhod123 fluorescent images of CA1 area before (baseline, BL) and after FCCP application obtained from representative control slice, and slices subjected to OGD followed by placebo or Ru265 treatment. *Lower* - Traces represent Rhod123 $\Delta F/F_0$ (%) changes in slices shown on the upper images (**black** – control, **red**

- OGD, **blue** – OGD+Ru265). Arrows indicate the time points at which fluorescent images were taken. **B** - Bar graph show average maximum FCCP-evoked Rhod123 $\Delta F/F_0$ *100% ($\pm$ SE) increases, symbols represent individual data points for each slice. The symbols depicting Rhod123 increases in slices presented on A are marked with an X. control: n=15 slices from 11 mice, OGD: n=7 slices from 7 mice, OGD+Ru265: n=8 slices from 7 mice. * - $p < 0.01$, ANOVA.

Figure 4. Treatment with Ru265 considerably improves Ca^{2+} homeostasis in CA1 neurons. Acute slices were prepared from the brains of Salsa6f/CaM mice. **A**, Upper - *GCamp6f* and *tdTomato* images of CA1 region at baseline and after administration of KCl (90 mM for 3 min) in a representative slice. Lower - trace illustrates average ($\pm$ SE) *Gcamp6f* / *tdTomato* ratio changes in n=5 neurons. Arrows show time points at which upper images were taken. **B**, upper - *GCamp6f* and *tdTomato* images of CA1 region before and after subjecting slice to prolonged OGD. Trace shows average changes ($\pm$ SE) in *Gcamp6f* / *tdTomato* ratio in n=4 neurons. **C** - Brain slices from Salsa6f mice were subjected to OGD (8 min)/reperfusion, and Ru265 (or placebo treatment) was applied 3 min after OGD withdrawal for 30 min. Upper - *GCamp6f* and *tdTomato* fluorescent images of CA1 region in representative control untreated slice, 1 and 3 h after OGD alone, and 3 h after OGD+treatment with Ru265, as indicated. Lower – Bar graph represents average ($\pm$ SE) *Gcamp6f* / *tdTomato* ratios, and symbols show *Gcamp6f* / *tdTomato* ratios in individual neurons (n=46 neurons from 6 mice in control, n=59 cells from 6 mice at 1 hour post-OGD, n=95 neurons from 8 mice at 3 hour post-OGD and n=67 neurons from 8 mice at 3 h after OGD+Ru265), * - $p < 0.01$, ANOVA

Figure 5. Mitochondrial swelling in CA1 pyramidal neurons 24 h after cardiac arrest is mitigated by MCU blockade with Ru265. Rats were subjected to sham treatment, transient global ischemia induced by CA (8 min), or CA followed by intravenous treatment with Ru265 (1 mg/kg). After 24 hours of recovery, rats were euthanized, and brain tissue was collected for histological examination. 20 μ m slices were cut with a vibratome and stained with TOM-20 antibody. **A** - *Appearance of mitochondria in CA1 neurons*. Representative confocal images show TOM-20 labeled mitochondria. Bottom - Enlarged images of regions indicated on the top panel show representative mitochondria at high magnifications. Inset lines illustrate how measurements of lengths and widths were performed. Note that ischemia caused fragmentation and visible swelling of mitochondria, with a decrease in length and increase in width; and that this change was attenuated by delayed treatment with Ru265. **B-D** - *Quantitative analysis*. Bar graphs show mean ($\pm$ SD) mitochondrial length (B), width (C) and length/width ratio (D). Mitochondrial length and width were averaged for each neuron (>200 cells per animal analyzed). Symbols represent mitochondrial measures averaged per individual neuron. **E** – *Length/width distribution*. Bar graphs show L/W ratio distribution for single mitochondria (n>500 for each group) with bin size 5. Data obtained from 3 sham rats, 4 ischemia rats and 3 ischemia+Ru265 rats. * - $p < 0.01$, ANOVA

Figure 6. MCU inhibition considerably decreases neuronal injury in CA1 pyramidal neurons after cardiac arrest. Rats were subjected to sham treatment, transient global ischemia induced by CA (8 min), or CA followed by intravenous injection of Ru265 (1 mg/kg). After 24-hour recovery, rats were euthanized, and brain tissue collected for histological examination. To assess injury, slices were subjected to a modified acid fuchsin labeling procedure (VAF). **A** - *Representative images of VAF stained CA1 and CA3 regions in brain slices*. Note that the majority of neurons appear healthy (yellow arrows) in slices from sham treated animals, but after ischemia most neurons are injured (red arrows) and have VAF labelling in cytoplasm or are swollen with no clear cellular membrane. Note that the injury is considerably less in the CA3 region and also that the injury was largely reversed in both regions, in animals treated with Ru265. **B** - *Quantitative assessment*. All neurons were blindly analyzed and rated as healthy or injured, and the percentages of neurons determined to be injured were calculated for each animal. Note that the number of injured cells was far greater in animals subjected to ischemia than after sham treatment, and that Ru265 administration greatly diminished neuronal injury. Bars represent mean ($\pm$ SE) from 7 rats sham, 7 animals after CA, and 8 rats after CA+Ru265. Symbols show number of injured cells calculated per individual animal. * - $p < 0.01$, ANOVA

Figure 7. MCU inhibition with Ru265 after CA/CPR considerably decreases neuronal injury in cortical layer III and V neurons. **A** - Representative images of cortical layer III and V. Note that the majority of neurons appear healthy (yellow arrows) in slices from sham treated animals, but after ischemia most neurons are injured (red arrows) and have VAF labelling in cytoplasm. Note that injury was considerably diminished in cortex from Ru265 treated animals. **B** - Quantitative assessment. All neurons were blindly analyzed and rated as intact or injured, and the percentages of neurons determined to be injured within each animal were calculated. Bars represent mean ($\pm$ SE) from 7 animals sham, 7 animals after CA, and 8 animals after CA+Ru265. Symbols show numbers of injured cells calculated per individual animal. * - $p < 0.01$, ANOVA

Figure 8. MCU inhibition reverses neurological deficit caused by transient brain ischemia. The Neurologic Deficit Scale (NDS) - a cumulative scores for multiple tests evaluating arousal, brainstem function, motor and sensory assessment, - was used to evaluate the effect of CA and the treatment with Ru265 after CA on animal behavior. Bar graphs represent average NDS in animals 4 and 20 h after CA, as indicated, and symbols show individual data points for each group; $n=7$ sham treated animals, $n=10$ animals subjected to CA and $n=11$ rats subjected to CA+Ru265, * - $p < 0.05$, Mann-Whitney U test.

References

- Hosseini M, Wilson RH, Crouzet C, Amirhekmat A, Wei KS and Akbari Y, (2020) *Resuscitating the Globally Ischemic Brain: TTM and Beyond*. Neurotherapeutics. 17: 539-562. DOI: 10.1007/s13311-020-00856-z <http://www.ncbi.nlm.nih.gov/pubmed/32367476>
- Holgersson J, Meyer MAS, Dankiewicz J, Lilja G, Ullen S, Hassager C, et al., (2022) *Hypothermic versus Normothermic Temperature Control after Cardiac Arrest*. NEJM Evid. 1: EVIDoA2200137. DOI: 10.1056/EVIDoA2200137 <http://www.ncbi.nlm.nih.gov/pubmed/38319850>
- Wechsler LR, Adeoye O, Alemseged F, Bahr-Hosseini M, Deljkich E, Favilla C, et al., (2023) *Most Promising Approaches to Improve Stroke Outcomes: The Stroke Treatment Academic Industry Roundtable XII Workshop*. Stroke. 54: 3202-3213. DOI: 10.1161/STROKEAHA.123.044279 <http://www.ncbi.nlm.nih.gov/pubmed/37886850>
- Petito CK, Feldmann E, Pulsinelli WA and Plum F, (1987) *Delayed hippocampal damage in humans following cardiorespiratory arrest*. Neurology. 37: 1281-6. <http://www.ncbi.nlm.nih.gov/pubmed/3614648>
- Kirino T, (1982) *Delayed neuronal death in the gerbil hippocampus following ischemia*. Brain Res. 239: 57-69. <http://www.ncbi.nlm.nih.gov/pubmed/7093691>
- Ordy JM, Wengenack TM, Bialobok P, Coleman PD, Rodier P, Baggs RB, et al., (1993) *Selective vulnerability and early progression of hippocampal CA1 pyramidal cell degeneration and GFAP-positive astrocyte reactivity in the rat four-vessel occlusion model of transient global ischemia*. Exp Neurol. 119: 128-39. DOI: 10.1006/exnr.1993.1014 <http://www.ncbi.nlm.nih.gov/pubmed/8432346>
- Sugawara T, Fujimura M, Morita-Fujimura Y, Kawase M and Chan PH, (1999) *Mitochondrial release of cytochrome c corresponds to the selective vulnerability of hippocampal CA1 neurons in rats after transient global cerebral ischemia*. J Neurosci. 19: RC39. <http://www.ncbi.nlm.nih.gov/pubmed/10559429>
- Pulsinelli WA, Brierley JB and Plum F, (1982) *Temporal profile of neuronal damage in a model of transient forebrain ischemia*. Ann Neurol. 11: 491-8. DOI: 10.1002/ana.410110509 <http://www.ncbi.nlm.nih.gov/pubmed/7103425>
- Baron JC, Yamauchi H, Fujioka M and Endres M, (2014) *Selective neuronal loss in ischemic stroke and cerebrovascular disease*. J Cereb Blood Flow Metab. 34: 2-18. DOI: 10.1038/jcbfm.2013.188 <http://www.ncbi.nlm.nih.gov/pubmed/24192635>
- Chan PH, (2001) *Reactive oxygen radicals in signaling and damage in the ischemic brain*. J Cereb Blood Flow Metab. 21: 2-14. DOI: 10.1097/00004647-200101000-00002 <http://www.ncbi.nlm.nih.gov/pubmed/11149664>

11. Fiskum G, Murphy AN and Beal MF, (1999) *Mitochondria in neurodegeneration: acute ischemia and chronic neurodegenerative diseases*. J Cereb Blood Flow Metab. 19: 351-69. DOI: 10.1097/00004647-199904000-00001 <http://www.ncbi.nlm.nih.gov/pubmed/10197505>
12. Piantadosi CA and Zhang J, (1996) *Mitochondrial generation of reactive oxygen species after brain ischemia in the rat*. Stroke. 27: 327-31; discussion 332. DOI: 10.1161/01.str.27.2.327 <http://www.ncbi.nlm.nih.gov/pubmed/8571432>
13. Ji SG, Medvedeva YV, Wang HL, Yin HZ and Weiss JH, (2019) *Mitochondrial Zn(2+) Accumulation: A Potential Trigger of Hippocampal Ischemic Injury*. Neuroscientist. 25: 126-138. DOI: 10.1177/1073858418772548 <http://www.ncbi.nlm.nih.gov/pubmed/29742958>
14. Shuttleworth CW and Weiss JH, (2011) *Zinc: new clues to diverse roles in brain ischemia*. Trends Pharmacol Sci. 32: 480-6. DOI: 10.1016/j.tips.2011.04.001 <http://www.ncbi.nlm.nih.gov/pubmed/21621864>
15. Medvedeva YV, Ji SG, Yin HZ and Weiss JH, (2017) *Differential Vulnerability of CA1 versus CA3 Pyramidal Neurons After Ischemia: Possible Relationship to Sources of Zn2+ Accumulation and Its Entry into and Prolonged Effects on Mitochondria*. J Neurosci. 37: 726-737. DOI: 10.1523/JNEUROSCI.3270-16.2016 <http://www.ncbi.nlm.nih.gov/pubmed/28100752>
16. Medvedeva YV, Yin HZ, Bazrafkan A, Yeromin A, Ji SG, Weiss-Hung EJ, et al., (2022) *Blocking Mitochondrial Zn(2+) Accumulation after Ischemia Reduces Mitochondrial Dysfunction and Neuronal Injury*. J Neurosci. 42: 5281-5292. DOI: 10.1523/JNEUROSCI.0874-21.2022 <http://www.ncbi.nlm.nih.gov/pubmed/35623885>
17. Tapia R and Velasco I, (1997) *Ruthenium red as a tool to study calcium channels, neuronal death and the function of neural pathways*. Neurochem Int. 30: 137-47. <http://www.ncbi.nlm.nih.gov/pubmed/9017661>
18. Woods JJ, Lovett J, Lai B, Harris HH and Wilson JJ, (2020) *Redox Stability Controls the Cellular Uptake and Activity of Ruthenium-Based Inhibitors of the Mitochondrial Calcium Uniporter (MCU)*. Angew Chem Int Ed Engl. 59: 6482-6491. DOI: 10.1002/anie.202000247 <http://www.ncbi.nlm.nih.gov/pubmed/32039530>
19. Woods JJ, Nemani N, Shanmughapriya S, Kumar A, Zhang M, Nathan SR, et al., (2019) *A Selective and Cell-Permeable Mitochondrial Calcium Uniporter (MCU) Inhibitor Preserves Mitochondrial Bioenergetics after Hypoxia/Reoxygenation Injury*. ACS Cent Sci. 5: 153-166. DOI: 10.1021/acscentsci.8b00773 <http://www.ncbi.nlm.nih.gov/pubmed/30693334>
20. Nathan SR, Pino NW, Arduino DM, Perocchi F, MacMillan SN and Wilson JJ, (2017) *Synthetic Methods for the Preparation of a Functional Analogue of Ru360, a Potent Inhibitor of Mitochondrial Calcium Uptake*. Inorg Chem. 56: 3123-3126. DOI: 10.1021/acs.inorgchem.6b03108 <http://www.ncbi.nlm.nih.gov/pubmed/28244741>
21. Azadian M, Tian G, Bazrafkan A, Maki N, Rafi M, Chetty N, et al., (2020) *Overnight Caloric Restriction Prior to Cardiac Arrest and Resuscitation Leads to Improved Survival and Neurological Outcome in a Rodent Model*. Front Neurosci. 14: 609670. DOI: 10.3389/fnins.2020.609670 <http://www.ncbi.nlm.nih.gov/pubmed/33510613>
22. Han S, Contreras MI, Bazrafkan A, Rafi M, Dara SM, Orujyan A, et al., (2022) *Cortical Anoxic Spreading Depolarization During Cardiac Arrest is Associated with Remote Effects on Peripheral Blood Pressure and Postresuscitation Neurological Outcome*. Neurocrit Care. 37: 139-154. DOI: 10.1007/s12028-022-01530-2 <http://www.ncbi.nlm.nih.gov/pubmed/35729464>
23. Medvedeva YV, Lin B, Shuttleworth CW and Weiss JH, (2009) *Intracellular Zn2+ accumulation contributes to synaptic failure, mitochondrial depolarization, and cell death in an acute slice oxygen-glucose deprivation model of ischemia*. J Neurosci. 29: 1105-14. http://www.ncbi.nlm.nih.gov/entrez/query.fcgi?cmd=Retrieve&db=PubMed&dopt=Citation&list_uids=19176819
24. Gee KR, Zhou ZL, Ton-That D, Sensi SL and Weiss JH, (2002) *Measuring zinc in living cells. A new generation of sensitive and selective fluorescent probes*. Cell Calcium. 31: 245-51. http://www.ncbi.nlm.nih.gov/entrez/query.fcgi?cmd=Retrieve&db=PubMed&dopt=Citation&list_uids=12098227


25. Sensi SL, Ton-That D, Sullivan PG, Jonas EA, Gee KR, Kaczmarek LK, et al., (2003) *Modulation of mitochondrial function by endogenous Zn²⁺ pools*. Proc Natl Acad Sci U S A. 100: 6157-62. http://www.ncbi.nlm.nih.gov/entrez/query.fcgi?cmd=Retrieve&db=PubMed&dopt=Citation&list_uids=12724524
26. Duchen MR, Surin A and Jacobson J, (2003) *Imaging mitochondrial function in intact cells*. Methods Enzymol. 361: 353-89. http://www.ncbi.nlm.nih.gov/entrez/query.fcgi?cmd=Retrieve&db=PubMed&dopt=Citation&list_uids=12624920
27. Crouzet C, Wilson RH, Bazrafkan A, Farahabadi MH, Lee D, Alcocer J, et al., (2016) *Cerebral blood flow is decoupled from blood pressure and linked to EEG bursting after resuscitation from cardiac arrest*. Biomed Opt Express. 7: 4660-4673. DOI: 10.1364/BOE.7.004660 <http://www.ncbi.nlm.nih.gov/pubmed/27896005>
28. Kang YJ, Tian G, Bazrafkan A, Farahabadi MH, Azadian M, Abbasi H, et al., (2017) *Recovery from Coma Post-Cardiac Arrest Is Dependent on the Orexin Pathway*. J Neurotrauma. 34: 2823-2832. DOI: 10.1089/neu.2016.4852 <http://www.ncbi.nlm.nih.gov/pubmed/28447885>
29. Crouzet C, Wilson RH, Lee D, Bazrafkan A, Tromberg BJ, Akbari Y, et al., (2020) *Dissociation of Cerebral Blood Flow and Femoral Artery Blood Pressure Pulsatility After Cardiac Arrest and Resuscitation in a Rodent Model: Implications for Neurological Recovery*. J Am Heart Assoc. 9: e012691. DOI: 10.1161/JAHA.119.012691 <http://www.ncbi.nlm.nih.gov/pubmed/31902319>
30. Victorov IV, Prass K and Dirnagl U, (2000) *Improved selective, simple, and contrast staining of acidophilic neurons with vanadium acid fuchsin*. Brain Res Brain Res Protoc. 5: 135-9. <http://www.ncbi.nlm.nih.gov/pubmed/10775832>
31. Baron KT and Thayer SA, (1997) *CGP37157 modulates mitochondrial Ca²⁺ homeostasis in cultured rat dorsal root ganglion neurons*. Eur J Pharmacol. 340: 295-300. http://www.ncbi.nlm.nih.gov/entrez/query.fcgi?cmd=Retrieve&db=PubMed&dopt=Citation&list_uids=9537826
32. Colegrove SL, Albrecht MA and Friel DD, (2000) *Dissection of mitochondrial Ca²⁺ uptake and release fluxes in situ after depolarization-evoked [Ca²⁺]_i elevations in sympathetic neurons*. J Gen Physiol. 115: 351-70. DOI: 10.1085/jgp.115.3.351 <http://www.ncbi.nlm.nih.gov/pubmed/10694263>
33. Garcia-Chacon LE, Nguyen KT, David G and Barrett EF, (2006) *Extrusion of Ca²⁺ from mouse motor terminal mitochondria via a Na⁺-Ca²⁺ exchanger increases post-tetanic evoked release*. J Physiol. 574: 663-75. DOI: 10.1113/jphysiol.2006.110841 <http://www.ncbi.nlm.nih.gov/pubmed/16613870>
34. Ji SG, Medvedeva YV, Wang HL, Yin HZ and Weiss JH, (2018) *Mitochondrial Zn(2+) Accumulation: A Potential Trigger of Hippocampal Ischemic Injury*. Neuroscientist. 1073858418772548. DOI: 10.1177/1073858418772548 <http://www.ncbi.nlm.nih.gov/pubmed/29742958>
35. Weiss JH, Sensi SL and Koh JY, (2000) *Zn(2+): a novel ionic mediator of neural injury in brain disease*. Trends Pharmacol Sci. 21: 395-401. http://www.ncbi.nlm.nih.gov/entrez/query.fcgi?cmd=Retrieve&db=PubMed&dopt=Citation&list_uids=11050320
36. Bodalia A, Li H and Jackson MF, (2013) *Loss of endoplasmic reticulum Ca²⁺ homeostasis: contribution to neuronal cell death during cerebral ischemia*. Acta Pharmacol Sin. 34: 49-59. DOI: 10.1038/aps.2012.139 <http://www.ncbi.nlm.nih.gov/pubmed/23103622>
37. Rahi V and Kaundal RK, (2024) *Exploring the intricacies of calcium dysregulation in ischemic stroke: Insights into neuronal cell death and therapeutic strategies*. Life Sci. 347: 122651. DOI: 10.1016/j.lfs.2024.122651 <http://www.ncbi.nlm.nih.gov/pubmed/38642844>
38. Tymianski M and Tator CH, (1996) *Normal and abnormal calcium homeostasis in neurons: a basis for the pathophysiology of traumatic and ischemic central nervous system injury*. Neurosurgery. 38: 1176-95. DOI: 10.1097/00006123-199606000-00028 <http://www.ncbi.nlm.nih.gov/pubmed/8727150>
39. Dong TX, Othy S, Jairaman A, Skupsky J, Zavala A, Parker I, et al., (2017) *T-cell calcium dynamics visualized in a ratiometric tdTomato-GCaMP6f transgenic reporter mouse*. Elife. 6. DOI: 10.7554/eLife.32417 <http://www.ncbi.nlm.nih.gov/pubmed/29239725>

40. Novorolsky RJ, Nichols M, Kim JS, Pavlov EV, J JW, Wilson JJ, et al., (2020) *The cell-permeable mitochondrial calcium uniporter inhibitor Ru265 preserves cortical neuron respiration after lethal oxygen glucose deprivation and reduces hypoxic/ischemic brain injury*. J Cereb Blood Flow Metab. 40: 1172-1181. DOI: 10.1177/0271678X20908523 <http://www.ncbi.nlm.nih.gov/pubmed/32126877>
41. Yin HZ, Wang HL, Ji SG, Medvedeva YV, Tian G, Bazrafkan AK, et al., (2019) *Rapid Intramitochondrial Zn²⁺ Accumulation in CA1 Hippocampal Pyramidal Neurons After Transient Global Ischemia: A Possible Contributor to Mitochondrial Disruption and Cell Death*. J Neuropathol Exp Neurol. 78: 655-664. DOI: 10.1093/jnen/nlz042 <http://www.ncbi.nlm.nih.gov/pubmed/31150090>
42. Jia X, Koenig MA, Shin HC, Zhen G, Pardo CA, Hanley DF, et al., (2008) *Improving neurological outcomes post-cardiac arrest in a rat model: immediate hypothermia and quantitative EEG monitoring*. Resuscitation. 76: 431-42. DOI: 10.1016/j.resuscitation.2007.08.014 <http://www.ncbi.nlm.nih.gov/pubmed/17936492>
43. Jia X, Koenig MA, Shin HC, Zhen G, Yamashita S, Thakor NV, et al., (2006) *Quantitative EEG and neurological recovery with therapeutic hypothermia after asphyxial cardiac arrest in rats*. Brain Res. 1111: 166-75. DOI: 10.1016/j.brainres.2006.04.121 <http://www.ncbi.nlm.nih.gov/pubmed/16919609>
44. Medvedeva YV, Sharman E and Weiss JH, (2025) *Mechanisms of delayed ischemia/reperfusion evoked ROS generation in the hippocampal CA1 zone of adult mouse brain slices*. Res Sq. DOI: 10.21203/rs.3.rs-5640324/v1 <http://www.ncbi.nlm.nih.gov/pubmed/40313775>
45. Hoyte L, Barber PA, Buchan AM and Hill MD, (2004) *The rise and fall of NMDA antagonists for ischemic stroke*. Curr Mol Med. 4: 131-6. <http://www.ncbi.nlm.nih.gov/pubmed/15032709>
46. Ikonomidou C and Turski L, (2002) *Why did NMDA receptor antagonists fail clinical trials for stroke and traumatic brain injury?* Lancet Neurol. 1: 383-6. <http://www.ncbi.nlm.nih.gov/pubmed/12849400>
47. Goyal M, Menon BK, van Zwam WH, Dippel DW, Mitchell PJ, Demchuk AM, et al., (2016) *Endovascular thrombectomy after large-vessel ischaemic stroke: a meta-analysis of individual patient data from five randomised trials*. Lancet. 387: 1723-31. DOI: 10.1016/S0140-6736(16)00163-X <http://www.ncbi.nlm.nih.gov/pubmed/26898852>
48. Koh JY, Suh SW, Gwag BJ, He YY, Hsu CY and Choi DW, (1996) *The role of zinc in selective neuronal death after transient global cerebral ischemia*. Science. 272: 1013-6. http://www.ncbi.nlm.nih.gov/entrez/query.fcgi?cmd=Retrieve&db=PubMed&dopt=Citation&list_uids=8638123
49. Tonder N, Johansen FF, Frederickson CJ, Zimmer J and Diemer NH, (1990) *Possible role of zinc in the selective degeneration of dentate hilar neurons after cerebral ischemia in the adult rat*. Neurosci Lett. 109: 247-52. http://www.ncbi.nlm.nih.gov/entrez/query.fcgi?cmd=Retrieve&db=PubMed&dopt=Citation&list_uids=2330128
50. Calderone A, Jover T, Mashiko T, Noh KM, Tanaka H, Bennett MV, et al., (2004) *Late calcium EDTA rescues hippocampal CA1 neurons from global ischemia-induced death*. J Neurosci. 24: 9903-13. http://www.ncbi.nlm.nih.gov/entrez/query.fcgi?cmd=Retrieve&db=PubMed&dopt=Citation&list_uids=15525775
51. Medvedeva YV and Weiss JH, (2014) *Intramitochondrial Zn(2+) accumulation via the Ca(2+) uniporter contributes to acute ischemic neurodegeneration*. Neurobiol Dis. 68: 137-44. DOI: 10.1016/j.nbd.2014.04.011 <http://www.ncbi.nlm.nih.gov/pubmed/24787898>
52. Jiang LJ, Vasak M, Vallee BL and Maret W, (2000) *Zinc transfer potentials of the alpha - and beta-clusters of metallothionein are affected by domain interactions in the whole molecule*. Proc Natl Acad Sci U S A. 97: 2503-8. http://www.ncbi.nlm.nih.gov/entrez/query.fcgi?cmd=Retrieve&db=PubMed&dopt=Citation&list_uids=10716985
53. Maret W and Vallee BL, (1998) *Thiolate ligands in metallothionein confer redox activity on zinc clusters*. Proc Natl Acad Sci U S A. 95: 3478-82. http://www.ncbi.nlm.nih.gov/entrez/query.fcgi?cmd=Retrieve&db=PubMed&dopt=Citation&list_uids=9520391

54. Maret W, (2011) *Redox biochemistry of mammalian metallothioneins*. J Biol Inorg Chem. 16: 1079-86. DOI: 10.1007/s00775-011-0800-0 <http://www.ncbi.nlm.nih.gov/pubmed/21647775>
55. Ji SG and Weiss JH, (2018) *Zn(2+)-induced disruption of neuronal mitochondrial function: Synergism with Ca(2+), critical dependence upon cytosolic Zn(2+) buffering, and contributions to neuronal injury*. Exp Neurol. 302: 181-195. DOI: 10.1016/j.expneurol.2018.01.012 <http://www.ncbi.nlm.nih.gov/pubmed/29355498>
56. Lee JY, Kim JH, Palmiter RD and Koh JY, (2003) *Zinc released from metallothionein-iii may contribute to hippocampal CA1 and thalamic neuronal death following acute brain injury*. Exp Neurol. 184: 337-47. http://www.ncbi.nlm.nih.gov/entrez/query.fcgi?cmd=Retrieve&db=PubMed&dopt=Citation&list_uids=14637104
57. Ji SG, Medvedeva YV and Weiss JH, (2020) *Zn(2+) entry through the mitochondrial calcium uniporter is a critical contributor to mitochondrial dysfunction and neurodegeneration*. Exp Neurol. 325: 113161. DOI: 10.1016/j.expneurol.2019.113161 <http://www.ncbi.nlm.nih.gov/pubmed/31881218>
58. Jiang D, Sullivan PG, Sensi SL, Steward O and Weiss JH, (2001) *Zn(2+) induces permeability transition pore opening and release of pro-apoptotic peptides from neuronal mitochondria*. J Biol Chem. 276: 47524-9. http://www.ncbi.nlm.nih.gov/entrez/query.fcgi?cmd=Retrieve&db=PubMed&dopt=Citation&list_uids=11595748
59. Sensi SL, Yin HZ, Carriedo SG, Rao SS and Weiss JH, (1999) *Preferential Zn²⁺ influx through Ca²⁺-permeable AMPA/kainate channels triggers prolonged mitochondrial superoxide production*. Proc Natl Acad Sci U S A. 96: 2414-9. http://www.ncbi.nlm.nih.gov/entrez/query.fcgi?cmd=Retrieve&db=PubMed&dopt=Citation&list_uids=10051656
60. Sensi SL, Yin HZ and Weiss JH, (2000) *AMPA/kainate receptor-triggered Zn²⁺ entry into cortical neurons induces mitochondrial Zn²⁺ uptake and persistent mitochondrial dysfunction*. Eur J Neurosci. 12: 3813-8. http://www.ncbi.nlm.nih.gov/entrez/query.fcgi?cmd=Retrieve&db=PubMed&dopt=Citation&list_uids=11029652
61. Gazaryan IG, Krasinskaya IP, Kristal BS and Brown AM, (2007) *Zinc irreversibly damages major enzymes of energy production and antioxidant defense prior to mitochondrial permeability transition*. J Biol Chem. 282: 24373-80. http://www.ncbi.nlm.nih.gov/entrez/query.fcgi?cmd=Retrieve&db=PubMed&dopt=Citation&list_uids=17565998
62. Gazaryan IG, Krasnikov BF, Ashby GA, Thorneley RN, Kristal BS and Brown AM, (2002) *Zinc is a potent inhibitor of thiol oxidoreductase activity and stimulates reactive oxygen species production by lipoamide dehydrogenase*. J Biol Chem. 277: 10064-72. http://www.ncbi.nlm.nih.gov/entrez/query.fcgi?cmd=Retrieve&db=PubMed&dopt=Citation&list_uids=11744691
63. Bernardi P, (1996) *The permeability transition pore. Control points of a cyclosporin A-sensitive mitochondrial channel involved in cell death*. Biochim Biophys Acta. 1275: 5-9. http://www.ncbi.nlm.nih.gov/entrez/query.fcgi?cmd=Retrieve&db=PubMed&dopt=Citation&list_uids=8688451
64. Bernardi P, Petronilli V, Di Lisa F and Forte M, (2001) *A mitochondrial perspective on cell death*. Trends Biochem Sci. 26: 112-7. http://www.ncbi.nlm.nih.gov/entrez/query.fcgi?cmd=Retrieve&db=PubMed&dopt=Citation&list_uids=11166569
65. Crompton M, (1999) *The mitochondrial permeability transition pore and its role in cell death*. Biochem J. 341 (Pt 2): 233-49. <http://www.ncbi.nlm.nih.gov/pubmed/10393078>
66. Fatokun AA, Dawson VL and Dawson TM, (2014) *Parthanatos: mitochondrial-linked mechanisms and therapeutic opportunities*. Br J Pharmacol. 171: 2000-16. DOI: 10.1111/bph.12416 <http://www.ncbi.nlm.nih.gov/pubmed/24684389>

67. Juan CA, Perez de la Lastra JM, Plou FJ and Perez-Lebena E, (2021) *The Chemistry of Reactive Oxygen Species (ROS) Revisited: Outlining Their Role in Biological Macromolecules (DNA, Lipids and Proteins) and Induced Pathologies*. Int J Mol Sci. 22. DOI: 10.3390/ijms22094642 <http://www.ncbi.nlm.nih.gov/pubmed/33924958>
68. Herz J, Sabellek P, Lane TE, Gunzer M, Hermann DM and Doeppner TR, (2015) *Role of Neutrophils in Exacerbation of Brain Injury After Focal Cerebral Ischemia in Hyperlipidemic Mice*. Stroke. 46: 2916-25. DOI: 10.1161/STROKEAHA.115.010620 <http://www.ncbi.nlm.nih.gov/pubmed/26337969>
69. Iadecola C and Anrather J, (2011) *The immunology of stroke: from mechanisms to translation*. Nat Med. 17: 796-808. DOI: 10.1038/nm.2399 <http://www.ncbi.nlm.nih.gov/pubmed/21738161>
70. Li J, Baud O, Vartanian T, Volpe JJ and Rosenberg PA, (2005) *Peroxynitrite generated by inducible nitric oxide synthase and NADPH oxidase mediates microglial toxicity to oligodendrocytes*. Proc Natl Acad Sci U S A. 102: 9936-41. DOI: 10.1073/pnas.0502552102 <http://www.ncbi.nlm.nih.gov/pubmed/15998743>
71. Mander PK, Jekabsone A and Brown GC, (2006) *Microglia proliferation is regulated by hydrogen peroxide from NADPH oxidase*. J Immunol. 176: 1046-52. DOI: 10.4049/jimmunol.176.2.1046 <http://www.ncbi.nlm.nih.gov/pubmed/16393992>
72. Rojo AI, McBean G, Cindric M, Egea J, Lopez MG, Rada P, et al., (2014) *Redox control of microglial function: molecular mechanisms and functional significance*. Antioxid Redox Signal. 21: 1766-801. DOI: 10.1089/ars.2013.5745 <http://www.ncbi.nlm.nih.gov/pubmed/24597893>
73. Smith AN, Shaughnessy M, Collier S, Hopkins D and Byrnes KR, (2022) *Therapeutic targeting of microglia mediated oxidative stress after neurotrauma*. Front Med (Lausanne). 9: 1034692. DOI: 10.3389/fmed.2022.1034692 <http://www.ncbi.nlm.nih.gov/pubmed/36405593>
74. Brennan AM, Suh SW, Won SJ, Narasimhan P, Kauppinen TM, Lee H, et al., (2009) *NADPH oxidase is the primary source of superoxide induced by NMDA receptor activation*. Nat Neurosci. 12: 857-63. DOI: 10.1038/nn.2334 <http://www.ncbi.nlm.nih.gov/pubmed/19503084>
75. Clausen A, McClanahan T, Ji SG and Weiss JH, (2013) *Mechanisms of rapid reactive oxygen species generation in response to cytosolic Ca²⁺ or Zn²⁺ loads in cortical neurons*. PLoS One. 8: e83347. DOI: 10.1371/journal.pone.0083347 <http://www.ncbi.nlm.nih.gov/pubmed/24340096>

A. *Ex vivo* mouse slice OGD



B. *In vivo* rat cardiac arrest

