UC Irvine

UC Irvine Previously Published Works

Title

Complete mitochondrial genome of the white char *Salvelinus albus* (Salmoniformes, Salmonidae).

Permalink

<https://escholarship.org/uc/item/03r1z2sw>

Journal

Mitochondrial DNA. Part A, DNA mapping, sequencing, and analysis, 27(5)

ISSN

2470-1394

Authors

Balakirev, Evgeniy S
Parensky, Valery A
Kovalev, Mikhail Yu
et al.

Publication Date

2016-09-01

DOI

10.3109/19401736.2015.1079890

Peer reviewed

Mitochondrial DNA

The Journal of DNA Mapping, Sequencing, and Analysis

ISSN: 1940-1736 (Print) 1940-1744 (Online) Journal homepage: <http://www.tandfonline.com/loi/imdn20>

Complete mitochondrial genome of the white char *Salvelinus albus* (Salmoniformes, Salmonidae)

Evgeniy S. Balakirev, Valery A. Parensky, Mikhail Yu. Kovalev & Francisco J. Ayala

To cite this article: Evgeniy S. Balakirev, Valery A. Parensky, Mikhail Yu. Kovalev & Francisco J. Ayala (2015): Complete mitochondrial genome of the white char *Salvelinus albus* (Salmoniformes, Salmonidae), Mitochondrial DNA

To link to this article: <http://dx.doi.org/10.3109/19401736.2015.1079890>

Published online: 10 Sep 2015.

Submit your article to this journal [↗](#)

View related articles [↗](#)

View Crossmark data [↗](#)

MITOGENOME ANNOUNCEMENT

Complete mitochondrial genome of the white char *Salvelinus albus* (Salmoniformes, Salmonidae)Evgeniy S. Balakirev^{1,2,3}, Valery A. Parensky², Mikhail Yu. Kovalev², and Francisco J. Ayala¹¹Department of Ecology and Evolutionary Biology, University of California, Irvine, CA, United States of America, ²A.V. Zhirmunsky Institute of Marine Biology, Far Eastern Branch, Russian Academy of Science, Vladivostok, Russia, and ³Far Eastern Federal University, Vladivostok, Russia

Abstract

The complete mitochondrial genome was sequenced in two individuals of white char *Salvelinus albus*. The genome sequences are 16 653 bp in size, and the gene arrangement, composition, and size are very similar to the salmonid fish genomes published previously. The low level of sequence divergence detected between the genome of *S. albus* and the GenBank complete mitochondrial genomes of the Northern Dolly Varden char *S. malma* (KJ746618) and the Arctic char *S. alpinus* (AF154851) may likely be due to recent divergence of the species and/or historical hybridization and interspecific replacement of mtDNA.

Keywords

Arctic char *S. alpinus*, complete mitochondrial genome, hybridization, mtDNA introgression, Northern Dolly Varden char *S. malma*, salmonids, white char *Salvelinus albus*

History

Received 28 July 2015

Accepted 2 August 2015

Published online 3 September 2015

The white char *Salvelinus albus* Glubokovsky is an endemic of the Kamchatka River, Russia (Glubokovsky, 1977). The species status of the white char was supported with morphological (Glubokovsky, 1995; Parensky et al., 2004; Romanov et al., 2011; Viktorovskii, 1978) and karyological (Frolov, 2001) data. However, genetic markers including allozymes, myogens, mitochondrial DNA, nuclear ribosomal DNA, ITS sequences, and microsatellites failed to confirm the identity of *S. albus* (review in Salmenkova & Omelchenko, 2013).

We have sequenced two complete mitochondrial (mt) genomes of *S. albus* (GenBank accession numbers KT266870 and KT266871) from the Azabachye lake creek (Kamchatka, Russia) to increase the power of phylogenetic analysis of this complex salmonid group, using primers designed with the program mitoPrimer_V1 (Yang et al., 2011). The size of the genome is 16 653 bp and the gene arrangement, composition, and size are very similar to the salmonid fish genomes published previously. The overall base composition was 28.0% for A, 26.4% for T, 17.0% for G, and 28.6% for C. The 54.4% A + T base

composition was higher than G + C, 45.6%. There were no any nucleotide differences between the two isolates (SA1 and SA2) studied.

The comparison of mt genomes now obtained with other complete mt genomes available in GenBank for the family Salmonidae including genera *Salvelinus*, *Parahucho*, *Salmo*, *Hucho*, and *Brachymystax* (Figure 1) reveals a close affinity of *S. albus* to other *Salvelinus* species with a very low level of sequence divergence ($D_{xy} = 0.0020 \pm 0.0003$) between our specimens SA1 and SA2 and the complete mt genome of the Northern Dolly Varden char *S. malma* (KJ746618; Balakirev et al., 2014) (Figure 1). The divergence was higher ($D_{xy} = 0.0079 \pm 0.0006$) between *S. albus* and *S. alpinus* (AF154851; Doiron et al., 2002), but still too low for considering them as separate species. The low level of sequence divergence among *S. albus*, *S. malma*, and *S. alpinus* could be explained by recent divergence and/or historical hybridization and interspecific replacement of mtDNA, as it has been found for other char species (e.g., Bernatchez et al. 1995; Shedko et al. 2007).

Figure 1. Maximum likelihood tree based on 12 protein-coding genes (11 026 bp) for the *Salvelinus albus* specimens SA1 and SA2, and the GenBank representatives of the family Salmonidae. The tree is based on the General Time Reversible + gamma + invariant sites (GTR + G+I) model of nucleotide substitution. The numbers at the nodes are bootstrap percent probability values based on 1000 replications.

Acknowledgements

The authors thank Elena Balakireva for encouragement and help.

Declaration of interest

The authors report that they have no conflicts of interest. The authors alone are responsible for the content and writing of the paper. The work on mitochondrial genome sequencing was supported by the Bren Professor Funds at the University of California Irvine to F. J. Ayala. The analysis of the data was supported by the Russian Science Foundation (RSF; Grant no. 14-50-00034) to E. S. Balakirev

References

- Balakirev ES, Romanov NS, Ayala FJ. (2014). Complete mitochondrial genomes of the Northern (*Salvelinus malma*) and Southern (*Salvelinus curilus*) Dolly Varden charrs (Salmoniformes, Salmonidae). Mitochondrial DNA. [Epub ahead of print]. doi:10.3109/19401736.2014.926531.
- Bernatchez L, Glémet H, Wilson CC, Danzmann R. (1995). Introgression and fixation of Arctic charr (*Salvelinus alpinus*) mitochondrial genome in an allopatric population of brook trout (*Salvelinus fontinalis*). Can J Fish Aquat Sci 52:179–85.
- Doiron S, Bernatchez L, Blier PU. (2002). A comparative mitogenomic analysis of the potential adaptive value of Arctic charr mtDNA introgression in brook charr populations (*Salvelinus fontinalis* Mitchell). Mol Biol Evol 19:1902–9.
- Frolov SV. (2001). Karyological differences between Northern Dolly Varden *Salvelinus malma malma* and white char *Salvelinus albus* from the Kamchatka River basin. Russ J Genet 37:269–75.
- Glubokovsky MK. (1977). *Salvelinus albus* sp. n. from the Kamchatka River basin. Biol Morya 4:49–56.
- Glubokovsky MK. (1995). Evolutionary biology of salmonid fishes (Evolutsionnaya biologiya lososevykh ryb). Moscow: Nauka.
- Parensky VA, Romanov NS, Frolov SV, Nikanorov AP, Repin MYu. (2004). Craniological analysis of Kronotskoe Lake's (Kamchatka) charrs (*Salvelinus*). In: Biodiversity preservation of Kamchatka and adjacent seas (materials of the fifth scientific conference). November 2004. Petropavlovsk-Kamchatsky: Kamchat Press. p 280–3.
- Romanov NS, Frolov SV, Nikanorov AP, Repin MYu. (2011). Some aspects of morphological variability of Kronotskoe Lake's (Kamchatka) charrs. In: Vladimir Ya. Levanidov's biennial memorial meetings, vol. 5. Vladivostok: Dal'nauka. p 458–63.
- Salmenkova EA, Omelchenko VT. (2013). Genetic divergence and taxonomic status of charrs of the genus *Salvelinus*. Biol Bull Rev 3: 481–92.
- Shedko SV, Ginatulina LK, Miroshnichenko IL, Nemkova GA. (2007). Phylogeography of mitochondrial DNA in South Asian Dolly Varden charr *Salvelinus curilis* (Pallas, 1814) (Salmoniformes, Salmonidae): mediated gene introgression? Russ J Genet 43:165–76.
- Viktorovskii RM. (1978). The mechanisms of Charr's Speciation from the Kronotskoe Lake (Mekhanizmy vidobrazovaniya u gol'tsov Kronotskogo ozera). Moscow: Nauka. 112 p.
- Yang CH, Chang HW, Ho CH, Chou YC, Chuang LY. (2011). Conserved PCR primer set designing for closely-related species to complete mitochondrial genome sequencing using a sliding window-based PSO algorithm. PLoS One 6:e17729.