

# UC Irvine

## UC Irvine Previously Published Works

### Title

Mechanosensor Piezo1 mediates bimodal patterns of intracellular calcium and FAK signaling

### Permalink

<https://escholarship.org/uc/item/046748bj>

### Journal

The EMBO Journal, 41(17)

### ISSN

0261-4189

### Authors

Pan, Yijia  
Shi, Linda Zhixia  
Yoon, Chi Woo  
[et al.](#)

### Publication Date

2022-09-01

### DOI




10.15252/emj.2022111799

### Copyright Information

This work is made available under the terms of a Creative Commons Attribution License, available at <https://creativecommons.org/licenses/by/4.0/>

Peer reviewed

# Mechanosensor Piezo1 mediates bimodal patterns of intracellular calcium and FAK signaling

Yijia Pan<sup>1</sup>, Linda Zhixia Shi<sup>2</sup>, Chi Woo Yoon<sup>1</sup>, Daryl Preece<sup>2</sup>, Veronica Gomez-Godinez<sup>2</sup>, Shaoying Lu<sup>1</sup>, Christopher Carmona<sup>1</sup>, Seung-Hyun Woo<sup>3,4</sup>, Shu Chien<sup>1,2,5,\*</sup> , Michael W Berns<sup>2,6,\*\*</sup> , Longwei Liu<sup>1,2,\*\*\*</sup>  & Yingxiao Wang<sup>1,2,\*\*\*\*</sup> 

## Abstract

Piezo1 belongs to mechano-activatable cation channels serving as biological force sensors. However, the molecular events downstream of Piezo1 activation remain unclear. In this study, we used biosensors based on fluorescence resonance energy transfer (FRET) to investigate the dynamic modes of Piezo1-mediated signaling and revealed a bimodal pattern of Piezo1-induced intracellular calcium signaling. Laser-induced shockwaves (LIS) and its associated shear stress can mechanically activate Piezo1 to induce transient intracellular calcium ( $Ca_{[i]}$ ) elevation, accompanied by an increase in FAK activity. Interestingly, multiple pulses of shockwave stimulation caused a more sustained calcium increase and a decrease in FAK activity. Similarly, tuning the degree of Piezo1 activation by titrating either the dosage of Piezo1 ligand Yoda1 or the expression level of Piezo1 produced a similar bimodal pattern of FAK responses. Further investigations revealed that SHP2 serves as an intermediate regulator mediating this bimodal pattern in Piezo1 sensing and signaling. These results suggest that the degrees of Piezo1 activation induced by both mechanical LIS and chemical ligand stimulation may determine downstream signaling characteristics.

**Keywords** calcium; FAK; laser-induced shockwaves; Piezo1; SHP2

**Subject Categories** Membrane & Trafficking; Signal Transduction

**DOI** 10.15252/emj.2022111799 | Received 31 May 2022 | Revised 19 June 2022 | Accepted 28 June 2022 | Published online 17 July 2022

**The EMBO Journal (2022) 41: e111799**

## Introduction

Cells constantly encounter mechanical forces in the forms of shear stress, compression, and stretching. Mechanotransduction is crucial

for cells to perceive the local microenvironments to maintain cellular function and tissue homeostasis (Vogel & Sheetz, 2006; Liang & Boppart, 2010; Park *et al*, 2010; Egan *et al*, 2015). Alterations of mechanotransduction can lead to pathological conditions (Jaalouk & Lammerding, 2009). However, it remains unclear on the detailed mechanism by which cells perceive mechanical cues of the external microenvironment and transduce them into molecular signals and genetic regulations for the coordination of cellular responses.

Several families of molecular mechanoreceptors have been reported to transduce mechanical stimulation into ion currents to modulate physiological functions, particularly the transmembrane ion channel proteins (Ranade *et al*, 2015). Piezo1 has been identified as a component of mechanically activated cation channels in both vertebrates and non-vertebrates. It induces currents in various cell types upon the application of mechanical pressure, tension, or deformation to the cell membrane (Coste *et al*, 2010, 2012; Liao *et al*, 2019). Most of the mechanical stimulations resulted in transient calcium influx, although sustained calcium influx upon prolonged mechanical stretch was also reported (Miyamoto *et al*, 2014). Consequently, Piezo1 plays different roles in physiology, including angiogenesis, vasculogenesis (Li *et al*, 2014), and erythrocyte volume regulation (Cahalan *et al*, 2015). Piezo1 is also involved in regulating cancer biology in complicated manners. For instance, blocking Piezo1 decreased the motility of the breast cancer cell line MCF-7 (Li *et al*, 2015), but increased the migration speed of small cell lung cancer cell lines (McHugh *et al*, 2012). These results suggest a complex role of Piezo1 in regulating cancer cell migration under different biological and environmental contexts.

Due to the important role of Piezo1 in regulating physiological and pathological cell migration, an emerging topic of interest has been focused on the dynamics and effects of Piezo1 activation on cytoskeleton (Nourse & Pathak, 2017). Focal adhesion zones are hubs for cytoskeletal structures that connect the extracellular matrix

1 Department of Bioengineering, University of California, San Diego, La Jolla, CA, USA

2 Institute of Engineering in Medicine, University of California, San Diego, La Jolla, CA, USA

3 Department of Cell Biology, Dorris Neuroscience Center, The Scripps Research Institute, La Jolla, CA, USA

4 Genomic Institute of the Novartis Research Foundation, San Diego, CA, USA

5 Department of Medicine, University of California, San Diego, La Jolla, CA, USA

6 Beckman Laser Institute and Medical Clinic, University of California, Irvine, Irvine, CA, USA

\*Corresponding author. Tel: 1 (858) 534 5195; E-mail: shuchien@ucsd.edu

\*\*Corresponding author. Tel: 1 (858) 822 2954; E-mail: mwberns17@gmail.com

\*\*\*Corresponding author. Tel: 1 (858) 999 7657; E-mail: lol001@eng.ucsd.edu


\*\*\*\*Corresponding author. Tel: 1 (858) 882 2450; E-mail: yiw015@eng.ucsd.edu

to the intracellular cytoskeleton, and they can detect and transmit external mechanical forces. Focal adhesion kinase (FAK) belongs to focal adhesion zone proteins and is a key regulator of focal adhesion dynamics and cytoskeletal remodeling. Therefore, studying the effect of Piezo1 activation on FAK dynamics can give insights into how Piezo1 affects cellular migration. In this study, we applied laser-induced shockwaves (LIS) and chemical stimulations to activate the Piezo1 in live cells and studied the mechanism by which Piezo1 regulates molecular events, for example, intracellular calcium ( $Ca_{[i]}$ ) and the activity of FAK. Laser-induced shockwaves have been applied in mechanical destruction of cancer cells; at the same time, it serves as a useful tool to study cancer cell mechanical properties (Quinto-Su *et al*, 2011; Steinhauser & Schmidt, 2014; Shannon *et al*, 2017). LIS are produced by the sudden buildup of energy at the focus of a short-pulsed laser beam (Lokhandwalla *et al*, 2001). The subsequent microplasma cavitation, followed by microbubble expansion and contraction, and consequently shear stress, can exert a large mechanical force in the surrounding area (Rau *et al*, 2006). The force is controllable in magnitude by tuning the laser power and can exert instantaneous pressures at the Mega Pascal range to induce intracellular calcium responses (Suhr *et al*, 1996; Tsukamoto *et al*, 2011; Fan *et al*, 2012; Compton *et al*, 2014; Gomez-Godinez *et al*, 2015; Li *et al*, 2018). Therefore, laser-induced cavitation and the consequent shear stress can serve as a mechanical stimulator of Piezo1 on single live cells with high precision in space and time (Li *et al*, 2013, 2018; Compton *et al*, 2014; Yuan *et al*, 2015).

Genetically encoded biosensors based on FRET have enabled the visualization of signaling events in live cells with high spatiotemporal resolution (Lu *et al*, 2013). They can be applied at the same time to monitor in live cells the Piezo1 activation and its associated  $Ca_{[i]}$  dynamics, as well as the FAK activation, a crucial signaling component that regulates cell adhesion and motility (Mitra *et al*, 2005). In this study, by combining mechanical LIS- or chemical ligand-induced Piezo1 activation and live-cell FRET imaging, we revealed a bimodal pattern of Piezo1-induced action of  $Ca_{[i]}$  dynamics (transient vs. sustained) and downstream FAK molecular events (activation vs. inhibition), dependent on the degree of Piezo1 activation. These results advance our understanding of Piezo1 activation and its role in perceiving mechanical and chemical cues and regulating intracellular molecular signals.

## Results

To explore whether LIS-induced mechanical stimulation can activate the Piezo1 channel, HEK 293T cells co-transfected with Piezo1-tdTomato and a CFP/YFP-based FRET calcium biosensor (Palmer & Tsien, 2006) were examined in a LIS-FRET imaging system (Gomez-Godinez *et al*, 2015) (Fig 1). To create a shockwave sufficient to mechanically stimulate the target cells (Rau *et al*, 2006) without inducing cell damage, the laser power was precisely controlled in a range between 160 and 180  $\mu$ W (Appendix Figs S1 and S2A and B). In a previous study, we showed that cells subjected to LIS treatment with powers at the level up to 342  $\mu$ W and at a distance of 70  $\mu$ m remained healthy and did not uptake PI (Gomez-Godinez *et al*, 2015), a reliable marker for cell necrosis and membrane portion (Fan *et al*, 2012; Li *et al*, 2018). Upon one pulse of LIS stimulation




**Figure 1. Diagram of an integrated system of laser-induced shockwave (LIS) and FRET imaging.**

The CCD camera captures fluorescent images of FRET biosensors with 480/30 nm and 535/25 nm bandpass filters integrated into the dual view system. The ablation laser is guided through the side port of the microscope. The arc lamp excitation light enters through the backport of the microscope and is filtered with a 440/20 nm bandpass filter.

of the Piezo1-expressing cells, calcium influx was immediately detected in cells cultured in calcium-containing HBSS media, but not in HBSS without calcium (Fig 2A and B). In control HEK cells transfected only with a calcium biosensor but not Piezo1, there was no significant calcium influx in HBSS media with or without calcium (Fig 2A and B, Appendix Fig S3 and Movie EV1). In HBSS media without calcium, there was a minor calcium increase in the Piezo1-transfected cells (Appendix Fig S3), possibly due to the ER calcium release as Piezo1 can be expressed in ER (McHugh *et al*, 2012). Intermittent shockwave pulses elicited multiple calcium waves (Fig 2C), verifying a healthy response and a lack of shockwave-induced damage on the target cell.

We next examined the effect of this LIS on FAK activity visualized by a FAK FRET biosensor developed earlier by our group (Seong *et al*, 2011). HEK 293T cells co-transfected with Piezo1-tdTomato and the ECFP/YFP-based FAK FRET biosensor were exposed to LIS in HBSS media containing calcium. One pulse of shockwave stimulation caused an increase of FAK activity in HEK cells transfected with Piezo1, but no significant change in the control HEK cells without Piezo1 expression (Fig 3A–C and Movie EV2). These results indicate that LIS-induced Piezo1 activation could lead to FAK activation, potentially due to the transient calcium influx.

Aside from mechanical stimulation, chemical agonists can also activate the Piezo1 ion channel. To further gain molecular insights into the role of Piezo1 in FAK activation, Yoda1, a synthetic small molecule that has been reported to act as a specific agonist for



**Figure 2. The Piezo1-dependent calcium influx upon LIS stimulation in HEK cells.**


- A The time courses of normalized FRET ratio (Mean  $\pm$  SEM) of calcium biosensor before and after shockwave stimulation in HBSS Ca<sup>2+</sup> medium with (yellow line,  $n = 6$ ,  $N = 3$ ) or without the Piezo1 expression (black line,  $n = 6$ ,  $N = 3$ ).
- B The representative FRET/ECFP ratio images of calcium biosensor in Piezo1-expressed HEK cells before (top left) and after (top right) LIS stimulation. Phase image with laser initiating point (left) and fluorescent image indicating Piezo1 expression (right) were also shown on the bottom panels (Scale bar, 40  $\mu$ m). Color scale bars in the figures are to show the FRET/ECFP ratios, with cold and hot colors representing low and high ratios, respectively.
- C The time course of normalized FRET ratio (Mean  $\pm$  SEM) of calcium biosensor in the Piezo1-expressing HEK cells before and after multiple shockwave stimulations in HBSS Ca<sup>2+</sup> medium ( $n = 6$ ,  $N = 3$ ).

Data information: “ $n$ ” means the total cell number. “ $N$ ” means the number of individual experiments.

activating human and mouse Piezo1 (Syeda *et al*, 2015), was applied to stimulate Piezo1. We stimulated HEK cells that have been co-transfected with Piezo1 and the FAK FRET biosensor with 25  $\mu$ M Yoda1 as a standard dosage (Syeda *et al*, 2015; Lacroix *et al*, 2018). Surprisingly, Yoda1 stimulation of Piezo1 caused a significant decrease in FAK activity in the HEK cells expressing exogenous Piezo1 (Fig 4A and B); this is opposite to the increase in FAK activity in response to mechanical stimulation of Piezo1 by one pulse of LIS. Hence, it was hypothesized that the LIS-induced mechanical and 25  $\mu$ M Yoda1-based chemical stimulations result in different modes of the Piezo1 activation. To test this hypothesis, we co-transfected HEK cells with Piezo1 and the calcium FRET biosensor. Upon Yoda1 stimulation at 25  $\mu$ M, a sustained calcium influx was observed in the cells transfected with Piezo1 (Fig 4C and D), which is significantly different from the transient Ca<sub>[i]</sub> dynamics observed in the one-pulse LIS stimulated cells (Fig 2A). We also measured the cell viability by calcein-AM dye after 25  $\mu$ M Yoda1 stimulation, and we did not observe cell damage (Appendix Fig S4A), suggesting the sustained Ca<sup>2+</sup> did not affect cell health. To further exclude the influence of reporting artifacts of FRET biosensors, we expressed in cells a soluble nonfunctional FRET biosensor containing ECFP and YPet pair but with its binding domain mutated and tracked its FRET change after stimulated with 25  $\mu$ M Yoda1. No significant FRET

ratio change was observed (Appendix Fig S4B). These results suggest that our biosensor-reported biochemical signals induced by Piezo1 activation are not due to the cell morphology changes nor the reporting artifacts of FRET biosensors. The differential FAK responses may hence be attributed to the differences in Ca<sub>[i]</sub> dynamic patterns upon the mechanical and chemical stimulations.

To further examine this hypothesis, we titrated the concentration of Yoda1 used to stimulate cells transfected with Piezo1. We observed that a decrease in Yoda1 concentration from 25 to 0.5  $\mu$ M caused a shift in Ca<sub>[i]</sub> dynamics pattern from a sustained increase in a transient rise (Fig 5A and B). Correspondently, the FAK response to Yoda1 stimulation changed from a decrease to an increase when the Yoda1 dosage was reduced from 25 to 0.5  $\mu$ M in the Piezo1-expressing cells (Fig 5C and D). Consistent with these results, increases in the Piezo1 expression level in cells under an intermediate dosage of 2  $\mu$ M Yoda1 stimulation caused a shift in the dynamic pattern of Ca<sub>[i]</sub> signaling from transient to sustained (Fig 5E and Appendix Fig S5). We also tested additional concentrations of Yoda1 at higher or lower ranges. Indeed, when we increased Yoda1 concentration from 25  $\mu$ M to 40  $\mu$ M, a greater reduction in FAK activity was observed. At a lower range, we did not observe a significant change in the degree of FAK activity increase, potentially due to less sensitive responses of FAK activities at low levels (Appendix Fig S6A–E).



**Figure 3. The Piezo1-dependent FAK activation upon LIS stimulation in HEK cells.**

**A** The representative ECFP/FRET ratio images of FAK biosensor in Piezo1 overexpressed HEK cells before (left) and after (right) shockwave stimulation (Scale bar, 40  $\mu$ m). Color scale bars in the figures are to show the ECFP/FRET ratios, with cold and hot colors representing low and high ratios, respectively.

**B** The time courses of normalized ECFP/FRET ratio (Mean  $\pm$  SEM) of FAK FRET biosensor before and after shockwave stimulation in HBSS  $\text{Ca}^{2+}$  media with (green line,  $n = 16$ ,  $N = 3$ ) or without Piezo1 (black line,  $n = 10$ ,  $N = 3$ ).

**C** The percentage change of FRET ratio of the FAK FRET biosensor in HEK cells with or without Piezo1 after shockwave stimulation in HBSS  $\text{Ca}^{2+}$  medium. \* $P < 0.05$  from Student's two-tailed  $t$ -test.


Data information: "n" means the total cell number. "N" means the number of individual experiments.

These results suggest that the level of Piezo1 activation upon Yoda1 stimulation may have led to differential dynamic patterns of calcium influx to result in FAK activation versus inhibition.

We thereafter examined whether the bimodal pattern of Piezo1 action observed by chemical stimulation can also be induced by LIS mechanical stimulation with different strength: comparing the effect of multiple versus single pulse(s) of LIS stimulation. This multiple-pulse LIS stimulation was created by firing laser repeatedly three times within a short time interval of 5 s (Fig 6A). Multiple LIS stimulation caused a more sustained calcium increase and a decrease of FAK activity in cells transfected with Piezo1, compared with a more transient calcium increase and a slight increase in FAK activity upon single-pulsed stimulation (Fig 6B–D). These results are consistent with Piezo1 activation under chemical stimulation, suggesting that different modes of Piezo1 activation can also be triggered by mechanical LIS stimulation. As such, there exists a bimodal pattern of Piezo1 activation-induced  $\text{Ca}_{[i]}$  dynamics (transient vs. sustained) and downstream FAK (activation vs. inhibition), with the transient

mode causing a transient  $\text{Ca}_{[i]}$  rise and a FAK activation while the sustained mode causing a sustained  $\text{Ca}_{[i]}$  rise and a FAK suppression (Fig 6E).

Next, we investigated the molecular mechanism by which the degree of Piezo1 activation and calcium dynamics lead to the FAK activation or inhibition. The Src-homology-2 (SH2) domain-containing protein tyrosine phosphatase (SHP2), which can be regulated by calcium influx (Rusanescu *et al*, 2005), is a well-known regulator of cell adhesion (Yu *et al*, 1998; Oh *et al*, 1999; Inagaki *et al*, 2000) and FAK activity through its tyrosine phosphatase catalytic domain (PTP domain) directly (Yamauchi *et al*, 1995; Ouwens *et al*, 1996; Mañes *et al*, 1999). SHP2 can also work as an adaptor molecule that binds to phospho-tyrosine containing activators via its two SH2 domains, including insulin receptor substrate 1 (IRS1), GRB2-associated binding protein 1 (GAB1), and non-tyrosine containing substrates such as p53 and fatty acid synthase (FASN) (Noguchi *et al*, 1994; Yu *et al*, 2013; Guo & Xu, 2020). This function of SHP2 is independent of its phosphatase activity



**Figure 4. The FAK and calcium activation upon Yoda1 stimulation in the Piezo1-expressing HEK cells.**

- A The representative ECFP/FRET ratio images of FAK biosensor in Piezo1 expressing HEK cells before and after 25  $\mu$ M Yoda1 stimulation (Scale bar, 10  $\mu$ m). Color scale bars in the figures indicate ECFP/FRET ratios, with cold and hot colors representing low and high ratios, respectively.
- B The time courses of normalized FRET ratio (Mean  $\pm$  SEM) of FAK FRET biosensor before and after Yoda1 stimulation with (pink line,  $n = 10$ ,  $N = 3$ ) or without Piezo1 (black line,  $n = 11$ ,  $N = 3$ ). The gray line represents the DMSO control group ( $n = 8$ ,  $N = 3$ ). Color scale bars in the figures indicate FRET/ECFP ratios, with cold and hot colors representing low and high ratios, respectively.
- C The representative FRET/ECFP ratio images of the calcium biosensor in the Piezo1-expressing HEK cells before and after 25  $\mu$ M Yoda1 stimulation (Scale bar, 10  $\mu$ m). Color scale bars in the figures indicate FRET/ECFP ratios, with cold and hot colors representing low and high ratios, respectively.
- D The time courses of normalized FRET ratio (Mean  $\pm$  SEM) of calcium biosensor in the Piezo1-expressing HEK cells (purple line,  $n = 9$ ,  $N = 3$ ) before and after 25  $\mu$ M Yoda1 stimulation in culture medium.

Data information: “ $n$ ” means the total cell number. “ $N$ ” means the number of individual experiments.

(Noguchi *et al*, 1994; Yu *et al*, 2013; Guo & Xu, 2020). Thus, we hypothesized that SHP2 may work as an intermediate regulator in Piezo1 signaling to regulate FAK differentially in response to different stimulations, through its adaptor or phosphatase function under different conditions. To test this hypothesis and investigate whether and how SHP2 regulates FAK activity downstream of Piezo1 activation and calcium dynamics, we first examined the SHP2 activation pattern following Piezo1 agonist stimulation utilizing a FRET-based SHP2 biosensor (Sun *et al*, 2017). A rapid and sustained SHP2 activation was observed in Piezo1 overexpressing cells after being stimulated with 25  $\mu$ M Yoda1, while no apparent SHP2 activation in cells stimulated with 0.5  $\mu$ M Yoda1 (Fig 7A and B). This SHP2 activation is Piezo1 dependent, as no obvious response was observed in Piezo1-negative cells (Fig 7A and B). Such SHP2 activation can lead to the FAK dephosphorylation and suppression (Fig 4B) (Yamauchi *et al*, 1995; Ouwens *et al*, 1996; Mañes *et al*, 1999). In contrast, in cells overexpressing a catalytically dead SHP2 (C495S) mutant, FAK was activated when stimulated with 25  $\mu$ M Yoda1 in cells expressing Piezo1 (Fig 7C), to the same extent as in the control cells

stimulated by 0.5  $\mu$ M Yoda1 stimulation (Fig 7D). Similar results were obtained in cells pretreated with a potent, highly selective SHP2 targeting inhibitor (SHP 099) (Yuan *et al*, 2020) that stabilized the auto-inhibited SHP2 conformation and inhibited its catalytic function (Fig 7C and D). These results suggest that SHP2 can lead to FAK activation independent of its catalytic function upon Piezo1 activation. As such, SHP2 may function as a molecular switch to regulate FAK activity in a Piezo1-dependent manner (Fig 7E). SHP2 can be activated under sustained calcium influx to enzymatically dephosphorylate and suppress the FAK kinase; in contrast, under transient calcium, SHP2 serves as an adaptor protein with its enzymatic domain kept in an autoinhibited form to facilitate the activation of FAK.

## Discussion

In this study, we revealed a bimodal pattern of Piezo1 activation-induced  $\text{Ca}_{\text{[i]}}$  dynamics (transient vs. sustained) and downstream

FAK (activation vs. inhibition), with the transient mode causing a transient  $Ca_{[i]}$  rise and a FAK activation while the sustained mode causing a sustained  $Ca_{[i]}$  rise and a FAK suppression. By combining

FRET imaging with laser-based photonics, we have successfully monitored the distinct molecular signaling events with high spatiotemporal resolutions in live cells following the mechanosensor




Figure 5.



**Figure 5. The titration of Yoda1 stimulation in the Piezo1-expressing HEK cells.**

- A The time courses of normalized FRET ratio (Mean  $\pm$  SEM) of calcium biosensor in the Piezo1-expressing HEK cells upon 0.5  $\mu$ M (Orange line,  $n = 10$ ,  $N = 3$ ) or 25  $\mu$ M (purple line,  $n = 8$ ,  $N = 3$ ) Yoda1 stimulation.
- B The percentage changes of FRET ratio (Mean  $\pm$  SEM) decrease of calcium biosensor in Piezo1-expressing HEK cells upon 0.5  $\mu$ M (Orange bar,  $n = 8$ ,  $N = 3$ ) or 25  $\mu$ M (purple bar,  $n = 10$ ,  $N = 3$ ) Yoda1 stimulation at different time points after they reached the peak. \*\*\*\* $P < 0.0001$  from Student's two-tailed  $t$ -test.
- C The time courses of normalized FRET ratio (Mean  $\pm$  SEM) of FAK FRET biosensor before and after 0.5  $\mu$ M (green line,  $n = 8$ ,  $N = 3$ ) or 25  $\mu$ M (pink line,  $n = 10$ ,  $N = 3$ ) Yoda1 stimulation.
- D The percentage change of FRET ratio (Mean  $\pm$  SEM) changes of FAK biosensor in the Piezo1-expressing HEK cells after 0.5  $\mu$ M (green bar,  $n = 8$ ,  $N = 3$ ) or 25  $\mu$ M (pink bar,  $n = 10$ ,  $N = 3$ ) Yoda1 stimulation at different time points. \*\*\* $P = 0.0005$  and \*\*\*\* $P < 0.0001$  from Student's two-tailed  $t$ -test.
- E The time courses of normalized FRET ratio (Mean  $\pm$  SEM) of calcium biosensor before and after 2  $\mu$ M Yoda1 stimulation in cells with high (intensity  $>4,650$  au., light red line,  $n = 14$ ,  $N = 3$ ) or low (intensity  $<1,000$  au., Blue line,  $n = 5$ ,  $N = 2$ ) Piezo1 expression.
- Data information: "n" means the total cell number. "N" means the number of individual experiments.

Piezo1 activation by LIS mechanical stimulation and the ligand chemical stimulation.

Interestingly, high levels of Piezo1 activation, either attributed to high dosages of Yoda1 application or to high expressions of Piezo1, caused sustained calcium signaling and FAK suppression. Tuning down the Piezo1 activation level can lead to a switch of calcium and FAK responses, with the dynamic  $Ca_{[i]}$  signaling becoming more transient, accompanied by an activation of FAK. This is consistent with previous reports that transient calcium influx could lead to increased FAK phosphorylation (Giannone *et al*, 2002; Kaczmarek *et al*, 2005). These results indicate that the strength of Piezo1 activation can lead to two distinctive activation patterns (bimodal) of downstream signaling, with low levels of Piezo1 activation triggering transient calcium signaling and FAK activation (transient mode) while high levels of Piezo1 activation led to sustained calcium signaling and FAK suppression (sustained mode) (Fig 6E). We believe that calcium plays a central role in the bimodal actions of FAK. It is well established that the kinetics and amplitude of  $Ca^{2+}$  signals allow the activation of specific  $Ca^{2+}$  targets and potentially the differential focal adhesion (FA) characteristics (Berridge, 1997). Specifically, the process of cellular adhesion involves the dynamic regulation of FAs complexes (Giannone *et al*, 2004; Wolfenson *et al*, 2014). FAK can phosphorylate local substrate molecules and create docking sites for the molecular recruitment and further assembly of the FA complex (Yu *et al*, 2011; Oakes *et al*, 2012). We indeed examined FAK dynamics by tracking a focal adhesion-associated adaptor protein paxillin (Schaller, 2001). We stimulated Piezo1-expressing HEK cells with 25  $\mu$ M or 0.5  $\mu$ M Yoda1, and observed a decrease in paxillin intensity in 25  $\mu$ M Yoda1-treated cells, but a slight increase in paxillin intensity with 0.5  $\mu$ M Yoda1 treatment

(Appendix Fig S6A–D). These observations are also consistent with the FAK suppression and activation caused by 25  $\mu$ M and 0.5  $\mu$ M Yoda1, respectively.

Further investigations suggest that SHP2 may serve as a molecular switch downstream of Piezo1 to regulate FAK activity, which leads to different FAK responses upon the different levels of Piezo1 activation. SHP2 can be activated under sustained calcium influx to enzymatically dephosphorylate and suppress the FAK kinase, while serving as an adaptor protein independent of its enzymatic function to facilitate the activation of FAK under transient calcium. The difference in the time course of calcium signal in imaging experiments compared with the time course of Piezo1 activation revealed in electrophysiological recordings (Syeda *et al*, 2015) raises the possibility that there may exist a signaling amplifier to mediate the actions of Piezo1 and facilitate the bimodal downstream calcium and other signaling cascades. However, this difference could also be attributed to the distinct detecting methods, in which a fluorescent biosensor typically monitors accumulated calcium in the cytoplasm, and patch-clamps measure the membrane potential or electrophysiological currents passing through ionic channels on the plasma membrane. Although correlated, the transformation from the electrophysiological signal to calcium is inherently nonlinear due to the dynamics of calcium concentrations at different subcellular locations (Scheuss *et al*, 2006; Akerboom *et al*, 2012; Wei *et al*, 2020). In addition, the calcium response illustrated by the biosensor represents the accumulative net outcome of the activation of the whole Piezo1 population as a group on the cell surface, in which the large number of Piezo1 may become activated in an unsynchronized fashion, which contributes to the relatively slow kinetics of observed calcium signals. In contrast,

**Figure 6. Calcium and FAK activation upon multiple pulsed LIS stimulation in Piezo1 overexpressing HEK cells.**

- A The stimulation pattern of multiple-pulsed laser stimulation (3 pulses, 5 s interval).
- B The time courses of normalized FRET ratio (Mean  $\pm$  SEM) of d3cpv biosensor before and after three-coupled multiple pulsed or single-pulsed LIS stimulation with 5-s interval (blue line,  $n = 6$ ,  $N = 3$ ; red line,  $n = 15$ ,  $N = 3$ ).
- C The percentage change of normalized FRET integral change ratio (See materials and methods, Mean  $\pm$  SEM) of calcium biosensor in HEK cells with (red bar,  $n = 15$ ,  $N = 3$ ) or without (blue bar,  $n = 12$ ,  $N = 3$ ) Piezo1 expression before and after multiple-pulsed LIS pulsed stimulations. For each cell, the FRET integral ratio was calculated as the ratio of the area under the time course of the FRET/ECFP ratio (blue area A) to the area under the maximum ratio (patterned area B) over the whole time course after LIS stimulation. \*\*\*\* $P < 0.0001$ , two-tailed unpaired Student's  $t$ -test.
- D The time courses of normalized FRET ratio (Mean  $\pm$  SEM) of FAK biosensor before and after multiple-pulsed LIS stimulation at 5-s interval (blue line,  $n = 16$ , red line,  $n = 36$ ,  $N = 3$ ).
- E A cartoon diagram depicting the activation patterns of Piezo1 on downstream calcium (blue or red represents transient or sustained calcium increase pattern, respectively) and FAK signaling (biphasic: blue or red represents a suppression or activation of FAK activity, respectively) upon stimulation. LIS specifically elicits the lower phase (transient mode) of Piezo1 actions (indicated by the pink shade).
- Data information: "n" means the total cell number. "N" means the number of individual experiments.






Figure 6.

electrophysiological signals represent the instantaneous response of localized and limited copies of Piezo1 on the cell membrane at the proximity of patch clamp, which may better reflect single Piezo1 behavior and be much faster. Thus, the transient and sustained calcium response patterns that we observed here, whose

time courses are different from the reported electrophysiological signals, do not necessarily suggest a signal amplifier downstream of Piezo1 in regulating the calcium signals. It is possible that a stronger stimulation can cause the opening of multiple channel clusters with more accumulative and sustained signaling, in

contrast to the limited copies of channel opening induced by a weaker stimulation which can result in a quick reversible and transient signaling. Nevertheless, we cannot rule out the possibility that there is a signal amplifier downstream of Piezo1 that is required to lead to both the transient and the sustained calcium

signals presented in this study. While further investigations will be warranted to reveal more molecular insights, this study should shed new light on how the mechanosensitive Piezo1 regulates cell adhesion and paves the way to a better understanding of Piezo1-related mechanotransduction processes.




Figure 7.

**Figure 7. SHP2 works as a molecular switch downstream of Piezo1 to regulate FAK.**

- A, B The time courses (Mean  $\pm$  SEM) (A) and representative images (Scale bar, 10  $\mu$ m) of FRET ratio of SHP2 biosensor (Y542F) before and after 25  $\mu$ M or 0.5  $\mu$ M Yoda1 stimulation in cells with or without Piezo1 overexpression (red bar,  $n = 5$ ,  $N = 3$ ; blue bar,  $n = 5$ ,  $N = 3$ ; purple bar,  $n = 11$ ,  $N = 3$ ; green bar,  $n = 5$ ,  $N = 3$ ).
- C The time courses of normalized FRET ratio (Mean  $\pm$  SEM) of FAK FRET biosensor under different treatments. (From top to the bottom,  $n = 8$ , 10, 14, and 14, respectively; SHP 099, the cells were pretreated with 10  $\mu$ M SHP099; SHP2 C495S, the cells overexpressing SHP2 protein with a non-functional catalytic domain).
- D The percentage change of FRET ratio (Mean  $\pm$  SEM) changes of FAK biosensor in the Piezo1-expressing cells in C at 5 min. (From left to right,  $n = 8$ , 10, 14, and 14 respectively; one-way ANOVA, \*\* $P = 0.006$ , \*\*\*\* $P < 0.0001$ , NS = not significant.)
- E Proposed working model of FAK regulation by Piezo1 in which SHP2 works as a molecular switch.
- Data information: “ $n$ ” means the total cell number. “ $N$ ” means the number of individual experiments.

Our findings of the bimodality of Piezo1-induced signaling should also help to advance our understanding of the opposite effects of Piezo1 knockdown on the migration of two cell lines: a decrease for gastric tumor cell (Yang *et al*, 2014), but an increase for lung epithelial cells (McHugh *et al*, 2012). As stomach and lung cells have a low and high level of Piezo1 expression, respectively (Miyamoto *et al*, 2014), the knockdown of Piezo1 on these cells may have tuned their Piezo1 levels and hence sensing strengths differentially in response to environmental cues. This may ultimately lead to the different outcomes in perturbing cell migration, in which FAK plays a crucial role. As such, our results provide some insights into the paradox that the same manipulation of Piezo1 under different cell types can lead to differential outcomes in cell migration (Bagriantsev *et al*, 2014; Li *et al*, 2015).

## Materials and Methods

### DNA plasmid

The construct of the mPiezo1-tdTomato was obtained from Dr. Patapoutian's lab (Cahalan *et al*, 2015). The D3cpv calcium FRET biosensor was previously described (Palmer & Tsien, 2006), the cytosolic FAK FRET biosensor was previously developed and described (Seong *et al*, 2011).

### Cell culture and transfection

HEK 293T cell line was a kind gift from Dr. Patapoutian's lab (Coste *et al*, 2010) and HeLa cells were purchased from American Type Culture Collection (ATCC). Cell culture reagents were obtained from Invitrogen. Cells were maintained in Dulbecco's modified Eagle medium (DMEM) supplemented with 10% fetal bovine serum (FBS), 2 mM L-glutamine, 1 unit/ml penicillin, 100  $\mu$ g/ml streptomycin, and 1 mM sodium pyruvate. Cells were cultured in a humidified 95% air, 5% CO<sub>2</sub> incubator at 37 °C. FUGENE 6 (Invitrogen) was used for transfection of DNA plasmids. Cells were tested 36–48 h post-transfection. The cells were tested mycoplasma negative.

During imaging, cells were cultured in fibronectin-coated coverglass-bottom dishes and maintained in Hanks buffered saline solution (HBSS) with or without calcium and magnesium (Invitrogen, Carlsbad, CA). Chemical stimulant, Yoda1, was purchased from Thermo fisher scientific. Calcein AM was purchased from Thermo fisher scientific. Cells were stained with Calcein AM according to the manufacturer's manual.

### Laser-shockwave system

The laser used for the shockwave stimulation is a 532 nm wavelength, 100 Hz repetition rate, and 2 ns pulse width Coherent Flare system (Spectra-Physics). Laser power is attenuated by rotating an optical polarizer mounted in a stepper-motor-controlled rotating mount (New Port). A mechanical shutter (Vincent Associates) with a 10–15 ms duty cycle is gated to allow 1–2 laser pulses to pass through when firing. The beam expanders are used to adjust the beam diameter of the lasers to fill the back aperture of the objective. The custom laser entry port (CLEP) is placed underneath the objective (phase III, 40 $\times$ , NA 1.3, oil immersion) of a Zeiss Axiovert 200M microscope. The filter on the CLEP was chosen to direct the laser lights onto the lower right corner of the image screen.

### Laser-induced shockwave (LIS)

To understand precisely the forces presented in the shockwave system, we followed a similar method in a previous publication to measure and calculate forces (Rau *et al*, 2006). The time-resolved shockwave signal was determined by monitoring the size of each bubble created by the laser focus plasma. To achieve this, a low power measurement laser at 633 nm was arranged to be coaxial with the shockwave beam. The transmitted power of the beam was monitored using a fast photodiode. Laser focus was at 10  $\mu$ m above the substrate. Upon the creation of a bubble during the measurement, laser was deflected and the signal was captured.

A series of measurements at different powers between time-averaged 160  $\mu$ W and 293  $\mu$ W measured at the back aperture were conducted. The peak beam deflection and the time of bubble rise and collapse were recorded for calculation.

By utilizing the Rayleigh Equation for bubble growth:

$$R_{max} = \frac{T}{0.915} \left[ \frac{p_{\infty} - p_v}{\rho} \right]^{\frac{1}{2}}$$

where  $R_{max}$  is the maximum bubble radius,  $T$  is the time for bubble collapse. Here  $p_{\infty}$  is the atmospheric pressure,  $p_v$  is the vapor pressure within the bubble, and  $\rho$  is the density of the fluid. This information can then be fit into the Gilmore Equation and applied to a Logarithmic fit function to calculate the time-dependent bubble expansions.

We calculated the radial shear forces in our system by applying the methods shown previously (Lokhandwalla *et al*, 2001; Lokhandwalla & Sturtevant, 2001; Gomez-Godinez *et al*, 2015).

The radial distribution of shear forces was shown below.

Time averaged power ( $\mu\text{W}$ )	Optical intensity ( $\text{GW}/\text{m}^2$ )	Pulse Energy ( $\mu\text{J}$ )	Bubble Energy ( $\mu\text{J}$ )	Peak Shear Stress (MPa)
$164 \pm 0.1$	$1.24\text{e}+4 \pm 1\text{e}+3$	$1.6 \pm 0.04$	$0.21 \pm 0.02$	$21.10 \pm 4$
$170 \pm 0.1$	$1.28\text{e}+4 \pm 1\text{e}+3$	$1.7 \pm 0.04$	$0.22 \pm 0.02$	$21.38 \pm 4$
$180 \pm 0.1$	$1.36\text{e}+4 \pm 1\text{e}+3$	$1.8 \pm 0.04$	$0.23 \pm 0.02$	$21.84 \pm 4$

We noted that cells have been shown to survive through pressures up to 120 MPa (Rau *et al.*, 2004), as cell lysis is not only dependent on the peak shear but also on the duration of the applied shear stress, as well as cell types and bubble expansion radius (Yuan *et al.*, 2015). The cell lysis was observed to occur at distances below 120  $\mu\text{m}$  away from the epicenter of the cavitation with our current experimental configuration. In our mechano-stimulation experiments, cells were positioned more than 190  $\mu\text{m}$  away from the cavitation epicenter to avoid cell damages.

### Image acquisition

Images of shockwave stimulation were collected by ORCA-Flash4.0 V2 Digital CMOS camera (Hamamatsu) passing a 440DF20 excitation filter, a 455DRLP dichroic mirror, and two emission filters placed in a dual view adaptor (480DF30 for ECFP and 535DF25 for EYFP). Images of agonist stimulation were collected from a Nikon microscope with a charge-coupled device (CCD) camera with the same filter setting. For FAK biosensor, the pixel-by-pixel ratio images of ECFP/FRET were calculated based on the background-subtracted fluorescence intensity images of CFP and YFP by the Metafluor program to allow quantification and statistical analysis of FRET responses. For calcium biosensor, the emission ratio of FRET/ECFP was used to represent calcium activity. For each cell, the FRET integral ratio was calculated as the ratio of the area under the time course of the FRET/ECFP ratio (blue area A) to the area under the maximum ratio (patterned area B) over the whole time course after LIS stimulation. The emission ratio images were shown in the intensity modified display (IMD) mode. In terms of emission ratios quantifying the biosensor activations, we used ECFP/FRET for FAK biosensor, and FRET/ECFP for calcium biosensor. The reason is that the two biosensors are designed with different FRET activation mechanisms. For FAK biosensor, FAK activation causes a decrease in FRET efficiency, while for calcium biosensor, calcium activation causes an increase in FRET efficiency. Therefore, we used the indicated emission ratios described above to represent biosensor activations, so that an increase in the emission ratio indicates an activity increase in the target molecule.

### Focal adhesion analysis

The detection method of FA locations based on the GFP fluorescence intensity (FI) images was as previously described (Lu *et al.*, 2014). In brief, the image analysis for focal adhesion was conducted using our customized software fluocell developed in MATLAB. The source code of *fluocell* is published at Github (<https://github.com/lu6007/fluocell>). All the images were background-subtracted and smoothed using a median-filter with a window size of  $3 \times 3$  pixels. The GFP-

paxillin images were used to compute the masks of the cells. The cell masks were then divided into four layers with the outer layer representing the lamellipodium region. After that, a fan-shaped region was selected in the first image of the video sequence to identify the location of active lamellipodia (Appendix Fig S6A and B). The interception of this fan region and the outer layer of the cell mask were utilized for all the images in the sequence to represent the active lamellipodium region of interest (ROI). As such, the paxillin-GFP images were processed with high-pass filter which removed diffusive cytosolic background (Appendix Fig S6B) and a threshold value was used to detect paxillin-labeled focal adhesions in the filtered image. Total FI of the detected paxillin focal adhesions were quantified in the ROI over time and normalized such that the average value before treatment was 1. The normalized time courses of total FI were then collected for further analysis and statistical comparison.

### Statistical analysis

Sample sizes were chosen on the basis of previous literature in this field. Before plasmid transfection, cell manipulation, and drug treatment, the cells were randomized. Investigators were not blinded to group allocation during data collection and analysis. The sample size, statistical significance value, and error bar graphs were listed in figure legends. Pairwise comparisons were performed using a two-tailed unpaired Student's *t*-test. A *P*-value  $< 0.05$  was considered statistically significant. Comparisons between more than two groups were performed using one-way ANOVA followed by Tukey's multiple comparisons test to compare individual means as indicated in the figure legends.

### Data availability

This study includes no data deposited in external repositories.

**Expanded View** for this article is available online.

### Acknowledgments

The authors thank Dr. Ardem Patapoutian for his generosity in providing materials and advice. This work is supported by grants from NIH GM140929, EB029122, HD107206 (Y. Wang), and HL121365 (Y. Wang and SC), NSF CBET1360341 (Y. Wang and SL), the Beckman Laser Institute Foundation, AFOSR FA9550-08-1-0284, Air Force Office of Scientific Research under award number FA9550-17-1-0193, Beckman Laser Institute Inc. Foundation (MWB). The funding agencies had no role in study design, data collection and analysis, decision to publish, or preparation of the manuscript.

### Author contributions

**Yijia Pan:** Conceptualization; data curation; formal analysis; validation; visualization; methodology; writing – original draft. **Linda Zhixia Shi:** Resources; data curation; investigation; visualization; methodology; project administration; writing – review and editing. **Chi Woo Yoon:** Resources; formal analysis; validation; investigation; methodology; writing – review and editing. **Daryl Preece:** Resources; data curation; investigation; methodology; writing – review and editing. **Veronica Gomez-Godinez:** Resources; data curation; investigation; methodology. **Shaoying Lu:** Data curation; software; formal analysis; funding acquisition; investigation; visualization; methodology; writing – review and editing. **Christopher Carmona:** Resources; investigation; methodology;

writing – review and editing. **Seung-Hyun Woo:** Resources; investigation; methodology; writing – review and editing. **Shu Chien:** Conceptualization; resources; formal analysis; supervision; funding acquisition; project administration; writing – review and editing. **Michael W Berns:** Conceptualization; resources; supervision; funding acquisition; investigation; project administration; writing – review and editing. **Longwei Liu:** Conceptualization; resources; data curation; formal analysis; supervision; investigation; visualization; methodology; writing – original draft; project administration; writing – review and editing. **Yingxiao Wang:** Conceptualization; resources; data curation; supervision; funding acquisition; methodology; writing – original draft; project administration; writing – review and editing.

In addition to the CRediT author contributions listed above, the contributions in detail are:

YP, SC, MWB, LL, and YW designed research; YP, LL, LZS, DP, CY, CC, and VG performed research; SW contributed new reagents; YP and SL analyzed data; YP, DP, SC, MWB, LL, and YW wrote the paper.

### Disclosure and competing interests statement

Yingxiao Wang and Shaoying Lu are scientific co-founders of Cell E&G Inc and Acoustic Cell Therapy LLC. However, these financial interests do not affect the design, conduct or reporting of this research.

## References

- Akerboom J, Chen T-W, Wardill TJ, Tian L, Marvin JS, Mutlu S, Calderón NC, Esposti F, Borghuis BG, Sun XR *et al* (2012) Optimization of a GCaMP calcium indicator for neural activity imaging. *J Neurosci* 32: 13819–13840
- Bagrientsev SN, Gracheva EO, Gallagher PG (2014) Piezo proteins: regulators of mechanosensation and other cellular processes. *J Biol Chem* 289: 31673–31681
- Berridge MJ (1997) The AM and FM of calcium signalling. *Nature* 386: 759–760
- Cahalan SM, Lukacs V, Ranade SS, Chien S, Bandell M, Patapoutian A (2015) Piezo1 links mechanical forces to red blood cell volume. *Elife* 4: e07370
- Compton JL, Luo JC, Ma H, Botvinick E, Venugopalan V (2014) High-throughput optical screening of cellular mechanotransduction. *Nat Photonics* 8: 710–715
- Coste B, Mathur J, Schmidt M, Earley TJ, Ranade S, Petrus MJ, Dubin AE, Patapoutian A (2010) Piezo1 and Piezo2 are essential components of distinct mechanically activated cation channels. *Science* 330: 55–60
- Coste B, Xiao B, Santos JS, Syeda R, Grandl J, Spencer KS, Kim SE, Schmidt M, Mathur J, Dubin AE *et al* (2012) Piezo proteins are pore-forming subunits of mechanically activated channels. *Nature* 483: 176–181
- Egan P, Sinko R, LeDuc PR, Keten S (2015) The role of mechanics in biological and bio-inspired systems. *Nat Commun* 6: 7418
- Fan Z, Liu H, Mayer M, Deng CX (2012) Spatiotemporally controlled single cell sonoporation. *Proc Natl Acad Sci USA* 109: 16486–16491
- Giannone G, Dubin-Thaler BJ, Dobreiner HG, Kieffer N, Bresnick AR, Sheetz MP (2004) Periodic lamellipodial contractions correlate with rearward Actin waves. *Cell* 116: 431–443
- Giannone G, Ronde P, Gaire M, Haiech J, Takeda K (2002) Calcium oscillations trigger focal adhesion disassembly in human U87 astrocytoma cells. *J Biol Chem* 277: 26364–26371
- Gomez-Godinez V, Preece D, Shi L, Khatibzadeh N, Rosales D, Pan Y, Lei L, Wang Y, Berns MW (2015) Laser-induced shockwave paired with FRET: a method to study cell signaling. *Microsc Res Tech* 78: 195–199
- Guo W, Xu Q (2020) Phosphatase-independent functions of SHP2 and its regulation by small molecule compounds. *J Pharmacol Sci* 144: 139–146
- Inagaki K, Noguchi T, Matozaki T, Horikawa T, Fukunaga K, Tsuda M, Ichihashi M, Kasuga M (2000) Roles for the protein tyrosine phosphatase SHP-2 in cytoskeletal organization, cell adhesion and cell migration revealed by overexpression of a dominant negative mutant. *Oncogene* 19: 75–84
- Jaalouk DE, Lammerding J (2009) Mechanotransduction gone awry. *Nat Rev Mol Cell Biol* 10: 63–73
- Kaczmarek E, Erb L, Koziak K, Jarzyna R, Wink MR, Guckelberger O, Blusztajn JK, Trinkaus-Randall V, Weisman GA, Robson SC (2005) Modulation of endothelial cell migration by extracellular nucleotides: involvement of focal adhesion kinase and phosphatidylinositol 3-kinase-mediated pathways. *Thromb Haemost* 93: 735–742
- Lacroix JJ, Botello-Smith WM, Luo Y (2018) Probing the gating mechanism of the mechanosensitive channel Piezo1 with the small molecule Yoda1. *Nat Commun* 9: 2029
- Li F, Chan CU, Ohl CD (2013) Yield strength of human erythrocyte membranes to impulsive stretching. *Biophys J* 105: 872–879
- Li J, Hou B, Tumova S, Muraki K, Bruns A, Ludlow MJ, Sedo A, Hyman AJ, McKeown L, Young RS *et al* (2014) Piezo1 integration of vascular architecture with physiological force. *Nature* 515: 279–282
- Li C, Rezaia S, Kammerer S, Sokolowski A, Devaney T, Gorischek A, Jahn S, Hackl H, Groschner K, Windpassinger C *et al* (2015) Piezo1 forms mechanosensitive ion channels in the human MCF-7 breast cancer cell line. *Sci Rep* 5: 8364
- Li F, Yang C, Yuan F, Liao D, Li T, Guilak F, Zhong P (2018) Dynamics and mechanisms of intracellular calcium waves elicited by tandem bubble-induced jetting flow. *Proc Natl Acad Sci USA* 115: E353–E362
- Liang X, Boppart SA (2010) Biomechanical properties of *in vivo* human skin from dynamic optical coherence elastography. *IEEE Trans Biomed Eng* 57: 953–959
- Liao D, Li F, Lu D, Zhong P (2019) Activation of Piezo1 mechanosensitive ion channel in HEK293T cells by 30MHz vertically deployed surface acoustic waves. *Biochem Biophys Res Commun* 518: 541–547
- Lokhandwalla M, McAteer JA, Williams JC Jr, Sturtevant B (2001) Mechanical haemolysis in shock wave lithotripsy (SWL): II. *In vitro* cell lysis due to shear. *Phys Med Biol* 46: 1245–1264
- Lokhandwalla M, Sturtevant B (2001) Mechanical haemolysis in shock wave lithotripsy (SWL): I. analysis of cell deformation due to SWL flow-fields. *Phys Med Biol* 46: 413–437
- Lu SY, Seong J, Wang Y, Chang SC, Eichorst JP, Ouyang MX, Li JYS, Chien S, Wang YX (2014) Decipher the dynamic coordination between enzymatic activity and structural modulation at focal adhesions in living cells. *Sci Rep* 4: 5756
- Lu S, Wang Y, Huang H, Pan Y, Chaney EJ, Boppart SA, Ozer H, Strongin AY, Wang Y (2013) Quantitative FRET imaging to visualize the invasiveness of live breast cancer cells. *PLoS One* 8: e58569
- Mañes S, Mira E, Gómez-Mouton C, Zhao ZJ, Lacalle RA, Martínez-A C (1999) Concerted activity of tyrosine phosphatase SHP-2 and focal adhesion kinase in regulation of cell motility. *Mol Cell Biol* 19: 3125–3135
- McHugh BJ, Murdoch A, Haslett C, Sethi T (2012) Loss of the integrin-activating transmembrane protein Fam38A (Piezo1) promotes a switch to a reduced integrin-dependent mode of cell migration. *PLoS One* 7: e40346
- Mitra SK, Hanson DA, Schlaepfer DD (2005) Focal adhesion kinase: in command and control of cell motility. *Nat Rev Mol Cell Biol* 6: 56–68
- Miyamoto T, Mochizuki T, Nakagomi H, Kira S, Watanabe M, Takayama Y, Suzuki Y, Koizumi S, Takeda M, Tominaga M (2014) Functional role for

- Piezo1 in stretch-evoked Ca<sup>2+</sup>(+) influx and ATP release in urothelial cell cultures. *J Biol Chem* 289: 16565–16575
- Noguchi T, Matozaki T, Horita K, Fujioka Y, Kasuga M (1994) Role of SH-PTP2, a protein-tyrosine phosphatase with Src homology 2 domains, in insulin-stimulated Ras activation. *Mol Cell Biol* 14: 6674–6682
- Nourse JL, Pathak MM (2017) How cells channel their stress: interplay between Piezo1 and the cytoskeleton. *Semin Cell Dev Biol* 71: 3–12
- Oakes PW, Beckham Y, Stricker J, Gardel ML (2012) Tension is required but not sufficient for focal adhesion maturation without a stress fiber template. *J Cell Biol* 196: 363–374
- Oh E-S, Gu H, Saxton TM, Timms JF, Hausdorff S, Frevert EU, Kahn BB, Pawson T, Neel BG, Thomas SM (1999) Regulation of early events in integrin signaling by protein tyrosine phosphatase SHP-2. *Mol Cell Biol* 19: 3205–3215
- Ouwens DM, HMM M, van der GCM Z, Stein Gerlach M, Ullrich A, Maassen AJ (1996) Insulin-induced tyrosine dephosphorylation of paxillin and focal adhesion kinase requires active phosphotyrosine phosphatase 1D. *Biochem J* 318: 609–614
- Palmer AE, Tsien RY (2006) Measuring calcium signaling using genetically targetable fluorescent indicators. *Nat Protoc* 1: 1057–1065
- Park Y, Best CA, Badizadegan K, Dasari RR, Feld MS, Kuriabova T, Henle ML, Levine AJ, Popescu G (2010) Measurement of red blood cell mechanics during morphological changes. *Proc Natl Acad Sci USA* 107: 6731–6736
- Quinto-Su PA, Kuss C, Preiser PR, Ohi CD (2011) Red blood cell rheology using single controlled laser-induced cavitation bubbles. *Lab Chip* 11: 672–678
- Ranade SS, Syeda R, Patapoutian A (2015) Mechanically activated ion channels. *Neuron* 87: 1162–1179
- Rau KR, Guerra A, Vogel A, Venugopalan V (2004) Investigation of laser-induced cell lysis using time-resolved imaging. *Appl Phys Lett* 84: 2940–2942
- Rau KR, Quinto-Su PA, Hellman AN, Venugopalan V (2006) Pulsed laser microbeam-induced cell lysis: time-resolved imaging and analysis of hydrodynamic effects. *Biophys J* 91: 317–329
- Rusanescu G, Yang W, Bai A, Neel BG, Feig LA (2005) Tyrosine phosphatase SHP-2 is a mediator of activity-dependent neuronal excitotoxicity. *EMBO J* 24: 305–314
- Schaller MD (2001) Paxillin: a focal adhesion-associated adaptor protein. *Oncogene* 20: 6459–6472
- Scheuss V, Yasuda R, Sobczyk A, Svoboda K (2006) Nonlinear [Ca<sup>2+</sup>] signaling in dendrites and spines caused by activity-dependent depression of Ca<sup>2+</sup> extrusion. *J Neurosci* 26: 8183–8194
- Seong J, Ouyang M, Kim T, Sun J, Wen PC, Lu S, Zhuo Y, Llewellyn NM, Schlaepfer DD, Guan JL et al (2011) Detection of focal adhesion kinase activation at membrane microdomains by fluorescence resonance energy transfer. *Nat Commun* 2: 406
- Shannon EK, Stevens A, Edrington W, Zhao Y, Jayasinghe AK, Page-McCaw A, Hutson MS (2017) Multiple mechanisms drive calcium signal dynamics around laser-induced epithelial wounds. *Biophys J* 113: 1623–1635
- Steinhauser MO, Schmidt M (2014) Destruction of cancer cells by laser-induced shock waves: recent developments in experimental treatments and multiscale computer simulations. *Soft Matter* 10: 4778–4788
- Suhr D, Brummer F, Irmer U, Hulser DF (1996) Disturbance of cellular calcium homeostasis by *in vitro* application of shock waves. *Ultrasound Med Biol* 22: 671–679
- Sun J, Lei L, Tsai CM, Wang Y, Shi Y, Ouyang M, Lu S, Seong J, Kim TJ, Wang P et al (2017) Engineered proteins with sensing and activating modules for automated reprogramming of cellular functions. *Nat Commun* 8: 477
- Syeda R, Xu J, Dubin AE, Coste B, Mathur J, Huynh T, Matzen J, Lao J, Tully DC, Engels IH et al (2015) Chemical activation of the mechanotransduction channel Piezo1. *Elife* 4: e07369
- Tsukamoto A, Higashiyama S, Yoshida K, Watanabe Y, Furukawa KS, Ushida T (2011) Stable cavitation induces increased cytoplasmic calcium in L929 fibroblasts exposed to 1-MHz pulsed ultrasound. *Ultrasonics* 51: 982–990
- Vogel V, Sheetz M (2006) Local force and geometry sensing regulate cell functions. *Nat Rev Mol Cell Biol* 7: 265–275
- Wei Z, Lin B-J, Chen T-W, Daie K, Svoboda K, Druckmann S (2020) A comparison of neuronal population dynamics measured with calcium imaging and electrophysiology. *PLoS Comput Biol* 16: e1008198
- Wolfenson H, Iskratsch T, Sheetz MP (2014) Early events in cell spreading as a model for quantitative analysis of biomechanical events. *Biophys J* 107: 2508–2514
- Yamauchi K, Milarski KL, Saltiel AR, Pessin JE (1995) Protein-tyrosine-phosphatase SHPTP2 is a required positive effector for insulin downstream signaling. *Proc Natl Acad Sci* 92: 664–668
- Yang XN, Lu YP, Liu JJ, Huang JK, Liu YP, Xiao CX, Jazag A, Ren JL, Guleng B (2014) Piezo1 is as a novel trefoil factor family 1 binding protein that promotes gastric cancer cell mobility *in vitro*. *Dig Dis Sci* 59: 1428–1435
- Yu J, Deng R, Zhu HH, Zhang SS, Zhu C, Montminy M, Davis R, Feng GS (2013) Modulation of fatty acid synthase degradation by concerted action of p38 MAP kinase, E3 ligase COP1, and SH2-tyrosine phosphatase Shp2. *J Biol Chem* 288: 3823–3830
- Yu CH, Law JB, Suryana M, Low HY, Sheetz MP (2011) Early integrin binding to Arg-Gly-asp peptide activates Actin polymerization and contractile movement that stimulates outward translocation. *Proc Natl Acad Sci USA* 108: 20585–20590
- Yu D-H, Qu C-K, Henegariu O, Lu X, Feng G-S (1998) Protein-tyrosine phosphatase Shp-2 regulates cell spreading, migration, and focal adhesion \*. *J Biol Chem* 273: 21125–21131
- Yuan X, Bu H, Zhou J, Yang C-Y, Zhang H (2020) Recent advances of SHP2 inhibitors in cancer therapy: current development and clinical application. *J Med Chem* 63: 11368–11396
- Yuan F, Yang C, Zhong P (2015) Cell membrane deformation and bioeffects produced by tandem bubble-induced jetting flow. *Proc Natl Acad Sci USA* 112: E7039–E7047