UC Irvine

UC Irvine Previously Published Works

Title

3T hippocampal glutamate-glutamine complex reflects verbal memory decline in aging.

Permalink

<https://escholarship.org/uc/item/0472x9qm>

Authors

Nikolova, Simona
Stark, Shauna M
Stark, Craig EL

Publication Date

2017-06-01

DOI

10.1016/j.neurobiolaging.2017.01.026

Copyright Information

This work is made available under the terms of a Creative Commons Attribution License, available at <https://creativecommons.org/licenses/by/4.0/>

Peer reviewed

3T hippocampal glutamate-glutamine complex reflects verbal memory decline in aging

Simona Nikolova^{a,b}, Shauna M. Stark^{a,b}, Craig E.L. Stark^{a,b,*}

^a Department of Neurobiology and Behavior, University of California, Irvine, CA, USA

^b Center for the Neurobiology of Learning and Memory, University of California, Irvine, CA, USA

ARTICLE INFO

Article history:

Received 3 October 2016

Received in revised form 5 December 2016

Accepted 6 January 2017

Available online 18 March 2017

Keywords:

3T MRI

Aging

MRS

Memory

Imaging

Glx

ABSTRACT

The hippocampus is a critical site for alterations that are responsible for age-related changes in memory. Here, we present a relatively novel approach of examining the relationship between memory performance and glutamate-glutamine levels using short echo time magnetic resonance spectroscopy. Specifically, we investigated the relationship between Glx (a composite of glutamate and glutamine) levels in the hippocampus, performance on a word-recall task, and resting-state functional connectivity. While there was no overall difference in Glx intensity between young and aging adults, we identified a positive correlation between delayed word-list recall and Glx, bilaterally in older adults, but not in young adults. Collapsed across age, we also discovered a negative relationship between Glx intensity and resting-state functional connectivity between the anterior hippocampus and regions in the subcallosal gyrus. These findings demonstrate the possible utility of Glx in identifying age-related changes in the brain and behavior and provide encouragement that magnetic resonance spectroscopy can be useful in predicting age-related decline before any physical abnormalities are present.

© 2017 Elsevier Inc. All rights reserved.

1. Introduction

One of the key concerns of older adults is the experience of memory loss, both in the normal course of aging and as it is associated with neural degenerative disease. Many of the types of memory that demonstrate age-related decline are mediated by the hippocampus and surrounding medial temporal lobe structures (Morrison and Baxter, 2012; Small, 2001; Yassa et al., 2011a). While there are many laboratory tasks that are designed to tax-specific aspects of hippocampal function, standardized neuropsychological tests are broadly available and extensively used in both research and clinical settings. The Rey Auditory Verbal Memory Task (RAVLT; Rey, 1941) has been shown to be highly effective at detecting age-related memory decline (Mitrushina et al., 1991; Schoenberg et al., 2006). In this task, memory is assessed based on the acquisition of a list of 15 unrelated words, followed by free recall, and repeated over the course of 5 presentations. Participants are then given an interference list of 15 new words, followed by free recall of the original list and another

free recall 15 minutes later. Age-related decline has been observed on all recall trials (Mitrushina et al., 1991), with performance on delayed recall correlating with functional scales of memory in those with subjective memory complaints (Estevez-Gonzalez et al., 2003). It is currently used by a number of research groups to identify healthy agers who are particularly “successful” (Mapstone et al., 2014), “super” (Rogalski et al., 2013), or “unimpaired” (Stark et al., 2013) in their memory performance relative to other, healthy participants. Likewise, a delayed word-recall performance on a related task is particularly sensitive to early decline associated with Alzheimer’s disease (Albert, 1996).

Although the aging brain has been associated with declines in hippocampal volume (Raz et al., 2005), there is strong evidence that this neuronal loss is not simply due to fewer neurons (Burke and Barnes, 2006). Instead, there are evidence of synaptic alterations (Nicholson et al., 2004) and decreases in gene expression (Berchtold et al., 2013) in aging. Likewise, there are neuromodulatory changes in the hippocampus, including decreased cholinergic input to the hippocampus (Sugaya et al., 1998) and a reduction in dopamine that correlates with memory performance (Stemmelin et al., 2000). While these findings have been reported in rodents, we now have the technology to investigate some of these neurotransmitters in humans using magnetic resonance spectroscopy (MRS). Instead of looking at MRS related to disease, the goal of this work is to link it to behavioral performance.

* Corresponding author at: Center for Neurobiology of Learning and Memory, Department of Neurobiology and Behavior, Francisco J. Ayala School of Biological Sciences, University of California, Irvine, CA, USA. Tel.: (949) 824-4201; fax: (949) 824-8439.

E-mail address: cestark@uci.edu (C.E.L. Stark).

MRS quantifies the concentration of various metabolites (intermediary and byproducts of metabolism) in the brain, with many of these accessible via conventional proton (^1H) imaging. Numerous studies have utilized the efficiency of MRS in providing biological information on cellular/metabolic changes in the aging brain or diseased brain (see Wang et al., 2015 for a review). Specifically, our focus is on Glutamate (Glu), the most abundant metabolite in the brain, a dominant excitatory neurotransmitter. Glutamate is released during neuronal excitation and converted to Glutamine (Gln) with the Glu-Gln cycle requiring high energy demands. Alterations in concentrations of Glu and Gln have been reported in numerous neurological and psychiatric diseases, such as depression and mood disorders, epilepsy, alcohol and drug abuse, schizophrenia, and neurodegenerative disorders (see Ramadan et al., 2013 for a review).

MRS studies have shown reduced Glu levels in a mouse model of Alzheimer's disease (AD; Chen et al., 2012b) and in humans with AD (Rupsingh et al., 2011). They have also shown a decrease in combined Glu and Gln (referred to as Glx) in the cingulate cortices and posterior cingulate gyrus of AD patients (Antuono et al., 2001; Hattori et al., 2002). Little is known about any such alterations in healthy aging or in the hippocampus per se, despite the clear potential for age-related changes in hippocampal activity (Miller et al., 2008; Wilson et al., 2006; Yassa et al., 2011b). In one report, Wagner et al. (2016) investigated the relationship between hippocampal Glx, functional connectivity with the hippocampus, and verbal memory performance (on a different task) in healthy young males. They reported that lower Glx levels were associated with higher functional connectivity of the hippocampus to prefrontal cortex and anterior cingulate. In addition, they found that Glx concentration in the posterior hippocampus predicted verbal memory performance. These relationships emphasize the utility of MRS as a functional measure of neuronal activity, related to both behavior and brain connectivity.

Given the relationship of Glx to aging and AD, and the possible link to memory performance, the purpose of this study was to investigate age-related changes in Glx levels in the hippocampus using short echo time MRS in young and older adults. We hypothesized that Glx levels would correlate with age-related decline on the RAVLT. Following the findings of Wagner et al. (2016), we also investigated the relationship between Glx and resting-state functional connectivity (rs-fMRI) in a subset of our sample for which we were fortunate enough to also have rs-fMRI scans. We hypothesized that we would replicate the negative relationship found between the resting-state functional connectivity (RSFC) anterior hippocampus to regions in the prefrontal cortex and Glx levels in the right anterior hippocampus. These findings would significantly extend those of Wagner et al. (2016) by showing the behavioral relevance of Glx in the hippocampus to healthy aging and link them to the highly-used RAVLT.

2. Materials and methods

2.1. Participant recruitment

Forty-one participants, consisting of 21 older (ages 59–84) and 20 younger adults (ages 20–38) were recruited for this study from the community surrounding the University of California in Irvine (UCI). Two participants were removed from the analysis: 1 young adult was excluded because one of their MRS voxels contained only 20% overlap with the hippocampus (motion between scan preparation and acquisition) and one older adult was excluded for RAVLT scores that fell outside the age-thresholded norms. Thus, we analyzed data from 20 older (8 males, 12 females, mean age of 70 ± 6 years) and 19 younger (8 males, 11 females, mean age of $27 \pm$

5 years) adults. All participants were right-handed, had no history of neurological disease or psychiatric illness, and had normal or corrected-to-normal vision. In addition, they all performed within the normal range for their age on standardized neuropsychological testing and scored in the normal to mild range on the Geriatric Depression Scale. All participants gave informed consent approved by the UCI Internal Review Board before participation and were compensated for their time.

Verbal memory was evaluated using the Rey Auditory Verbal Learning Test (RAVLT; Rey, 1941). The RAVLT involved learning a list of 15 unrelated words, repeated 5 times, with a recall test after each administration of the list. Following the 5 presentations, a new interferences list of 15 words was administered, along with a recall task. After the interference list, participants were then asked to recall the original list for an immediate recall test and assessed again 15-minutes later with a delayed recall test.

2.2. MRS data acquisition

MRS scans were collected using a 3.0T Philips whole-body MRI (Best, the Netherlands), using a 32-channel head coil. High-resolution T1-weighted and T2-weighted images were acquired parallel to the hippocampus.

T1-weighted whole-brain anatomical images were acquired using a sagittal magnetization-prepared rapid gradient echo scan (MP-RAGE) at a resolution of 0.75 mm (isotropic), repetition time (TR)/echo time (TE) = 11/4.6 ms, field of view = 320×264 , and flip angle = 18° . In addition, a T2-weighted whole-brain anatomical (TR/TE = 8/4.6 seconds) scan was used as an anatomical reference to localize the hippocampal voxel placement for each participant. The rectangular voxel of interest was placed along the long axis of the hippocampus on the sagittal T2 image and contained mainly the anterior hippocampus (Fig. 1).

^1H spectroscopy data were acquired using PRESS (TR/TE = 2000/37 ms), from a nominal $2.5 \times 1.5 \times 1.0 \text{ cm}^3$ voxel in the left and right hippocampi of all the subjects. The acquisition of MRS spectra took about 8 minutes of total MR time of 60 minutes per subject. Voxel dimensions were chosen to maximize the volume while largely contained within the anterior hippocampus of each subject. Hippocampal volumes were calculated using Freesurfer (see MRS Data Analysis section). For the right hippocampus, the values ranged between 3.0 cm^3 to 3.75 cm^3 (mean and SD $3.3 \pm 0.6 \text{ cm}^3$ in aged subjects and $3.70 \pm 0.2 \text{ cm}^3$ for young participants). For the left hippocampi, the volumes varied between 2.4 cm^3 to 4.06 cm^3 (mean and SD $3.3 \pm 0.6 \text{ cm}^3$ in aged subjects and $3.1 \pm 0.6 \text{ cm}^3$ for young participants). Shimming was accomplished using the iterative VOI method provided by Philips. Only shims below 10 Hz were included in the study. MRS data for 5 participants were excluded because either of the linewidths (left or right hippocampus) exceeded 10 Hz.

2.3. Rs-fMRI data acquisition

We were fortunate to have resting-state data from 22 participants (13 young and 9 older adults), collected on a separate day. Echo-planar images were collected on a separate day from the MRS scans using the same scanning equipment. Whole-brain resting-state data were acquired with a T_2^* weighted echo-planar imaging scan, at a resolution of 2.5 mm (isotropic), 46 axial slices, TE = 26 ms, flip angle = 70° , TR = 2500 ms, dynamics = 188. The resting-state scan took 7 minutes 27 seconds, during that time the participants were instructed to relax with their eyes open. Using AFNI (<http://afni.nimh.nih.gov/afni>; Analysis of Functional NeuroImages). The data were aligned to the participant's MP-RAGE with the script align_epi_anat.py (Saad et al., 2009). Each participant's structural

Fig. 1. Representative spectra and voxel locations from 1 participant. Left hippocampus voxel is shown on axial, sagittal, and coronal view (yellow). The right hippocampus MRS voxel is shown in red. Abbreviations: A, anterior; L, left; P, posterior; R, right. (For interpretation of the references to color in this figure legend, the reader is referred to the Web version of this article.)

scan and functional data (statistical maps) were aligned to a model template using Advanced Normalization Tools (Avants et al., 2008).

2.4. MRS data analysis

Quantitative MRS data were initially analyzed with TARQUIN v.4.3.6 (Wilson et al., 2011), using a non-negative least squares projection to estimate signal amplitudes. Initial point truncation and HSVD water removal was used to reduce baseline interferences (Wilson et al., 2011).

The fractional contributions of gray matter (GM), white matter (WM), and cerebrospinal fluid (CSF) were determined for each voxel by segmenting the T_1 -weighted images using SPM8 (Ashburner and Friston, 1997; SPM - Statistical Parametric Mapping, 2016) and MATLAB (the Mathworks, INC. Natick, MA, USA). The fractional tissue contributions were used to correct the estimated metabolite concentrations for partial volume effects (Gussew et al., 2012) using a customized MATLAB code. The partial volume correction required relaxation rates (T_1 and T_2) of water and metabolites in both WM and GM (Stanisz et al., 2005).

Freesurfer was used to segment the T_1 -weighted images into hippocampal volumes. The percent of each voxel that overlapped with the hippocampus was calculated and reported in Table 1. The percent of MRS voxel GM consisting of hippocampal tissue was recorded as well. The mean MNI coordinates of each voxel were determined first by warping each anatomical T_1 to MNI152 template using Advanced Normalization Tools (Avants et al., 2008), and subsequently by applying the warp transform to each voxels 3D binary mask (1 mm isotropic resolution). The mean MNI coordinates were calculated from the warped binary mask in MNI space. Hippocampal volumes and ventricular volumes were also calculated on the MP-RAGE for each participant using Freesurfer's automatic software for volumetric measures (Fischl et al., 2002; FreeSurfer, 2016). These volumes were normalized by dividing by total intracranial volume for each participant.

2.5. Rs-fMRI data analysis

Data were preprocessed using AFNI and included detrended using 2nd-order Legendre polynomials. The time series was then normalized to have zero mean and unit variance and the motion vectors and first derivatives were regressed out of the signal. The first 6 principal components of the variance were computed using CompCor (not including global signal) and regressed out of the signal (Behzadi et al., 2007). The data were then temporally band-pass filtered (0.009–0.08 Hz) and TRs with framewise displacement >0.5 mm, as well as those 1 TR before and 2 TRs after, were removed from analyses by censoring as recommended by Power et al. (2012). Thus, the remaining data for both the young and older adults contained similar, very low levels of motion. Note, all rs-fMRI analyses to identify regions showing reliable connectivity to the hippocampus were done on the entire set of participants, regardless of age.

We estimated the inherent blur using 3dFWHMx (AFNI) for each participant and used the mean of these values in 3dClustSim¹ (AFNI: June 2016) to determine the combined voxel-wise ($p < 0.005$) and cluster-size (nearest-neighbor connectivity of 10 contiguous voxels) thresholds needed to arrive at a multiple-comparison corrected alpha of $p < 0.05$.

2.6. RAVLT data analysis

Free recall from the 5 acquisition tests in the RAVLT was summed to produce a RAVLT total score. In addition, the RAVLT interference score was calculated (total recalled from the interference word list administered after these). Finally, we also scored the total recalled

¹ 3dClustSim was updated in May 2015 to address an error in underestimating multiple corrections, resulting in an overestimation of significance for a given cluster (Eklund et al., 2016). Here, we have used the updated version of 3dClustSim, which corrected this error.

Table 1

Age-group contrasts for RAVLT performance, Glx values, brain volumes with values reported as mean (standard deviation)

	Young	Aging	Independent <i>t</i> test
Age (y)	27 (5)	70 (6)	
Sex (M, F)	8, 11	8, 12	
Behavior			
RAVLT total	63 (1.3)	56 (1.8)	$t(37) = 3.14, p < 0.01$
RAVLT immediate	13.9 (0.3)	12.4 (0.5)	$t(38) = 2.62, p < 0.05$
RAVLT delay	14.1 (0.3)	12.1 (0.6)	$t(38) = 3.01, p < 0.01$
RAVLT interference	7.1 (2.0)	5.6 (2.0)	$t(37) = 2.30, p < 0.05$
Metabolites			
Left Glx	18.4 (4.8)	16.7 (7.4)	$t(37) = 0.67, p = 0.51$
Right Glx	14.9 (5.6)	14.0 (4.0)	$t(37) = 0.63, p = 0.53$
Volumes			
Left hippocampus	0.28% (0.02%)	0.25% (0.03%)	$t(37) = 3.15, p < 0.01$
Right hippocampus	0.27% (0.02%)	0.24% (0.04%)	$t(37) = 2.71, p < 0.05$
Ventricular volume	1.0% (0.3%)	2.1% (1.1%)	$t(37) = 4.54, p < 0.01$
Left GM fraction	70% (3%)	71% (4%)	$t(37) = 0.99, p = 0.32$
Right GM fraction	69% (4%)	69% (4%)	$t(37) = 0.71, p = 0.48$
Left WM fraction	22% (4%)	22% (4%)	$t(37) = 0.32, p = 0.75$
Right WM fraction	26% (4%)	25% (4%)	$t(37) = 0.37, p = 0.71$
Left CSF fraction	8% (2%)	7% (2%)	$t(37) = 1.11, p = 0.27$
Right CSF fraction	6% (2%)	5% (2%)	$t(37) = 0.61, p = 0.55$
Left voxel volume [c.c.]	3.7 (0.2)	3.1 (0.6)	$t(37) = 3.48, p < 0.01$
Right voxel volume [c.c.]	3.7 (0.2)	3.2 (0.6)	$t(37) = 3.05, p < 0.01$
% of left voxel = hipp	59% (6%)	61% (8%)	$t(37) = 1.16, p = 0.25$
% of right voxel = hipp	58% (4%)	59% (8%)	$t(37) = 0.32, p = 0.75$
Left %GM = hipp	84% (9%)	86% (8%)	$t(37) = 0.77, p = 0.44$
Right %GM = hipp	85% (6%)	84% (9%)	$t(37) = 0.24, p = 0.81$
Avg. MNI left voxel [X]	−26 (1)	−27 (1)	
Avg. MNI left voxel [Y]	−20 (4)	−19 (2)	
Avg. MNI left voxel [Z]	−18 (2)	−20 (6)	
Avg. MNI right voxel [X]	27 (2)	26 (1)	
Avg. MNI right voxel [Y]	−19 (3)	−19 (2)	
Avg. MNI right voxel [Z]	−18 (2)	−19 (2)	

The bold values reflect comparisons between the Young and Aging groups that pass a statistical threshold of *p* values.

Key: CSF, cerebrospinal fluid; Glx, a composite of glutamate and glutamine; GM, gray matter; RAVLT, Rey Auditory Verbal Memory Task; WM, white matter.

from the immediate recall and the delayed recall, administered 15 minutes later. We used Prism 6 (www.graphpad.com) to calculate unpaired *t* tests for between group comparisons and correlations between behavior and Glx values.

3. Results

3.1. Age-related differences in RAVLT performance

One older adult performed more than 2 standard deviations below standardized normative values for his age. As this may represent either exceptionally inattentive performance or potential early clinical impairment, this individual was excluded from further analysis. All other young and older participants performed within 1 standard deviation of the standardized norms for their age.

In the resulting data set, we examined the effect of aging on the RAVLT using planned independent samples *t* tests. As expected, we found that older adults performed worse than young adults on all 4 measures of the RAVLT: total acquisition, interference, and immediate and delayed recall (see Table 1).

3.2. No age-related differences in Glx concentration

Tissue fractions and normalized hippocampal and ventricular volumes for both groups are shown in Table 1. There was no evidence for differences in the proportion of GM, WM, and CSF in our MRS voxels across age groups, reducing potential concerns for artefactual differences in metabolite concentrations resulting from varying partial volume effects across groups. Consistent with a wide range of studies, both left and right hippocampal volume, as well as ventricular volume, were lower in the aging group. Not surprisingly, the average volume of the MRS voxels was also smaller in aged than young participants. However, the average Glx, presented in absolute concentrations (mmol/L), did not differ between the two groups. Thus, although hippocampal atrophy was observed in the aging group, we were still able to obtain a robust measure of Glx.

3.3. Age-related differences in MRS and RAVLT performance

To reduce noise and as we had no *a priori* hypotheses concerning laterality, we began investigating the relationship between Glx and memory performance by averaging the left and right Glx values for each participant, correlating the result with RAVLT performance. We found significant positive correlations between Glx and RAVLT total, immediate, and delayed recall for the aged group (Fig. 2). There was no evidence for such relationship in the young group.

Since MRS was collected from a single, often unilateral voxel, our combination of the 2 hemispheres' Glx values in our initial analysis is somewhat atypical. Thus, as a secondary analysis, we analyzed these relationships separately for each hemisphere to determine if the results were consistent and replicable in each. Similarly, independent correlations were observed in each hemisphere in the aging group between Glx and RAVLT total (left: $r = 0.62, p < 0.01$; right: $r = 0.39, p = 0.09$), immediate (left: $r = 0.54, p < 0.05$; right: $r = 0.52, p < 0.05$), and delayed recall (left: $r = 0.65, p < 0.01$; right: $r = 0.42, p = 0.07$). No relationship between Glx and RAVLT performance was observed in the young group (all *p*'s > 0.5). We note that this lack of a relationship in our anterior hippocampal voxels is consistent with the findings of Wagner et al. (2016). Although they observed a relationship between hippocampal Glx and performance on a test that provides a similar measure of memory as the RAVLT Interference score, this relationship was only found for posterior and not anterior hippocampal voxels.

3.4. Relationship between RSFC and Glx

RSFC analyses were carried out by correlating the regional signal-time courses extracted from the left and right anterior hippocampus (corresponding to the regions where the MRS scans were collected), to all other voxels in the brain. We applied Fisher *z*-transformation to the linear correlation maps before performing statistical analyses. Due to the reduced number of participants in the young and aging groups who had both MRS and rs-fMRI, we collapsed the young and older adults into a single group.

Using 3dRegAna (AFNI), we tested the relationship between hippocampal RSFC (separately for the left and the right anterior hippocampus) and Glx intensity. There were 14 regions that revealed a relationship between left or right RSFC and Glx concentration presented in Table 2 (multiple-comparison corrected). Of these, 2 of the regions (Fig. 3) were located in the left subcallosal gyrus, near the orbitofrontal cortex (OFC) location identified by Wagner et al. (2016). In both of these regions, we replicated the Wagner et al. (2016) finding of a negative relationship between Glx concentration and RSFC from the anterior hippocampus to the OFC. Interestingly, although the Wagner group found this relationship in young adult males, we extended these findings in a group of

Fig. 2. Regression analyses of the average Glx with RAVLT total acquisition, immediate recall, and delay recall memory scores for young and aging participants. The aging group demonstrated a positive relationship between average Glx and each memory measure, yet there is no evidence of this relationship in the young group. Abbreviations: Glx, a composite of glutamate and glutamine; RAVLT, Rey Auditory Verbal Memory Task.

healthy, mixed gender, older, and younger adults. We (post hoc) plotted the young and older adults separately in Fig. 3 to demonstrate the consistency in both groups but, given our small sample size, we did not contrast them directly. In addition to the 2 OFC regions that overlapped with those reported by Wagner et al. (2016), we also identified 12 other regions, displaying a mix of positive and negative relationships between Glx intensity and RSFC.

4. Discussion

In this study, we measured Glx (a combined measure of glutamate and glutamine) in the anterior hippocampus of young and older adults. We found that greater Glx concentration in older

adults was correlated with higher memory performance in a word recall on the RAVLT, both at encoding and retrieval. Moreso, we showed that this effect was observed in both the left and the right hippocampus, providing an internal replication of this relationship. Interestingly, this relationship was not evident in young subjects.

These findings extend those of Wagner et al. (2016), who reported a positive correlation in young males between hippocampal Glx levels and retroactive interference on a task similar to the RAVLT. Our results are largely consistent and complementary here, as the studies contain both overlapping and independent measures. Briefly, Wagner et al. (2016) observed their Glx-verbal memory correlation in posterior but not anterior Glx measures and only in the interference memory measure. In addition, they

Table 2
Regions that show a relationship between Glx intensity and RSFC with the left and right hippocampus as seed regions

Region name	Glx	X	Y	Z	Dir.
Left subcallosal gyrus (24)	Left	−13	19	−12	−
Left subcallosal gyrus (7)	Right	−20	18	−9	−
Left superior frontal gyrus (11)	Right	−4	16	55	−
Right middle frontal gyrus (8)	Left	28	24	38	−
Left medial frontal gyrus (20)	Right	−4	0	54	+
Left postcentral gyrus (3)	Right	−40	−11	32	−
Right angular gyrus (36)	Left	37	−58	31	−
Left cuneus (19)	Right	−4	−77	17	−
Left inferior parietal lobe (5)	Right	−55	−35	26	−
Right lingual gyrus (6)	Right	6	−62	−3	−
Left precuneus (5)	Left	−37	−74	36	−
Right lingual gyrus (1)	Right	11	−86	−26	+
Right inferior occipital (10)	Right	22	−89	0	+
Right inferior occipital gyrus (15)	Right	32	−82	−3	+
Right supramarginal gyrus (8)	Right	38	−45	24	+
Right precentral gyrus (9)	Right	57	−14	26	−
Right middle temporal gyrus (18)	Right	50	−52	6	−
Right cerebellum (29)	Left	6	−60	−27	+
Right posterior cingulate (34)	Left	13	−54	11	−
Left parahippocampal gyrus (16)	Right	−27	−56	2	−

Region names are defined as the focal point in the Tailairach-Tournoux Atlas. Glx refers to the left or right hippocampal seed that showed RSFC to that region. X (right/left), Y (anterior/posterior), and Z (inferior/superior) coordinates given in Tailairach space (TLRC).
Key: Glx, a composite of glutamate and glutamine; RSFC, resting-state functional connectivity.

only scanned young males. Our data from the anterior hippocampus failed to show a relationship between Glx and verbal memory consistent with their observations. As we do not have MRS data from the posterior hippocampus, we cannot address whether young individuals show a similar relationship with the interference score of the RAVLT. While it is by no means clear, there is growing

evidence for some form of anterior-posterior distinction in the hippocampus (Strange et al., 2014). For example, activity in the posterior hippocampus has been shown to predict the degree of retroactive interference in a paired-associates task (Kuhl et al., 2010), highlighting the relevance of this region for retroactive interference specifically. We collected data from the anterior hippocampus, which has been associated with novelty detection (Daselaar et al., 2006) and a relationship to the dorsal attention network (Kim, 2015). Thus, the anterior hippocampus may be more sensitive to the recall measures of the RAVLT than to retroactive interference. We caution, however, that such interpretations are entirely speculative and based on reverse inferences at this stage.

We did, however, replicate their basic effect by demonstrating a relationship between hippocampal Glx and RAVLT performance. In our study, we found this relationship in the anterior hippocampus in a heterogeneous group of older adults for several of the verbal learning measures. Note, Wagner et al. (2016) did not test older individuals or women. Thus, we view our results as being consistent with and complementary to those of Wagner et al. (2016) in this regard.

Our observation of a relationship between Glx levels and RAVLT performance in aging adults suggests a possible modulatory effect of glutamate and/or glutamine on memory performance. Higher levels of glutamate in the striatum have been associated with better performance on tests of executive functioning in a group of older adults, provided a potential mechanism for age-related decline in the frontostriatal system (Zahr et al., 2008). N-methyl-D-aspartate (NMDA) receptors represent one of the glutamate receptors that are present in high density in the hippocampus and cerebral cortex, even in aging and AD (Cotman et al., 1989), but there is mounting evidence that NMDA receptor function declines with age (Barnes et al., 1997). Likewise, an age-related deficiency in NMDA receptor function contributes to impairments in spatial learning and

Fig. 3. Two regions in the left subcallosal gyrus replicate the negative relationship observed by Wagner et al. (2016) between Glx intensity and RSFC to the right and left anterior hippocampus. The young and aging groups are plotted separately here but were analyzed as a single group, irrespective of age. Abbreviations: Glx, a composite of glutamate and glutamine; RSFC, resting-state functional connectivity.

memory in older rodents (Magnusson, 1998; Zhao et al., 2009), providing a mechanism for glutamate-related dysfunction in aging that may be contributing to a decline in verbal memory.

As noted earlier, in our current data, we are unable to separate the Glu and Gln components of the signal and, like many studies, must combine this into a Glx value. Glu and Gln are complementary, such that Glu is taken up in the synaptic cleft and then converted to Gln within the astrocytes. This cycle is rapid and highly dynamic, with Glu located primarily in neurons and Gln located predominantly in the glial cells (Ottersen et al., 1992). Although we found no age-related decrease in Glx concentration between young and older adults in the hippocampus, other studies have found evidence for an age-related decline in Glu, accompanied by an age-related increase in Gln in the hippocampus (Hadel et al., 2013; Schubert et al., 2004). These opposing effects may have mitigated any age-related difference in overall Glx concentration here, but an underlying age-related imbalance may still account for the relationship with RAVLT performance. Outside of the hippocampus, age-related decreases in glutamate have been reported in the striatum (Zahr et al., 2008; but see Choi et al. (2014) for the opposite pattern), but not the cerebellum (Zahr et al., 2008, 2013), emphasizing the possible regional specificity of these metabolic imbalances. In addition, the opposing dynamics of the Glu/Gln cycle may be completely masked by the combined Glx metric. Future studies using acquisition techniques that can isolate Glu and Gln (e.g., Mega-PRESS sequences) are needed to identify the opposing contributions of these 2 metabolites.

The relationship between Glu and Gln concentrations in aging is a complicated one. These compounds reflect both neuronal and mitochondrial energy production, with Glu serving as a metabolic precursor of γ -aminobutyric acid (GABA), the main inhibitory neurotransmitter in the cerebral cortex. Mitochondrial dysfunction in normal aging may lead to an increase in Glu (Boumezbeur et al., 2009; Lin and Beal, 2006) and be a major contributor to the underlying cause of AD (Lin et al., 2003). Further, ameliorating this mitochondrial dysfunction has been shown to improve cognitive performance in rodent models of AD (Chen et al., 2012a). Increased in hippocampal Glu levels have also been observed in a number of disorders, including medial temporal lobe epilepsy (Simister et al., 2002; Woermann et al., 1999) and schizophrenia (Kraguljac et al., 2014). Based on these findings, we might have predicted that greater Glu levels would be associated with worse memory performance on the RAVLT in older adults. In contrast, we found that for healthy older adults, greater Glx concentration is associated with greater memory performance. One might speculate that higher Glx values reflect greater regulation of the Glu/Gln cycle, since the combined Glx measure does not allow us to analyze the individual contributions of Glu and Gln to memory performance. Yet, any speculation here must be tempered by the fact that overall Glx concentration did not differ between the older and young adults, therefore the relationship between Glx and memory performance in the older adults may not reflect a global shift in Glu associated with clinical conditions such as schizophrenia, epilepsy, and AD. Notably, in these diseased states, the decrease of Glu is consistently accompanied by a decrease in NAA (Barth et al., 2015; Jessen et al., 2001; Rupasingh et al., 2011), signifying neuronal loss or dysfunction. In this study, a decrease in NAA was not observed or expected as healthy aging is not associated with overt neuronal loss in the hippocampus (Burke and Barnes, 2006).

In this study, we also replicated another finding reported by Wagner et al. (2016) of a negative relationship between hippocampal Glx in the anterior hippocampus and the RSFC from the anterior hippocampus to regions in the OFC. The OFC is one of the regions of the default mode network (DMN), a network of highly interconnected brain structures that show synchronous

low-frequency fluctuations during a resting state condition (Raichle et al., 2001). Previous correlations between Glx and nodes of the DMN have been observed, particularly with increased glutamate levels in the anterior cingulate cortex (Enzi et al., 2012) and the posterior cingulate cortex (Duncan et al., 2013; Hu et al., 2013). Thus, the inter-regional synchrony in the DMN may be modulated, at least in part, by the excitation-inhibition balances based on the ratio of Glu and Gln.

In contrast to the findings of Wagner et al. (2016), it is worth noting that we also found both negative and positive relationships between Glx and RSFC between the anterior hippocampus and other regions outside of the DMN, including regions in the parahippocampal gyrus, parietal cortex, and cerebellum. Only a few studies have investigated the relationship between glutamate levels and task-induced BOLD signal or RSFC in humans. Using MRS in the dorsal anterior cingulate cortex, Falkenberg et al. (2012) categorized young healthy adults into high- and low-glutamate individuals and showed that those with low-glutamate levels showed an increased BOLD response during a task that demanded low cognitive control, whereas individuals with high-Glu levels showed the opposite effect. Similarly, a positive relationship between RSFC and Glu levels in the anterior cingulate cortex has been shown in young adults. These findings emphasize an interaction between Glu levels and the activation levels of these regions, though they have been limited to very specific regions of interest.

Glu, as an excitatory neurotransmitter, may result in high levels of RSFC in regions that are task activated, whereas a reduction in this glutamatergic activity, or a counteraction of it by an inhibitory neurotransmitter such as GABA, may lead to task-induced deactivation. These relationships are highly dependent on the task and the regions involved in the task, but reflect the utility of these metabolites as functional measures of neuronal activity. In this study, the rs-fMRI data were collected during “rest”, which is a cognitively uncontrolled condition that results in high BOLD activity in the hippocampus (Stark and Squire, 2001). The complex relationship between how glutamate levels in the hippocampus modulates rest-based RSFC is beyond the scope of this investigation. Instead, we report these relationships to (1) report on a replication with Wagner et al. (2016); (2) demonstrate a relationship between Glx and RSFC across many regions; and (3) reveal the complexity of this relationship across regions. With a larger sample size or multiple MRS voxels, it would be interesting for future studies to investigate any age-related modulation of these relationships.

This study was limited by a small sample size for the rs-fMRI data set, which prevented us from examining any age-related differences in the relationship between Glx and resting state activity. We were also limited in our ability to contrast Glx and network relationships because we collected MRS from only the anterior hippocampus. In future studies, collecting MRS voxels from the anterior and posterior hippocampus, or from other memory-related regions such as the parahippocampal gyrus, would provide an opportunity to compare the relationship between Glx and functional activity. Finally, since Glx is a composite measure, we are unable to distinguish the individual contributions of glutamate and glutamine to the behavior or the brain activity. The use of a MEGA-PRESS sequence may be able to differentiate these 2 metabolites in the future. GABAergic inhibition plays a key role in glutamatergic excitation (McCormic et al., 1989), and the balance of GABA and Glu is essential to consider with respect to hippocampal function and memory. In MRS, the Glu and GABA peaks however overlap, making it difficult to separate those metabolites without the use of J-editing spectral techniques. We were unable to investigate the effects of GABA here since special sequences are needed to resolve this peak but would like to address the role of GABA in future work.

5. Conclusions

We report that decreases in Glx in the anterior hippocampus were accompanied by decreased performance in verbal memory in aged adults. These preliminary findings suggest a potential age-related modulatory role of Glu and/or Glu in the hippocampal regulation of memory. Future studies are needed to focus on how a dysregulation of the excitation-inhibition balance in the hippocampus may contribute to age-related memory decline.

Disclosure statement

The authors have no conflicts of interest to disclose.

Acknowledgements

The authors would like to thank Samantha Rutledge for her assistance with data collection and Derek Huffman and Veronique Boucquey for assistance with preprocessing the resting state functional connectivity data. Funding for this project was provided in part by a grant from the National Institutes on Aging R01-AG034613 and R21-AG053040.

References

- Albert, M.S., 1996. Cognitive and neurobiologic markers of early Alzheimer's disease. *Proc. Natl. Acad. Sci. U. S. A.* 93, 13547–13551.
- Antuono, P.G., Jones, J.L., Wang, Y., Li, S.J., 2001. Decreased glutamate + glutamine in Alzheimer's Disease detected in vivo with (1)H-MRS at 0.5 T. *Neurology* 2001, 737–742.
- Ashburner, J., Friston, K., 1997. Multimodal image coregistration and partitioning: a unified framework. *NeuroImage* 6, 209–217.
- Avants, B.B., Epstein, C.L., Grossman, M., Gee, J.C., 2008. Symmetric diffeomorphic image registration with cross-correlation: evaluating automated labeling of elderly and neurodegenerative brain. *Med. Image Anal.* 12, 26–41.
- Barnes, C.A., Rao, G., Shen, J., 1997. Age-related decrease in the N-methyl-D-aspartate-mediated excitatory postsynaptic potential in the hippocampal region CA1. *Neurobiol. Aging* 18, 445–452.
- Barth, C., Villringer, A., Sacher, J., 2015. Sex hormones affect neurotransmitters and shape the adult female brain during hormonal transition periods. *Front. Neurosci.* 9, 37.
- Behzadi, Y., Restom, K., Liu, J., Liu, T.T., 2007. A component based noise correction method (CompCor) for BOLD and perfusion based fMRI. *Neuroimage* 37, 90–101.
- Berchtold, N.C., Coleman, P.D., Cribbs, D.H., Rogers, J., Gillen, D.L., Cotman, C.W., 2013. Synaptic genes are extensively downregulated across multiple brain regions in normal human aging and Alzheimer's disease. *Neurobiol. Aging* 34, 1653–1661.
- Boumezeur, F., Mason, G.F., de Graaf, R.A., Behar, K.L., Cline, G.W., Shulman, G.I., Rothman, D.L., Peterson, K.F., 2009. Altered brain mitochondrial metabolism in healthy aging as assessed by in vivo magnetic resonance spectroscopy. *J. Cereb. Blood Flow Metab.* 30, 211–221.
- Burke, S.N., Barnes, C.A., 2006. Neural plasticity in the ageing brain. *Nat. Rev. Neurosci.* 7, 30–40.
- Chen, L., Yoo, S.E., Na, R., Liu, Y., Ran, Q., 2012a. Cognitive impairment and increased Aβ levels induced by paraquat exposure are attenuated by enhanced removal of mitochondrial H₂O₂. *Neurobiol. Aging* 33, 432.e15–432.e26.
- Chen, S.Q., Cai, Q., Shen, Y.Y., Wang, P.J., Teng, G.J., Zhang, W., Zang, F.C., 2012b. Age-related changes in brain metabolites and cognitive function in APP/PS1 transgenic mice. *Behav. Brain Res.* 235, 1–6.
- Choi, I.Y., Lee, P., Wang, W.T., Hui, D., Wang, X., Brooks, W.M., Michaelis, E.K., 2014. Metabolism changes during aging in the hippocampus and striatum of Glut1 (glutamate dehydrogenase 1) transgenic mice. *Neurochem. Res.* 39, 446–455.
- Cotman, C.W., Geddes, J.W., Bridges, R.J., Monaghan, D.T., 1989. N-methyl-D-aspartate receptors and Alzheimer's disease. *Neurobiol. Aging* 10, 603–605.
- Daselaar, S.M., Fleck, M.S., Cabeza, R., 2006. Triple dissociation in the medial temporal lobes: recollection, familiarity, and novelty. *J. Neurophysiol.* 96, 1902–1911.
- Duncan, N.W., Wiebking, C., Tiet, B., Marjanska, M., Hayes, D.J., Lyttleton, O., Doyon, J., Northoff, G., 2013. Glutamate concentrations in the medial prefrontal cortex predicts resting-state cortical-subcortical functional connectivity in humans. *PLoS One* 8, e60312.
- Eklund, A., Nichols, T.E., Knutsson, H., 2016. Cluster failure: why fMRI inferences for spatial extent have inflated false-positive rates. *Proc. Natl. Acad. Sci. U. S. A.* 113, 7900–7905.
- Enzi, B., Duncan, N.W., Kaufmann, J., Tempelmann, C., Wiebking, C., Northoff, G., 2012. Glutamate modulates resting state activity in the perigenual anterior cingulate cortex—a combined fMRI-MRS study. *Neuroscience* 227, 102–109.
- Estevez-Gonzalez, A., Kulisevsky, J., Boltes, A., Otermer, P., Garcia-Sanchez, C., 2003. Rey verbal learning is a useful tool for differential diagnosis in the preclinical phase of Alzheimer's disease: comparison with mild cognitive impairment and normal aging. *Int. J. Geriatr. Psychiatry* 18, 1021–1028.
- Falkenberg, L.E., Westerhausen, R., Specht, K., Hugdahl, K., 2012. Resting-state glutamate level in the anterior cingulate predicts blood-oxygen level-dependent response to cognitive control. *Proc. Natl. Acad. Sci. U. S. A.* 109, 5069–5073.
- Fischl, B., Salat, D.H., Busa, E., Albert, M., Dieterich, M., Haselgrove, C., van der Kouwe, A., Killiany, R., Kennedy, D., Klaveness, S., Montillo, A., Makris, N., Rosen, B., Dale, A., 2002. Whole brain segmentation: automated labeling of neuroanatomical structures in the human brain. *Neuron* 33, 341–355.
- FreeSurfer, 2016. Available at: <http://surfer.nmr.mgh.harvard.edu/>. Accessed March 3, 2016.
- Gussew, A., Erdtel, M., Hiepe, P., Rzanny, R., Reichenbach, J.R., 2012. Absolute quantitation of brain metabolites with respect to heterogeneous tissue compositions in (1)H-MR spectroscopic volumes. *Magma* 25, 321–333.
- Hadel, S., Wirth, C., Rapp, M., Gallinat, J., Schubert, F., 2013. Effects of age and sex on the concentrations of glutamate and glutamine in the human brain. *J. Magn. Reson. Imaging* 38, 1480–1487.
- Hattori, N., Abe, K., Sakoda, S., Sawada, T., 2002. Proton MR spectroscopic study at 3 Tesla on glutamate/glutamine in Alzheimer's Disease. *Neuroreport* 13, 183–186.
- Hu, Y., Chen, X., Gu, H., Yang, Y., 2013. Resting-state glutamate and GABA concentrations predict task-induced deactivation in the default mode network. *J. Neurosci.* 33, 18566–18573.
- Jessen, F., Block, W., Traber, F., Keller, E., Flacke, S., Lamerichs, R., Schild, H.H., Huen, R., 2001. Decrease of N-acetylaspartate in the MTL correlates with cognitive decline of AD patients. *Neurology* 57, 930–932.
- Kim, H., 2015. Encoding and retrieval along the long axis of the hippocampus and their relationships with dorsal attention and default mode networks: the HERNET model. *Hippocampus* 25, 500–510.
- Kraguljac, N.V., White, D.M., Hadley, J., Reid, M.A., Lahti, A.C., 2014. Hippocampal-parietal dysconnectivity and glutamate abnormalities in unmedicated patients with schizophrenia. *Hippocampus* 24, 1524–1532.
- Kuhl, B.A., Shah, A.T., DuBrow, S., Wagner, A.D., 2010. Resistance to forgetting associated with hippocampus-mediated reactivation during new learning. *Nat. Neurosci.* 13, 501–506.
- Lin, A.P., Shic, F., Enriquez, C., Ross, B.D., 2003. Reduced glutamate transmission in patients with Alzheimer's disease - in vivo ¹³C magnetic resonance spectroscopy study. *Mag. Res. Mater. Phys. Bio Med.* 16, 29–42.
- Lin, M.T., Beal, M.F., 2006. Mitochondrial dysfunction and oxidative stress in neurodegenerative diseases. *Nature* 443, 787–795.
- Magnusson, K.R., 1998. Aging of glutamate receptors: correlations between binding and spatial memory performance in mice. *Mech. Ageing Dev.* 104, 227–248.
- Mapstone, M., Cheema, A.K., Fiandaca, M.S., Zhong, X., Mhyre, T.R., MacArthur, L.H., Hall, W.J., Fisher, S.G., Peterson, D.R., Haley, J.M., Nazar, M.D., Rich, S.A., Berlau, D.J., Peltz, C.B., Tan, M.T., Kawas, C.H., Federhoff, H.J., 2014. Plasma phospholipids identify antecedent memory impairment in older adults. *Nat. Med.* 4, 415–418.
- McCormick, D.A., Williamson, A., 1989. Convergence and divergence of neurotransmitter action in human cerebral cortex. *Proc. Natl. Acad. Sci. U. S. A.* 86, 8098–8102.
- Miller, S.L., Celone, K., DePeau, K., Diamond, E., Dickerson, B.C., Rentz, D., Pihlajamaki, M., Sperling, R.A., 2008. Age-related memory impairment associated with loss of parietal deactivation but preserved hippocampal activation. *Proc. Natl. Acad. Sci. U. S. A.* 105, 2181–2186.
- Mitrushina, M., Satz, P., Chervinsky, A., D'Elia, L., 1991. Performance of four age groups of normal elderly on the Rey Auditory-Verbal Learning Test. *J. Clin. Psych.* 47, 351–357.
- Morrison, J.H., Baxter, M.G., 2012. The ageing cortical synapse: hallmarks and implications for cognitive decline. *Nature Rev. Neurosci.* 13, 240–250.
- Nicholson, D.A., Yoshida, R., Berry, R.W., Gallagher, M., Geinisman, Y., 2004. Reduction in size of perforated postsynaptic densities in hippocampal axospinous synapses and age-related spatial learning impairments. *J. Neurosci.* 24, 7648–7653.
- Ottersen, O., Zhang, N., Walberg, F., 1992. Metabolic compartmentation of glutamate and glutamine: morphological evidence obtained by quantitative immunocytochemistry in rat cerebellum. *Neuroscience* 46, 519–534.
- Power, J.D., Snyder, A.Z., Schlagger, B.L., Peterson, S.E., 2012. Spurious but systematic correlations in functional connectivity MRI networks arise from subject motion. *Neuroimage* 59, 2142–2154.
- Raichle, M.E., MacLeod, A.M., Snyder, A.Z., Powers, W.J., Gusnard, D.A., Shulman, G.L., 2001. A default mode of brain function. *Proc. Natl. Acad. Sci. U. S. A.* 98, 676–682.
- Ramadan, S., Lin, A., Stanwell, P., 2013. Glutamate and glutamine: a review of in vivo MRS in the human brain. *NMR Biomed.* 26, 1630–1646.
- Raz, N., Lindenberger, U., Rodrigue, K.M., Kennedy, K.M., Head, D., Williamson, A., Dahle, C., Gerstorf, D., Acker, J.D., 2005. Regional brain changes in aging healthy adults: general trends, individual differences and modifiers. *Cereb. Cortex* 15, 1676–1689.
- Rey, A., 1941. L'examen psychologique dans les cas d'encephalopathie traumatique. *Arch. Psychol.* 28, 286–340.
- Rogalski, E.J., Gefen, T., Shi, J., Samimi, M., Bigio, E., Weintraub, S., Geula, C., Mesulam, M.M., 2013. Youthful memory capacity in old brains: anatomic and genetic clues from the Northwestern SuperAging Project. *J. Cogn. Neurosci.* 25, 29–36.

- Rupasingh, R., Borrie, M., Smith, M., Wells, J.L., Bartha, R., 2011. Reduced hippocampal glutamate in Alzheimer disease. *Neurobiol. Aging* 32, 802–810.
- Saad, Z.S., Glen, D.R., Chen, G., Beauchamp, M.S., Desai, R., Cox, R.W., 2009. A new method for improving functional-to-structural MRI alignment using local Pearson correlation. *Neuroimage* 44, 839–848.
- Schoenberg, M.R., Dawson, K.A., Duff, K., Patton, D., Scott, J.G., Adams, R.L., 2006. Test performance and classification statistics for the Rey Auditory Verbal Learning Test in selected clinical samples. *Arch. Clin. Neuropsychol.* 21, 693–703.
- Schubert, F., Gallinat, J., Seifert, F., Rinneberg, H., 2004. Glutamate concentrations in human brain using single voxel proton magnetic resonance spectroscopy at 3 Tesla. *NeuroImage* 21, 1762–1771.
- Simister, R.J., Woermann, F.G., McLean, M.A., Bartlett, P.A., Barker, G.J., Duncan, J.S., 2002. A short-echo-time proton magnetic resonance spectroscopic imaging study of temporal lobe epilepsy. *Epilepsia* 43, 12021–12031.
- Small, S.A., 2001. Age-related memory decline: current concepts and future directions. *Neurol. Rev.* 58, 360–364.
- SPM - Statistical Parametric Mapping, 2016. Available at: <http://www.fil.ion.ucl.ac.uk/spm/>. Accessed March 3, 2016.
- Stanisz, G.J., Odobina, E.E., Pun, J., Escaravage, M., Graham, S.J., Bronskill, M.J., Henkelman, R.M., 2005. T1, T2 relaxation and magnetization transfer in tissue at 3T. *Magn. Reson. Med.* 54, 507–512.
- Stark, C.E.L., Squire, L.R., 2001. When zero is not zero: the problem of ambiguous baseline condition in fMRI. *Proc. Natl. Acad. Sci. U. S. A.* 98, 12760–12766.
- Stark, S.M., Yassa, M.A., Lacy, J.W., Stark, C.E.L., 2013. A task to assess behavioral pattern separation (BPS) in humans: data from healthy aging and mild cognitive impairment. *Behav. Neurosci.* 51, 2442–2449.
- Stemmelin, J., Lazarus, C., Cassel, S., Kelche, C., Cassel, J.C., 2000. Immunohistochemical and neurochemical correlates of learning deficits in aged rats. *Neuroscience* 24, 275–289.
- Strange, B.A., Witter, M.P., Lein, E.S., Moser, E.I., 2014. Functional organization of the hippocampal longitudinal axis. *Nat. Rev. Neurosci.* 15, 655–669.
- Sugaya, K.R.G., Personett, D., Robbins, M., Kent, C., Bryan, D., Skiba, E., Gallagher, M., McKinney, M., 1998. Septo-hippocampal cholinergic and neurotrophin markers in age-induced cognitive decline. *Neurobiol. Aging* 19, 351–361.
- Wagner, G., Gussew, A., Köhler, S., de la Cruz, F., Smesny, S., Reichenbach, J.R., Bär, K.J., 2016. Resting state functional connectivity of the hippocampus along the anterior–posterior axis and its association with glutamatergic metabolism. *Cortex* 81, 104–117.
- Wang, H., Tan, L., Wang, H.F., Liu, Y., Yin, R.H., Wang, W.Y., Chang, X.L., Jiang, T., Yu, J.T., 2015. Magnetic resonance spectroscopy in Alzheimer's disease: systematic review and meta-analysis. *J. Alzheimer's Dis.* 46, 1049–1070.
- Wilson, I.A., Gallagher, M., Eichenbaum, H., Tanila, H., 2006. Neurocognitive aging: Prior memories hinder new hippocampal encoding. *Trends Neurosci.* 29, 662–670.
- Wilson, M., Reynolds, G., Kauppinen, R.A., Arvanitis, T.N., Peet, A.C., 2011. A constrained least-squares approach to the automated quantitation of in vivo (1)H magnetic resonance spectroscopy data. *Magn. Reson. Med.* 65, 1–12.
- Woermann, F.G., McLean, M.A., Bartlett, P.A., Parker, G.J., Duncan, J.S., 1999. Short-echo-time single-voxel 1H magnetic resonance spectroscopy in magnetic resonance imaging negative temporal lobe epilepsy: different biochemical profile compared with hippocampal sclerosis. *Ann. Neurol.* 45, 369–376.
- Yassa, M.A., Lacy, J.W., Stark, S.M., Albert, M.S., Gallagher, M., Stark, C.E., 2011b. Pattern separation deficits associated with increased hippocampal CA3 and dentate gyrus activity in nondemented older adults. *Hippocampus* 21, 968–979.
- Yassa, M.A., Mattfeld, A.T., Stark, S.M., Stark, C.E.L., 2011a. Age-related memory deficits linked to circuit-specific disruptions in the hippocampus. *Proc. Natl. Acad. Sci. U. S. A.* 108, 8873–8878.
- Zahr, N.M., Mayer, D., Pfefferbaum, A., Sullivan, E.V., 2008. Low striatal glutamate levels underlie cognitive decline in the elderly: evidence from in vivo molecular spectroscopy. *Cereb. Cortex* 18, 2241–2250.
- Zahr, N.M., Mayer, D., Rohlfing, T., Chanraud, S., Gu, M., Sullivan, E.V., Pfefferbaum, A., 2013. In vivo glutamate measured with MR spectroscopy: behavioral correlates in aging. *Neurobiol. Aging* 34, 1265–1276.
- Zhao, X., Rosenke, R., Kronemann, D., Brim, B., Das, S.R., Dunah, A.W., Magnusson, K.R., 2009. The effects of aging on N-methyl-D-aspartate receptor subunits in the synaptic membrane and relationships to long-term spatial memory. *Behav. Neurosci.* 162, 933–945.