

UC Davis

UC Davis Previously Published Works

Title

Direct and indirect parental exposure to endocrine disruptors and elevated temperature influences gene expression across generations in a euryhaline model fish.

Permalink

<https://escholarship.org/uc/item/04d788jx>

Authors

DeCourten, Bethany

Brander, Susanne

Connon, Richard

Publication Date

2019

DOI

10.7717/peerj.6156

Copyright Information

This work is made available under the terms of a Creative Commons Attribution License, available at <https://creativecommons.org/licenses/by/4.0/>

Peer reviewed



Direct and indirect parental exposure to endocrine disruptors and elevated temperature influences gene expression across generations in a euryhaline model fish

Bethany M. DeCourten^{1,2}, Richard E. Connon³ and Susanne M. Brander^{1,2}

¹Department of Environmental and Molecular Toxicology, Oregon State University, Corvallis, OR, United States of America

²Department of Biology and Marine Biology, University of North Carolina at Wilmington, Wilmington, NC, United States of America

³Department of Anatomy, Physiology and Cell Biology, University of California, Davis, CA, United States of America

ABSTRACT

Aquatic organisms inhabiting polluted waterways face numerous adverse effects, including physiological disruption by endocrine disrupting compounds (EDCs). Little is known about how the temperatures associated with global climate change may influence the response of organisms exposed to EDCs, and the effects that these combined stressors may have on molecular endpoints such as gene expression. We exposed *Menidia beryllina* (inland silversides) to environmentally relevant concentrations (1 ng/L) of two estrogenic EDCs (bifenthrin and 17 α -ethinylestradiol; EE2) at 22 °C and 28 °C. We conducted this experiment over multiple generations to better understand the potential effects to chronically exposed populations in the wild. We exposed adult parental fish (F0) for 14 days prior to spawning of the next generation. F1 larvae were then exposed from fertilization until 21 days post hatch (dph) before being transferred to clean water tanks. F1 larvae were reared to adulthood, then spawned in clean water to test for further effects of parental exposure on offspring (F2 generation). Gene expression was quantified by performing qPCR on F0 and F1 gonads, as well as F1 and F2 larvae. We did not detect any significant differences in the expression of genes measured in the parental or F1 adult gonads. We found that the 28 °C EE2 treatment significantly decreased the expression of nearly all genes measured in the F1 larvae. This pattern was transferred to the F2 generation for expression of the follicle-stimulating hormone receptor (FSHR) gene. Expression of 17 β -hydroxysteroid dehydrogenase (17 β -HSD) and G protein-coupled receptor 30 (GPR30) revealed changes not measured in the previous generation. Effects of the bifenthrin treatments were not observed until the F2 generation, which were exposed to the chemicals indirectly as germ cells. Our results indicate that effects of EDCs and their interactions with abiotic factors, may not be adequately represented by singular generation testing. These findings will contribute to the determination of the risk of EDC contamination to organisms inhabiting contaminated waterways under changing temperature regimes.

Submitted 26 June 2018

Accepted 24 November 2018

Published 8 January 2019

Corresponding author

Bethany M. DeCourten,
bethany.decourten@oregonstate.edu

Academic editor

Mark Hahn

Additional Information and
Declarations can be found on
page 14

DOI 10.7717/peerj.6156

© Copyright

2019 DeCourten et al.

Distributed under

Creative Commons CC-BY 4.0

OPEN ACCESS

Subjects Genomics, Toxicology, Climate Change Biology, Ecotoxicology, Environmental Impacts

Keywords Ecotoxicology, Gene expression, Climate change, Endocrine disruption, Multiple stressors, Multigenerational, Temperature, *Menidia beryllina*, Bifenthrin, Ethinylestradiol

INTRODUCTION

Many chemicals intended for human use, such as pharmaceuticals and pesticides, are found to pollute adjacent waterways and some are also known to disrupt endocrine function (Brander *et al.*, 2013; Kuivila *et al.*, 2012; Weston & Lydy, 2012; Brander *et al.*, 2016a). Endocrine disrupting compounds (EDCs) influence hormone excretion and signaling in exposed organisms. For example, estrogenic EDCs can alter the expression of genes involved in reproduction and development (Scholz & Gutzeit, 2000; Tu *et al.*, 2016; Brander *et al.*, 2016b). These changes in gene expression may result in higher order consequences potentially impacting organismal function (Brander *et al.*, 2013; Brander *et al.*, 2016b). For example, the synthetic estrogen found in oral contraceptives, 17 α -ethinylestradiol (EE2), has been shown to reduce egg production, influence sex ratios, and influence development in fishes exposed to environmentally relevant concentrations (Kidd *et al.*, 2007; Vajda *et al.*, 2008; Schwindt *et al.*, 2014; Bhandari, Vom Saal & Tillitt, 2015). Many pesticides, such as the commonly used pyrethroid pesticide, bifenthrin, can produce estrogenic effects at low doses that deviates from their intended mode of action on the sodium channels of presynaptic neurons of target pests (Brander *et al.*, 2013). Like EE2, bifenthrin has been shown to impact fishes at multiple levels of biological organization; e.g., gene expression, leading to impaired development and reproduction (Brander *et al.*, 2016a; Brander *et al.*, 2016b; Tu *et al.*, 2016).

Estuaries are typically highly impacted by pollution because they are heavily used by growing coastal communities (Cloern & Jassby, 2012). Protection of estuaries is vital for maintaining healthy food webs and ecosystem services. The model species, *Menidia beryllina* (common name: inland silverside), is a small forage fish that is a food source to a number of estuarine taxa, including commercially important species. *Menidia* are tolerant of a wide range of salinities but are sensitive to chemical stressors, making them an excellent model species to test the effects of anthropogenic stressors that are common in estuaries (Hemmer, Middaugh & Moore, 1990; Brander *et al.*, 2013). Past studies have shown that the expression of genes involved in reproduction, growth and immune function can be altered following EDC exposure in *M. beryllina*. *Menidia* species may be especially susceptible to increasing temperatures and pollutants as their sex determination can be influenced by both temperature and EDC exposure (Duffy, McElroy & Conover, 2009; DeCourten & Brander, 2017). Understanding how organisms with sensitive physiology, such as *M. beryllina*, respond to EDC exposure will help determine the potential risk of these stressors on estuarine ecosystems.

As our planet continues to warm at an unprecedented rate (Stocker *et al.*, 2013), it becomes increasingly important to understand the impacts pollution may have in the context of global climate change. Increases in temperature can inhibit growth, reproduction and early-life stage survival of marine fishes (Munday *et al.*, 2008; Pankhurst & Munday, 2011). Expression of genes along the hypothalamus-pituitary-gonad (HPG) axis can be

reduced by exposure to increased temperatures, inhibiting gonadal function (*Miranda et al., 2013*). The impacts of increased temperature on fishes could lead to changes to overall community structure and trophic dynamics (*Jeppesen et al., 2010*). Studies suggest that the effects of estrogenic endocrine disruptors may be influenced by other abiotic factors such as ambient temperature (*Hemmer, Middaugh & Moore, 1990; Luzio et al., 2016; DeCourten & Brander, 2017*). Testing effects of EDCs under different temperature regimes will help determine the risk they pose to wildlife in the context of global climate change. Elevated temperature may exacerbate the effects of EDCs by increasing the uptake and metabolism of these chemicals in ectothermic organisms (*Clarke & Fraser, 2004*). Additionally, some chemicals such as bifenthrin, which is converted into a metabolite (4-hydroxybifenthrin), have more potent metabolites than their parent compounds (*DeGroot & Brander, 2014*). In this study, we investigated the effects of two estrogenic EDCs (EE2 and bifenthrin) on genes involved in development and reproduction of *M. beryllina*, at two temperatures: 22 °C and 28 °C, to understand how rising temperatures may influence the toxicity of these chemicals. The temperatures chosen for this study represent the most extreme possibility from the worst-case scenario temperature increase (RCP 8.5) expected by the year 2100, put forth by the International Panel for Climate Change (IPCC; *Stocker et al., 2013*). The RCP 8.5 model predicts a surface temperature increase of 1.4–2.6 °C between 2046 and 2065 and a 2.6–4.8 °C 2081–2100 (*Stocker et al., 2013*). Assuming these warming periods are additive, we might expect a 4–7.4 °C increase over the next century, illustrating the importance for testing a 6 °C temperature increase. This is especially important for organisms with plasticity in development, such as *M. beryllina*, whose sex determination can be influenced by temperature and EDC exposure independently (*Duffy, McElroy & Conover, 2009; DeCourten & Brander, 2017*). The 28 °C upper temperature limit was chosen to allow us to test for effects on TSD without reaching the upper breeding limit of *M. beryllina* (30 °C), at which spawning is curtailed (*Hubbs, 1982*). This experiment was conducted over three generations to determine effects at different life stages and effects of parental exposure on offspring. Understanding the effects of concurrent stressors across multiple generations on the gene expression of exposed organisms is necessary to determine the potential for long-term population impacts.

METHODS

Menidia beryllina rearing and exposures

A subset of individuals used in a previous study (*DeCourten & Brander, 2017*) investigating the effects of elevated temperature and EDC exposure on reproduction, development and survival, were sampled for analysis of molecular endpoints. The following methods pertaining to fish rearing, exposures and water chemistry were initially described in *DeCourten & Brander (2017)*. The parental generation of *M. beryllina* were obtained from Aquatic Biosystems (Fort Collins, CO, USA) as juveniles and reared to adulthood at 25 °C, at a salinity of 15 ppt at the University of North Carolina, Wilmington's Center for Marine Science. Fish were fed Hikari tropical micro pellets (Kyorin Food Industries, Ltd, Japan) and live *Artemia nauplii* supplemented with Selcon™ (vitamin supplement; American

Marine Inc., Ridgefield, CT), during rearing and experimental exposures. Adult fish (341 dph) were moved to experimental tanks and acclimated to either 22 °C or 28 °C for 3 weeks prior to experimental exposures under a continuous 16:8 light-dark cycle.

Parental exposures were initiated by replacing approximately 90 percent of the water in experimental tanks with water containing either 1 ng/L of bifenthrin (purity: 99.5%, Chem Service, West Chester, PA, USA) or 1ng/L 17 α -ethinylestradiol (EE2) (purity: \geq 98%, Sigma-Aldrich, St. Louis, MO, USA) suspended in 1 μ L/L of methanol due to the hydrophobic nature of the EDCs. We used 1 μ L/L concentration of methanol in our control group to account for any adverse effects that may be attributed to the solvent. Adult fish (341 dph) were exposed for 14 days in 7.6 L glass jars, containing approximately 6 L of artificial seawater. Exposure jars were housed in 60 L water baths fitted with aquarium heaters set to maintain experimental temperatures. Fish were maintained at a constant salinity target of 15 ppt; optimal for growth and survival of *M. beryllina* (Middaugh, Hemmer & Lamadrid-Rose, 1986). Artificial seawater was made daily by adding Instant Ocean (Spectrum Brands, Blacksburg, VA) to distilled water, which was aerated for 24 h prior to use. Solutions with experimental EDCs were mixed in glass containers immediately prior to daily water changes. Jars were continuously aerated and water quality parameters including temperature (21.9 °C \pm S.D. 0.41 and 27.7 °C \pm S.D. 0.89, respectively), dissolved oxygen (5.81 \pm S.D. 0.98), pH (7.94 \pm S.D. 0.56), salinity (16.26 \pm S.D. 1.59), and ammonia (0.22 \pm S.D. 0.76) were measured with a YSI Professional Plus Quatro water quality meter daily before and after water changes in two representative control jars at each temperature. Water bath temperatures were measured once daily throughout the duration of the experiment. Debris was removed, and sixty percent of the water replaced daily in experimental chambers. All air pumps and heaters were connected to an emergency power source to ensure continuous function. Adult fish were fed a diet of Hikari fish food pellets, freeze-dried blood worms and live *Artemia nauplii* supplemented with SelconTM. Larvae were fed live rotifers (*Brachionus rotundiformis*) from 0–14 dph, and a combination of live *Artemia* and ground Hikari pellets from 10 dph to adulthood. Fish were fed daily and allowed to feed for one hour prior to water change. Following exposures, spawning trials were initiated with the parental generation adults.

Spawning trials were initiated by the addition of spawning substrate (6-inch strands of bleached and rinsed dye-free yarn), as described in Brander *et al.* (2016b). The fish were allowed to spawn for 48 h to produce the subsequent generation. Following the spawning period adult fish were euthanized in an overdose solution of buffered tricaine methanesulfonate (MS-222; approximately 200 mg/L) and immediately weighed and measured, and gonads of three males and three females were collected from each replicate jar. Spawning substrate with attached embryos remained in experimental jars allowing embryos to hatch without transfer. F1 larvae were reared with chemical exposure until 21 dph at which point a subset of five larvae were sampled from each of the 5 replicate jars, to be used for gene expression analysis. Sampled adult tissue and 21dph larvae were immediately frozen in liquid nitrogen and stored at –80 °C. All remaining fish were transferred to clean 37.5 L round black polypropylene tanks and reared to maturity at either 22 °C or 28 °C. Each tank was equipped with Marineland Penguin PF0100B filter,

one Eheim Jager aquarium thermostat heater (75 watt) and Resun AF-2009D automatic feeder. F1 adults were spawned, and the F2 larvae produced were then reared and spawned in clean water (without EDC contamination) at experimental temperatures until 21 dph, to test the effect of parental exposure on offspring. At this time, all larvae were sampled, and the experiment was terminated. All experiments were conducted in accordance with UNCW Institutional Animal Care and Use protocol #A1314-010 and #A1415-010.

Analytical chemistry

Water samples for analytical chemistry were made using the same methods used to make water for the experimental treatments, poured into 950 mL sterile amber bottles, transported on wet ice and stored in the dark at 4 °C for no longer than 24 h. Extraction was performed using conditioned 6 mL solid phase-extraction C₁₈ cartridges (Superclean™ 500 mg; Sigma Aldrich) at a slow drip. Columns were rinsed twice with 5 mL of a 1:1 solution of hexane:ethyl acetate to elute the bifenthrin. The solvent elution (10 mL) was concentrated to 0.4 mL under a gentle stream of nitrogen. Final extract was analyzed using gas chromatography negative chemical ionization mass spectrometry (GC-NCI-MS) on an Agilent 5973 series gas chromatograph (Agilent Technologies, Palo Alto, CA, USA) equipped with a split-splitless injector (280 °C, splitless, 1.5 min purge time). The columns used were Supelco DB-5MS column (30 mm × 0.25 mm with a 0.3 μm film thickness). The instrument was calibrated using nine sets of standard bifenthrin solutions, the surrogate trans-permethrin D6 (EQ Laboratories, Atlanta, GA, USA), and the internal standard dibromooctafluorobiphenyl (Chem Service, West Chester, PA, USA) in hexane. Controls were conducted by analyzing a method blank of deionized water (Milli-Q) to ensure samples were not contaminated. A surrogate was added to each sample before extraction to monitor matrix effects and method performance. The detection limit of bifenthrin in whole water was 0.6 ng/L. No bifenthrin was detected in the controls or method blanks. Measured concentrations of bifenthrin were 1.3 ng/L. Surrogate recovery was 133.84%, with corrected value of bifenthrin concentration being 0.89 ng/L. EE2 treatment water was prepared, in a similar matter to the bifenthrin, according to previously established protocols (*DeGroot & Brander, 2014; Brander et al., 2016b*). All stock and experimental solutions were mixed in the same manner as bifenthrin to ensure like concentrations.

Gene expression quantification

A random subset of 2–3 larvae and 1–2 male and female gonads were selected from each experimental replicate ($n = 3–5$) and used for quantification of gene expression ([Table S1](#)). RNA was extracted using RNeasy Mini Kits (Qiagen, Hilden, Germany) Total RNA concentration and 260/280 ratios (acceptable values > 1.6) were quantified using a Nanodrop 1000 spectrophotometer (ThermoFisher Scientific, Waltham, MA). RNA purity was assessed via electrophoresis on a 1% agarose gel and visualized on a Gel Doc™ XR + Gel Documentation system (BioRad, Hercules, CA, USA). Samples containing trace amounts of genomic DNA were treated with DNase I (Invitrogen™, Waltham, MA) and cleaned up using RNeasy MinElute Cleanup Kit (Qiagen, Hilden, Germany). RNA concentrations were adjusted to a concentration of 100 ng/μl in 96-well plates, randomizing samples within each

generation and tissue type prior to cDNA synthesis. Complimentary DNA (cDNA) was synthesized using Superscript III (Invitrogen, Carlsbad, CA, USA), following manufacture protocols and methods in *Jeffries et al. (2015)*. Twelve genes were selected for quantitative polymerase chain reaction (qPCR), including two reference genes (*Table 1*). We chose to examine a suite of genes involved in reproduction, growth and development based on previous findings from *DeCourten & Brander (2017)*, as well as *Brander et al. (2016b)*. Within the larval samples we chose genes primarily involved in steroidogenesis (17 β -HSD, 3 β -HSD, CYP19b), reproductive pathways (ERs, ARx, FSHR, GnRHR) and growth (IGF2) due to the differences in sex ratio and developmental deformities after early life stage exposure observed in our previous study (*DeCourten & Brander, 2017*). Within the adult gonadal tissues, we selected genes involved in reproductive pathways (ERs, ARx, FSHR, HSP90), and steroidogenesis (17 β -HSD, 3 β -HSD, CYP19a, INHA) also due to our previous observations of skewed sex ratios and reduced reproductive success (*DeCourten & Brander, 2017*). Larval cDNA samples were diluted 2:3 and adult gonad samples were diluted 1:4 with nuclease free water to optimize the reaction. We conducted qPCRs using previously established protocols in *Brander et al. (2016b)*, with Maxima Probe/ROX qPCR Master Mix (ThermoFisher Scientific). Primers and probes were obtained from Eurofins Genomics (Louisville, KY) and Roche LifeScience (Indianapolis, IN), respectively. We conducted qPCRs using 8 μ L cDNA and 12 μ L of mastermix containing 83.3% Maxima Probe/ROX qPCR Master Mix (Thermo Fisher Scientific), 8.3% primers, and 8.3% probe. qPCRs were run on a BioRad CFX96. Reaction protocols included a 10 min hot start incubation at 95 °C, and 40 cycles of 15s at 95 °C, and 30s at 60 °C, during which optical data for the plate was recorded. The environmentally relevant concentrations used elicit small changes in gene expression, thus high concentrations of cDNA were required. Technical replication (x3) was conducted for qPCR efficiency tests when validating primers which demonstrated a high level of accuracy between replicates. Efficiency values for the majority of genes used in this study ranged from 95.1%–107.8%, with the exception of CYP19b (119.2%) and INHA (123.6%), both of which have been used successfully in a previous study (*Brander et al., 2016b*). We did not run technical replicates for experimental qPCRs due to the small amount RNA available from small tissue and larval samples. We opted instead to have a greater number of gene responses examined instead of including technical replicates as previously done in *Connon et al. (2011)*, *Connon et al. (2012)*, *Jeffries et al. (2015)*, *Brander et al. (2016b)*, *Frank et al. (2019)* and *Goff et al. (2017)*. Protocols were developed based on manufacturer's recommendations and previously established protocols from *Brander et al. (2016b)*.

Data analysis

Data was collected from 1–2 individual male and female gonads and 2–3 individual larvae in each replicate tank ($n = 3–5$). Thresholds were set using CFX manager software (BioRad) and calculated CT values were exported to Microsoft Excel 2016. CT values were normalized by the geometric mean of the two housekeeping genes (EF1a and RPL7) used in this study. Housekeeping genes were confirmed to be stable across treatments using GeNorm (*Vandesompele et al., 2002*). Normalized CT values were then converted to

Table 1 Genes used for qPCR analysis. Information on genes selected for qPCR assays.

Gene	Symbol	Roche UPL #	Primer	Function
17-beta-hydroxysteroid dehydrogenase 14	HSD17B14	142	F: gatctgctcaacctcaatttgc R: caggtgtggcaacgcaaat	Steroid synthesis
3-beta-hydroxysteroid dehydrogenase delta 5	HSD3B2	133	F: caccatgctcaacaccacctt R: ctctgaccccaggtctttcttg	Glucocorticoid and androgen synthesis
Gonadal aromatase (<i>adults only</i>)	CYP19a	3	F: gcctcccacagaccaacaat R: gccatgctgaggtttcagtc	Steroid metabolism
Brain aromatase (<i>larvae only</i>)	CYP19b	25	F: cagaaccagatgtggagcag R: cacagactttcccctgaacacc	Steroid metabolism
Estrogen receptor alpha	ESR1	15	F: ctccattgtgccagtgacaga R: acgcttcgcatgctca	Hormone receptor
Estrogen receptor beta a	ESR2	52	F: gaccatcctgggaaactgatctt R: cattatgccctccacgcact	Hormone receptor
Estrogen receptor beta b	ESR3	130	F: gattttattcaaccggagcagtg R: catcgctcgtctgatgaact	Hormone receptor
Androgen receptor	Arx	31	F: atccgcatgcagtgctcata R: cccagacctcgtattcaacg	Hormone receptor
Follicle stimulating hormone receptor	FSHR	67	F: tgggatctcaacaatgccttc R: gatccgctctgcttaatttg	Hormone receptor
Gonadotropin-releasing hormone receptor 1	GnRHR	108	F: cgggtggcgtggacaatg R: ggtgatggtcacgtttgtgga	Hormone receptor
G-protein coupled estrogen receptor (<i>larvae only</i>)	GPR30	48	F: tcttcgtcgtctgttgcacat R: cgtcctcctccgcctctac	Hormone receptor
Insulin-like growth factor 2 (<i>larvae only</i>)	IGF2	38	F: gcaggtcatacccgatgacg R: ggctgccttctattccacac	Growth factor
Heat shock protein 90 beta (<i>adults only</i>)	HSP90AB1	55	F: gaaaaggtagaccgtgtccaac R: gccgtaagtgtggtcacaat	Molecular Chaperone
Inhibin alpha subunit (<i>adults only</i>)	INHA	24	F: gaggaggatcagggggactg R: cgggtggacgatccacttt	Negative Feedback
60s ribosomal protein l7	RPL_7	31	F: aacttctgtggccgttcaag R: tegcctccctccacaaagt	Reference gene
Elongation factor 1-alpha-like	EF1a	65	F: catcgctgcaagttcagc R: cccagacttcagcctt	Reference gene

log₂ fold change values using methods described in [Livak & Schmittgen \(2001\)](#). Further information is provided in [Data S1](#). Values from multiple fish in each replicate were averaged to obtain one value for each replicate tank. We tested for significant differences ($p < 0.05$) among all treatments using a two-way ANOVA and Tukey's post hoc multiple comparisons with false discovery rate p -value corrections applied ($p < 0.05$; [Benjamini & Hochberg, 1995](#)). All statistical tests were performed in R ([R Core Team, 2017](#); v 3.4.3) and Tukey's post hoc multiple comparisons were made using the *multcomp* package ([Hothorn, Bretz & Westfall, 2008](#)).




Figure 1 Relative gene expression of F1 larvae. Gene expression in fold change (log₂) of the ESR1 (A), ESR2 (B), ESR3 (C) GPR30 (D), 3β-HSD (E), 17β-HSD (F), FSHR (G), GnRHR(H), CYP19b (I), and IGF2 (J) genes. Values normalized to the geometric mean of the housekeeping genes in the F1 larvae (21 dph) at 22 °C and 28 °C in the control (C), bifenthrin (BF) and ethinylestradiol (EE2) treatments. Boxes denote inner quartile range, whiskers represent upper and lower quartiles, and median is denoted by solid bars. Significance is denoted by lowercase letters (two-way ANOVA, Tukey's HSD, $p < 0.05$, $n = 5$).

Full-size [DOI: 10.7717/peerj.6156/fig-1](https://doi.org/10.7717/peerj.6156/fig-1)

RESULTS

Parental generation

There were no significant differences in expression of genes tested in the parental generation adult ovaries (Fig. S1) or testes (Fig. S2) following experimental exposures. A full list of pairwise comparisons and p -values across all generations is presented in Table S2.

F1 generation

Within the F1 larvae, (21 dph; Fig. 1) we found that the 28 °C EE2 treatment significantly reduced the expression of all genes except for ESR1, relative to all other experimental treatments with the exception of 28 °C bifenthrin. There was reduced expression of most of the estrogen receptors tested including: ESR2 (two-way ANOVA, Tukey's HSD, $p \leq 0.05$), ESR3 (two-way ANOVA, Tukey's HSD, $p \leq 0.05$) and GPR30 (two-way ANOVA, Tukey's HSD, $p \leq 0.05$), in the 28 °C EE2 treatment. There was reduced expression of receptors for hormones secreted by the pituitary: GnRHR (two-way ANOVA, Tukey's HSD, $p \leq 0.05$) and FSHR (two-way ANOVA, Tukey's HSD, $p \leq 0.05$). Expression of genes involved in

steroidogenesis was reduced in the 28 °C EE2 treatment including: CYP19b, (two-way ANOVA, Tukey's HSD, $p \leq 0.05$), 17 β -HSD (two-way ANOVA, Tukey's HSD, $p \leq 0.05$) and 3 β -HSD (two-way ANOVA, Tukey's HSD, $p \leq 0.05$). We also determined a decrease in expression of IGF2 (two-way ANOVA, Tukey's HSD, $p \leq 0.05$), which is involved in growth regulation. For a full list of p -values for all pairwise comparisons refer to [Table S2](#). There were no significant differences in gene expression of the F1 larvae among any of the 22 °C treatments, and we found there were no significant changes in gene expression in any of the genes tested in the adult F1 ovaries ([Fig. S3](#)) or testes ([Fig. S4](#)) in any of the chemical or temperature treatments.

F2 generation

Within the F2 larvae ([Fig. 2](#)), we found that GPR30 expression was reduced by the 22 °C bifenthrin and EE2 treatments (two-way ANOVA, Tukey's HSD, $p \leq 0.01$). There were no differences in expression in the other estrogen receptors (ESR1, ESR2, ESR3). The pattern of expression of FSHR observed in the F1 generation was carried over into the F2 generation ([Fig. 2](#)), with the 28 °C EE2 treatment significantly reducing the expression of this gene when compared to all other treatments (two-way ANOVA, Tukey's HSD, $p \leq 0.01$). We did not observe any differences in expression of GnRHR, indicating that receptors of hormones secreted by the pituitary responded differently in the F2 generation. The expression of 17 β -HSD was significantly altered by the 28 °C bifenthrin treatment when compared to the 22 °C control (two-way ANOVA, Tukey's HSD, $p = 0.0249$). Other genes involved in steroidogenesis (3 β -HSD and CYP19b) in the F2 generation were not significantly affected by the experimental exposures. As observed in the previous generation, we did not observe differences in expression of IGF2 among the experimental treatments.

DISCUSSION

Many EDCs have been detected in estuarine waterways, including many commonly used pharmaceuticals and pesticides. Bifenthrin is a pyrethroid pesticide that enters watersheds primarily via runoff following application for urban and agricultural pest control ([Kuivila et al., 2012](#)). This compound is found in California surface waters at concentrations up to 106 ng/L ([Weston & Lydy, 2012](#)) and higher concentrations have been found in sediments throughout the world ([Li et al., 2017](#)). EE2 is a common component of oral contraceptives, which enters the waterways via wastewater effluent, at concentrations up to 7.0 ng/L ([Adeel et al., 2017](#); [King et al., 2016](#)). While waste water effluents are primarily found in riverine environments, these rivers discharge into downstream estuarine and marine systems, introducing EE2 at low (ng/L) concentrations ([Aris, Shamsuddin & Praveena, 2014](#)). Environmentally relevant concentrations of these estrogenic EDCs have been shown to elicit adverse effects in fish including disruption of homeostasis, reproduction and development ([Brander et al., 2016a](#); [Brander et al., 2016b](#); [DeCourten & Brander, 2017](#)). Our results support previous findings illustrating the importance of testing for adverse effects of estrogenic EDCs at low doses, which can produce stronger effects than those observed at moderate doses ([Brander et al., 2012](#); [Vandenberg et al., 2012](#)).

Relative gene expression of F2 larvae




Figure 2 Relative gene expression of F2 larvae. Gene expression in fold change (log₂) of the ESR1 (A), ESR2 (B), ESR3 (C), GPR30 (D), 3β-HSD (E), 17β-HSD (F), FSHR (G), GnRHR (H), CYP19b (I), and IGF2 (J) genes. Values normalized to the geometric mean of the housekeeping genes in the F2 larvae (21 dph) at 22 °C and 28 °C in the control (C), bifenthrin (BF) and ethinylestradiol (EE2) treatments. Boxes denote inner quartile range, whiskers represent upper and lower quartiles, and median is denoted by solid bars. Significance is denoted by lowercase letters (two-way ANOVA, Tukey's HSD, $p < 0.05$, $n = 5$ in the 22 °C EE2 and 22 °C Control, $n = 4$ in 22 °C BF, 28 °C BF and 28 °C Control, $n = 3$ in 28 °C EE2).

Full-size [DOI: 10.7717/peerj.6156/fig-2](https://doi.org/10.7717/peerj.6156/fig-2)

We found that the effects of EE2 exposure may be influenced by temperature in the F1 larvae, with the 28 °C EE2 treatment significantly altering expression of most genes tested. Many of the genes selected in this study are involved in the hypothalamic-pituitary-gonadal (HPG) axis and are useful in determining the effects of elevated temperature and EDC exposure on development and reproduction. We found that 28 °C EE2 treatment reduced the expression of the aromatase gene, CYP19b, which is primarily expressed in the brain and plays a pivotal role in sex determination by the conversion of androgens into estrogens (Diotel et al., 2010). Opposite of our findings with EE2, previous studies found that CYP19b expression was increased following exposure to environmental androgens or estrogens (Baumann et al., 2014; Filby et al., 2007). We found a similar pattern of expression in ESR2 and ESR3, which are nuclear, ligand-dependent estrogen receptors that can regulate estrogenic responses via direct mechanisms or by binding to estrogen response elements in the promoter regions of some genes (Filby & Tyler, 2005). Our findings in the 28 °C EE2 group are contrary to previous studies that reported an upregulation of estrogen receptors

following exposure to estrogenic EDCs (Filby & Tyler, 2005; Filby et al., 2007; Rhee et al., 2011) indicating that these effects may be dependent on temperature. These discrepancies may also be attributed to the length of chemical exposure (0 hpf to 21 dph, including a 7–10 day embryonic incubation), and responses regulated via ligand–receptor feedback loops can be dependent on exposure period (Golshan et al., 2014). There is evidence that genes upregulated during earlier timepoints in early life stage exposure to EDCs can begin to down regulate as the exposure continues past 30 days post fertilization (Shved et al., 2008). Analogous to the expression of nuclear estrogen receptors, we found a reduction in the expression of GPR30, an estrogen receptor coupled with seven transmembrane G proteins that plays a pivotal role in reproductive success including activating oocyte maturation (Maggiolini & Picard, 2010; Pang, Dong & Thomas, 2008). The production of the enzymes 17 β -HSD and 3 β -HSD, which act as catalysts in the steroidogenesis pathway, is essential in the synthesis pathway of biologically active sex steroids (Marchais-Oberwinkler et al., 2011; Liu et al., 2012). We found that EE2 (28 °C) downregulated the expression of genes coding for the production of these enzymes, corroborating previous findings that production of this class of enzymes can be influenced by exposure to EE2 (Liu et al., 2012; Filby et al., 2007). Similarities between the expression of the above-mentioned genes indicate that the response to EE2 can be exacerbated by temperature along the HPG axis following exposure at the larval stage. Changes in the expression of these genes may influence the sexual development and reproductive capacity of exposed fish. At the adult stage, F1 fish in this experiment were observed to have a more female biased sex ratio in the 28 °C EE2 than any other treatment (DeCourten & Brander, 2017). The observed changes in gene expression may explain the skewed sex ratios of the F1 adults in the 28 °C EE2 treatment (DeCourten & Brander, 2017) as sex of *M. beryllina* is determined in the larval stage (Conover & Fleisher, 1986; Strüssmann et al., 2010). Sex ratios deviating from the evolutionarily stable 50:50 males to females have the potential to result in population declines in *M. beryllina* (White et al., 2017).

We investigated the response of additional genes in larval tissue essential to reproductive function at later life stages. We found that the 28 °C EE2 treatment downregulated the production of GnRHR, which indicates that less gonadotropin releasing hormone (GnRH) is being produced in the hypothalamus (Levavi-Sivan et al., 2010). GnRH stimulates gonadotropin production in the gonads and is essential for gonad maturation, reproduction, and reproductive behaviors (Jodo, Ando & Urano, 2003). A similar pattern of expression was found with the FSHR gene indicating that EE2 (28 °C) downregulated the production of follicle stimulating hormone (FSH) in the pituitary of the larval fish. FSH levels can vary based on reproductive status of the individual and mediates steroidogenesis, gametogenesis and ovulation in the gonads (Rocha et al., 2009). A previous study using fish exposed to BPA, found an upregulation on GnRHR and FSHR, opposite of our finding (Rhee et al., 2011). However, Rhee et al. (2011) found similar expression patterns across GnRHR, FSHR, ERs, and CYP19a, indicating that these genes respond to EDCs similarly. Our findings align with Rhee et al. (2011) as the downregulation observed in the 28 °C EE2 treatment was similar across all genes we tested, indicating a similar response to EE2 exposure throughout the HPG axis. These results could explain the reduction

in egg production by the F1 adults in the 28 °C EE2 treatment in this experiment as multiple genes within the HPG axis were downregulated within the 28 °C EE2 treatment, potentially impacting the development and reproduction (*DeCourten & Brander, 2017*). However, these changes in gene expression were not observed in the F1 adult gonad tissues, suggesting that gene expression altered immediately after EE2 exposure was not measurable later in life after rearing in clean water. Despite the lack of altered gene expression at the adult stage, we were able to measure changes to higher order processes (e.g., sex ratio, egg production) that could have been mediated through the expression of genes at the larval stage, which highlights the importance of testing across levels of biological organization (*Brander, Hecht & Kuivila, 2015*).

Previous literature has established that exposure to estrogenic EDCs has been shown to alter the expression of many genes involved in the reproductive and developmental pathways in gonadal tissues (*Scholz & Gutzeit, 2000; Filby et al., 2007; Zhang et al., 2008*). Likewise, increased temperature has been shown to influence expression of genes such as gonadal aromatase, which is the mechanism through which animals are able to achieve environmental sex determination (*Kitano et al., 1999; Uchida et al., 2004*). However, we were unable to detect any significant differences in the expression of genes in the parental ovaries (*Fig. S1*), parental testes (*Fig. S2*), F1 adult ovaries (*Fig. S3*), F1 adult testes (*Fig. S4*) in any treatments tested in this study. This is likely due to high variability of measurements between our samples and the loss of statistical power within our tests resulting from a high number of pairwise comparisons. These discrepancies with existing literature may also be explained by exposing our fish to EDCs at a salinity of 15 ppt because EE2 uptake can be influenced by changes in salinity (*Blewett et al., 2013*). Additionally, because we were only able to sample fish at one time point in each life stage we may have missed an opportunity to detect changes in gene expression that had occurred earlier on or there may have been unmeasured differences in protein or hormone concentrations that were not represented by gene expression quantifications at the end of the exposure. This may explain why these exposures resulted in phenotypic changes such as decreases in egg production within 48 h of the sampling time (*DeCourten & Brander, 2017*).

Like the genes involved in the HPG axis, we found that the IGF2 expression was reduced by exposure to EE2 at 28 °C. Production of IGFs are regulated by growth hormones, mediating the uptake of glucose and amino acids, and protein synthesis (*Shamblott et al., 1995*). A study with fathead minnows (*Pimephales promelas*) found that bifenthrin downregulated the expression of IGF2, when using higher concentrations than those used in our study (*Beggel et al., 2011*). This supports our findings that high doses or highly estrogenic EDCs, such as EE2 in our study, can downregulate the production of IGF2, although we did not find a significant difference between low concentrations of bifenthrin and the controls. Previous literature has shown that IGF2 transcription is not influenced by temperature (*Gabillard et al., 2003*), despite the growth of ectothermic organisms being heavily influenced by elevated temperatures (*Moyano et al., 2016*). Our results support the findings that temperature alone does not influence IGF2 transcription. However, we found that EE2 exposure, in combination with elevated temperature, had a greater effect on IGF2 regulation than either stressor alone.

It has been shown that temperature can exacerbate the feminizing effects of EE2 exposure and delay the development of zebrafish testes (Luzio *et al.*, 2016). Previous studies have found that effects of estrogenic EDCs on molecular endpoints and development can be influenced by abiotic factors such as temperature and photoperiod (Brian *et al.*, 2008; Jin *et al.*, 2009). For example, Brian *et al.* (2008) found that the vitellogenin production increased at higher temperatures in fish exposed to a mixture of estrogenic EDCs. In the case of 17 β -HSD, we found that increased temperature can also influence effects of bifenthrin exposure in the F2 generation. Bifenthrin has been shown to alter the expression of genes involved in hormone regulation producing higher order effects on reproduction (Brander *et al.*, 2016b). It has been suggested that effects produced by concurrent exposure to EDCs and elevated temperature can negatively impact survival, development and reproduction, resulting in population declines over time (Brown *et al.*, 2015).

Our results indicate that organisms exposed at early life stages are more susceptible to molecular effects of EDCs, as found with previous studies (Guillette Jr *et al.*, 1995; Van Aerle *et al.*, 2002), as we determined a greater effect of EDCs and temperature at the larval life stage than adult. In the case of FSHR expression, we found that the same pattern measured in the previous generation was also observed in their offspring. Parental exposure to environmental factors can result in phenotypic changes to offspring to allow for rapid response to environmental stressors (Burgess & Marshall, 2014). It has previously been shown that the effects of EDCs can be observed across multiple generations following exposure (Corrales *et al.*, 2014). Transgenerational effects have been linked to lowered survival and reproduction in generations following parental exposure, including generations only exposed as germ cells (Schwindt *et al.*, 2014; Bhandari, Vom Saal & Tillitt, 2015). However, similar effects between the F1 and F2 larvae were not seen in all genes with the expression of 17 β -HSD and GPR30 differing between generations. For example, we saw a greater influence of bifenthrin on the expression of 17 β -HSD and GRP30 in the F2 larvae, which were exposed indirectly as germ cells, than the previous generation. Our previous findings suggest that EDCs can have more deleterious effects on development in indirectly exposed larvae (DeCourten & Brander, 2017). Our results indicate that the full impact of some EDCs may not be realized when conducting exposures on a single generation. Multigenerational testing is important to determine the effects of parental exposure and potential epigenetic modifications caused by EDC exposure that may result in persistent effects to exposed populations (Brander, Biales & Connon, 2017).

CONCLUSIONS

We found that increases in temperature can increase the effects of EDC exposure, as we observed more alterations of gene expression in higher temperatures. Based off predictions of temperature increases over the next 100 years by the IPCC 2013 report (Stocker *et al.*, 2013), we can expect to see rising temperatures influence how organisms respond to exposure to EDCs. Our findings confirm the importance for research incorporating multiple stressors to assess the risk of anthropogenic stressors to aquatic organisms. We found that the changes in larval gene expression resulting from exposure to a combination

of EDCs and temperature can result in higher order effects that are measurable later in life after molecular endpoints are no longer affected. These findings imply that organisms exposed briefly as larvae can be affected by these stressors throughout their lives regardless of time allotted for recovery. Additionally, we found that effects of EDCs can be carried over to the next generation or in some cases produce different effects in subsequent generations. These findings suggest mechanisms dictating generational transfer of effects following parental exposure may not affect all molecular endpoints equally. Further research is needed to determine the effects of parental exposure on subsequent generations. We found that larval life stages were more sensitive to EDC exposure than adults, and that changes in gene expression following EDC exposure can be further influenced by temperature. Our findings will inform the long-term management of aquatic systems vulnerable to pollution under future climate change scenarios, many of which provide critical habitat for fishes during early life stages.

ACKNOWLEDGEMENTS

We thank the staff at the UNCW Center for Marine Science and Department of Biology and Marine Biology for support during the experiment, as well as our collaborators in the Connon laboratory at the University of California, Davis for informing the laboratory protocols used. We thank Mr. Nathan Burns for assistance with RNA extraction and the many undergraduate assistants for their contribution to the project, and Mr. Christian Commander for his advice and assistance with statistical software.

ADDITIONAL INFORMATION AND DECLARATIONS

Funding

This work was supported by the California Department of Fish and Wildlife (contract # E1183010 to Susanne M. Brander and Richard E. Connon) and the US Environmental Protection Agency (EPA STAR #835799 to Susanne M. Brander). Bethany M. DeCourten received additional support from the Ahuja Water Quality Fellowship and the Francis Peter Fensel Memorial Fellowship. Additional research funds were acquired from a 2011 UNCW Center for Marine Science pilot grant. The funders had no role in study design, data collection and analysis, decision to publish, or preparation of the manuscript.

Grant Disclosures

The following grant information was disclosed by the authors:

California Department of Fish and Wildlife: #E1183010.

US Environmental Protection Agency: EPA STAR #835799.

Ahuja Water Quality Fellowship.

Francis Peter Fensel Memorial Fellowship.

2011 UNCW Center for Marine Science.

Competing Interests

Susanne M. Brander is an Academic Editor for PeerJ.

Author Contributions

- Bethany M. DeCourten conceived and designed the experiments, performed the experiments, analyzed the data, prepared figures and/or tables, authored or reviewed drafts of the paper, approved the final draft.
- Richard E. Connon contributed reagents/materials/analysis tools, authored or reviewed drafts of the paper, approved the final draft.
- Susanne M. Brander conceived and designed the experiments, contributed reagents/materials/analysis tools, authored or reviewed drafts of the paper, approved the final draft.

Animal Ethics

The following information was supplied relating to ethical approvals (i.e., approving body and any reference numbers):

Our research was approved by the Institutional Animal Care and Use Committee at the University of North Carolina, Wilmington (protocols #A1314-010 and #A1415-010).

Data Availability

The following information was supplied regarding data availability:

The raw data is provided in the [Supplementary Files](#).

Supplemental Information

Supplemental information for this article can be found online at <http://dx.doi.org/10.7717/peerj.6156#supplemental-information>.

REFERENCES

- Adeel M, Song X, Wang Y, Francis D, Yang Y. 2017.** Environmental impact of estrogens on human, animal and plant life: a critical review. *Environment International* **99**:107–119 DOI [10.1016/j.envint.2016.12.010](https://doi.org/10.1016/j.envint.2016.12.010).
- Aris AZ, Shamsuddin AS, Praveena SM. 2014.** Occurrence of 17 α -ethynylestradiol (EE2) in the environment and effect on exposed biota: a review. *Environment International* **69**:104–119 DOI [10.1016/j.envint.2014.04.011](https://doi.org/10.1016/j.envint.2014.04.011).
- Baumann L, Knörr S, Keiter S, Nagel T, Rehberger K, Volz S, Oberrauch S, Schiller V, Fenske M, Holbech H, Segner H, Braunbeck T. 2014.** Persistence of endocrine disruption in zebrafish (*Danio rerio*) after discontinued exposure to the androgen 17 β -trenbolone. *Environmental Toxicology and Chemistry* **33(11)**:2488–2496 DOI [10.1002/etc.2698](https://doi.org/10.1002/etc.2698).
- Beggel S, Connon R, Werner I, Geist J. 2011.** Changes in gene transcription and whole organism responses in larval fathead minnow (*Pimephales promelas*) following short-term exposure to the synthetic pyrethroid bifenthrin. *Aquatic Toxicology* **105(1–2)**:180–188 DOI [10.1016/j.aquatox.2011.06.004](https://doi.org/10.1016/j.aquatox.2011.06.004).
- Benjamini Y, Hochberg Y. 1995.** Controlling the false discovery rate: a practical and powerful approach to multiple testing. *Journal of the Royal Statistical Society. Series B (Methodological)* **57(1)**:289–300.

- Bhandari RK, Vom Saal FS, Tillitt DE. 2015.** Transgenerational effects from early developmental exposures to bisphenol A or 17 α -ethynylestradiol in medaka, *Oryzias latipes*. *Scientific Reports* 5:9303 DOI 10.1038/srep09303.
- Blewett TA, Robertson LM, MacLatchy DL, Wood CM. 2013.** Impact of environmental oxygen, exercise, salinity, and metabolic rate on the uptake and tissue-specific distribution of 17 α -ethynylestradiol in the euryhaline teleost *Fundulus heteroclitus*. *Aquatic toxicology* 138:43–51 DOI 10.1016/j.aquatox.2013.04.006.
- Brander SM, Biales AD, Connon RE. 2017.** The role of epigenomics in aquatic toxicology. *Environmental Toxicology and Chemistry* 36(10):2565–2573 DOI 10.1002/etc.3930.
- Brander SM, Cole BJ, Cherr GN. 2012.** An approach to detecting estrogenic endocrine disruption via choriogenin expression in an estuarine model fish species. *Ecotoxicology* 21(4):1272–1280.
- Brander SM, Connon RE, He G, Hobbs JA, Smalling KL, Teh SJ, White JW, Werner I, Denison MS, Cherr GN. 2013.** From ‘omics to otoliths: responses of an estuarine fish to endocrine disrupting compounds across biological scales. *PLOS ONE* 8(9):e74251 DOI 10.1371/journal.pone.0074251.
- Brander SM, Gabler MK, Fowler NL, Connon RE, Schlenk D. 2016a.** Pyrethroid pesticides as endocrine disruptors: molecular mechanisms in vertebrates with a focus on fishes. *Environmental Science & Technology* 50(17):8977–8992 DOI 10.1021/acs.est.6b02253.
- Brander S, Hecht S, Kuivila K. 2015.** The Challenge: “Bridging the gap” with fish: advances in assessing exposure and effects across biological scales. *Environmental Toxicology and Chemistry* 34(3):459–459.
- Brander SM, Jeffries KM, Cole BJ, DeCourten BM, White JW, Hasenbein S, Fangue NA, Connon RE. 2016b.** Transcriptomic changes underlie altered egg protein production and reduced fecundity in an estuarine model fish exposed to bifenthrin. *Aquatic Toxicology* 174:247–260 DOI 10.1016/j.aquatox.2016.02.014.
- Brian JV, Harris CA, Runnalls TJ, Fantinati A, Pojana G, Marcomini A, Booy P, Lamoree M, Kortenkamp A, Sumpter J. 2008.** Evidence of temperature-dependent effects on the estrogenic response of fish: implications with regard to climate change. *Science of the Total Environment* 397(1–3):72–81 DOI 10.1016/j.scitotenv.2008.02.036.
- Brown AR, Owen SF, Peters J, Zhang Y, Soffker M, Paull GC, Hosken DJ, Wahab MA, Tyler CR. 2015.** Climate change and pollution speed declines in zebrafish populations. *Proceedings of the National Academy of Sciences of the United States of America* 112(11):E1237–E1246 DOI 10.1073/pnas.1416269112.
- Burgess SC, Marshall DJ. 2014.** Adaptive parental effects: the importance of estimating environmental predictability and offspring fitness appropriately. *Oikos* 123(7):769–776 DOI 10.1111/oik.01235.
- Clarke A, Fraser KPP. 2004.** Why does metabolism scale with temperature? *Functional Ecology* 18(2):243–251 DOI 10.1111/j.0269-8463.2004.00841.x.

- Cloern JE, Jassby AD. 2012.** Drivers of change in estuarine-coastal ecosystems: discoveries from four decades of study in San Francisco Bay. *Reviews of Geophysics* **50(4)**:RG4001 DOI [10.1029/2012RG000397](https://doi.org/10.1029/2012RG000397).
- Connon RE, Beggel S, D'Abronzio LS, Geist JP, Pfeiff J, Loguinov AV, Vulpe CD, Werner I. 2011.** Linking molecular biomarkers with higher level condition indicators to identify effects of copper exposures on the endangered delta smelt (*Hypomesus transpacificus*). *Environmental Toxicology and Chemistry* **30(2)**:290–300 DOI [10.1002/etc.400](https://doi.org/10.1002/etc.400).
- Connon RE, D'Abronzio LS, Hostetter NJ, Javidmehr A, Roby DD, Evans AF, Loge FJ, Werner I. 2012.** Transcription profiling in environmental diagnostics: health assessments in Columbia River basin steelhead (*Oncorhynchus mykiss*). *Environmental Science and Technology* **46(11)**:6081–6087 DOI [10.1021/es3005128](https://doi.org/10.1021/es3005128).
- Conover DO, Fleisher MH. 1986.** Temperature-sensitive period of sex determination in the Atlantic silverside, *Menidia*. *Canadian Journal of Fisheries and Aquatic Sciences* **43(3)**:514–520 DOI [10.1139/f86-061](https://doi.org/10.1139/f86-061).
- Corrales J, Thornton C, White M, Willett KL. 2014.** Multigenerational effects of benzo [a] pyrene exposure on survival and developmental deformities in zebrafish larvae. *Aquatic Toxicology* **148**:16–26 DOI [10.1016/j.aquatox.2013.12.028](https://doi.org/10.1016/j.aquatox.2013.12.028).
- DeCourten BM, Brander SM. 2017.** Combined effects of increased temperature and endocrine disrupting pollutants on sex determination, survival, and development across generations. *Scientific Reports* **7(1)**:9310 DOI [10.1038/s41598-017-09631-1](https://doi.org/10.1038/s41598-017-09631-1).
- DeGroot BC, Brander SM. 2014.** The role of P450 metabolism in the estrogenic activity of bifenthrin in fish. *Aquatic Toxicology* **156**:17–20 DOI [10.1016/j.aquatox.2014.07.007](https://doi.org/10.1016/j.aquatox.2014.07.007).
- Diotel N, Le Page Y, Mouriec K, Tong SK, Pellegrini E, Vaillant C, Anglade I, Brion F, Pakdel F, Chung BC, Kah O. 2010.** Aromatase in the brain of teleost fish: expression, regulation and putative functions. *Frontiers in Neuroendocrinology* **31(2)**:172–192 DOI [10.1016/j.yfrne.2010.01.003](https://doi.org/10.1016/j.yfrne.2010.01.003).
- Duffy TA, McElroy AE, Conover DO. 2009.** Variable susceptibility and response to estrogenic chemicals in *Menidia menidia*. *Marine Ecology Progress Series* **380**:245–254 DOI [10.3354/meps07928](https://doi.org/10.3354/meps07928).
- Filby AL, Thorpe KL, Maack G, Tyler CR. 2007.** Gene expression profiles revealing the mechanisms of anti-androgen- and estrogen-induced feminization in fish. *Aquatic Toxicology* **81(2)**:219–231 DOI [10.1016/j.aquatox.2006.12.003](https://doi.org/10.1016/j.aquatox.2006.12.003).
- Filby AL, Tyler CR. 2005.** Molecular characterization of estrogen receptors 1, 2a, and 2b and their tissue and ontogenic expression profiles in fathead minnow (*Pimephales promelas*). *Biology of Reproduction* **73(4)**:648–662 DOI [10.1095/biolreprod.105.039701](https://doi.org/10.1095/biolreprod.105.039701).
- Frank DF, Brander SM, Hasenbein S, Harvey DJ, Lein PJ, Geist J, Connon RE. 2019.** Developmental exposure to environmentally relevant concentrations of bifenthrin alters transcription of mTOR and ryanodine receptor-dependent signaling molecules

- and impairs predator avoidance behavior across early life stages in inland silversides (*Menidia beryllina*). *Aquatic Toxicology* **206**:1–13.
- Gabillard JC, Weil C, Rescan PY, Navarro I, Gutiérrez J, Le Bail PY. 2003.** Effects of environmental temperature on IGF1, IGF2, and IGF type I receptor expression in rainbow trout (*Oncorhynchus mykiss*). *General and Comparative Endocrinology* **133**(2):233–242 DOI [10.1016/S0016-6480\(03\)00167-9](https://doi.org/10.1016/S0016-6480(03)00167-9).
- Goff AD, Saranjampour P, Ryan LM, Hladik ML, Covi JA, Armbrust KL, Brander SM. 2017.** The effects of fipronil and the photodegradation product fipronil desulfinyl on growth and gene expression in juvenile blue crabs, *Callinectes sapidus*, at different salinities. *Aquatic Toxicology* **186**:96–104 DOI [10.1016/j.aquatox.2017.02.027](https://doi.org/10.1016/j.aquatox.2017.02.027).
- Golshan M, Hatf A, Zare A, Socha M, Milla S, Gosiewski G, Fontaine P, Sokołowska-Mikołajczyk M, Habibi HR, Alavi SMH. 2014.** Alternations in neuroendocrine and endocrine regulation of reproduction in male goldfish (*Carassius auratus*) following an acute and chronic exposure to vinclozolin, *in vivo*. *Aquatic Toxicology* **155**:73–83 DOI [10.1016/j.aquatox.2014.06.004](https://doi.org/10.1016/j.aquatox.2014.06.004).
- Guillette Jr LJ, Crain DA, Rooney AA, Pickford DB. 1995.** Organization versus activation: the role of endocrine-disrupting contaminants (EDCs) during embryonic development in wildlife. *Environmental Health Perspectives* **103**(Suppl 7):157–164 DOI [10.1289/ehp.95103s7157](https://doi.org/10.1289/ehp.95103s7157).
- Hemmer MJ, Middaugh DP, Moore JC. 1990.** Effects of temperature and salinity on *Menidia beryllina* embryos exposed to terbufos. *Diseases of Aquatic Organisms* **8**(2):127–136 DOI [10.3354/dao008127](https://doi.org/10.3354/dao008127).
- Hothorn T, Bretz F, Westfall P. 2008.** Simultaneous inference in general parametric models. *Biometrical Journal* **50**(3):346–363 DOI [10.1002/bimj.200810425](https://doi.org/10.1002/bimj.200810425).
- Hubbs C. 1982.** Life history dynamics of *Menidia beryllina* from Lake Texoma. *American Midland Naturalist* **107**(1):1–12 DOI [10.2307/2425184](https://doi.org/10.2307/2425184).
- Jeffries KM, Brander SM, Britton MT, Fangue NA, Connon RE. 2015.** Chronic exposures to low and high concentrations of ibuprofen elicit different gene response patterns in a euryhaline fish. *Environmental Science and Pollution Research* **22**(22):17397–17413 DOI [10.1007/s11356-015-4227-y](https://doi.org/10.1007/s11356-015-4227-y).
- Jeppesen E, Meerhoff M, Holmgren K, González-Bergonzoni I, Teixeira-de Mello F, Declerck SA, De Meester L, Søndergaard M, Lauridsen TL, Bjerring R, Conde-Porcuna JM. 2010.** Impacts of climate warming on lake fish community structure and potential effects on ecosystem function. *Hydrobiologia* **646**(1):73–90 DOI [10.1007/s10750-010-0171-5](https://doi.org/10.1007/s10750-010-0171-5).
- Jin M, Zhang X, Wang L, Huang C, Zhang Y, Zhao M. 2009.** Developmental toxicity of bifenthrin in embryo-larval stages of zebrafish. *Aquatic Toxicology* **95**(4):347–354 DOI [10.1016/j.aquatox.2009.10.003](https://doi.org/10.1016/j.aquatox.2009.10.003).
- Jodo A, Ando H, Urano A. 2003.** Five different types of putative GnRH receptor gene are expressed in the brain of masu salmon (*Oncorhynchus masou*). *Zoological Science* **20**(9):1117–1125 DOI [10.2108/zsj.20.1117](https://doi.org/10.2108/zsj.20.1117).

- Kidd KA, Blanchfield PJ, Mills KH, Palace VP, Evans RE, Lazorchak JM, Flick RW. 2007.** Collapse of a fish population after exposure to a synthetic estrogen. *Proceedings of the National Academy of Sciences* **104**(21):8897–8901 DOI [10.1073/pnas.0609568104](https://doi.org/10.1073/pnas.0609568104).
- King OC, Van de Merwe JP, McDonald JA, Leusch FD. 2016.** Concentrations of levonorgestrel and ethinylestradiol in wastewater effluents: is the progestin also cause for concern? *Environmental Toxicology and Chemistry* **35**(6):1378–1385 DOI [10.1002/etc.3304](https://doi.org/10.1002/etc.3304).
- Kitano T, Takamune K, Kobayashi T, Nagahama Y, Abe SI. 1999.** Suppression of P450 aromatase gene expression in sex-reversed males produced by rearing genetically female larvae at a high water temperature during a period of sex differentiation in the Japanese flounder (*Paralichthys olivaceus*). *Journal of Molecular Endocrinology* **23**(2):167–176 DOI [10.1677/jme.0.0230167](https://doi.org/10.1677/jme.0.0230167).
- Kuivila KM, Hladik ML, Ingersoll CG, Kemble NE, Moran PW, Calhoun DL, Nowell LH, Gilliom RJ. 2012.** Occurrence and potential sources of pyrethroid insecticides in stream sediments from seven US metropolitan areas. *Environmental Science & Technology* **46**(8):4297–4303 DOI [10.1021/es2044882](https://doi.org/10.1021/es2044882).
- Levavi-Sivan B, Bogerd J, Mañanós EL, Gómez A, Lareyre JJ. 2010.** Perspectives on fish gonadotropins and their receptors. *General and Comparative Endocrinology* **165**(3):412–437 DOI [10.1016/j.ygcen.2009.07.019](https://doi.org/10.1016/j.ygcen.2009.07.019).
- Li H, Cheng F, Wei Y, Lydy MJ, You J. 2017.** Global occurrence of pyrethroid insecticides in sediment and the associated toxicological effects on benthic invertebrates: an overview. *Journal of Hazardous Materials* **324**:258–271 DOI [10.1016/j.jhazmat.2016.10.056](https://doi.org/10.1016/j.jhazmat.2016.10.056).
- Liu S, Qin F, Wang H, Wu T, Zhang Y, Zheng Y, Li M, Wang Z. 2012.** Effects of 17 α -ethinylestradiol and bisphenol A on steroidogenic messenger ribonucleic acid levels in the rare minnow gonads. *Aquatic Toxicology* **122**:19–27 DOI [10.1016/j.aquatox.2012.05.010](https://doi.org/10.1016/j.aquatox.2012.05.010).
- Livak KJ, Schmittgen TD. 2001.** Analysis of relative gene expression data using real-time quantitative PCR and the 2 $^{-\Delta\Delta CT}$ method. *Methods* **25**(4):402–408 DOI [10.1006/meth.2001.1262](https://doi.org/10.1006/meth.2001.1262).
- Luzio A, Santos D, Fontainhas-Fernandes AA, Monteiro SM, Coimbra AM. 2016.** Effects of 17 α -ethinylestradiol at different water temperatures on zebrafish sex differentiation and gonad development. *Aquatic Toxicology* **174**:22–35 DOI [10.1016/j.aquatox.2016.02.003](https://doi.org/10.1016/j.aquatox.2016.02.003).
- Maggiolini M, Picard D. 2010.** The unfolding stories of GPR30, a new membrane-bound estrogen receptor. *Journal of Endocrinology* **204**(2):105–114 DOI [10.1677/JOE-09-0242](https://doi.org/10.1677/JOE-09-0242).
- Marchais-Oberwinkler S, Henn C, Möller G, Klein T, Negri M, Oster A, Spadaro A, Werth R, Wetzel M, Xu K, Frotscher M. 2011.** 17 β -Hydroxysteroid dehydrogenases (17 β -HSDs) as therapeutic targets: protein structures, functions, and recent progress in inhibitor development. *The Journal of Steroid Biochemistry and Molecular Biology* **125**(1-2):66–82 DOI [10.1016/j.jsbmb.2010.12.013](https://doi.org/10.1016/j.jsbmb.2010.12.013).

- Middaugh DP, Hemmer MJ, Lamadrid-Rose Y. 1986.** Laboratory spawning cues in *Menidia beryllina* and *M. peninsulae* (Pisces, Atherinidae) with notes on survival and growth of larvae at different salinities. *Environmental Biology of Fishes* **15**(2):107–117.
- Miranda LA, Chalde T, Elisio M, Strüssmann CA. 2013.** Effects of global warming on fish reproductive endocrine axis, with special emphasis in pejerrey *Odontesthes bonariensis*. *General and Comparative Endocrinology* **192**:45–54
DOI [10.1016/j.ygcen.2013.02.034](https://doi.org/10.1016/j.ygcen.2013.02.034).
- Moyano M, Illing B, Peschutter P, Huebert KB, Peck MA. 2016.** Thermal impacts on the growth, development and ontogeny of critical swimming speed in Atlantic herring larvae. *Comparative Biochemistry and Physiology Part A: Molecular & Integrative Physiology* **197**:23–34 DOI [10.1016/j.cbpa.2016.02.020](https://doi.org/10.1016/j.cbpa.2016.02.020).
- Munday PL, Jones GP, Pratchett MS, Williams AJ. 2008.** Climate change and the future for coral reef fishes. *Fish and Fisheries* **9**(3):261–285
DOI [10.1111/j.1467-2979.2008.00281.x](https://doi.org/10.1111/j.1467-2979.2008.00281.x).
- Pang Y, Dong J, Thomas P. 2008.** Estrogen signaling characteristics of Atlantic croaker G protein-coupled receptor 30 (GPR30) and evidence it is involved in maintenance of oocyte meiotic arrest. *Endocrinology* **149**(7):3410–3426
DOI [10.1210/en.2007-1663](https://doi.org/10.1210/en.2007-1663).
- Pankhurst NW, Munday PL. 2011.** Effects of climate change on fish reproduction and early life history stages. *Marine and Freshwater Research* **62**(9):1015–1026
DOI [10.1071/MF10269](https://doi.org/10.1071/MF10269).
- R Core Team. 2017.** R: a language and environment for statistical computing. Vienna: R Foundation for Statistical Computing. Available at <https://www.R-project.org/>.
- Rhee JS, Kim BM, Lee CJ, Yoon YD, Lee YM, Lee JS. 2011.** Bisphenol A modulates expression of sex differentiation genes in the self-fertilizing fish, *Kryptolebias marmoratus*. *Aquatic Toxicology* **104**(3-4):218–229
DOI [10.1016/j.aquatox.2011.04.020](https://doi.org/10.1016/j.aquatox.2011.04.020).
- Rocha A, Zanuy S, Carrillo M, Gómez A. 2009.** Seasonal changes in gonadal expression of gonadotropin receptors, steroidogenic acute regulatory protein and steroidogenic enzymes in the European sea bass. *General and Comparative Endocrinology* **162**(3):265–275 DOI [10.1016/j.ygcen.2009.03.023](https://doi.org/10.1016/j.ygcen.2009.03.023).
- Scholz S, Gutzeit HO. 2000.** 17- α -ethinylestradiol affects reproduction, sexual differentiation and aromatase gene expression of the medaka (*Oryzias latipes*). *Aquatic Toxicology* **50**(4):363–373 DOI [10.1016/S0166-445X\(00\)00090-4](https://doi.org/10.1016/S0166-445X(00)00090-4).
- Schwindt AR, Winkelman DL, Keteles K, Murphy M, Vajda AM. 2014.** An environmental oestrogen disrupts fish population dynamics through direct and transgenerational effects on survival and fecundity. *Journal of Applied Ecology* **51**(3):582–591
DOI [10.1111/1365-2664.12237](https://doi.org/10.1111/1365-2664.12237).
- Shamblott MJ, Cheng CM, Bolt D, Chen TT. 1995.** Appearance of insulin-like growth factor mRNA in the liver and pyloric ceca of a teleost in response to exogenous growth hormone. *Proceedings of the National Academy of Sciences* **92**(15):6943–6946
DOI [10.1073/pnas.92.15.6943](https://doi.org/10.1073/pnas.92.15.6943).

- Shved N, Berishvili G, Baroiller JF, Segner H, Reinecke M. 2008. Environmentally relevant concentrations of 17 α -ethinylestradiol (EE2) interfere with the growth hormone (GH)/insulin-like growth factor (IGF)-I system in developing bony fish. *Toxicological Sciences* **106**(1):93–102 DOI [10.1093/toxsci/kfn150](https://doi.org/10.1093/toxsci/kfn150).
- Stocker TF, Qin D, Plattner GK, Tignor M, Allen SK, Boschung J, Nauels A, Xia Y, Bex V, Midgley PM (eds.) 2013. IPCC, 2013: summary for policymakers in climate change 2013: the physical science basis, contribution of working group I to the fifth assessment report of the intergovernmental panel on climate change. Available at www.ipcc.ch/pdf/assessmentreport/ar5/wg1/WG1AR5_SPM_FINAL.pdf.
- Strüssmann CA, Conover DO, Somoza GM, Miranda LA. 2010. Implications of climate change for the reproductive capacity and survival of New World silversides (family Atherinopsidae). *Journal of Fish Biology* **77**(8):1818–1834 DOI [10.1111/j.1095-8649.2010.02780.x](https://doi.org/10.1111/j.1095-8649.2010.02780.x).
- Tu W, Xu C, Lu B, Lin C, Wu Y, Liu W. 2016. Acute exposure to synthetic pyrethroids causes bioconcentration and disruption of the hypothalamus—pituitary—thyroid axis in zebrafish embryos. *Science of the Total Environment* **542**:876–885 DOI [10.1016/j.scitotenv.2015.10.131](https://doi.org/10.1016/j.scitotenv.2015.10.131).
- Uchida D, Yamashita M, Kitano T, Iguchi T. 2004. An aromatase inhibitor or high water temperature induce oocyte apoptosis and depletion of P450 aromatase activity in the gonads of genetic female zebrafish during sex-reversal. *Comparative Biochemistry and Physiology Part A: Molecular & Integrative Physiology* **137**(1):11–20 DOI [10.1016/S1095-6433\(03\)00178-8](https://doi.org/10.1016/S1095-6433(03)00178-8).
- Vajda AM, Barber LB, Gray JL, Lopez EM, Woodling JD, Norris DO. 2008. Reproductive disruption in fish downstream from an estrogenic wastewater effluent. *Environmental Science & Technology* **42**(9):3407–3414 DOI [10.1021/es0720661](https://doi.org/10.1021/es0720661).
- Van Aerle R, Pounds N, Hutchinson TH, Maddix S, Tyler CR. 2002. Window of sensitivity for the estrogenic effects of ethinylestradiol in early life-stages of fathead minnow, *Pimephales promelas*. *Ecotoxicology* **11**(6):423–434 DOI [10.1023/A:1021053217513](https://doi.org/10.1023/A:1021053217513).
- Vandenberg LN, Colborn T, Hayes TB, Heindel JJ, Jacobs Jr DR, Lee DH, Shioda T, Soto AM, Vom Saal FS, Welshons WV, Zoeller RT. 2012. Hormones and endocrine-disrupting chemicals: low-dose effects and nonmonotonic dose responses. *Endocrine Reviews* **33**(3):378–455 DOI [10.1210/er.2011-1050](https://doi.org/10.1210/er.2011-1050).
- Vandesompele J, De Preter K, Pattyn F, Poppe B, Van Roy N, De Paepe A, Speleman F. 2002. Accurate normalization of real-time quantitative RT-PCR data by geometric averaging of multiple internal control genes. *Genome Biology* **3**(7):research0034–1.
- Weston DP, Lydy MJ. 2012. Stormwater input of pyrethroid insecticides to an urban river. *Environmental Toxicology and Chemistry* **31**(7):1579–1586 DOI [10.1002/etc.1847](https://doi.org/10.1002/etc.1847).
- White JW, Cole BJ, Cherr GN, Cannon RE, Brander SM. 2017. Scaling up endocrine disruption effects from individuals to populations: outcomes depend on how many males a population needs. *Environmental Science & Technology* **51**(3):1802–1810 DOI [10.1021/acs.est.6b05276](https://doi.org/10.1021/acs.est.6b05276).

Zhang X, Hecker M, Park JW, Tompsett AR, Newsted J, Nakayama K, Jones PD, Au D, Kong R, Wu RS, Giesy J. 2008. Real-time PCR array to study effects of chemicals on the hypothalamic–pituitary–gonadal axis of the Japanese medaka. *Aquatic Toxicology* 88(3):173–182 DOI [10.1016/j.aquatox.2008.04.009](https://doi.org/10.1016/j.aquatox.2008.04.009).