

UC Irvine

UC Irvine Previously Published Works

Title

Rearrangements of azidoquinones. IX. Pyrolysis of 2,3-diazido-1,4-naphthoquinone-dibenzenesulfonimide

Permalink

<https://escholarship.org/uc/item/0529f30p>

Journal

Tetrahedron Letters, 12(39)

ISSN

0040-4039

Authors

Moore, Harold W

Lee, Mei-Sie

Publication Date

1971

DOI

10.1016/s0040-4039(01)97252-1

Copyright Information

This work is made available under the terms of a Creative Commons Attribution License, available at

<https://creativecommons.org/licenses/by/4.0/>

Peer reviewed

REARRANGEMENTS OF AZIDOQUINONES. IX. PYROLYSIS OF 2,3-DIAZIDO-1,4-NAPHTHOQUINONE-DIBENZENESULFONIMIDE

Harold W. Moore and Mei-Sie Lee


Department of Chemistry
University of California
Irvine, California 92664

(Received in USA 2 June 1971; received in UK for publication 1 September 1971)


Azidoquinones have been shown to be very versatile synthetic intermediates. γ -cyano-alkylidene- $\Delta^{\alpha, \beta}$ -butenolides,¹ 2-cyano-1,3-cyclopentenediones,² cyanoketenes,³ azapinediones,⁴ aminoquinones⁵ and acylcyanides⁶ can be prepared from azidoquinones by employing the appropriate reaction conditions. The synthetic utility of this class of compounds suggested an investigation of certain azidoquinone derivatives to see if analogous transformations could be accomplished. Reported here is a study of the thermal decomposition of 2,3-diazido-1,4-naphthoquinone-dibenzenesulfonimide, **2**. This vicinal diazide cleaves pyrolytically in refluxing benzene to give phthaloyl cyanide-dibenzenesulfonimide, **3**. Compound **3** undergoes subsequent hydrolytic ring closure upon reaction with alcohols to 1,3-substituted isoindolines and phthalimidine-benzenesulfonimides.

The purple crystalline diazide, **2**, was prepared in nearly quantitative yield from 2,3-dichloro-1,4-naphthoquinone-dibenzenesulfonimide,⁷ **1**, upon reaction with excess sodium azide. A solution of 2.1 mmol of **1** in 40 ml of acetone was reacted with 6.1 mmol of sodium azide in 5 ml of water at 0-5°. The reaction solution immediately turned from yellow to purple upon addition of the azide. Water (20 ml) was added to the reaction solution which induced the precipitation of the diazide, **2**.

Phthaloyl cyanide-dibenzenesulfonimide, **3**, was obtained by refluxing a suspension of 725 mg (1.4 mmol) of **2** in 7 ml of anhydrous benzene for 6 hrs. A small amount (40 mg) of an unidentified insoluble precipitate was collected by filtration and the solvent removed in vacuo to give **3** as a brownish oil. The structure of **3** is in complete agreement with its spectral data (Table I) and its chemical properties. The thermal cleavage, **2** \rightarrow **3**, has good precedent; 2,3-diazido-1,4-naphthoquinone loses nitrogen in refluxing toluene and forms phthaloyl cyanide,⁶ and o-diazidobenzene thermally fragments to cis-1,4-dicyanobutadiene.⁸



The α -cyanobenzesulfonyl moieties in **3** are very reactive toward hydroxylic reagents. Compound **3** reacts readily with water to give the substituted N-benzenesulfonyllactam, **4**, in 85% yield. The formation of **4** is envisaged as arising *via* the isoindoline, **4**, which then eliminates hydrogen cyanide. Hydrolysis of the lactam **5** with 5% NaOH converted it to benzenesulfonamide, **6**, and N-benzenesulfonylphthalimide, **7**,⁹ which were identified by comparison (ir and mixed mp) with authentic samples.



The hydrolytic ring closure of $\underline{3}$ could also be accomplished with alcohols. Methanol and isopropanol react with $\underline{3}$ to give respectively the 1,3-substituted isoindolines, $\underline{8}$ and $\underline{9}$. These compounds are heat labile and lose hydrogen cyanide upon attempted recrystallization from acetone-diethyl ether to give respectively the substituted N-benzenesulfonylphthalimidine-benzenesulfonimides, $\underline{10}$ and $\underline{11}$. The spectral data (Table I) for compounds $\underline{8}$ - $\underline{11}$ are in complete accord with their proposed structures. Of particular utility in the structural assignments is the nmr spectrum of $\underline{11}$ which shows nonequivalent methyl absorptions (δ , 1.38 and 1.10) for the isopropyl group. This is consistent with structure $\underline{11}$ which has the isopropoxide moiety directly substituted to an asymmetric center.




TABLE I
Spectral Data of New Compounds¹⁰

Compound	Yield	mp °C	ir(Nujol) cm ⁻¹	nmr (CDCl ₃) ppm
$\underline{2}$	>95%	126-28°	2137 2105 1558	8.83-7.26 m
$\underline{3}$		oil	N ₃ C=N- 2208 1585	
$\underline{5}$	85%	158-60	>NH C=O 3247 1754	7.90-6.40 m
$\underline{8}$	72%	175-185 dec	>NH 3205	
$\underline{9}$	74%	165-185 dec	>NH 3200	
$\underline{10}$		187-89	C=N- 1621	3.29 s (3) OCH ₃ 8.20-7.30 m (14)
$\underline{11}$		166-68	C=N 1630	1.10 d (3) CH ₃ 1.38 d (3) CH ₃ 4.08 m (1) CH 8.00-7.08 m (14)

Adams and coworkers ¹⁰ have published an extensive investigation concerning the synthesis and chemistry of variously substituted quinonedimides. The availability of the corresponding azidoquinonedimides via standard routes gives a class of previously unexplored azides. The chemistry of these compounds as compared to that of the corresponding azidoquinones is currently under study.

ACKNOWLEDGEMENT

The authors are grateful to the National Science Foundation (GP 19268) for financial support of this project.

REFERENCES

1. H. W. Moore, H. R. Shelden, D. W. Deters and R. J. Wikholm, J. Am. Chem. Soc., 92, 1675 (1969).
2. H. W. Moore, H. R. Shelden and W. Weyler, Jr., Tetrahedron Lett., 3947 (1969).
3. H. W. Moore, and W. Weyler, Jr., J. Am. Chem. Soc., 92, 4132 (1970).
4. H. W. Moore, H. R. Shelden and W. Weyler, Jr., Tetrahedron Lett., 1243 (1969).
5. H. W. Moore and H. R. Shelden, J. Org. Chem., 33, 4019 (1968).
6. J. A. Van Allen, W. J. Priest, A. S. Marshall and G. A. Reynolds, J. Org. Chem., 33, (1968).
7. R. Adams and R. A. Wankel, J. Am. Chem. Soc., 73, 131 (1951).
8. J. H. Hall and E. Patterson, J. Am. Chem. Soc., 89, 5856 (1967).
9. T. W. Evans and R. A. Wankel, J. Am. Chem. Soc., 73, 131 (1951).
10. Combustion analysis for the new compounds 2, 10 and 11 are all in complete accord with their proposed formulations.
11. R. Adams and W. Reifschneider, Bull. Soc., Chim. France, 23 (1958).